

---

## Submission to Clinical Epidemiology and Global Health - manuscript number

4 messages

---

**Clinical Epidemiology and Global Health** <em@editorialmanager.com>  
Reply-To: Clinical Epidemiology and Global Health <support@elsevier.com>  
To: Santi Martini <santi-m@fkm.unair.ac.id>

Mon, Sep 6, 2021 at 1:50 PM

\*This is an automated message.\*

Manuscript Number: CEGH-D-21-00557

Assessing Quality of Life and Associated Factors in Post-Stroke Patients Using the World Health Organization Abbreviated Generic Quality Of Life Questionnaire (WHOQOL-BREF)

Dear Dr. Martini,

Your above referenced submission has been assigned a manuscript number: CEGH-D-21-00557.

To track the status of your manuscript, please log in as an author at <https://www.editorialmanager.com/cegh/>, and navigate to the "Submissions Being Processed" folder.

Thank you for submitting your work to this journal.

Kind regards,  
Clinical Epidemiology and Global Health

More information and support

You will find information relevant for you as an author on Elsevier's Author Hub: <https://www.elsevier.com/authors>

FAQ: How can I reset a forgotten password?

[https://service.elsevier.com/app/answers/detail/a\\_id/28452/supporthub/publishing/](https://service.elsevier.com/app/answers/detail/a_id/28452/supporthub/publishing/)

For further assistance, please visit our customer service site: <https://service.elsevier.com/app/home/supporthub/publishing/>

Here you can search for solutions on a range of topics, find answers to frequently asked questions, and learn more about Editorial Manager via interactive tutorials. You can also talk 24/7 to our customer support team by phone and 24/7 by live chat and email

#AU\_CEGH#

To ensure this email reaches the intended recipient, please do not delete the above code

---

In compliance with data protection regulations, you may request that we remove your personal registration details at any time. (Use the following URL: <https://www.editorialmanager.com/cegh/login.asp?a=r>). Please contact the publication office if you have any questions.

---

## Decision on submission to Clinical Epidemiology and Global Health

4 messages

---

Clinical Epidemiology and Global Health <em@editorialmanager.com>  
Reply-To: Clinical Epidemiology and Global Health <support@elsevier.com>  
To: Santi Martini <santi-m@fkm.unair.ac.id>

Tue, Dec 7, 2021 at 7:43 PM

Manuscript Number: CEGH-D-21-00557

Assessing Quality of Life and Associated Factors in Post-Stroke Patients Using the World Health Organization Abbreviated Generic Quality Of Life Questionnaire (WHOQOL-BREF)

Dear Dr. Martini,

Thank you for submitting your manuscript to Clinical Epidemiology and Global Health.

I have completed my evaluation of your manuscript. The reviewers recommend reconsideration of your manuscript following minor revision and modification. I invite you to resubmit your manuscript after addressing the comments below. Please resubmit your revised manuscript by Dec 21, 2021.

When revising your manuscript, please consider all issues mentioned in the reviewers' comments carefully: please outline every change made in response to their comments and provide suitable rebuttals for any comments not addressed. Please note that your revised submission may need to be re-reviewed.

To submit your revised manuscript, please log in as an author at <https://www.editorialmanager.com/cegh/>, and navigate to the "Submissions Needing Revision" folder under the Author Main Menu.

Clinical Epidemiology and Global Health values your contribution and I look forward to receiving your revised manuscript.

Kind regards,

Shally Awasthi

Editor-in-Chief

Clinical Epidemiology and Global Health

Editor and Reviewer comments:

Reviewer #1: I have attached my review. Please revised it base on my review

More information and support

FAQ: How do I revise my submission in Editorial Manager?

[https://service.elsevier.com/app/answers/detail/a\\_id/28463/supporthub/publishing/](https://service.elsevier.com/app/answers/detail/a_id/28463/supporthub/publishing/)

You will find information relevant for you as an author on Elsevier's Author Hub: <https://www.elsevier.com/authors>

FAQ: How can I reset a forgotten password?

[https://service.elsevier.com/app/answers/detail/a\\_id/28452/supporthub/publishing/](https://service.elsevier.com/app/answers/detail/a_id/28452/supporthub/publishing/)

For further assistance, please visit our customer service site: <https://service.elsevier.com/app/home/supporthub/publishing/>

Here you can search for solutions on a range of topics, find answers to frequently asked questions, and learn more about Editorial Manager via interactive tutorials. You can also talk 24/7 to our customer support team by phone and 24/7 by live chat and email

#AU\_CEGH#

To ensure this email reaches the intended recipient, please do not delete the above code

# Clinical Epidemiology and Global Health

## Assessing Quality of Life and Associated Factors in Post-Stroke Patients Using the World Health Organization Abbreviated Generic Quality Of Life Questionnaire (WHOQOL-BREF) --Manuscript Draft--

<b>Manuscript Number:</b>	CEGH-D-21-00557R1
<b>Article Type:</b>	Original article
<b>Keywords:</b>	educational level; internal consistency; Post-Stroke; QOL; WHOQOL-BREF
<b>Corresponding Author:</b>	Santi Martini, Dr Airlangga University Faculty of Public Health: Universitas Airlangga Fakultas Kesehatan Masyarakat INDONESIA
<b>First Author:</b>	Santi Martini, Dr
<b>Order of Authors:</b>	Santi Martini, Dr Defi Amalia Setia Ningrum, master Khadizah H. Abdul-Mumin, Ph.D Chung Yi-Li, Professor
<b>Abstract:</b>	<p><b>Introduction</b></p> <p>This study piloted the World Health Organization Quality of Life short version form (WHOQOL-BREF) to compare the quality of life (QoL) in post-stroke patients who followed the medical stroke rehabilitation program and those whom did not. This study determined which quality of life domains were mostly affected by stroke, and whether there are influential factors other than rehabilitation.</p> <p><b>Methods</b></p> <p>A cross-sectional study was conducted at Public Hospital in Surabaya. Patients with stroke less than two years since the first attack were included in this study (n=52). Cronbach's alpha test was performed to assess the internal consistency of WHOQOL-BREF questionnaire. The independent t test and ANOVA were used to compare the differences between patient's characteristics and the six domains of QoL (i.e., perception of QoL, perception of health, physical of health, psychological health, social relationship, and environmental health). Multiple linear regression was performed to assess the influential factors of QoL.</p> <p><b>Results</b></p> <p>post-stroke patients aged more than 50 years old (76.9%), male (55.8%). But, only 48.1% of those patients participating in medical rehabilitation programs. Age group was significantly associated with QoL in the psychological health domain (p=0.021); participating in medical stroke rehabilitation programs was significantly associated with social relationship domain (p=0.026) and the education level was significantly associated with physical health (p=0.005), psychological health (p=0.035) and perception of health (p=0.003)..</p> <p><b>Conclusion</b></p> <p>Lower education level was significantly associated with a low perception of health, one of QoL domains; and it was the most influential factor of QoL among post-stroke patients.</p>
<b>Suggested Reviewers:</b>	Muhammad Aziz Rahman, Ph.D Associate Dean, Federation University Australia ma.rahman@federation.edu.au

	<p>Rahayu Lubis, Ph.D Lecturer, University of Sumatera Utara: Universitas Sumatera Utara lubis@usu.ac.id</p>
	<p>Muhammad Atoillah Isfandiari, Dr Vice Dean, Airlangga University: Universitas Airlangga muhammad-a-i@fkm.unair.ac.id</p>
	<p>Kurnia Dwi Artanti, MPH Lecturer, Universitas Airlangga kurnia-d-a@fkm.unair.ac.id She's an editor of a journal in Indonesia</p>
<b>Opposed Reviewers:</b>	
<b>Response to Reviewers:</b>	

Dear Chief of Editor Clinical Epidemiology and Global Health

We are writing to request that you consider our manuscript, titled “Assessing Quality Of Life And Associated Factors In Post-Stroke Patients Using The World Health Organization Abbreviated Generic Quality of Life Questionnaire (WHOQOL-BREF)”

I declare that the work submitted for publication is original, previously unpublished and not under consideration for publication elsewhere.

All Author declare there is no conflict of Interest.

Thank you .

Sincerely,

Dr. Santi Martini, MD, MPH

Respond to Reviewer

Dear Editor and Reviewer, we already made revision as requested.

Please find the revised part in highlighted Green.

# **Assessing Quality of Life and Associated Factors in Post-Stroke Patients Using the World Health Organization Abbreviated Generic Quality Of Life Questionnaire (WHOQOL-BREF)**

Authors :

1. Dr. Santi Martini [[santi-m@fkm.unair.ac.id](mailto:santi-m@fkm.unair.ac.id)]  
Department of Epidemiology, Faculty of Public Health, Universitas Airlangga
2. Defi Amalia Setia Ningrum [[defi.amalia@gmail.com](mailto:defi.amalia@gmail.com)]  
FETP graduate, University of Indonesia
3. Dr. Khadizah H. Abdul-Mumin [[khadizah.mumin@ubd.edu.bn](mailto:khadizah.mumin@ubd.edu.bn)]  
Institute of Health Sciences (PAPRSB, IHS), Universiti Brunei Darussalam (UBD),
4. Co-author : Prof. Chung Yi-Li, BSc, MSc., PhD. [[cyli99@mail.ncku.edu.tw](mailto:cyli99@mail.ncku.edu.tw)]  
Department of Public Health, College of Medicine, National Cheng Kung University,  
Taiwan

Corresponding author: Dr. Santi Martini,

Dean Faculty of Public Health

Department of Epidemiology, Faculty of Public Health, Universitas Airlangga

Kampus C Unair, Surabaya 60115, East Java indonesia.

[santi-m@fkm.unair.ac.id](mailto:santi-m@fkm.unair.ac.id)

## **Acknowledgements**

Faculty of Public Health Universitas Airlangga

## **Declaration**

All Author declare there is no conflict of Interest

This study was approved by the Institutional Health Research Ethics Committee of the primary investigator in Surabaya








Figure 1. Bar chart demonstrating the influence of medical stroke rehabilitation on QoL. Patients were divided into two groups based on whether they participate or not in the medical stroke rehabilitation. The bar shows average domain score of WHOQOL-BREF.

Table 1. Characteristic of the sample (n=52)

Characteristic	n (%)
Age (years)	58.27 ± 9.5
≤ 50	12 (23.1)
> 50	40 (76.9)
Gender	
Male	29 (55.8)
Female	23 (44.2)
Education level	
Elementary	15 (28.8)
Secondary	21 (40.4)
University	16 (30.8)
Marital status	
Single/divorced	10 (19.2)
Married	42 (80.8)
Medical rehabilitation	
Yes	25 (48.1)
No	27 (51.9)

Table 2. Comparison of the WHOQOL-BREF mean scores according to gender, age, education level, marital status, and medical rehabilitation.

	Domain					
	Perception of QOL Mean ± SD	Perception of health Mean ± SD	Physical health Mean ± SD	Psychological health Mean ± SD	Social relationship Mean ± SD	Enviromental health Mean ± SD
Total	62.50 ± 16.789	49.03 ± 17.823	62.85 ± 19.031	75.44 ± 20.165	66.06 ± 1.311	72.11 ± 12.607
Gender						
Male	62.93 ± 15.840	52.59 ± 16.830	62.79 ± 21.222	76.59 ± 20.916	65.76 ± 12.858	73.17 ± 11.142
Female	61.96 ± 18.262	44.56 ± 18.397	62.91 ± 16.320	74.00 ± 19.544	66.43 ± 13.720	70.78 ± 14.390
<i>p</i> value	0.838	0.108	0.982	0.648	0.856	0.503
Age (years)						
≤ 50	66.67 ± 16.283	56.25 ± 15.539	63.17 ± 18.473	85.00 ± 12.778	70.92 ± 6.127	72.42 ± 12.479
> 50	61.25 ± 16.937	46.87 ± 18.07	62.75 ± 19.425	72.57 ± 21.194	64.60 ± 14.318	72.025 ± 12.801
<i>p</i> value	0.332	0.111	0.947	0.018*	0.033*	0.926
Education level						
Elementary	58.33 ± 18.094	36.67 ± 15.100	56.00 ± 17.708	67.53 ± 18.909	62.20 ± 16.772	68.87 ± 13.400
Secondary	60.71 ± 14.940	51.19 ± 16.726	64.48 ± 15.725	79.86 ± 17.316	66.14 ± 9.040	73.04 ± 10.500
University	68.75 ± 17.078	57.81 ± 15.052	67.12 ± 23.215	77.06 ± 23.589	69.56 ± 13.623	73.94 ± 14.507
<i>p</i> value	0.186	0.002*	0.238	0.183	0.301	0.494
Marital status						
Single/divorced	61.31 ± 16.750	50.00 ± 18.313	62.88 ± 19.215	76.43 ± 19.067	66.59 ± 12.172	71.714 ± 12.377
Married	67.50 ± 16.874	45.00 ± 15.811	62.70 ± 19.241	71.30 ± 24.989	71.71 ± 12.377	73.800 ± 14.101
<i>p</i> value	0.299	0.431	0.979	0.475	0.550	0.643
Medical Stroke rehabilitation						
Yes	60.18 ± 14.310	44.44 ± 14.434	67.88 ± 17.871	79.60 ± 20.279	69.80 ± 11.965	74.36 ± 13.801
No	65.00 ± 19.094	54.00 ± 20.000	58.19 ± 19.201	71.59 ± 19.647	62.59 ± 113.397	70.037 ± 11.250
<i>p</i> value	0.306	0.52	0.66	0.154	0.047*	0.220

QOL=Quality of life, SD=Standard deviations, \*The mean difference is significant at  $\leq 0.05$  level

Table 3: Backward multiple linear regression analyses of significant factors associated with QOL

QOL domains	Variables	Unstandardized Coefficients		Standardized Coefficients	t	P value
		B	SE	Beta		
Physical Health	Education level	0.806	0.276	0.382	0.923	0.005
Psychological Health	Education level	0.651	0.300	0.288	0.175	0.035
	Age Category	-1.973	0.826	-0.415	-0.388	0.021
Social Relationship	Participation of Medical Rehabilitation	1.229	0.536	0.306	0.294	0.026
Environmental Health	None	-	-	-	-	-
Perception of Health	Education level	0.218	0.071	0.399	0.080	0.003

QOL = Quality of life

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.



## *Clinical Epidemiology and Global Health*

### COPYRIGHT TRANSFER AGREEMENT



Clinical Epidemiology and Global Health will be pleased to publish your article (“the Work”), tentatively entitled

Assessing Quality of Life and Associated Factors in Post-Stroke Patients Using the World Health Organization Abbreviated Generic Quality Of Life Questionnaire (WHOQOL-BREF)

---

---

in the *Clinical Epidemiology and Global Health* (“the Journal”) if the Work is accepted for publication. The authors transfer all copyright ownership in and relating to the Work, in all forms and media, to the Clinical Epidemiology and Global Health in the event that the Work is published in the Journal. However, this agreement will be null and void if the Work is not published in the Journal.

The authors warrant that the Work is original, is not under consideration by another journal, and has not been previously published.

This statement is signed by all the listed authors in the manuscript. The authors certify that all the authors listed in the manuscript understand and agree with the terms in this Copyright Transfer Agreement.

Defi Amalia Setia Ningrum, Khadizah H. Abdul-Mumin, Chung Yi-Li

Co-authors

Dr. Santi Martini, dr., M.Kes

Corresponding author’s name (typed)

Santi-m@fkm.unair.ac.id


Corresponding author’s signature & E-mail address

August 31, 2021

Date

Defi Amalia Setia Ningrum

Co-authors name (typed)



defi.amalia@gmail.com

Co-authors signature & E-mail address

September 2, 2021

Date

Prof. Chung Yi-Li, BSc, MSc., PhD

Co-authors name (typed)

cyli99@mail.ncku.edu.tw

Co-authors signature & E-mail address

September 2, 2021

Date

Co-authors name (typed)

Co-authors signature & E-mail address

Date

Co-authors name (typed)

Co-authors signature & E-mail address

Date

Dr. Khadizah H. Abdul-Mumin

Co-authors name (typed)



khadizah.mumin@ubd.edu.bn

Co-authors signature & E-mail address

September 1, 2021

Date

Chung-Yi Li

Co-authors name (typed)



Co-authors signature & E-mail address

September 2, 2021

Date

Co-authors name (typed)

Co-authors signature & E-mail address

Date

Co-authors name (typed)

Co-authors signature & E-mail address

Date



# *Clinical Epidemiology and Global Health*

## AUTHORSHIP & CONFLICTS OF INTEREST STATEMENT



Manuscript title: Assessing Quality of Life and Associated Factors in Post-Stroke Patients Using the World Health Organization Abbreviated Generic Quality Of Life Questionnaire (WHOQOL-BREF)

---

**AUTHORSHIP** All persons who meet authorship criteria are listed as authors, and all authors certify that they have participated sufficiently in the work to take public responsibility for the content, including participation in the concept, design, analysis, writing, or revision of the manuscript. Furthermore, each author certifies that this material or similar material has not been and will not be submitted to or published in any other publication.

Please indicate the specific contributions made by each author (list the authors' initials followed by their surnames, e.g., Y.L. Chang). The name of each author must appear at least once in each of the three categories below.

### *Category 1*

Conception and design of study: S. Martini, D. Ningrum,

acquisition of data: S. Martini, D. Ningrum,

; analysis and/or interpretation of data: S. Martini, D. Ningrum, C.Yi Li, Khadizah,;

### *Category 2*

Drafting the manuscript: S. Martini, D. Ningrum,

revising the manuscript critically for important intellectual content S. Martini, D. Ningrum, C.Yi Li, Khadizah;

### *Category 3*

Approval of the version of the manuscript to be published (the names of all authors must be listed):

S. Martini, D. Ningrum, C.Yi Li, Khadizah,;

### **Acknowledgments**

All persons who have made substantial contributions to the work reported in the manuscript (e.g., technical help, writing and editing assistance, general support), but who do not meet the criteria for authorship, are named in the Acknowledgments and have given us their written permission to be named. If we have not included an Acknowledgments, then that indicates that we have not received substantial contributions from non-authors.

**CONFLICTS OF INTEREST**

A conflict of interest occurs when an individual’s objectivity is potentially compromised by a desire for financial gain, prominence, professional advancement or a successful outcome. *CEGH* Editors strive to ensure that what is published in the Journal is as balanced, objective and evidence-based as possible. Since it can be difficult to distinguish between an actual conflict of interest and a perceived conflict of interest, the Journal requires authors to disclose all and any potential conflicts of interest.

**Section I**

The authors whose names are listed immediately below certify that they have NO affiliations with or involvement in any organization or entity with any financial interest (such as honoraria; educational grants; participation in speakers’ bureaus; membership, employment, consultancies, stock ownership, or other equity interest; and expert testimony or patent-licensing arrangements), or non-financial interest (such as personal or professional relationships, affiliations, knowledge or beliefs) in the subject matter or materials discussed in this manuscript.

Author names:

Santi Martini, Defi Amalia Setia Ningrum, Khadizah H. Abdul-Mumin, Chung Yi-Li

---

---

---

**Section II**

The authors whose names are listed immediately below report the following details of affiliation or involvement in an organization or entity with a financial or non-financial interest in the subject matter or materials discussed in this manuscript. Please specify the nature of the conflict on a separate sheet of paper if the space below is inadequate.


Author names & E-mail address	Details of the conflict(s) of interest:
<u>Santi Martini, <a href="mailto:santi-m@fkm.unair.ac.id">santi-m@fkm.unair.ac.id</a></u>	none of conflict of interest
<u>Defi Amalia Setia Ningrum, <a href="mailto:defi.amalia@gmail.com">defi.amalia@gmail.com</a></u>	none of conflict of interest
<u>Khadizah H. Abdul-Mumin, <a href="mailto:khadizah.mumin@ubd.edu.bn">khadizah.mumin@ubd.edu.bn</a></u>	none of conflict of interest
<u>Chung Yi-Li, <a href="mailto:cyl99@mail.ncku.edu.tw">cyl99@mail.ncku.edu.tw</a></u>	none of conflict of interest

This Authorship & Conflicts of Interest Statement is signed by the corresponding author on behalf of all the listed authors in the manuscript. The corresponding author certifies that all the information contained in this statement is true, correct and agreed on by all the listed authors.

Author(s) name (typed)

Santi Martini

Author(s) signature



Date

August 31, 2021



## **Assessing Quality of Life and Associated Factors in Post-Stroke Patients Using the World Health Organization Abbreviated Generic Quality of Life Questionnaire (WHOQoL-Bref)**

### **Abstract**

**Introduction:** This study piloted the World Health Organization Quality of Life short version form (WHOQOL-BREF) to compare the quality of life (QoL) in post-stroke patients who followed the medical stroke rehabilitation program and those whom did not. This study determined which quality of life domains were mostly affected by stroke, and whether there are influencing factors other than rehabilitation. **Methods:** A cross-sectional study was conducted at Public Hospital in Surabaya. Patients with stroke less than two years since the first attack were included in this study (n=52). Cronbach's alpha test was performed to assess the internal consistency of WHOQOL-BREF questionnaire. The independent *t* test and ANOVA were used to compare the differences between patient's characteristics and the six domains of QoL (i.e., perception of QoL, perception of health, physical of health, psychological health, social relationship, and environmental health). Multiple linear regression was performed to assess the influential factors of QoL. **Results:** post-stroke patients aged more than 50 years old (76.9%), male (55.8%). But, only 48.1% of those patients participating in medical rehabilitation programs. Age group was significantly associated with QoL in the psychological health domain ( $p=0.021$ ); participating in medical stroke rehabilitation programs was significantly associated with social relationship domain ( $p=0.026$ ) and the education level was significantly associated to physical health ( $p=0.005$ ), psychological health ( $p=0.035$ ) and perception of health ( $p=0.003$ ). **Conclusion:** Lower education level was significantly associated with a low perception of

health, one of QoL domains; and it was the most influential factor of QoL among post-stroke patients.

Keywords: educational level; internal consistency; Post-Stroke; quality of life; WHOQOL-BREF

## Introduction

Epidemiological studies evident that approximately 5.5 million people worldwide died annually due to stroke with the estimation that the global prevalence of stroke will gradually increase to as high as 21.9% by 2030 (1-2). Stroke is one of the most frequent causes of disability worldwide with 4.4 million disability-adjusted life-years lost (1). Most of stroke is ischemic stroke and based on CT scan of ischemic stroke patients, there are 58,6% subject with lacunar stroke, 20,7% with large artery atherosclerotic (LAA) stroke, and 20,7% with cardio-emboli stroke (3). The outcome of stroke is often affecting all aspect of individual's life. Motoric function disability is not the only impairment caused by stroke. Beyond those, reduction of social contact, lower self-esteem, and depression after stroke, also contributing factors causing deterioration quality of life (4-6). As well, stroke cause depression, the WHOQOL-BREF scores were significantly and persistently lower 1 year after stroke in patients with post-stroke depression (7).

The management of post-stroke patient is often sequentially complex. Stroke patients have considerably undergone ongoing and changing difficulties related to their disability, their self perception and coping with a new life (8). Participation in medical stroke rehabilitation is an important determinant of the adaption process in individuals after stroke (9). Medical stroke rehabilitation enables the patients to be less likely to experience worsening functional ability to perform daily activities (10). It helps not only to recover physical health but also recover psychosocial aspects.

Measuring the quality of life is the best way to assess the efficacy of intervention such as the medical stroke rehabilitation (11). Quality of life among post-stroke patients are lower compared to quality of life before stroke attack and also lower than general population, as shown study in Singapore. One of the widely used tools for assessing quality of life is the World Health Organization Quality of Life (WHO-QOL). The short version of the WHO-QOL is the WHOQOL-BREF. The tool asses quality of life for the general population that focused on individual's perception of their well-being in the domain of perception of quality of life, perception of health, physical health, psychological health, social relationship,

environment, and quality of life (12-14). The tool is useful in epidemiological studies or clinical trials, as well as to evaluate the effectiveness of treatment and disease control (15). All domains of the quality-of-life measurement assessed in this study because we would like to know in which domain the most affected among post stroke patients. In order to prioritize the domains intervened. In other hand, if we didn't measure all aspects of quality of life it can't describe validity of the measurement and as well, the quality of life in a whole.

WHOQOL-BREF had used to measure the quality of life for elderly diabetic patients. WHOQOL-BREF could measure the quality of intervention programs to increase the quality of life for elderly diabetic (16). The WHO-QOL-100 also had conducted to measure quality of life after stroke rehabilitation (17). Previous study on WHO-QOL carried on stroke patients to measure the quality of life after visiting Post-stroke clinics and also identify various factors were associated with improving quality of live such as unemployment or reduced income can cause poor quality of life anxiety or depression also cause it. Furthermore, it's very important to identify which factor influencing quality of life. The results become evidence for public health policy.

This study aimed to assess which of domains of quality of life influenced the most among post-stroke patients with use of WHOQOL-BREF scale and to investigate the influencing factors to quality of life. As well, to calculate internal consistency of WHOQOL-BREF after translation into Bahasa Indonesia in order to have a valid answer and response among Indonesian people.

## **Material and Methods**

This research was a cross-sectional study using the WHOQOL-BREF scale. Data were collected for two months from January to February 2016 from stroke patients who discharged from one hospital in Surabaya within two years after first stroke attack. This study was approved by the Institutional Health Research Ethics Committee of the primary investigator in Surabaya. All eligible subjects were invited to

the study and were adequately acquainted about the purpose and specific details of the study. Only subjects who consented to participate in the research were included in the study.

The population in this study were all stroke patients who discharge from the hospital within two years from the first attack as many as 128 patients. The study included 52 post-stroke patients (23 females and 29 males) who had first stroke attack within two years. Selected patient randomly based on the list of the patients from the hospital. These participants were those who did not have cognitive and communication problems. Post-stroke patients had Transient Ischemic Attack (TIA) patients were excluded from the study.

The World Health Organization WHOQOL-BREF questionnaire and those questions divided into six domains were used to assess the quality of life, as the dependent variables. The original version of the questionnaire was provided by the WHO and approved for utilization and translation into Bahasa Indonesia. WHOQOL-BREF consists of 26 questions: two questions assess the domains on perception of quality of life and perception of health of the patients; and the rest of the 24 questions were categorized into another 4 domains: Physical Health with 7 items; psychological health with 6 items; social relationship with 3 items; and environmental health with 8 items. All questions were scored by Likert scale of 5-points. The total score of each domain was calculated according to the scores tables guidelines provided by WHO in the original WHOQOL-BREF. A higher score means a higher quality of life.

Other additional variables collected included gender, age, education level, marital status, and participation in medical stroke rehabilitation programs as independent variables. The age of participants was categorized into two groups: 50 years old and less or over 50 years old. The youngest age was 37 years old and the oldest was 78 years old. The education level was represented by categories into three groups: elementary, secondary, or university. The marital status was categorized into two groups: single or divorced and married. The participation in medical stroke rehabilitation was categorized into two groups: following or not following medical stroke rehabilitation programs.

Statistical analysis was performed using descriptive analysis that included frequencies, percentages, means, and standard deviation (SD). Cronbach's alpha (-C, internal consistency index) was used to estimate the reliability of the WHOQOL-BREF questionnaire in our study sample. The acceptable value of Cronbach's alpha was  $0.70 < -C < 0.95$ . The associations and differences of the patients' characteristics and their QoL was examined and compared by using independent *t* test and ANOVA. Linear multiple regression analysis was performed to assess the associated factors to quality of life and level of significance was set at  $p \leq 0.05$ .

## Results

The baseline demographic characteristics of the 52 post-stroke patients are shown in Table 1. Participants were primarily male (55.8%), the mean age of the sample was 58.72 years with range 48.77 till 67.77 years old and dominated by 50 years old group and over (76.9%). Most of those patients had at least a secondary education level (40.4%), married (80.8%), and had approximately ratio 1:1 of those whom followed medical stroke rehabilitation as compared to those whom do not. In this study, Cronbach's alpha value was applied to examine the internal consistency of the WHOQOL-BREF scale (26 items) and the value was 0.797, which is considered acceptable.

Amongst all the different domains of WHOQOL-BREF, the highest and the lowest mean was found for psychological health (mean = 75.44) and perception of health (mean = 49.03), respectively as seen in Table 2. It means that post-stroke patients had the best quality of life in psychological health domain. On the other hand, amongst those patients had the lowest of quality of life for perception of health domain. According to age group, there's significant difference in psychological health ( $p= 0.018$ ) and social relationship ( $p=0.033$ ) between patients of 50 years or less and those aged over 50 years. And the score was higher amongst post-stroke patients aged 50 years or less. For educational level, there's significant difference for quality of life in domain of perception of health ( $p=0.002$ ). The highest score was 67.12

for post-stroke patients who graduated from University. For following medical rehabilitation, it's found significant difference in the domain of social relationship ( $p = 0.047$ ) and the score for post-stroke patients who followed medical rehabilitation was higher than those patients who did not follow any medical rehabilitation programs, namely 69.80 and 62.59 respectively (table 2). There's no significant difference in quality of life between patients with different marital status.

Figure 1 illustrated the comparison of the quality of life between the group of patients who followed medical stroke rehabilitation with those who did not. The results indicated that patients whom followed medical stroke rehabilitation scored higher than the group of patients whom did not. The results also showed that the group of patients who followed medical stroke rehabilitation scored higher in all the four domains of the questionnaire, as well as two additional domains of general quality of life. The highest domain score was psychological health which was evident in those patients whom followed medical stroke rehabilitation. But, the highest deviation domain score between the two groups was physical health score, with 9.69 deviation score.

Cronbach alpha coefficient was performed to examine internal consistency of all of the questions in the WHOQOL-BREF (26 items) and all domains of the questionnaire. The Cronbach alpha coefficient of WHOQOL-BREF was in the range of acceptable value, it's 0.791. Cronbach alpha analysis of each domain indicated the following values: physical health domain (0.747), psychological health domain (0.744), social relationship domain (0.466) and environmental health domain (0.401).

Table 3 illustrates the results of Backward Multiple Linear Regression. It's found that psychological health domain of QoL was significantly associated to both age group ( $p=0.021$ ) and education level ( $p=0.035$ ). Physical health domain of QoL was also significantly associated to education level ( $p=0.005$ ). Similarly, perception of health of QoL was associated to education level ( $p=0.003$ ). For another domain, social relationship was associated to participation of medical rehabilitation ( $p=0.026$ ).

## Discussion

One of the objectives of this study was to evaluate the internal consistency (reliability) of WHOQOL-BREF questionnaire in post-stroke patients. Reliability analysis in this study indicated an acceptable internal consistency of WHOQOL-BREF scale ( $\alpha = 0.791$ ) and for each of its domains were high, except for social relationships domain that is partly low ( $\alpha = 0.466$ ) and environmental health domain ( $\alpha = 0.401$ ). Using the same instrument, Lee et al, investigated validation WHOQOL-BREF in patients with stroke and spinal cord injury. In this study, the value of Cronbach's alpha ranged from 0.603 to 0.875, values that are higher than the present study. Lower internal consistency in social relationships domain can be caused by the small number of questions (3 items) (18).

Meanwhile, the other objectives of this study were to evaluate the quality of life in patients two years after stroke and to determine which of these factors influence QOL most. Using the WHOQOL-BREF tool, we can observe that in the four main domains, the highest mean satisfaction rating was found for psychological health (mean = 75.44), implying good self-esteem, spiritually, cognition, and positive feeling. Moreover, the lowest mean satisfaction was showed for physical health (mean = 62.85), indicating not very good activity of daily living, dependence on medication and medicinal aids, less energy and fatigue, pain and discomfort (14).

In some cases, the coefficient of correlation is not statistically significant when comparing the WHOQOL-BREF mean score between patient's characteristic and their QOL. The results indicated a significance association between the mean score of social relationship and psychological health in patients which was evidently higher score among patients who aged 50 years old and less than those who aged over 50 years old. The results of this study are consistent with the results of the study by Deborah et al and Celeb et al. Increasing age of post-stroke patients was followed by a decrease in the total score of quality of life, which can be interpreted as low quality of life (6,19). The finding also showed that every one year increase in age causing a decrease in the score of quality of life by 0.226 (20). Older post-



stroke patients are more at risk of having depression and anterior circulation stroke (7, 21). Indeed, depression has long been the most common psychological disorder reported in stroke survivors, and might affect physiological defects (22).

The mean score also showed that the social health of patients whom followed medical stroke rehabilitation was significantly better than those whom do not. This is echoed in research conducted by Rice et al where there were significant improvements in physical function, mobility, and confidence in post-stroke patients who attended medical rehabilitation (23). It was consistent with this study as seen in Figure 1. There are some significant association amongst post-stroke patient whom followed medical stroke rehabilitation with better quality of life in all WHOQOL-BREF domains. The most significant results was “social relations”, which refers to intimate emotional interactions and a sense of social support, for example, from family or friends (24). Medical stroke rehabilitation were evident to be a means of post-stroke patients to connect with others who had the same disabilities. Group exercises may be more effective and enjoyable. A local pilot study showed that group exercises supervised by a physiotherapist in a social center setting were effective in improving social relation, mobility and stability in elderly people who had been chronically disabled by stroke (25).

The results of multiple linear regression found that education level was an associated factor affecting perception of health, physical health, and psychological health. The existing knowledge often highlighted health disparities (26-27). The level of education is a condition that influences a person to accept or reject new knowledge that they received. Hence, a person whom have adequate educational background is likely to be positively influence in their behavior in seeking care and treatment of a disease, as well as to decide action to be and should be taken in order to treat their health problems. Patients whom have higher level of education had higher quality of life status. This is related to their positive level of adequacy of knowledge to understand and comply with doctor's suggestions related to improving their quality of life (28)

### **Study strengths and limitations**

Result of this study identified limitations of the National Institutes of Health Stroke Scale (NIHSS) and Beck Depression Inventory (BDI) which were developed in the United States of America (US), particularly in terms of clinical factors measure of stroke such as stroke severity. Other areas also may need exploration to better explain factors affecting post-stroke QOL.

However, this study evidence that the translated version of the WHOQOL-BREF into Bahasa Indonesia is found to be cross-culturally adaptable and is usable in Indonesia. It has potential to be used in the Asian and Asia Pacific region that used Bahasa Indonesia as the mode of communication. WHOQOL-BREF can be safely and effectively applied to post-stroke patients. Older post-stroke patients have been associated with negative psychological and social relationships. Yet, the findings of this study emphasize the impact of medical stroke rehabilitation which is evident to increase all WHOQOL-BREF domains score, in comparison with post-stroke patients whom do not follow such rehabilitation.

### **Conclusion**

The most **associated** factor for quality of life based on WHOQOL-BREF among post stroke patients was education level. Higher of education level cause increased score for quality of life in domain of physical health, psychological health, and perception of health. It means high education level cause good quality of life. Those patients aged 50 years old or younger was associated with good quality of life mainly psychological health domain. Participating in medical stroke rehabilitation program was associated with social relationship domain of quality of life. Internal consistency was good and adequate for all 26 questions of WHOQOL-BREF, its value was 0.791.

Education, specifically in terms of knowledge about medical stroke rehabilitation is identified as one of the most associated factors for quality of life amongst post-stroke patient. This study highlights the importance of improving communication and provision of information concerning the benefits of medical

stroke rehabilitation, which consequently, facilitate improvement of the quality of life of post-stroke patients. Structured planning of post-stroke rehabilitation programs and discharge preparation needs also be taken into consideration.

## References:

1. Mukherjee, D. and Patil, C.G.(2011). Epidemiology and the global burden of stroke. *World neurosurgery*, 76(6), pp.S85-S90.
2. Go AS, Mozaffarian D, Roger VI, Benjamin EJ, Berry JD, Blaha MJ, et al (2020). Heart disease and stroke Statistics–2013 update: a report from the American heart association. *Circulation. J Am Heart Assoc*: 129.
3. Amalia L, Furqani MA, Parwati I, Rizal A, Panigoro R (2020). Correlation between Alberta Stroke Programme Early Computed Tomography Score (ASPECTS) and National Institute of Health Stroke Score (NIHSS) in Ischemic Stroke. *Sains Malaysiana* 49 (5): 1115-1120
4. Marchant NL, Reed BR, Sanossian N, Madison CM, Kriger S, Dhada R, Mack WJ, DeCarli C, Weiner MW, Mungas DM, Chui HC (2013). The aging brain and cognition: contribution of vascular injury and aβ to mild cognitive dysfunction. *JAMA neurology*. Apr 1;70(4):488-95.
5. Marta Dabrowska-Bender, Magdalena Milewska, Aleksandra Golabek, Aneta Duda-Zalewska, Anna Staniszevska (2017). The impact of ischemic cerebral stroke on the quality of life of patients based on clinical, social, and psychoemotional factors. *Journal of Stroke and Cerebrovascular Disease*, 26: 101-107
6. De Wit L, Theuns P, Dejaeger E, Devos S, Gantenbein AR, Kerckhofs E, Schuback B, Schupp W, Putman K (2017). Long-term impact of stroke on patients' health-related quality of life. *Disability and rehabilitation*. Jul 3;39(14):1435-40.
7. Kim E-S, Kim J-W, Kang H-J, Bae K-Y, Kim S-W, Kim J-T, Park M-S, Cho K-Y, Kim J-M (2018). Longitudinal Impact of Depression on Quality of Life in Stroke Patients. *Psychiatry Investig*, 15(2): 141-146

8. Allesen H. Body, coping and self-identity. A qualitative 5-year follow-up study of stroke (2015). *Disabil Rehabil* [serial on i2014. Available from: <http://dx.doi.org/10.3109/09638288.2013.788217>
9. Brands IM, Wade DT, Stapert SZ, van Heugten CM (2012). The adaptation process following acute onset disability: an interactive two dimensional approach applied to acquired brain injury. *Clin Rehabil*, 26: 840-852.
10. Chen Y, Abel KT, Janecek JT, Chen Y, Zheng K, Cramer SC (2019). Home-based technologies for stroke rehabilitation: A systematic review. *International journal of medical informatics*. Mar 1;123:11-22.
11. Schindel D, Schneider A, Grittner U, Jöbges M, Schenk L (2021). Quality of life after stroke rehabilitation discharge: a 12-month longitudinal study. *Disability and rehabilitation*. Jul 31;43(16):2332-41.
12. Purba FD, Hunfeld JA, Iskandarsyah A, Fitriana TS, Sadarjoen SS, Passchier J, Busschbach JJ (2018). Quality of life of the Indonesian general population: Test-retest reliability and population norms of the EQ-5D-5L and WHOQOL-BREF. *PLoS One*. May 11;13(5):e0197098.
13. Reba K, Birhane BW, Gutema H (2019). Validity and reliability of the Amharic version of the World Health Organization's quality of life questionnaire (WHOQOL-BREF) in patients with diagnosed type 2 diabetes in Felege Hiwot referral hospital, Ethiopia. *Journal of diabetes research*. May 6;2019.
14. Ilić I, Šipetić-Grujičić S, Grujičić J, Živanović Mačužić I, Kocić S, Ilić M (2019). Psychometric properties of the world health organization's quality of life (WHOQOL-BREF) questionnaire in medical students. *Medicina*. Dec;55(12):772.
15. Pinto JM, Arenillas JI, Martín Nogueras AM, Ramos González J, Gómez Gómez FP (2010). The quality of life of asthmatic patients evaluated by the WHOQOL-BREF and the SGRQ. *Fisioterapia (Madr)*, 32: 116-22.

16. Jahromi MK, Ramezanli S, Taheri L (2015). Effectiveness of diabetes self-management education on quality of life in diabetic elderly females. *Global journal of health science*, Jan 7(1): 10.
17. Giaquinto S, Giachetti I, Spiridigliozzi C, Nolfi G (2010). Quality of life after stroke in a rehabilitation setting. *Clinical and Experimental Hypertension*, Nov 1; 32 (7): 426-30
18. Lee Kuem Ju, Hye In Jan, Hyun Choi (2017). Korean translation and validation of the WHOQOL-DIS for people with spinal cord injury and stroke. *Disability and Health Journal*, 30: 1-5
19. Celeb AG., Aderonke O., Akinpelu (2012). Quality of life of nigerian stroke survivors during first 12 months post-stroke. *Hong Kong Physiotherapy Journal*, 30: 18-24.
20. She R, Yan Z, Hao Y, Zhang Z, Du Y, Liang Y, Vetrano DL, Dekker J, Bai B, Lau JT, Qiu C (2021). Health-related quality of life after first-ever acute ischemic stroke: associations with cardiovascular health metrics. *Quality of Life Research*. May 1:1-1.
21. Laurie Theeke, A. Noelle Lucke-Wold, Jennifer Mallow, Patricia Horstman (2017). Life After Stroke in Appalachia. *International Journal of Nursing Sciences*, 4: 105-111.
22. Kim ES, Kim JW, Kang HJ, Bae KY, Kim SW, Kim JT, Park MS, Cho KH, Kim JM (2018). Longitudinal impact of depression on quality of life in stroke patients. *Psychiatry investigation*. Feb;15(2):141.
23. Rice D, Janzen S, McIntyre A, Vermeer J, Britt E, Teasell R (2016). Comprehensive outpatient rehabilitation program: Hospital-based stroke patient rehabilitation. *Journal stroke cerebrovascular disease*, 25(5): 1158-64.

24. Marta Dabrowska-Bender, Magdalena Milewska, Aleksandra Golabek, Aneta Duda-Zalewska, Anna Staniszevska (2017). The impact of ischemic cerebral stroke on the quality of life of patients based on clinical, social, and psychoemotional factors. *Journal of Stroke and Cerebrovascular Disease*, 26: 101-107
25. Sarfo FS, Adamu S, Awuah D, Sarfo-Kantanka O, Ovbiagele B (2017). Potential role of tele-rehabilitation to address barriers to implementation of physical therapy among West African stroke survivors: A cross-sectional survey. *Journal of the neurological sciences*. Oct 15;381:203-8.
26. Odonkor CA, Esparza R, Flores LE, Verduzco-Gutierrez M, Escalon MX, Solinsky R, Silver JK (2021). Disparities in health Care for Black Patients in physical medicine and rehabilitation in the United States: a narrative review. *PM&R*. Feb;13(2):180-203.
27. Chuluunbaatar E, Chou Y-J, Pu C (2016). Quality of Life of Stroke Survivors and Their Informal Caregivers: A Prospective Study. *Disability and Health Journal* 9: 306-312
28. Pakpahan M, Siregar D, Susilawaty A, Tasnim T, Ramdany R, Manurung EI, Sianturi E, Tompunu MR, Sitanggang YF, Maisyarah M (2021). Promosi Kesehatan dan Perilaku Kesehatan. *Yayasan Kita Menulis*; Feb 19.

---

**Decision on submission to Clinical Epidemiology and Global Health**

4 messages

---

**Clinical Epidemiology and Global Health** <em@editorialmanager.com>  
Reply-To: Clinical Epidemiology and Global Health <support@elsevier.com>  
To: Santi Martini <santi-m@fkm.unair.ac.id>

Fri, Dec 24, 2021 at 4:39 PM

Manuscript Number: CEGH-D-21-00557R1

Assessing Quality of Life and Associated Factors in Post-Stroke Patients Using the World Health Organization Abbreviated Generic Quality Of Life Questionnaire (WHOQOL-BREF)

Dear Dr. Martini,

Thank you for submitting your manuscript to Clinical Epidemiology and Global Health.

I am pleased to inform you that your manuscript has been accepted for publication.

My comments, and any reviewer comments, are below.

Your accepted manuscript will now be transferred to our production department. We will create a proof which you will be asked to check, and you will also be asked to complete a number of online forms required for publication. If we need additional information from you during the production process, we will contact you directly.

We appreciate and value your contribution to Clinical Epidemiology and Global Health. We regularly invite authors of recently published articles to participate in the peer review process. You are now part of the Clinical Epidemiology and Global Health reviewer pool. We look forward to your continued participation in our journal, and we hope you will consider us again for future submissions.

Kind regards,  
Shally Awasthi  
Editor-in-Chief

Clinical Epidemiology and Global Health

Editor and Reviewer comments:

More information and support

FAQ: When and how will I receive the proofs of my article?  
[https://service.elsevier.com/app/answers/detail/a\\_id/6007/p/10592/supporthub/publishing/related/](https://service.elsevier.com/app/answers/detail/a_id/6007/p/10592/supporthub/publishing/related/)You will find information relevant for you as an author on Elsevier's Author Hub: <https://www.elsevier.com/authors>FAQ: How can I reset a forgotten password?  
[https://service.elsevier.com/app/answers/detail/a\\_id/28452/supporthub/publishing/](https://service.elsevier.com/app/answers/detail/a_id/28452/supporthub/publishing/)For further assistance, please visit our customer service site: <https://service.elsevier.com/app/home/supporthub/publishing/>  
Here you can search for solutions on a range of topics, find answers to frequently asked questions, and learn more about Editorial Manager via interactive tutorials. You can also talk 24/7 to our customer support team by phone and 24/7 by live chat and email

#AU\_CEGH#





## Original article

# Assessing quality of life and associated factors in post-stroke patients using the world health organization abbreviated generic quality of life questionnaire (WHOQOL-BREF)

Santi Martini <sup>a,\*</sup>, Defi Amalia Setia Ningrum <sup>b</sup>, Khadizah H. Abdul-Mumin <sup>c</sup>, Chung Yi-Li <sup>d</sup>

<sup>a</sup> Division of Epidemiology, Faculty of Public Health, Universitas Airlangga, Indonesia

<sup>b</sup> FETP Graduate, University of Indonesia, Indonesia

<sup>c</sup> Institute of Health Sciences (PAPRSB, IHS), Universiti Brunei Darussalam (UBD), Brunei

<sup>d</sup> Department of Public Health, College of Medicine, National Cheng Kung University, Taiwan



## ARTICLE INFO

## Keywords:

Educational level  
Internal consistency  
Post-stroke  
Quality of life  
WHOQOL-BREF

## ABSTRACT

**Introduction:** This study piloted the World Health Organization Quality of Life short version form (WHOQOL-BREF) to compare the quality of life (QoL) in post-stroke patients who followed the medical stroke rehabilitation program and those whom did not. This study determined which quality of life domains were mostly affected by stroke, and whether there are influencing factors other than rehabilitation.

**Methods:** A cross-sectional study was conducted at Public Hospital in Surabaya. Patients with stroke less than two years since the first attack were included in this study (n = 52). Cronbach's alpha test was performed to assess the internal consistency of WHOQOL-BREF questionnaire. The independent *t*-test and ANOVA were used to compare the differences between patient's characteristics and the six domains of QoL (i.e., perception of QoL, perception of health, physical of health, psychological health, social relationship, and environmental health). Multiple linear regression was performed to assess the influential factors of QoL.

**Results:** post-stroke patients aged more than 50 years old (76.9%), male (55.8%). But, only 48.1% of those patients participating in medical rehabilitation programs. Age group was significantly associated with QoL in the psychological health domain (p = 0.021); participating in medical stroke rehabilitation programs was significantly associated with social relationship domain (p = 0.026) and the education level was significantly associated to physical health (p = 0.005), psychological health (p = 0.035) and perception of health (p = 0.003).

**Conclusion:** Lower education level was significantly associated with a low perception of health, one of QoL domains; and it was the most influential factor of QoL among post-stroke patients.

## 1. Introduction

Epidemiological studies evident that approximately 5.5 million people worldwide died annually due to stroke with the estimation that the global prevalence of stroke will gradually increase to as high as 21.9% by 2030.<sup>1,2</sup> Stroke is one of the most frequent causes of disability worldwide with 4.4 million disability-adjusted life-years lost.<sup>1</sup> Most of stroke is ischemic stroke and based on CT scan of ischemic stroke patients, there are 58,6% subject with lacunar stroke, 20,7% with large artery atherosclerotic (LAA) stroke, and 20,7% with cardio-emboli stroke.<sup>3</sup> The outcome of stroke is often affecting all aspect of individual's life. Motoric function disability is not the only impairment

caused by stroke. Beyond those, reduction of social contact, lower self-esteem, and depression after stroke, also contributing factors causing deterioration quality of life.<sup>4-6</sup> As well, stroke cause depression, the WHOQOL-BREF scores were significantly and persistently lower 1 year after stroke in patients with post-stroke depression.<sup>7</sup>

The management of post-stroke patient is often sequentially complex. Stroke patients have considerably undergone ongoing and changing difficulties related to their disability, their self perception and coping with a new life.<sup>8</sup> Participation in medical stroke rehabilitation is an important determinant of the adaption process in individuals after stroke.<sup>9</sup> Medical stroke rehabilitation enables the patients to be less likely to experience worsening functional ability to perform daily

\* Corresponding author. Division of Epidemiology, Faculty of Public Health, Universitas Airlangga, Kampus C Unair, Surabaya, 60115, East Java, Indonesia.

E-mail addresses: [santi-m@fkm.unair.ac.id](mailto:santi-m@fkm.unair.ac.id) (S. Martini), [defi.amalia@gmail.com](mailto:defi.amalia@gmail.com) (D.A. Setia Ningrum), [khadizah.mumin@ubd.edu.bn](mailto:khadizah.mumin@ubd.edu.bn) (K.H. Abdul-Mumin), [cyli99@mail.ncku.edu.tw](mailto:cyli99@mail.ncku.edu.tw) (C. Yi-Li).

<https://doi.org/10.1016/j.cegh.2021.100941>

Received 4 September 2021; Received in revised form 19 December 2021; Accepted 24 December 2021

Available online 31 December 2021

2213-3984/© 2022 The Authors. Published by Elsevier B.V. on behalf of INDIACLEN. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

activities.<sup>10</sup> It helps not only to recover physical health but also recover psychosocial aspects.

Measuring the quality of life is the best way to assess the efficacy of intervention such as the medical stroke rehabilitation.<sup>11</sup> Quality of life among post-stroke patients are lower compared to quality of life before stroke attack and also lower than general population, as shown study in Singapore. One of the widely used tools for assessing quality of life is the World Health Organization Quality of Life (WHO-QOL). The short version of the WHO-QOL is the WHOQOL-BREF. The tool assess quality of life for the general population that focused on individual's perception of their well-being in the domain of perception of quality of life, perception of health, physical health, psychological health, social relationship, environment, and quality of life.<sup>12-14</sup> The tool is useful in epidemiological studies or clinical trials, as well as to evaluate the effectiveness of treatment and disease control.<sup>15</sup> All domains of the quality-of-life measurement assessed in this study because we would like to know in which domain the most affected among post stroke patients. In order to prioritize the domains intervened. In other hand, if we didn't measure all aspects of quality of life it can't describe validity of the measurement and as well, the quality of life in a whole.

WHOQOL-BREF had used to measure the quality of life for elderly diabetic patients. WHOQOL-BREF could measure the quality of intervention programs to increase the quality of life for elderly diabetic.<sup>16</sup> The WHO-QOL-100 also had conducted to measure quality of life after stroke rehabilitation.<sup>17</sup> Previous study on WHO-QOL carried on stroke patients to measure the quality of life after visiting Post-stroke clinics and also identify various factors were associated with improving quality of live such as unemployment or reduced income can cause poor quality of life anxiety or depression also cause it. Furthermore, it's very important to identify which factor influencing quality of life. The results become evidence for public health policy.

This study aimed to assess which of domains of quality of life influenced the most among post-stroke patients with use of WHOQOL-BREF scale and to investigate the influencing factors to quality of life. As well, to calculate internal consistency of WHOQOL-BREF after translation into Bahasa Indonesia in order to have a valid answer and response among Indonesian people.

## 2. Material and methods

This research was a cross-sectional study using the WHOQOL-BREF scale. Data were collected for two months from January to February 2016 from stroke patients who discharged from one hospital in Surabaya within two years after first stroke attack. This study was approved by the Institutional Health Research Ethics Committee of the primary investigator in Surabaya. All eligible subjects were invited to the study and were adequately acquainted about the purpose and specific details of the study. Only subjects who consented to participate in the research were included in the study.

The population in this study were all stroke patients who discharge from the hospital within two years from the first attack as many as 128 patients. The study included 52 post-stroke patients (23 females and 29 males) who had first stroke attack within two years. Selected patient randomly based on the list of the patients from the hospital. These participants were those who did not have cognitive and communication problems. Post-stroke patients had Transient Ischemic Attack (TIA) patients were excluded from the study.

The World Health Organization WHOQOL-BREF questionnaire and those questions divided into six domains were used to assess the quality of life, as the dependent variables. The original version of the questionnaire was provided by the WHO and approved for utilization and translation into Bahasa Indonesia. WHOQOL-BREF consists of 26 questions: two questions assess the domains on perception of quality of life and perception of health of the patients; and the rest of the 24 questions were categorized into another 4 domains: Physical Health with 7 items; psychological health with 6 items; social relationship with 3 items; and

environmental health with 8 items. All questions were scored by Likert scale of 5-points. The total score of each domain was calculated according to the scores tables guidelines provided by WHO in the original WHOQOL-BREF. A higher score means a higher quality of life.

Other additional variables collected included gender, age, education level, marital status, and participation in medical stroke rehabilitation programs as independent variables. The age of participants was categorized into two groups: 50 years old and less or over 50 years old. The youngest age was 37 years old and the oldest was 78 years old. The education level was represented by categories into three groups: elementary, secondary, or university. The marital status was categorized into two groups: single or divorced and married. The participation in medical stroke rehabilitation was categorized into two groups: following or not following medical stroke rehabilitation programs.

Statistical analysis was performed using descriptive analysis that included frequencies, means, and standard deviation (SD). Cronbach's alpha (-C, internal consistency index) was used to estimate the reliability of the WHOQOL-BREF questionnaire in our study sample. The acceptable value of Cronbach's alpha was  $0.70 < -C < 0.95$ . The associations and differences of the patients' characteristics and their QoL was examined and compared by using independent t-test and ANOVA. Linear multiple regression analysis was performed to assess the associated factors to quality of life and level of significance was set at  $p \leq 0.05$ .

## 3. Results

The baseline demographic characteristics of the 52 post-stroke patients are shown in Table 1. Participants were primarily male (55.8%), the mean age of the sample was 58.72 years with range 48.77 till 67.77 years old and dominated by 50 years old group and over (76.9%). Most of those patients had at least a secondary education level (40.4%), married (80.8%), and had approximately ratio 1:1 of those whom followed medical stroke rehabilitation as compared to those whom do not. In this study, Cronbach's alpha value was applied to examine the internal consistency of the WHOQOL-BREF scale (26 items) and the value was 0.797, which is considered acceptable.

Amongst all the different domains of WHOQOL-BREF, the highest and the lowest mean was found for psychological health (mean = 75.44) and perception of health (mean = 49.03), respectively as seen in Table 2. It means that post-stroke patients had the best quality of life in psychological health domain. On the other hand, amongst those patients had the lowest of quality of life for perception of health domain. According to age group, there's significant difference in psychological health ( $p = 0.018$ ) and social relationship ( $p = 0.033$ ) between patients of 50 years or less and those aged over 50 years. And the score was higher amongst post-stroke patients aged 50 years or less. For

**Table 1**  
Characteristic of the sample (n = 52).

Characteristic	n (%)
Age (years)	58.27 ± 9.5
≤ 50	12 (23.1)
> 50	40 (76.9)
Gender	
Male	29 (55.8)
Female	23 (44.2)
Education level	
Elementary	15 (28.8)
Secondary	21 (40.4)
University	16 (30.8)
Marital status	
Single/divorced	10 (19.2)
Married	42 (80.8)
Medical rehabilitation	
Yes	25 (48.1)
No	27 (51.9)


**Table 2**  
Comparison of the WHOQOL-BREF mean scores according to gender, age, education level, marital status, and medical rehabilitation.

	Domain					
	Perception of QOL Mean ± SD	Perception of health Mean ± SD	Physical health Mean ± SD	Psychological health Mean ± SD	Social relationship Mean ± SD	Enviromental health Mean ± SD
Total	62.50 ± 16.789	49.03 ± 17.823	62.85 ± 19.031	75.44 ± 20.165	66.06 ± 1.311	72.11 ± 12.607
Gender						
Male	62.93 ± 15.840	52.59 ± 16.830	62.79 ± 21.222	76.59 ± 20.916	65.76 ± 12.858	73.17 ± 11.142
Female	61.96 ± 18.262	44.56 ± 18.397	62.91 ± 16.320	74.00 ± 19.544	66.43 ± 13.720	70.78 ± 14.390
p value	0.838	0.108	0.982	0.648	0.856	0.503
Age (years)						
≤ 50	66.67 ± 16.283	56.25 ± 15.539	63.17 ± 18.473	85.00 ± 12.778	70.92 ± 6.127	72.42 ± 12.479
> 50	61.25 ± 16.937	46.87 ± 18.07	62.75 ± 19.425	72.57 ± 21.194	64.60 ± 14.318	72.025 ± 12.801
p value	0.332	0.111	0.947	0.018*	0.033*	0.926
Education level						
Elementary	58.33 ± 18.094	36.67 ± 15.100	56.00 ± 17.708	67.53 ± 18.909	62.20 ± 16.772	68.87 ± 13.400
Secondary	60.71 ± 14.940	51.19 ± 16.726	64.48 ± 15.725	79.86 ± 17.316	66.14 ± 9.040	73.04 ± 10.500
University	68.75 ± 17.078	57.81 ± 15.052	67.12 ± 23.215	77.06 ± 23.589	69.56 ± 13.623	73.94 ± 14.507
p value	0186	0.002*	0.238	0.183	0.301	0.494
Marital status						
Single/ divorced	61.31 ± 16.750	50.00 ± 18.313	62.88 ± 19.215	76.43 ± 19.067	66.59 ± 12.172	71.714 ± 12.377
Married	67.50 ± 16.874	45.00 ± 15.811	62.70 ± 19.241	71.30 ± 24.989	71.71 ± 12.377	73.800 ± 14.101
p value	0.299	0.431	0.979	0.475	0.550	0.643
Medical Stroke rehabilitation						
Yes	60.18 ± 14.310	44.44 ± 14.434	67.88 ± 17.871	79.60 ± 20.279	69.80 ± 11.965	74.36 ± 13.801
No	65.00 ± 19.094	54.00 ± 20.000	58.19 ± 19.201	71.59 ± 19.647	62.59 ± 113.397	70.037 ± 11.250
p value	0.306	0.52	0.66	0.154	0.047*	0.220

QOL = Quality of life, SD=Standard deviations, \*The mean difference is significant at ≤ 0.05 level.

educational level, there's significant difference for quality of life in domain of perception of health (p = 0.002). The highest score was 67.12 for post-stroke patients who graduated from University. For following medical rehabilitation, it's found significant difference in the domain of social relationship (p = 0.047) and the score for post-stroke patients who followed medical rehabilitation was higher than those patients who did not follow any medical rehabilitation programs, namely 69.80 and 62.59 respectively (Table 2). There's no significant difference in quality of life between patients with different marital status.

Fig. 1 illustrated the comparison of the quality of life between the group of patients who followed medical stroke rehabilitation with those who did not. The results indicated that patients whom followed medical stroke rehabilitation scored higher than the group of patients whom did not. The results also showed that the group of patients who followed medical stroke rehabilitation scored higher in all the four domains of the questionnaire, as well as two additional domains of general quality of life. The highest domain score was psychological health which was evident in those patients whom followed medical stroke rehabilitation.



**Fig. 1.** Bar chart demonstrating the influence of medical stroke rehabilitation on QoL. Patients were divided into two groups based on whether they participate or not in the medical stroke rehabilitation. The bar shows average domain score of WHOQOL-BREF.

But, the highest deviation domain score between the two groups was physical health score, with 9.69 deviation score.

Cronbach alpha coefficient was performed to examine internal consistency of all of the questions in the WHOQOL-BREF (26 items) and all domains of the questionnaire. The Cronbach alpha coefficient of WHOQOL-BREF was in the range of acceptable value, it's 0.791. Cronbach alpha analysis of each domain indicated the following values: physical health domain (0.747), psychological health domain (0.744), social relationship domain (0.466) and environmental health domain (0.401).

Table 3 illustrates the results of Backward Multiple Linear Regression. It's found that psychological health domain of QoL was significantly associated to both age group (p = 0.021) and education level (p = 0.035). Physical health domain of QoL was also significantly associated to education level (p = 0.005). Similarly, perception of health of QoL was associated to education level (p = 0.003). For another domain, social relationship was associated to participation of medical rehabilitation (p = 0.026).

#### 4. Discussion

One of the objectives of this study was to evaluate the internal consistency (reliability) of WHOQOL-BREF questionnaire in post-stroke patients. Reliability analysis in this study indicated an acceptable internal consistency of WHOQOL-BREF scale (α = 0.791) and for each of its domains were high, except for social relationships domain that is partly low (α = 0.466) and environmental health domain (α = 0.401). Using the same instrument, Lee et al., investigated validation WHOQOL-BREF in patients with stroke and spinal cord injury. In this study, the value of Cronbach's alpha ranged from 0.603 to 0.875, values that are higher than the present study. Lower internal consistency in social relationships domain can be caused by the small number of questions (3 items).<sup>18</sup>

Meanwhile, the other objectives of this study were to evaluate the quality of life in patients two years after stroke and to determine which of these factors influence QoL most. Using the WHOQOL-BREF tool, we can observe that in the four main domains, the highest mean satisfaction rating was found for psychological health (mean = 75.44), implying good self-esteem, spiritually, cognition, and positive feeling. Moreover,

**Table 3**  
Backward multiple linear regression analyses of significant factors associated with QOL.

QOL domains	Variables	Unstandardized Coefficients		Standardized Coefficients	t	P value
		B	SE	Beta		
Physical Health	Education level	0.806	0.276	0.382	0.923	0.005
Psychological Health	Education level	0.651	0.300	0.288	0.175	0.035
	Age Category	-1.973	0.826	-0.415	-0.388	0.021
Social Relationship	Participation of Medical Rehabilitation Program	1.229	0.536	0.306	0.294	0.026
Environmental Health	None	-	-	-	-	-
Perception of Health	Education level	0.218	0.071	0.399	0.080	0.003

QOL = Quality of life.

the lowest mean satisfaction was showed for physical health (mean = 62.85), indicating not very good activity of daily living, dependence on medication and medicinal aids, less energy and fatigue, pain and discomfort.<sup>14</sup>

In some cases, the coefficient of correlation is not statistically significant when comparing the WHOQOL-BREF mean score between patient's characteristic and their QOL. The results indicated a significance association between the mean score of social relationship and psychological health in patients which was evidently higher score among patients who aged 50 years old and less than those who aged over 50 years old. The results of this study are consistent with the results of the study by Deborah et al. and Celeb et al. Increasing age of post-stroke patients was followed by a decrease in the total score of quality of life, which can be interpreted as low quality of life.<sup>6,19</sup> The finding also showed that every one year increase in age causing a decrease in the score of quality of life by 0.226.<sup>20</sup> Older post-stroke patients are more at risk of having depression and anterior circulation stroke.<sup>7,21</sup> Indeed, depression has long been the most common psychological disorder reported in stroke survivors, and might affect physiological defects.<sup>22</sup>

The mean score also showed that the social health of patients whom followed medical stroke rehabilitation was significantly better than those whom do not. This is echoed in research conducted by Rice et al. where there were significant improvements in physical function, mobility, and confidence in post-stroke patients who attended medical rehabilitation.<sup>23</sup> It was consistent with this study as seen in Fig. 1. There are some significant association amongst post-stroke patient whom followed medical stroke rehabilitation with better quality of life in all WHOQOL-BREF domains. The most significant results was "social relations", which refers to intimate emotional interactions and a sense of social support, for example, from family or friends.<sup>24</sup> Medical stroke rehabilitation were evident to be a means of post-stroke patients to connect with others who had the same disabilities. Group exercises may be more effective and enjoyable. A local pilot study showed that group exercises supervised by a physiotherapist in a social center setting were effective in improving social relation, mobility and stability in elderly people who had been chronically disabled by stroke.<sup>25</sup>

The results of multiple linear regression found that education level was an associated factor affecting perception of health, physical health, and psychological health. The existing knowledge often highlighted health disparities.<sup>26,27</sup> The level of education is a condition that influences a person to accept or reject new knowledge that they received. Hence, a person whom have adequate educational background is likely to be positively influence in their behavior in seeking care and treatment of a disease, as well as to decide action to be and should be taken in order to treat their health problems. Patients whom have higher level of education had higher quality of life status. This is related to their positive level of adequacy of knowledge to understand and comply with doctor's suggestions related to improving their quality of life.<sup>28</sup>

#### 4.1. Study strengths and limitations

Result of this study identified limitations of the National Institutes of Health Stroke Scale (NIHSS) and Back Depression Inventory (BDI) which

were developed in the United States of America (US), particularly in terms of clinical factors measure of stroke such as stroke severity. Other areas also may need exploration to better explain factors affecting post-stroke QOL.

However, this study evidence that the translated version of the WHOQOL-BREF into Bahasa Indonesia is found to be cross-culturally adaptable and is useable in Indonesia. It has potential to be used in the Asian and Asia Pacific region that used Bahasa Indonesia as the mode of communication. WHOQOL-BREF can be safely and effectively applied to post-stroke patients. Older post-stroke patients have been associated with negative psychological and social relationships. Yet, the findings of this study emphasize the impact of medical stroke rehabilitation which is evident to increase all WHOQOL-BREF domains score, in comparison with post-stroke patients whom do not follow such rehabilitation.

## 5. Conclusion

The most associated factor for quality of life based on WHOQOL-BREF among post stroke patients was education level. Higher of education level cause increased score for quality of life in domain of physical health, psychological health, and perception of health. It means high education level cause good quality of life. Those patients aged 50 years old or younger was associated with good quality of life mainly psychological health domain. Participating in medical stroke rehabilitation program was associated with social relationship domain of quality of life. Internal consistency was good and adequate for all 26 questions of WHOQOL-BREF, its value was 0.791.

Education, specifically in terms of knowledge about medical stroke rehabilitation is identified as one of the most associated factors for quality of life amongst post-stroke patient. This study highlights the importance of improving communication and provision of information concerning the benefits of medical stroke rehabilitation, which consequently, facilitate improvement of the quality of life of post-stroke patients. Structured planning of post-stroke rehabilitation programs and discharge preparation needs also be taken into consideration.

## Declaration of competing interest

All Author declare there is no conflict of Interest.

This study was approved by the Institutional Health Research Ethics Committee of the primary investigator in Surabaya.

## Acknowledgements

Faculty of Public Health Universitas Airlangga.

## References

- Mukherjee D, Patil CG. Epidemiology and the global burden of stroke. *World neurosurgery*. 2011;76(6):S85-S90.
- Go AS, Mozaffarian D, Roger VL, et al. Heart disease and stroke Statistics-2013 update: a report from the American heart association. *Circulation. J Am Heart Assoc*. 2020;129.
- Amalia L, Furqani MA, Parwati I, Rizal A, Panigoro R. Correlation between Alberta stroke programme early computed tomography score (ASPECTS) and National

- Institute of health stroke score (NIHSS) in ischemic stroke. *Sains Malays.* 2020;49(5): 1115–1120.
- 4 Marchant NL, Reed BR, Sanossian N, et al. The aging brain and cognition: contribution of vascular injury and  $\alpha\beta$  to mild cognitive dysfunction. *JAMA Neurol.* 2013;70(4):488–495. Apr 1.
  - 5 Dabrowska-Bender Marta, Milewska Magdalena, Golabek Aleksandra, Duda-Zalewska Aneta, Anna Staniszevska. The impact of ischemic cerebral stroke on the quality of life of patients based on clinical, social, and psychoemotional factors. *J Stroke Cerebrovasc Dis.* 2017;26:101–107.
  - 6 De Wit L, Theuns P, Dejaeger E, et al. Long-term impact of stroke on patients' health-related quality of life. *Disabil Rehabil.* 2017;39(14):1435–1440. Jul 3.
  - 7 Kim E-S, Kim J-W, Kang H-J, et al. Longitudinal impact of depression on quality of life in stroke patients. *Psychiatry Investig.* 2018;15(2):141–146.
  - 8 Allesen H. *Body, Coping and Self-Identity. A Qualitative 5-year Follow-Up Study of Stroke.* *Disabil Rehabil* [serial on i2014]. 2015. <https://doi.org/10.3109/09638288.2013.788217>. Available from.
  - 9 Brands IM, Wade DT, Stapert SZ, van Heugten CM. The adaptation process following acute onset disability: an interactive two dimensional approach applied to acquired brain injury. *Clin Rehabil.* 2012;26:840–852.
  - 10 Chen Y, Abel KT, Janeczek JT, Chen Y, Zheng K, Cramer SC. Home-based technologies for stroke rehabilitation: a systematic review. *Int J Med Inf.* 2019;123:11–22. Mar 1.
  - 11 Schindel D, Schneider A, Grittner U, Jöbges M, Schenk L. Quality of life after stroke rehabilitation discharge: a 12-month longitudinal study. *Disabil Rehabil.* 2021;43(16):2332–2341. Jul 31.
  - 12 Purba FD, Hunfeld JA, Iskandarsyah A, et al. Quality of life of the Indonesian general population: test-retest reliability and population norms of the EQ-5D-5L and WHOQOL-BREF. *PLoS One.* 2018;13(5), e0197098. May 11.
  - 13 Reba K, Birhane BW, Gutema H. Validity and reliability of the Amharic version of the World Health Organization's quality of life questionnaire (WHOQOL-BREF) in patients with diagnosed type 2 diabetes in Felege Hiwot referral hospital, Ethiopia. *J Diabetes Res.* 2019. May 6;2019, 6 pages.
  - 14 Ilić I, Šipetić-Grujičić S, Grujičić J, Živanović Macuzić I, Kocić S, Ilić M. Psychometric properties of the world health organization's quality of life (WHOQOL-BREF) questionnaire in medical students. *Medicina.* 2019;55(12):772. Dec.
  - 15 Pinto JM, Arenillas JI, Martín Nogueiras AM, Ramos González J, Gómez Gómez FP. The quality of life of asthmatic patients evaluated by the WHOQOL-BREF and the SGRQ. *Fisioterapia (Madr).* 2010;32:116–122.
  - 16 Jahromi MK, Ramezanli S, Taheri L. Effectiveness of diabetes self-management education on quality of life in diabetic elderly females. *Global J Health Sci.* 2015;(1): 10. Jan 7.
  - 17 Giaquinto S, Giachetti I, Spiridigliozzi C, Nolfi G. Quality of life after stroke in a rehabilitation setting. *Clin Exp Hypertens.* 2010;32(7):426–430. Nov 1.
  - 18 Ju Lee Kuem, Jan Hye In, Choi Hyun. Korean translation and validation of the WHOQOL-DIS for people with spinal cord injury and stroke. *Disabil.Health J.* 2017; 30:1–5.
  - 19 Celeb AG, Aderonke O, Akinpelu. Quality of life of nigerian stroke survivors during first 12 months post-stroke. *Hong Kong Physiother J.* 2012;30:18–24.
  - 20 She R, Yan Z, Hao Y, et al. Health-related quality of life after first-ever acute ischemic stroke: associations with cardiovascular health metrics. *Qual Life Res.* 2021;(May 1), 1-1.
  - 21 Theeke Laurie, Lucke-Wold A Noelle, Mallow Jennifer, Horstman Patricia. Life after stroke in appalacia. *Int J Nurs Sci.* 2017;4:105–111.
  - 22 Kim ES, Kim JW, Kang HJ, et al. Longitudinal impact of depression on quality of life in stroke patients. *Psychiatry investigation.* Feb. 2018;15(2):141.
  - 23 Rice D, Janzen S, Mcintyre A, Vermeer J, Britt E, Teasell R. Comprehensive outpatient rehabilitation program: hospital-based stroke patient rehabilitation. *J Stroke Cerebrovasc Dis.* 2016;25(5):1158–1164.
  - 24 Dabrowska-Bender Marta, Milewska Magdalena, Golabek Aleksandra, Duda-Zalewska Aneta, Anna Staniszevska. The impact of ischemic cerebral stroke on the quality of life of patients based on clinical, social, and psychoemotional factors. *J Stroke Cerebrovasc Dis.* 2017;26:101–107.
  - 25 Sarfo FS, Adamu S, Awuah D, Sarfo-Kantanka O, Ovbiagele B. Potential role of tele-rehabilitation to address barriers to implementation of physical therapy among West African stroke survivors: a cross-sectional survey. *J Neurol Sci.* 2017;381:203–208. Oct 15.
  - 26 Odonkor CA, Esparza R, Flores LE, et al. Disparities in health Care for Black Patients in physical medicine and rehabilitation in the United States: a narrative review. *PM&R.* Feb. 2021;13(2):180–203.
  - 27 Chuluunbaatar E, Chou Y-J, Pu C. Quality of life of stroke survivors and their informal caregivers: a prospective study. *Disabil.Health J.* 2016;9:306–312.
  - 28 Pakpahan M, Siregar D, Susilawaty A, et al. *Promosi Kesehatan Dan Perilaku Kesehatan.* Yayasan Kita Menulis; 2021. Feb 19.