

The concentration of the potentially toxic element (PTEs) in black tea (*Camellia sinensis*) consumed in Iran: a systematic review, meta-analysis, and probabilistic risk assessment study

by Trias Mahmudiono

Submission date: 30-Mar-2023 01:22PM (UTC+0800)

Submission ID: 2050720066

File name: BELUM1.pdf (1.78M)

Word count: 5157

Character count: 25961



13

International Journal of Environmental Analytical Chemistry

 Taylor & Francis
Taylor & Francis GroupISSN: (Print) (Online) Journal homepage: <https://www.tandfonline.com/loi/geac20>

The concentration of the potentially toxic element (PTEs) in black tea (*Camellia sinensis*) consumed in Iran: a systematic review, meta-analysis, and probabilistic risk assessment study

Yadolah Fakhri, Hasti Daraei, Somayeh Hoseinvandtabar, Fereshteh Mehri, Trias Mahmudiono & Amin Mousavi Khaneghah

To cite this article: Yadolah Fakhri, Hasti Daraei, Somayeh Hoseinvandtabar, Fereshteh Mehri, Trias Mahmudiono & Amin Mousavi Khaneghah (2022): The concentration of the potentially toxic element (PTEs) in black tea (*Camellia sinensis*) consumed in Iran: a systematic review, meta-analysis, and probabilistic risk assessment study, International Journal of Environmental Analytical Chemistry, DOI: [10.1080/03067319.2022.2118596](https://doi.org/10.1080/03067319.2022.2118596)

To link to this article: <https://doi.org/10.1080/03067319.2022.2118596>

[View supplementary material](#)

Published online: 01 Sep 2022.

[Submit your article to this journal](#)

Article views: 44

[View related articles](#)[View Crossmark data](#)

Full Terms & Conditions of access and use can be found at
<https://www.tandfonline.com/action/journalInformation?journalCode=geac20>



15 The concentration of the potentially toxic element (PTEs) in black tea (*Camellia sinensis*) consumed in Iran: a systematic review, meta-analysis, and probabilistic risk assessment study

Yadolah Fakhri^a, Hasti Daraei^{b,c}, Somayeh Hoseinvandtabar^d, Fereshteh Mehri^e, Trias Mahmudiono^f and Amin Mousavi Khaneghah^g

^aFood Health Research Center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran;

^bEnvironmental Health Engineering Research Center, Kerman University of Medical Sciences, Kerman, Iran;

^cDepartment of Environmental Health Engineering, Faculty of Public Health, Kerman University of Medical Sciences, Kerman, Iran; ^dStudent Research Committee, School of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran;

^eNutrition Health Research Center, Hamadan University of Medical Sciences, Hamadan, Iran; ^fDepartment of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia; ^gDepartment of Fruit and Vegetable Product Technology, Prof. Waclaw Dąbrowski Institute of Agricultural and Food Biotechnology – State Research Institute, Warsaw, Poland

ABSTRACT

The contamination of tea as one most widely consumed beverage in Iran with the potentially toxic elements (PTEs) leads to adverse health effects. In the current study, a meta-analysis and probabilistic risk assessment regarding PTEs concentration in tea samples from Iran were performed using international databases (PubMed and Scopus) and national databases (SID, Irandoc, and Magiran) from 2005 to 1 June 2022. Data were analysed using the random-effects model based on Iranian and imported tea subgroups, and a probabilistic health risk assessment was conducted using the Monte Carlo simulation (MCS). The highest THQ in adults and children due to consumption of Iranian tea was related to As (0.0635) and As (0.2964), respectively, and the highest THQ in adults and children due to consumption of imported tea was related to Cu (0.0266) and Cu (0.1242), respectively. TTHQ in adults and children's consumers due to PTEs ingestion via consumption of Iranian tea was 0.21 and 1.01, respectively, and TTHQ due to PTEs ingestion via consumption of imported tea was 0.22 and 0.94, respectively. Therefore, non-carcinogenic risk in children due to consumption of Iranian tea was higher than 1 value. The mean CR for content of inorganic As in adults and children via consumption of Iranian tea was 2.89E-5 and 1.35E-4, respectively, and the mean CR for inorganic As in adults and children via consumption of imported tea was 9.44E-6 and 4.42E-5 respectively. Hence adults and children, due to both Iranian and imported tea consumption, are at considerable carcinogenic risk. It is recommended to carry out continuous monitoring plans at country customs and reduce sources of tea contamination to PTEs in agricultural fields.

30



ARTICLE HISTORY


Received 12 July 2022

Accepted 7 August 2022

KEYWORDS

Tea; potentially toxic elements; meta-analysis; risk assessment; Iran

42 CONTACT Trias Mahmudiono  trias-m@fkm.unair.ac.id; Amin Mousavi Khaneghah  MOUSAVU@UNICAMP.BR; mousavi.amin@gmail.com

 Supplemental data for this article can be accessed online at <https://doi.org/10.1080/03067319.2022.2118596>

© 2022 Informa UK Limited, trading as Taylor & Francis Group

1. Introduction

The chemical environment of contamination [1–3] followed by food contamination [9–13], has become a main global health concern. Tea is one of the most consumed and popular beverages in Asian countries, especially Iran. The average global tea consumption is 6.3 billion kg per year, estimated at 1.5 kg per year per each Iranian [14], which can be attributed to the health benefits of tea in the prevention of some diseases, including headache, depression, cancer, Alzheimer's, Parkinson's, vascular diseases, and enhancement of immune defence [15–17].

The benefits of tea consumption can be associated with the presence of compounds such as flavonoids, flavonols, caffeine, minerals, vitamins, trace elements, theobromine, and antioxidants [18]. However, tea can become contaminated with potentially toxic element, including heavy metals, during the growing, harvesting, processing, drying, processing, and packaging periods. In addition, soil, air, and water pollution, as well as pesticides, can affect the level of tea contamination [15,19]. In other words, the type and amount of potentially toxic element, especially heavy metals, in tea are different under the influence of the above factors [18]. Due to the high consumption of tea, the presence of PTEs in this drink can endanger human health. The most important heavy metals identified in tea are Iron (Fe), Zinc (Zn), Copper (Cu), Chromium (Cr), Lead (Pb), Cadmium (Cd), Nickel (Ni), and Mercury (Hg) [20–23]. Some PTEs, such as Fe, Zn, and Cu, are essential trace elements for human health; however, if their concentration exceeds the maximum allowable limit, they can accumulate in living systems [24–27]. In contrast, Cr, Pb, As, Cd, Ni, and Hg are toxic metals [28]. For example, exposure to Cd can lead to kidney disorders, high blood pressure, infertility, lung, bone marrow, stomach, and prostate cancers [29,30]. High concentrations of Pb in the body can prevent the synthesis of haemoglobin, kidney and cardiovascular system failure, autism, damage to the central nervous system, and high blood pressure [31,32]. Ni is most common adverse effects on humans are allergic reactions, shortness of breath, chronic bronchitis, headache, Irridium, kidney and blood problems, and lung and sinus cancer [33]. Hg can damage the liver, brain, respiratory system, skin, eyes, and central nervous system [34].

Despite numerous studies on the concentration of PTEs in tea, no systematic review and meta-analysis regarding the concentration of the potentially toxic elements (PTEs) in black tea in Iran were published. Therefore, this study was designed to meta-analyse the concentration of PTEs in Iranian and imported tea and probabilistic health risks.

5

2. Materials and methods

2.1. Search strategy

According to the PRISMA protocol [35], a search on international databases (Scopus and PubMed) and national databases (SID, Irandoc, and Magiran) were conducted for the period 2005 to 1 June 2022 to retrieve citations that investigated heavy metals in tea consumed in Iran. The search was conducted according to the following terms: 'Trace element' OR 'Metal OR 'Heavy metal' OR 'Elements' OR 'Potential toxic element' OR 'Toxic element' AND 'Black tea' OR 'Tea' AND 'Iran'. Disagreement between two authors for select of papers was resolved with the final opinion of the senior author [36].

8

2.2. Inclusion and exclusion criteria and extraction of data

Our criteria for inclusion of paper were 1) English and Persian language paper, 2) Descriptive studies, and 3) investigated concentration of heavy metals in tea. The experimental studies, unclear data studies, review, clinical trial, and thesis were excluded. The data, sample size, province, type of tea (Iranian or imported), and concentration (Mean, standard deviation, lowest and highest) were extracted [37].

2.3. Meta-analysis

A meta-analysis of the concentration of heavy metals in tea was conducted based on average and standard error. The data heterogeneity was detected by the I^2 index [38,39]. Depending on the I^2 index, random-effects model (> 50%) or fixed-effects model (< 50%) used to meta-analysis. Stata 14.00 software was used for meta-analysis.

23

2.4. Health risk assessment

In the current study, the non-carcinogenic risk of As, Cd, Pb, Ni, Hg, Cu, and Fe and the carcinogenic risk of As in consumers due to consumption of Iranian and imported tea were estimated using the Monte Carlo simulation (MCS). The distribution type of concentration and ingestion rate was selected as Log-normal and normal for body weight [40–42]. All equations of non-carcinogenic and carcinogenic risks were presented in Table 1.

3. Results and discussion

3.1. The level of PTEs based on metals types

Twenty-eight papers with 73 records reports were included in the current meta-analysis Figure 1 and Table S1. As seen at Table 2, the rank order of PTEs based on concentration was Fe (90.211 mg/kg) > Cu (16.881 mg/kg) > Ni (7.223 mg/kg) > Pb (1.305 mg/kg) > Cd (0.239 mg/kg) > As (0.228 mg/kg) > Hg (0.034 mg/kg). Findings obtained were lower than the standard limits suggested by Iran Iranian standard organisation (Pb: 10 mg/kg, Cd: 3 mg/

Table 1. Equations and description in our study.

Number	Equation	Description	References
1	$EDI = \frac{C_m \times IR}{BW}$	EDI = estimated daily intake C _m = concentration of metal (µg/Kg) IR = ingestion rate (4.13 ± 0.56 g/n-d) BW = Body weight (Adults = 70 kg and children = 15 kg)	[43,44]
2	$THQ = \frac{EDI}{RfD \text{ or } TDI}$	THQ = target hazard quotient RfD = oral reference dose: As (inorganic), Cd, Zn, Cu, Fe, Ni, methyl Hg, Mn and Zn is 0.0003, 0.001, 0.3, 0.04, 0.7, 0.011, 0.0001, 0.14, and 0.30 mg/kg-d, respectively. TDI: Tolerable daily intake TDI for Pb is 0.0036 mg/kg-d	[45,46]
3	$TTHQ = \frac{EDI_1}{RfD_1} + \frac{EDI_2}{RfD_2} + \dots + \frac{EDI_n}{RfD_n}$	EDI _n is the estimated daily intake for nth toxicant RfD _n is oral reference dose for nth toxicant	[47–49]
4	CR = EDI × SF	Slope factor of inorganic As is 1.5 (mg/kg-d) ⁻¹	[50]




Figure 1. Selection process of papers based on PRISMA.

Table 2. Meta-analysis concentration of PTEs in national and imported black tea in Iran (mg/kg).

PTEs	Type of tea	N ³⁷ per study	ES	Lower	Upper	Weight (%)	Heterogeneity statistic	Degrees of freedom	p-value	I ²
As	National	16	0.323	0.292	0.355	65.30	5233	15	<0.001	99.70%
	Imported	5	0.106	0.058	0.155	22.98	3154	4	<0.001	99.90%
	NM ¹	3	0.053	0.028	0.079	11.72	140	2	<0.001	98.60%
	Overall	24	0.228	0.209	0.247	100.00	9067	23	<0.001	99.70%
Cd	National	33	0.271	0.231	0.311	71.06	19,203	32	<0.001	99.80%
	Imported	9	0.243	0.191	0.295	21.45	3597	8	<0.001	99.80%
	NM	3	0.042	0.036	0.048	7.49	6	2	0.045	67.80%
	Overall	45	0.239	0.212	0.265	100.00	24,831	44	<0.001	99.80%
Pb	National	40	1.303	1.171	1.435	69.96	300,000	39	<0.001	100.00%
	Imported	13	1.427	0.977	1.877	22.82	180,000	12	<0.001	100.00%
	NM	4	0.856	0.114	1.598	7.22	1161	3	<0.001	99.70%
	Overall	57	1.305	1.183	1.426	100.00	580,000	56	<0.001	100.00%
Ni	National	20	7.991	6.894	9.088	79.80	750,000	19	<0.001	100.00%
	Imported	3	4.428	3.573	5.284	12.11	51	2	<0.001	96.10%
	NM	2	3.823	3.49	4.157	8.09	4	1	0.053	73.20%
	Overall	25	7.223	6.247	8.199	100.00	750,000	24	<0.001	100.00%
Methyl Hg	National	3	0.066	0.025	0.107	39.16	1598	2	<0.001	99.90%
	Imported	2	0.025	0.000	0.074	30.34	1586	1	<0.001	99.90%
	NM	2	0.012	0.008	0.016	30.50	49	1	<0.001	98.00%
	Overall	7	0.034	0.026	0.042	100.00	7368	6	<0.001	99.90%
Cu	National	37	16.275	15.736	16.815	76.92	170,000	36	<0.001	100.00%
	Imported	11	18.037	14.467	21.607	21.00	5112	10	<0.001	99.80%
	NM	1	19.46	18.435	20.485	2.08	0	0	.	%
	Overall	49	16.881	16.352	17.411	100.00	210,000	48	<0.001	100.00%
Fe	National	4	87.691	14.668	160.714	50.13	18,924	3	<0.001	100.00%
	Imported	3	95.898	14.906	176.89	37.32	3539	2	<0.001	99.90%
	NM	1	83.42	77.129	89.711	12.55	0	0	.	%
	Overall	8	90.211	38.764	141.658	100.00	36,470	7	<0.001	100.00%

kg, Cu: 50 mg/kg, As: 0.15 mg/kg, Hg: 0.02 mg/kg) [51]. Consistent with the findings, the level of PTEs in the studied black teas was very diverse between countries and compared to the standard level, which can be related to several factors such as the ability of the bioaccumulation and bioavailability of metals in the tea plant, nature of the tea plant species, biological form of cultivation, situation during plant growth of tea (humidity of soil, PH and water content), the composition of the irrigation water and soil, storage condition and processing technologies [52]. Each of these parameters can play an effective role in the content of metals in black tea consumed in Iran. Chemical properties of soil compounds, including pH level [53], level of carbon [54], amount of nitrogen [55], potassium sulphur [56], and type and amount of phosphate fertilisers, have an influence on metals level uptake via tea plants [57]. In a study conducted by Tuysuz and Yaylali-Abanuz, they showed a significant negative correlation between the pH of the soil and the uptake of various metals by tea plants. An increase in pH causes a reduction in the solubility of the Pb metal in soil [58]. It is observed that the higher level of Fe, Cu, and Zn elements, as compared with PTEs such as Pb, Cd, As, and Hg in black teas is related to their higher transfer rate [29,59,60]. PTEs such as Cu, Fe, and Cd have higher mobility properties than other PTEs such as Pb and Cd; hence, Cu, Fe, and Cd can quickly accumulate at the tea plant's roots [61,62]. Storage conditions and processing technologies are the significant factors in the concentration of PTEs. For example, the washing stage and pH environment in processing performed on tea leaves in industrial factories are important factors that affect the level of PTEs [63]. The content and age of tea plant leaves used in the production process, maturing, storage, and packing stage besides the fermentation methods have an effective role in contamination level in black tea [64]. The soil irrigated with wastewater had higher metals concentration than the soil irrigated with groundwater [65]. While the soil irrigated with wastewater contain a higher level of PTEs than the clean water and groundwater [66,67]. The different wastewater of domestic or industrial due to composition or, electrical conductivity can directly effect on bioavailability and then bioaccumulation of PTEs in crops [68]. Using various phosphate fertilisers and pesticides (fungicides) may lead to contamination of tea plants with PTEs [16]. Using ammonium sulphate fertiliser in agriculture can raise the content of the soil and increase the adsorption and mobility of the PTEs in the soil. In this regard, the soil fertilisation via ammonium sulphate is the important factor affecting the Cd absorption in the leaves of tea plants [58].

3.2. The concentration of PTEs based on origin of tea

The content of PTEs in the Iranian and imported teas was different (Table 2). Our findings indicated that the pooled concentration of Methyl Hg (0.066 mg/kg), As (0.323 mg/kg), Ni (7.991 mg/kg), Cu (18.037 mg/kg), Fe (95.898 mg/kg) in Iranian black tea was higher than imported tea (0.025 mg/kg), (0.106 mg/kg), (4.428 mg/kg), (16.275 mg/kg) and (87.691 mg/kg), respectively. While the pooled concentration of Cd and Pb in Iranian black tea (0.271 mg/kg) and (1.303 mg/kg) was almost equal to imported tea (0.243 mg/kg) and (1.427 mg/kg) respectively.

The mean level of As and Cd in Iranian black tea were 0.09 and 2.5 mg/kg, respectively [15]. In another investigation, the mean level of Cd and As in imported black tea were defined as 0.72 and 0.05 mg/kg, respectively [69]. Similarly, the mean level of Cd, Pb, Cu, and Fe in Iranian black tea were reported as 0.63, 1.72, 8.85, and 41.34 mg/kg, respectively [60].



The main sources of PTEs contamination in Iranian and imported teas are strongly dependent on geographic characteristics, including the weather conditions, annual rainfall, air pollution and as well as, and agricultural activities such as using of fertilisers and pesticides, different industrial and urban activities, proximity to roads, the mines, factories, and highways [70]. In this context, the low level of PTEs in some counties can be attributable to the agricultural soil's physicochemical properties and the region's weather conditions [71]. For example, various PTEs concentration teas plant mainly due to different climatic conditions in Japan (moister) compared with China and other Asian countries was noted [72]. The existence of mining and industry activities in the region close to planting plants, besides local dispersion in soil profiles can rise concentration of PTEs in tea plants. Cu contamination was higher than other PTEs in tea plants cultivated around a copper mine [72]. In another study, Argentina's rural or suburban regions have an extensive groundwater network polluted with unique As metals. Therefore, different plants that grow in these areas are more contaminated with As than in other areas [73]. A study conducted in Mengel, Konrad stated that Cd availability in soils was influenced completely by adding rock phosphate because rock phosphate is useful annually in several soils used for tea plants [74]. Contamination of pb in the soils of parts near the roads was noted because of emissio⁴⁷ of tetraethyl lead from exhaust vehicles was higher than in other areas [75]. Amouei et al. stated high level of Pb in Iranian black tea can be associated with the high traffic in this region, the alkalinity of the soil, the rise of cation exchange capacity, and the high volume of organic carbon and clay particles percentages of the soil [76].

4. Risk assessment

Non-carcinogenic risk

When, THQ and/or TTHQ > 1, exposed population are considerable non-carcinogenic risk [77,78]. The rank order of THQ of PTEs in Iranian tea for adults was As (0.0635) > Ni (0.0429) > Methyl Hg (0.0389) > Cu (0.0240) > Pb (0.0214) > Cd (0.0160) > Fe (0.0074) and for children was As (0.2964) > Ni (0.2000) > Methyl Hg (0.1817) > Cu (0.1120) > Pb (0.0997) > Cd (0.0746) > Fe (0.0345) (Figure S1). The rank order of THQ of PTEs in imported tea for adults was Cu (0.0266) > Ni (0.0238) > Pb (0.0234) > As (0.0208) > Methyl Hg (0.0148) > Cd (0.0143) > Fe (0.0081) and for children was Cu (0.1242) > Ni (0.1108) > Pb (0.1091) > As (0.0973) > Methyl Hg (0.0688) > Cd (0.0669) > Fe (0.0377) (Figure S2). Differences in concentrations of PTEs and their RfD⁹ were the reasons for the variation among THQ of Iranian and imported tea [79–81]. TTHQ in adults and children's consumers due to ingestion of PTEs in Iranian tea was 0⁶ and 1.01, respectively, and also TTHQ because of ingestion of PTEs in imported tea was 0.22 and 0.94, respectively (Figure 2). Hence, TTHQ in children due to consumption of Iranian tea is higher compared with 1 value.

5. Carcinogenic risk

The mean CR in children and adults because of ingestion of Iranian tea level of inorganic As was 2.89E-5 and 1.35E-4, respectively (Figure S3). The mean CR in adults and children due to inorganic As ingestion in imported tea was 9.44E-6 and 4.42E-5 respectively (Figure S4). If, CR > 1.00E-06, exposed population are considerable carcinogenic risk [82].




Figure 2. TTHQ in adults and children due to consumption of national and imported tea.

Therefore, CR for adults and children due to both Iranian and imported tea are at considerable carcinogenic risk.

6. Conclusion

This study was designed to meta-analysis the concentration of PTEs in black tea in Iran and probabilistic risk assessment. Pooled Fe, Cu, and Ni concentrations in consumed tea were higher than in other PTEs. The concentration of Methyl Hg, As, Ni, Cu, and Fe, in Iranian black tea, was higher than in imported tea. Consumption of Iranian tea resulted in non-carcinogenic risk to children as marginally. Adults and children were posed to non-acceptable carcinogenic risk due to consumption of Iranian and imported tea consumption. It is recommended to carry out continuous monitoring plans at country customs and agricultural fields to reduce the concentration of PTEs in black tea in Iran.

Note

1. Not mentioned

Disclosure statement

No potential conflict of interest was reported by the author(s).

ORCID

Yadolah Fakhri <http://orcid.org/0000-0002-3579-7641>

Amin Mousavi Khaneghah <http://orcid.org/0000-0001-5769-0004>

7

Consent to Participate

The authors declare their Consent to Participate in this article.

Consent to Publish

The authors declare their Consent to Publish this article.

Authors contributions

38

Search in databases was conducted by **Somayeh Hoseinvandtabar** and **Hasti Daraei**; Screening of obtained articles based on title, abstract and full text by **Somayeh Hoseinvandtabar**; Data extraction by **Somayeh Hoseinvandtabar**; Meta-analysis of data and risk assessment by **Yadolah Fakhri**; write and edit the manuscript by **Yadolah Fakhri**, **Trias Mahmudiono**, **Hasti Daraei**, **Fereshteh Mehri**, **Amin Mousavi Khaneghah**.

References

- [1] C. Shi et al. *Janus Particles with pH Switchable Properties for high-efficiency Adsorption of PPCPs in Water*, (Amsterdam: Solid State Sciences, 2021), Vol. 119, p. 106702
- [2] W. Liu et al., *Angewandte Chemie* **120**(30), 5701 (2008). doi:10.1002/ange.200800172.
- [3] W. Li et al., *Ecol. Indic.* **130**, 108031 (2021). doi:10.1016/j.ecolind.2021.108031.
- [4] X. Lin et al., *Ecol. Indic.* **126**, 107639 (2021). doi:10.1016/j.ecolind.2021.107639.
- [5] Q. Guan et al., *J. Environ. Manage.* **293**, 112961 (2021). doi:10.1016/j.jenvman.2021.112961.
- [6] D. Ge et al., *Sci. Total Environ.* **679**, 298 (2019). doi:10.1016/j.scitotenv.2019.05.060.
- [7] W. Liu et al., *Environ. Sci. Technol.* **54**(19), 11971 (2020). doi:10.1021/acs.est.0c01855.
- [8] N. Su et al., *Catena* **213**, 106163 (2022). doi:10.1016/j.catena.2022.106163.
- [9] T. Li et al., *Biosensors* **12**(5), 314 (2022). doi:10.3390/bios12050314.
- [10] G. Liu et al., *Sci. Total Environ.* **825**, 154058 (2022). doi:10.1016/j.scitotenv.2022.154058.
- [11] A.N. Munyasya et al., *Agric. Water Manage.* **269**, 107672 (2022). doi:10.1016/j.agwat.2022.107672.
- [12] Y. Wang et al., *J. Environ. Chem. Eng.* **10**(1), 107091 (2022). doi:10.1016/j.jece.2021.107091.
- [13] Y. Sun et al., *Curr. Opin. Food Sci.* **44**, 100813 (2022). doi:10.1016/j.cofs.2022.100813.
- [14] Statista. Volume of Tea Consumption Worldwide from 2012 to 2025. <https://www.statista.com/statistics/940102/global-tea-consumption/#:~:text=In%202020%2C%20global%20consumption%20of,7.4%20billion%20kilograms%20by%202025.&text=Tea%20is%20the%20second%20most,%2C%20yellow%2C%20oolong%20and%20black.> 2020.
- [15] G. Karimi et al. *Pharmacology*. **3**, 164 (2008).
- [16] F. Pourramezani et al., *Food Sci. Nutr.* **7**(12), 4021 (2019). doi:10.1002/fsn3.1267.
- [17] F. Vieux et al., *Nutrients* **11**(11), 2635 (2019). doi:10.3390/nu11112635.
- [18] E. Falahi and R. Hedayati, *Food Addit. Contam: Part B* **6**, 123 (2013). doi:10.1080/19393210.2013.764550.
- [19] J. Zhang et al., *Int. J. Environ. Res. Public Health* **15**(1), 133 (2018). doi:10.3390/ijerph15010133.
- [20] N. Yousefi, A. Jahangard and M.H. Mahmoudian, *Arch. Hygiene Sci.* **6**, 128 (2017). doi:10.29252/ArchHygSci.6.2.128.
- [21] Y. Fakhri et al., *Chemosphere* **279**, 130431 (2021). doi:10.1016/j.chemosphere.2021.130431.
- [22] A. Nematollahi et al., *Environ. Sci. Pollution Res.* **28**(39), 55186 (2021). doi:10.1007/s11356-021-14879-2.
- [23] M. Borzoei et al., *Biol. Trace Elem. Res.* **192**(2), 319 (2019). doi:10.1007/s12011-019-01662-1.
- [24] A. Bounar, K. Boukaka and E. Leghouchi, *Qual. Assur. Safety Crops & Foods* **12**, 76 (2020). doi:10.15586/QAS2019.639.

- [25] L. Gao et al., *Qual. Assur. Safety Crops & Foods* **14**(1), 82 (2022). doi:10.15586/qas.v14i1.1034.
- [26] A. Heshmati et al., *Qual. Assur. Safety Crops & Foods* **12**(3), 140 (2020). doi:10.15586/qas.v12i3.761.
- [27] M. Rezaei et al., *Qual. Assur. Safety Crops & Foods* **12**(3), 111 (2020). doi:10.15586/qas.v12i3.777.
- [28] A. RoyChowdhury, R. Datta and D. Sarkar, *Heavy Metal Pollution and Remediation, in Green Chemistry*, (Amsterdam: Elsevier, 2018), p. 359.
- [29] M.A. Zazouli et al., *Asian J. Chem.* **22** (2), 1387 (2010).
- [30] N.J. Vickers, *Curr. Biology* **27**, R713–R715 (2017). doi:10.1016/j.cub.2017.05.064.
- [31] A.T. Jan et al., *Int. J. Mol. Sci.* **16**(12), 29592 (2015). doi:10.3390/ijms161226183.
- [32] G. Barone, R. Giacomini-Stuffler and M.M. Storelli, *Food Chem. Toxicol.* **87**, 113 (2016). doi:10.1016/j.fct.2015.12.008.
- [33] A. Mudipalli and J.T. Zelikoff, *Essential and Non-Essential Metals* (New York City: Springer, 2017), p. 125–135.
- [34] E. Ha et al., *Environ. Res.* **152**, 419 (2017). doi:10.1016/j.envres.2016.06.042.
- [35] D. Moher et al., *Syst. Rev.* **4**(1), 1 (2015). doi:10.1186/2046-4053-4-1.
- [36] J.P. Aranega and C.A. Oliveira, *Qual. Assur. Safety Crops & Foods* **14**, 135 (2022). doi:10.15586/qas.v14i3.1079.
- [37] M. Mokhtarian et al., *Qual. Assur. Safety Crops & Foods* **12**(4), 15 (2020). doi:10.15586/qas.v12i4.784.
- [38] A. Atamaleki et al., *Int. J. Environ. Health Res.* **32**(7), 1 (2021).
- [39] J.P.T. Higgins et al. *Cochrane Handbook for Systematic Reviews of Interventions*, (Hoboken, New Jersey: Wiley, 2019), p. 125–155.
- [40] A. Heshmati, M. Khorshidi and A.M. Khaneghah, *Italian J. Food Sci.* **33**, 92 (2021). doi:10.15586/ijfs.v33iSP1.2065.
- [41] K. Jafari et al., *Italian J. Food Sci.* **33**(SP1), 103 (2021). doi:10.15586/ijfs.v33iSP1.2054.
- [42] M. Dadar et al., *Qual. Assur. Safety Crops & Foods* **14**(3), 55 (2022). doi:10.15586/qas.v14i3.1088.
- [43] EPA. *Exposure Factors Handbook - Chapter 15: Human Milk Intake.* <https://ofmpub.epa.gov/eims/eimscmm/getfile>. Vol. 5. 2011. 7.
- [44] Statista. *Annual per Capita Tea Consumption Worldwide as of 2016, by Leading Countries*(in Pounds). Statista. (23 January, 2020. Retrieved 10 February 2020. <https://www.statista.com/statistics/507950/global-per-capita-tea-consumption-by-country/>.
- [45] D.G. Barnes et al., *Regul. Toxicol. Pharmacol.* **8**(4), 471 (1988). doi:10.1016/0273-2300(88)90047-5.
- [46] EPA, *Risk-based Concentration Table*, (J United States Environmental Protection Agency, Washington DC , Philadelphia PA, 2000). p. 45–49.
- [47] L. Theodore and R.R. Dupont, *Environmental Health and Hazard Risk Assessment: Principles and Calculations* (Milton Park, Abingdon-on-Thames, Oxfordshire United Kingdom: CRC Press, 2017).
- [48] Y. Fakhri et al., *Toxin. Rev.* **24**, 1 (2019).
- [49] E.P.A. U, *Risk-based Concentration Table* (United States Environmental Protection Agency, Washington DC, Philadelphia PA, 2000).
- [50] EPA, *Exposure Factors Handbook*, (USA: EPA), 2011 (2011), p. 25–29. EPA/600/R-09
- [51] N.F. Soliman, *Health Econo. Outcome Res: Open Access* **2**, 1 (2016). doi:10.4172/2471-268X.1000109.
- [52] A. Heshmati et al., *J. Food Prot.* **83**(1), 101 (2020). doi:10.4315/0362-028X.JFP-19-312.
- [53] M. Li et al., *Catena* **101**, 11 (2013). doi:10.1016/j.catena.2012.09.012.
- [54] W. Lei et al., *Pedosphere* **23**(2), 177 (2013). doi:10.1016/S1002-0160(13)60005-3.
- [55] K. Oh, T. Kato and H. Xu, *Pedosphere* **18**, 222 (2008). doi:10.1016/S1002-0160(08)60010-7.
- [56] D. Kamau, J. Spiertz and O. Oenema, *Plant Soil* **307**, 29 (2008). doi:10.1007/s11104-008-9576-6.
- [57] S. Ananthacumaraswamy, L. Hettiarachchi and S. Dissanayake, *Commun. Soil Sci. Plant Anal.* **34**, 1481 (2003). doi:10.1081/CSS-120021291.
- [58] G. Yaylalı-Abanuz and N. Tüysüz, *Environ. Earth Sci.* **59**, 131 (2009). doi:10.1007/s12665-009-0011-y.

- [59] C.Y. Peng et al., *J. Food Sci.* **83**(4), 1165 (2018). doi:10.1111/1750-3841.14093.
- [60] M. Nejatolahi, S. Mortazavi, A. Ildoromi and Z. Levels of Cu, *J. Food Meas. Charact.* **8**, 277 (2014). doi:10.1007/s11694-014-9186-3.
- [61] E.T. Ghane et al., *Biol. Trace Elem. Res.* **200**(1), 437 (2022). doi:10.1007/s12011-021-02645-x.
- [62] S. Khazaei et al., *Int J Environ Anal Chem.*, 1 (2021). doi:10.1080/03067319.2021.1873309.
- [63] S. Hosseni, A. Shakerian and A. Moghimi, *J Food Biosci. Technol.* **3**, 67 (2013).
- [64] H. Matsuura et al., *Anal. Sci.* **17**(3), 391 (2001). doi:10.2116/analsci.17.391.
- [65] A. Al-Busaidi, P. Cookson and T. Yamamoto, *Soil Res.* **43**, 541 (2005). doi:10.1071/SR04102.
- [66] Z.-J. Xue et al., *Environ. Monit. Assess.* **184**(6), 3503 (2012). doi:10.1007/s10661-011-2204-6.
- [67] N. Gupta, D. Khan and S. Santra, *Environ. Monit. Assess.* **184**, 6673 (2012). doi:10.1007/s10661-011-2450-7.
- [68] K. Khanum et al., *Carpathian J. Earth Environ. Sci.* **12** (2), 403 (2017).
- [69] A. Rahmani et al., *Int J Environ Anal Chem.* **31**(1), (2022).
- [70] Y.-J. Cui et al., *Environ. Int.* **30**(6), 785 (2004). doi:10.1016/j.envint.2004.01.003.
- [71] A. Rahimi, E. Talebi Ghane and F. Mehri, *Int J Environ Anal Chem.* 1 (2021). doi:10.1080/03067319.2021.1974428
- [72] N. Tüysüz and M. Akçay, *Yıldönümü Yerbilimleri ve Madencilik Kongresi Bildiriler Kitabı* **2**, 625 (2000).
- [73] A.L. Pérez-Carrera, F.E. Arellano and A. Fernández-Cirelli, *Dairy Sci. Technol.* **96**, 591 (2016). doi:10.1007/s13594-016-0290-5.
- [74] K. Mengel and E.A. Kirkby, *Principles of Plant Nutrition* (New York City: Springer Science & Business Media, 2012).
- [75] M.E. Finster, K.A. Gray and H.J. Binns, *Sci. Total Environ.* **320**, 245 (2004). doi:10.1016/j.scitotenv.2003.08.009.
- [76] A.I. Amouei et al., *Environ. Justice* **5**(3), 153 (2012). doi:10.1089/env.2011.0038.
- [77] Y. Fakhri et al., *Biol. Trace Elem. Res.* **199**(8), 3089 (2021). doi:10.1007/s12011-020-02419-x.
- [78] V.N. Thai et al., *Biol. Trace Elem. Res.* **199**(9), 3497 (2021). doi:10.1007/s12011-020-02476-2.
- [79] G.R. Mostafaei et al., *Int J Environ Anal Chem.* **102**(5), 1192 (2022). doi:10.1080/03067319.2020.1734195.
- [80] M. Voltz et al., *Environmental Science and Pollution Research International* **29**(1), 1 (2022). doi:10.1007/s11356-021-17416-3.
- [81] P. Sadighara et al., *Int J Environ Anal Chem.*, 1 (2022). doi:10.1080/03067319.2022.2045589.
- [82] USEPA. United State Environmental Protection Agency. Quantitative Risk Assessment Calculations. <https://www.epa.gov/sites/production/files/2015-05/documents/13.pdf> 2015: p. 7.

The concentration of the potentially toxic element (PTEs) in black tea (*Camellia sinensis*) consumed in Iran: a systematic review, meta-analysis, and probabilistic risk assessment study

ORIGINALITY REPORT

19%

SIMILARITY INDEX

12%

INTERNET SOURCES

17%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1	repository.ju.edu.et Internet Source	2%
2	pericles.pericles-prod.literatumonline.com Internet Source	1%
3	jder.ssu.ac.ir Internet Source	1%
4	Mehrnoosh Abtahi, Yadolah Fakhri, Gea Oliveri Conti, Margherita Ferrante et al. "The Concentration of BTEX in the Air of Tehran: A Systematic Review-Meta Analysis and Risk Assessment", <i>International Journal of Environmental Research and Public Health</i> , 2018 Publication	1%
5	www.mdpi.com Internet Source	1%
6	Meghdad Pirsaeheb, Yadolah Fakhri, Mohammad Karami, Rokhsareh Akbarzadeh et al. "Measurement of permethrin,	1%

deltamethrin and malathion pesticide residues in the wheat flour and breads and probabilistic health risk assessment: a case study in Kermanshah, Iran", International Journal of Environmental Analytical Chemistry, 2019

Publication

7 Shima Jafarzadeh, Milad Hadidi, Mehrdad Forough, Abdorreza Mohammadi Nafchi, Amin Mousavi Khaneghah. "The control of fungi and mycotoxins by food active packaging: a review", Critical Reviews in Food Science and Nutrition, 2022

Publication

8 Yadolah Fakhri, Jamal Rahmani, Carlos Augusto Fernandes Oliveira, Larissa Tuanny Franco et al. "Aflatoxin M1 in human breast milk: A global systematic review, meta-analysis, and risk assessment study (Monte Carlo simulation)", Trends in Food Science & Technology, 2019

Publication

9 www.qascf.com

Internet Source

10 Hideaki Endo, Maki Igarashi, Akira Banba, Hitoshi Ohnuki, Hideki Ushio, Tetsuhito Hayashi, Huifeng Ren, Goro Yoshizaki. "Electrode-based immunologic assay system to monitor oocyte maturation-inducing hormone in fish", International

Journal of Environmental Analytical Chemistry, 2011

Publication

11 Ayat Rahmani, Safa Kalteh, Milad Derakhshan Jazari, Mansour Baziar, Mostafa Karimaei. "Trace elements in imported black tea products in Iran: quantification and probabilistic health risk assessment", International Journal of Environmental Analytical Chemistry, 2022
Publication 1 %

12 all-imm.com
Internet Source 1 %

13 ir.unimas.my
Internet Source 1 %

14 researchsystem.canberra.edu.au
Internet Source 1 %

15 Hamid Atabati, Hamid Kassiri, Ehsan Shamloo, Mitra Akbari et al. "The association between the lack of safe drinking water and sanitation facilities with intestinal Entamoeba spp infection risk: A systematic review and meta-analysis", PLOS ONE, 2020
Publication <1 %

16 uis.brage.unit.no
Internet Source <1 %

17 www.scienceinschool.org
Internet Source <1 %

18	Katia Lacasse. "Environmental considerations for textile processes and chemicals", Textile Chemicals, 2004 Publication	<1 %
19	iwaponline.com Internet Source	<1 %
20	iehpc.gdut.edu.cn Internet Source	<1 %
21	www.jchr.org Internet Source	<1 %
22	9dok.org Internet Source	<1 %
23	A. Domfeh, A. K. Anim, A. Asamoah. "Trace metals contamination in varieties of cereal-based pediatric foods sold in parts of Accra, Ghana", Research Square Platform LLC, 2023 Publication	<1 %
24	e-journal.unair.ac.id Internet Source	<1 %
25	researchers.usask.ca Internet Source	<1 %
26	Gamal A.E. Mostafa, Emad Alsarhani, Rashad AlSalahi. "Assessment of heavy metals in infused tea marketed in Riyadh, Saudi Arabia, using inductively coupled plasma-mass spectrometry: human health	<1 %

27

Lei Xu, Xiangyu Xing, Jianbiao Peng, Mingfei Ji. "Effects of In Situ Remediation on Copper Distribution and Soil Aggregate Adsorption-Desorption Characteristics in Smelter-Impacted Soil", Frontiers in Environmental Science, 2022

Publication

<1 %

28

Robel Hussen Kabthymer, Girum Gebremeskel Kanno, Mekonen Birhane Aregu, Susana Paixão, Tefera Belachew. "Prevalence and concentration of Aflatoxin M1 in human breast milk in sub-Saharan Africa: a systematic review and meta-analysis, and cancer risk assessment", International Journal of Environmental Health Research, 2022

Publication

<1 %

29

Zeinab Gholami, Mehrnoosh Abtahi, Mehdi Golbini, Iman Parseh et al. "The concentration and probabilistic health risk assessment of nitrate in Iranian drinking water: a case study of Ilam city", Toxin Reviews, 2019

Publication

<1 %

30

aura.abdn.ac.uk

Internet Source

<1 %

31

Internet Source

<1 %

32

Amin Mohammadpour, Enayat Berizi, Amin Allah Zarei, Mohadeseh Motamed-Jahromi et al. "The concentration of cadmium, lead, and nitrate in tomato and onion from Fars province, Iran: a health risk assessment study", International Journal of Environmental Analytical Chemistry, 2022

Publication

<1 %

33

Chammi P. Attanayake, Ganga M. Hettiarachchi, Sabine Martin, Gary M. Pierzynski. "Potential Bioavailability of Lead, Arsenic, and Polycyclic Aromatic Hydrocarbons in Compost-Amended Urban Soils", Journal of Environmental Quality, 2015

Publication

<1 %

34

Hamed Ghafari Farsani, Mohammad Hasan Gerami, Mehdi Naderi Farsani, Ghasem Rashidiyan et al. " Effect of different levels of essential oils () in diet on improvement growth, blood biochemical and immunity of Angelfish (Schultze, 1823) ", Natural Product Research, 2018

Publication

<1 %

35

Jian Zhang, Ruidong Yang, Yuncong C. Li, Yishu Peng, Xuefeng Wen, Xinran Ni. "Distribution, accumulation, and potential risks of heavy metals in soil and tea leaves

<1 %

from geologically different plantations",
Ecotoxicology and Environmental Safety,
2020

Publication

36

Kafeel Ahmad, Asma Ashfaq, Zafar Iqbal Khan, Humayun Bashir, Muhammad Sohail, Naunain Mehmood, Yunus Dogan. "Metal accumulation in Raphanus sativus and Brassica rapa: an assessment of potential health risk for inhabitants in Punjab, Pakistan", Environmental Science and Pollution Research, 2018

Publication

37

Maryam Dadar, Youcef Shahali, Sharaf S. Omar, Yadolah Fakhri, Jacques Godfroid, Amin Mousavi Khaneghah. "The prevalence of camel brucellosis and associated risk factors", Quality Assurance and Safety of Crops & Foods, 2022

Publication

38

Shabir Ahmad Mir, Manzoor Ahamd Shah, Mohammad Maqbool Mir, Tahira Sidiq et al. "Recent developments for controlling microbial contamination of nuts", Critical Reviews in Food Science and Nutrition, 2022

Publication

39

journal.fcisk.ru

Internet Source

40

journals.iau.ir

Internet Source

<1 %

<1 %

<1 %

<1 %

<1 %

41	trjfas.org Internet Source	<1 %
42	www.frontiersin.org Internet Source	<1 %
43	www.science.gov Internet Source	<1 %
44	"Development of recombinant vaccines to prevent the disease caused by the respiratory viruses SARS-CoV-2, hMPV, and hRSV", Pontificia Universidad Catolica de Chile, 2022 Publication	<1 %
45	Abdel-Rahim H. A. Hassan, Mohamed M. A. Zeinhom, Mariam A. Abdel-Wahab, Mona H. Tolba. "Heavy Metal Dietary Intake and Potential Health Risks for University Hostel Students", Biological Trace Element Research, 2015 Publication	<1 %
46	I. K. Mitsios, E. E. Golia, C. D. Tsadilas. "Heavy Metal Concentrations in Soils and Irrigation Waters in Thessaly Region, Central Greece", Communications in Soil Science and Plant Analysis, 2005 Publication	<1 %
47	Zahra Sarlak, Milad Rouhi, Adel Mirza Alizadeh, Ehsan Sadeghi, Hedayat Hosseini, Amin Mousavi Khaneghah. "Pb exposure from plant foods in Iran: a review",	<1 %

International Journal of Environmental Analytical Chemistry, 2021

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On