

ISSN 2455-3845
P-ISSN 2455-3845

AJPCR

An Open Access Peer Reviewed Journal

Asian Journal of Pharmaceutical & Clinical Research

Volume 10(1) 2024 | ISSN: 2455-3845
www.ajpcr.in



SMARTPHONE ADDICTION AND ITS RELATIONSHIP WITH PSYCHOLOGICAL HEALTH AMONG STUDENTS OF A MEDICAL SCHOOL IN EAST COAST MALAYSIA
AZWANIS ABDUL HADI, HAWARI MUSYIR MOHD NAWAWI, NURAFIFAH SHAMSURI, NURUL NAJIHAH RAHIM, HAFIZAH PASI 257-260

[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [10.22159/ajpcr.2019.v12i18.34396](#)

EPIDEMIOLOGIC ASPECTS OF TRAUMA IN AL-JOUF REGION, SAUDI ARABIA: A RETROSPECTIVE STUDY
DOAA M ABDEL-SALAM, RAWAN R ALRUWAILI, FARAH S ALHABLANI, NOUR H ALFAHEL, AMAL F ALMUNAH, ANWAR A ALBLAWI 261-264

[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [10.22159/ajpcr.2019.v12i18.33792](#)

THE ROLE OF SELENIUM MICRONUTRIENTS AS ANTIOXIDANTS IN EXPOSURE TO E-CIGARETTE SMOKE
RIVAN VIRLANDO SURYADINATA, MERRYANA ADRIANI, [SANTI MARTINI](#), SRI SUMARMI, BAMBANG WIRJATMADI 265-268

[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [10.22159/ajpcr.2019.v12i18.34454](#)

EVALUATION OF NUCLEAR CHANGES IN THE BUCCAL EPITHELIAL CELLS OF TOBACCO USERS IN NNEWI, SOUTH EAST NIGERIA
SAMUEL IFEDIORANMA OGENYI, ANTHONY AJULUCHUKWUNGOKERE, JONATHAN MADUKWE 269-273

[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [10.22159/ajpcr.2019.v12i18.34551](#)

EFFECTS EFFECTS OF PUMPKIN (CUCURBITA MOSCHATA DURCH) SEED POWDER ON BLOOD GLUCOSE LEVEL IN STREPTOZOTOCIN-INDUCED MICE
NOVARIANTI MARBUN, RUTH MAYANA RUMANTI 274-276

[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [10.22159/ajpcr.2019.v12i18.33730](#)



[HOME](#) / Editorial Board

Editorial Board

Editorial Board

AJPCR is committed to have dynamic and potential advisory-editorial board. Those established in the field can directly send their resume. New people are first needed to serve as referee before being considered member of advisory-editorial board. Email your resume to editor@ajpcr.com

Editor-in-Chief

- **Dr. Anurekha Jain**
Principal, Technocrats Institute of Technology and sciences-pharmacy, Bhopal
Email: anurekhajain1@gmail.com, editor@ajpcr.com

Associate Editor

- **Dr. Nuray Ari**
Department of Pharmacology, Faculty of Pharmacy, Ankara University, 06100 Ankara, Turkey. Email: ari@ankara.edu.tr
- **Dr. Idress Hamad Attitalla**
Department of Microbiology (Head) Faculty of Medical Technology (Dean) Box 919, Al-Bayda, Libya. Email: idress.hamad@omu.edu.ly

Assistant Editor

- **Dr. Omotoso Abayomi Ebenezer**
Department of Pharmaceutical & Medicinal Chemistry. Faculty of Pharmaceutical Sciences, University of Port Harcourt, Nigeria.
- **Dr. Vimal Kumar Jain**
Principal, ITM School of Pharmacy, ITM Universe, Vadodara & Associate Dean, Pharmacy, GTU, Ahmedabad, India
- **Dr. Adrianna Mostowska**
Department of Biochemistry and Molecular Biology Poznan University of Medical Sciences Poznan, Poland.

Editorial Board Members

- **Dr. Shivayogappa Teli**
Professor and Head, Department of Physiology Sri Manakula Vinayagar Medical College and Hospital Madagadipet, Puducherry 605107, India

- **Dr. Rupesh Kumar Gautam**
Associate Prof., ADINA Institute of Pharmaceutical Sciences, Sagar, MP, India
- **Dr. Tushar Treembak Shelke**
Head of Department of Pharmacology and Research Scholar, In Jspms Charak College of Pharmacy & Research, Pune, India
- **Dr. Subas Chandra Dinda**
Professor-cum-Director: School of Pharmaceutical Education & Research (SPER), Berhampur University, Berhampur, Orissa, India
- **Dr. Jagdale Swati Changdeo**
Professor and Head, Department of Pharmaceutics, MAEER's Maharashtra Institute of Pharmacy, S.No.124, MIT Campus, Kothrud, Pune, India
- **Dr. Biplab Kumar Dey**
Principal, Department of Pharmacy, Assam down town University, Sankar Madhab Path, Panikhaiti 781026, Guwahati, Assam, India
- **Dr. Durgesh Ranjan**
Pharmacy Global College of Pharmaceutical Technology Palpara, Krishnagar, Nadia West Bengal, India

Editorial office

Asian Journal of Pharmaceutical and Clinical Research
B-11, In front of Bima Hospital, Nai Abadi, Mandsaur 458001, MP, India
E-mail:editor@ajpcr.com



Online ISSN: 2455-3891

Print ISSN: 0974-2441

ICV 2021: 100.00

E-ISSN: 2455-3891
P-ISSN: 0974-2441

AJPCR

An Open Access Peer Reviewed Journal

Asian Journal of Pharmaceutical & Clinical Research

Citation Abbreviation: Asian J. Pharm. Clin. Res.
www.innovareacademics.in



Asian Journal of Pharmaceutical and Clinical Research

ISSN- 0974-2441

Vol 12, Issue 8, 2019

Editorial Board

Editor-in-Chief

Dr. Anurekha Jain

Dept. of Pharmaceutical Sciences, Jyoti Mahila Vidyapeeth University, Jaipur, Rajasthan Email: editor@ajpcr.com

Associate Editor

Dr. Nuray Ari

Prof., Department of Pharmacology, Faculty of Pharmacy, Ankara University, 06100 Ankara, Turkey

Dr. Neeraj Upmanyu

Peoples Institute of Pharmacy & Research Center, Bhopal, MP, India

Assistant Editor

Dr. Omotoso Abayomi Ebenezer

Prof., Department of Pharmaceutical & Medicinal Chemistry, Faculty of Pharmaceutical Sciences, University of Port Harcourt, Nigeria.

Dr. Vimal Kumar Jain

Institute of Pharmacy, Nirmal University, Ahemdabad, Gujarat, India

AJPCR Scope

AJPCR (Asian J Pharm Clin Res) is peer reviewed, Bi-monthly (Onward August 2014) open access Journal. This journal publishes original research in the field of Pharmaceutical and Clinical Sciences. The Journal has been designed to cover all the fields of research, which has any correlation and impact on Pharmaceutical Science. It aims to publish all the original research in field of science so a correlation can be made between these researches. Knowledge gained by such researches can be exposed to all and it can be brought in real utilization as all the branches of science are correlated and will assist all the researchers to potentiate their research capabilities.

Abstracting and Indexing

Google Scholar, Elsevier Products (EBSCO and EMBASE), SCImago(SJR), Index Copernicus (ICV: 92), CNKI (China Knowledge Resource Integrated Database), CAS, CASSI (American Chemical Society), ICAAP, Scientific commons, PSOAR, Open-J-Gate, Index Medicus for WHO South-East Asia (IMSEAR), OAI, LOCKKS, OCLC (World Digital Collection Gateway), UIUC.

Address

Published By

Innovare Academic Sciences Pvt Ltd
B-11 Housing Colony In Front Of Bima Hospital, Nai Abadi, Mandsaur M.P. 458001
Email: editor@ajpcr.com

Print By

Fun & Art
29-Nagar Palika Complex, Gandhi chouraha, Mandsaur M.P. 458001

Branch Office

Innovare Academic Sciences Pvt Ltd
T-8, Mahaveer Apartment, Near SIRT College, Ayodhya Bypaas, Bhopal-462041, MP, India
URL: <http://innovareacademics.in> | Email: iajournals@gmail.com | Contact: +91 810 972 5561

Asian Journal of Pharmaceutical and Clinical Research (AJPCR)

Vol 12, issue 8, 2019

P-ISSN-0974-2441

E-ISSN-2455-3891

Contents

SR. NO.	ARTICLE	PAGE NO.
	CASE STUDY(S)	
1	N-ACETYLCYSTEINE, A BOON FOR YELLOW PHOSPHORUS-INDUCED ACUTE LIVER FAILURE? A CASE REPORT AASHIQ AHAMED SHUKKOOR, NIMMY ELIZABETH GEORGE, SARAVANAN THANGAVELU	1-3
	REVIEW ARTICLE(S)	
2	A REVIEW OF BOTANY, MEDICINAL USES, AND BIOLOGICAL ACTIVITIES OF PENTANISIA PRUNELLOIDES (RUBIACEAE) ALFRED MAROYI	4-9
3	SCABIOSA COLUMBARIA: A REVIEW OF ITS MEDICINAL USES, PHYTOCHEMISTRY, AND BIOLOGICAL ACTIVITIES ALFRED MAROYI	10-14
4	A SYNTHESIS AND REVIEW OF MEDICINAL USES, PHYTOCHEMISTRY AND BIOLOGICAL ACTIVITIES OF HELICHRYSUM ODORATISSIMUM (L.) SWEET ALFRED MAROYI	15-23
5	ARSENIC: A HARMFUL AND DESECRATE COMPOUND FOR THE HUMANS SURESH CHANDRA, ARYENDU KUMAR SAINI, AKASH KUMAR GUPTA	24-29
6	ZANTEDESCHIA AETHIOPICA (L.) SPRENG.: A REVIEW OF ITS MEDICINAL USES, PHYTOCHEMISTRY, AND BIOLOGICAL ACTIVITIES ALFRED MAROYI	30-36
7	ANTI-ULCER AGENTS: A PHARMACOLOGICAL UPDATE OF THE PAST TEN YEARS SUDIP KUMAR MANDAL, SUBHOJIT DAWN, ANINDYA BOSE	37-41
	ORIGINAL ARTICLE(S)	
8	KNOWLEDGE AND PERCEPTIONS OF SELF-MEDICATION PRACTICES IN AN URBAN COMMUNITY VARGHESE SNEHA SUJA, SNEHA DUTTA, ANN MARY SWAROOP	42-45
9	BIOANALYTICAL METHOD DEVELOPMENT AND VALIDATION OF CANAGLIFLOZIN IN HUMAN PLASMA BY LIQUID CHROMATOGRAPHY-TANDEM MASS SPECTROMETRY DEEPAN T, BASAVESWARA RAO MV, DHANARAJU MD	46-51
10	PREPARATION AND EVALUATION OF IBUPROFEN EFFERVESCENT GRANULES JUTI RANI DEVI, BIDYUT DAS	52-55
11	STABILITY-INDICATING REVERSE-PHASE HIGH- PERFORMANCE LIQUID CHROMATOGRAPHIC METHOD DEVELOPMENT FOR SIMULTANEOUS ESTIMATION OF SOFOSBUVIR AND DACLATASVIR IN PURE AND PHARMACEUTICAL FORMULATION SWETHA A, RAMYA KUBER B	56-63
12	THE ANTIBIOTIC CONSUMPTION AT A PEDIATRIC WARD AT A PUBLIC HOSPITAL IN INDONESIA FAUNA HERAWATI, MUHAMAD SATRIA MANDALA PUA UP, RIKA YULIA, RETNOSARI ANDRAJATI	64-67
13	PROTECTIVE OF ETHANOLIC EXTRACT OF SAUSSUREA LAPPA AGAINST PARACETAMOL-INDUCED HEPATIC AND RENAL DAMAGE IN MALE RABBITS MARIAM A KADHEM	68-73
14	ENCAPSULATION OF IBUPROFEN INTO SOLID LIPID NANOPARTICLES FOR CONTROLLED AND SUSTAINED RELEASE USING EMULSIFICATION SOLVENT EVAPORATION TECHNIQUE WESLEY N OMWOYO, MAKWENA J MOLOTO	74-81
15	FREE RADICAL SCAVENGING, ANTIOXIDANT POTENTIAL, AND NITRIC OXIDE INHIBITION IN HUMAN THP1 DERIVED MACROPHAGES BY KOKILAKSHAM KASHAYAM TEENA MERLIN, PRAKASH KUMAR B	82-86
16	ANTIHYPERTENSIVE EFFECT OF RUTIN: PHARMACOLOGICAL AND COMPUTATIONAL APPROACH GANGA RAJU M, PREM PRASAD GOUD, SUVARCHALA REDDY NVL	87-92
17	AQUATIC/SEMI-AQUATIC MACROPHYTES USED IN HERBAL REMEDIES FROM THE WETLANDS OF WESTERN ASSAM, NORTH-EAST INDIA	93-96

18	UPEN DEKA, TAPAN DUTTA, SANJAY TALUKDAR EVALUATION OF CYTOTOXICITY AND APOPTOTIC POTENTIAL OF OPHIORRHIZA PECTINATA ARN. - A POTENT ANTICANCER AGENT	97-102
19	PREETHA MOL SN, JOHN E THOPPIL TO STUDY THE ULCEROPROTECTIVE EFFECT OF LEAVES OF MORINGA OLEIFERA ON EXPERIMENTALLY INDUCED INFLAMMATORY BOWEL DISEASE ON ANIMAL MODELS	103-106
20	SHIPRA KAUSHIK, SHOBHIT KAUSHIK IN SILICO DESIGN, SYNTHESIS AND EVALUATION OF IN VITRO GLUCOSE UPTAKE, GENE EXPRESSION, AND A-GLUCOSIDASE INHIBITORY ACTIVITY OF NOVEL 2-AMINOBENZIMIDAZOLE DERIVATIVES	107-114
21	SREEJA S, ANTON SMITH A, MATHAN S FORMULATION AND EVALUATION OF OINTMENT CONTAINING SUNFLOWER WAX	115-120
22	AVISH D MARU, SWAROOP R LAHOTI APPLICATION OF CENTRAL COMPOSITE DESIGN AND RESPONSE SURFACE METHODOLOGY FOR OPTIMIZATION OF METAL ORGANIC FRAMEWORK: NOVEL CARRIER FOR DRUG DELIVERY	121-127
23	PREETI KUSH, JITENDER MADAN, PARVEEN KUMAR A VALIDATED ANALYTICAL METHOD FOR THE SIMULTANEOUS ESTIMATION OF CYTARABINE AND DAUNORUBICIN IN BULK AND INFUSION FORMULATION BY REVERSE PHASE HIGH PERFORMANCE LIQUID CHROMATOGRAPHY	128-131
24	PRASANTHI CHENGALVA, LATHA LAVANYA PEDDAVENGARI, MADHAVI KUCHANA EVALUATION OF ANTIMICROBIAL ACTIVITY AND PHYTOCHEMICALS ANALYSIS OF WHOLE PLANT EXTRACT OF VINCA ROSEA	132-136
25	ANGELIN JEBAMALAR JAYARAJ, JOTHI UCHIMAHALI, THIYAGARAJAN GNANASUNDARAM, SIVAKUMAR THIRUMAL MOMORDICA CHARANTIA NANO DRY POWDER ORAL FORMULATION IN CONTROLLING HISTAMINE-INDUCED BRONCHOSPASM	137-140
26	KARTHIKA P, BHUVANESWARI K, BALACHANDER R PHYTOCHEMICAL EVALUATION, GC-MS ANALYSIS OF PHYTOACTIVE COMPOUNDS, AND ANTIBACTERIAL ACTIVITY STUDIES FROM LINUM USITATISSIMUM	141-149
27	THAMILMARAI SELVI B, SATHAMMAI PRIYA N, STEFFI PF, PRIYADARSHNI S SYNTHESIS AND ANTIBACTERIAL ACTIVITY OF NOVEL 5-ARYLIDENE-2-IMINO-3-(2-PHENYL-1,8-NAPHTHYRIDIN-3-YL)THIAZOLIDIN-4-ONES	150-153
28	RAMALINGAM KUNDENAPALLY, RAMESH DOMALA, SREENIVASULU BATHULA PROLONGED EXPOSURE TO DIFFERENT TYPES OF ENERGY RESTRICTED CALCIUM DIET MODULATES THE INFLAMMATORY AND OXIDATIVE STRESS RESPONSE IN HEALTHY MALE RATS	154-159
29	SANDEEP DAS, DIPAYAN CHOUDHURI THE THE TERATOGENIC EFFECTS OF ETHANOLIC EXTRACT OF BINTANGUR LEAVES (CALOPHYLLUM SOULATTRI BURM. F) ON FEMALE WHITE RATS	160-163
30	INARAH FAJRIATY, HAFRIZAL RIZA, FAJAR NUGRAHA, FRENCKI FRIANTO PROTECTIVE EFFECT OF ROTENONE AGAINST LIPOPOLYSACCHARIDE AND D-GALACTOSAMINE-INDUCED HEMATOTOXICITY	164-167
31	SAMRAT RAKSHIT, ANJANI VERMA, SATENDRA KUMAR NIRALA, MONIKA BHADAURIA IMPROVED CIPROFLOXACIN PENETRATION IN GOAT EYES USING COMPLEXATION TECHNIQUE	168-171
32	DURGA PANDEY, DEEPTI JAIN HYPOGLYCEMIC EFFECT OF HIGH-RESISTANT STARCH ANALOG RICE THROUGH GLP-1 AND INSULIN OR HIGH-RESISTANT STARCH ANALOG RICE ATTENUATES BLOOD GLUCOSE LEVEL THROUGH ENHANCEMENT OF GLP-1 AND INSULIN	172-175
33	HAIRRUDIN, SOETJIPTO, RETNO HANDAJANI DEVELOPMENT AND CHARACTERIZATION OF NEVIRAPINE LOADED AMORPHOUS SOLID DISPERSIONS FOR SOLUBILITY ENHANCEMENT	176-182
34	GAGANDEEP SINGH, NAVJOT SINGH, RANDEEP KUMAR, NEENA BEDI KNOWLEDGE AND PERCEPTIONS OF VITAMIN D DEFICIENCY AMONG THE UNITED ARAB EMIRATES POPULATION	183-186
35	OSAMA MOHAMED IBRAHIM, NOOR KIFAH AL-TAMEEMI, DALIA DAWOUD ATOMIC ABSORPTION SPECTROPHOTOMETER - A VERSATILE TOOL FOR THE ESTIMATION OF NICKEL CONTENT IN DRONEDARONE	187-190
36	VASAVI DEVI DASARI, SWARNALATHA DUGASANI, VENKATASUBBA REDDY GOPIREDDY PHYTOCHEMICAL INVESTIGATION OF TERMINALIA BELLIRICA FRUIT INSIDE	191-194

37	HAZRA K THE COMBINED EFFECT OF DEXAMETHASONE AND BACILLUS CALMETTE-GUÉRIN ON THE NEUROBEHAVIORAL ASPECT IN MALE MICE (MUS MUSCULUS)	195-201
38	ABDELKRIM HAOUJI, SAMIR DJEMLI, ABDELKRIM TAHRAOUI PHYTOCHEMICAL SCREENING AND ANTIOXIDANT POTENTIAL OF ANACARDIUM OCCIDENTALE, ACHYRANTHES ASPERA, AND AEGLE MARMELOS	202-205
39	BRIJYOG, LALITESHWAR PRATAP SINGH, SUSHIL KUMAR, SHWETA VERMA THE METAL COMPLEXES OF NOVEL SCHIFF BASE CONTAINING ISATIN: CHARACTERIZATION, ANTIMICROBIAL, ANTIOXIDANT AND CATALYTIC ACTIVITY STUDY	206-210
40	VAIRALAKSHMI M, PRINCESS R, JOHNSON RAJA S ASSESSMENT OF THE CURRENT GERIATRIC PHARMACEUTICAL CARE IN THE UNITED ARAB EMIRATES	211-215
41	OSAMA MOHAMED IBRAHIM, RANA IBRAHIM IN VITRO WOUND HEALING AND ANTIMICROBIAL PROPERTY OF COTTON FABRICS COATED OPTIMIZED SILVER NANOPARTICLES SYNTHESIZED USING PELTOPHORUM PTEROCARPUM LEAF EXTRACTS	216-222
42	ANNAMALAI P, BALASHANMUGAM P, KALAICHELVAN PT SYNTHESIS AND OPTIMIZATION OF GEMCITABINE-LOADED MIL-101NH ₂ (FE) NANOCARRIERS: RESPONSE SURFACE METHODOLOGY APPROACH	223-229
43	PREETI KUSH, JITENDER MADAN, PARVEEN KUMAR ANTIFUNGAL EFFICIENCY OF COPPER OXYCHLORIDE-CONJUGATED SILVER NANOPARTICLES AGAINST COLLETOTRICHUM GLOEOSPORIOIDES WHICH CAUSES ANTHRACNOSE DISEASE	230-233
44	RAGHAVENDRA SN, RAGHU HS, DIVYASHREE K, RAJESHWARA AN STUDIES ON GLUCOSE-LOWERING EFFICACY OF THE ACACIA SUMA ROXB. ROOT	234-238
45	JITENDRA DEBATA, SUNDEEP KUMAR HK AN IN-VITRO ANTIBIOFILM ACTIVITY OF CHLORELLA VULGARIS	239-242
46	SRIDEVI NS, DHANUSHA V, RAJESWARI M, SANTHI N STUDY OF ANTIMICROBIAL ACTIVITY OF SILVER-DOPED HYDROXYAPATITE	243-245
47	HIBU WAHID, ANJUVAN SINGH TERATOGENICITY TESTING OF SIDHA FORMULATION OF NILAVEMBU KUDINEER ON ZEBRAFISH (DANIO RERIO) EMBRYO	246-250
48	SHANMUGAPRIYA P, ELANSEKARAN S, RAMAMURTHY M, JEEVA GLADYS R, SIVARAMAN D, CHRISTIAN GJ STABILITY-INDICATING METHOD DEVELOPMENT AND VALIDATION FOR THE ESTIMATION OF ROSUVASTATIN CALCIUM IN BULK AND TABLET FORMULATION BY REVERSE-PHASE HIGH-PERFORMANCE LIQUID CHROMATOGRAPHY	251-256
49	SAILAJA B, SRAVANA KUMARI K SMARTPHONE ADDICTION AND ITS RELATIONSHIP WITH PSYCHOLOGICAL HEALTH AMONG STUDENTS OF A MEDICAL SCHOOL IN EAST COAST MALAYSIA	257-260
50	AZWANIS ABDUL HADI, HAWARI MUSYIR MOHD NAWAWI, NURAFIFAH SHAMSURI, NURUL NAJIAH RAHIM, HAFIZAH PASI EPIDEMIOLOGIC ASPECTS OF TRAUMA IN AL-JOUF REGION, SAUDI ARABIA: A RETROSPECTIVE STUDY	261-264
51	DOAA M ABDEL-SALAM, RAWAN R ALRUWAILI, FARAH S ALHABLANI, NOUR H ALFAHEL, ANWAR A ALBLAWI THE ROLE OF SELENIUM MICRONUTRIENTS AS ANTIOXIDANTS IN EXPOSURE TO E-CIGARETTE SMOKE	265-268
52	RIVAN VIRLANDO SURYADINATA, MERRYANA ADRIANI, SANTI MARTINI, SRI SUMARMI, BAMBANG WIRJATMADI EVALUATION OF NUCLEAR CHANGES IN THE BUCCAL EPITHELIAL CELLS OF TOBACCO USERS IN NNEWI, SOUTH EAST NIGERIA	269-273
53	SAMUEL IFEDIORANMA OGENYI1, ANTHONY AJULUCHUKWUNGOKERE, JONATHAN MADUKWE EFFECTS EFFECTS OF PUMPKIN (CUCURBITA MOSCHATA DURCH) SEED POWDER ON BLOOD GLUCOSE LEVEL IN STREPTOZOTOCIN-INDUCED MICE	274-276
54	NOVARIANTI MARBUN, RUTH MAYANA RUMANTI THERMODYNAMIC AND KINETIC STUDIES FOR THE BINDING OF LEAD ION BY CHELATING WITH THEOPHYLLINE 1,3-DIMETHYLXANTHINE (THERAPY DRUG)	277-283
55	HUDA N AL-ANI ANATOMICAL STUDIES OF THE FRUIT OF ZIZIPHUS RUGOSA	284-287
56	PREMA G, CHITRA M RESEARCH ARTICLE LEAVES EXTRACT-BASED BIOGENIC SYNTHESIS OF CUPRIC OXIDE NANOPARTICLES, CHARACTERIZATIONS, AND ANTIMICROBIAL ACTIVITY	288-291

57	YASHWANI PRAKASH A PREPARATION, CHARACTERIZATION, AND IN VITRO SKIN PENETRATION OF MORUS ALBA ROOT EXTRACT NANOEMULSION	297-301
58	MAZAYA FADHILA, ABDUL MUN IM, MAHDI JUFRI SPECTROPHOTOMETRIC DETERMINATION OF TOTAL PHENOLIC CONTENT FOR STANDARDIZATION OF VARIOUS PHYLLANTHUS SPECIES	297-301
59	RAKHI KHABIYA SOLUBILITY ENHANCEMENT OF GLIBENCLAMIDE USING MESOPOROUS SILICA	302-314
	SWATI MITTAL, AKSHAY SONAWANE, MANGESH KHUNE	

THE ROLE OF SELENIUM MICRONUTRIENTS AS ANTIOXIDANTS IN EXPOSURE TO E-CIGARETTE SMOKE

RIVAN VIRLANDO SURYADINATA^{1,2}, MERRYANA ADRIANI³, SANTI MARTINI⁴, SRI SUMARMI³,
BAMBANG WIRJATMADI^{3*}

¹Doctoral Program of Public Health, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia. ²Department of Public Health, Faculty of Medicine, Universitas Surabaya (UBAYA), Surabaya, Indonesia. ³Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia. ⁴Department of Epidemiology, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia
Email: b.wirjatmadi@fkm.unair.ac.id

Received: 09 July 2019, Revised and Accepted: 15 July 2019

ABSTRACT

Objective: E-cigarette products have resulted in various controversies concerning their posed impacts on health. Some argue that exposure to e-cigarette smoke could improve free radicals in the body; thus, it causes harming impacts on health. Peroral selenium (Se) administration can increase superoxide dismutase (SOD) and glutathione peroxidase (GPx) serving as antioxidants in the body.

Methods: This research is an experimental study aiming to analyze the effectiveness of Se to decrease free radical due to exposure to e-cigarette smoke as one of the preventive actions. The research was carried out to male Wistar rats with exposure to e-cigarette smoke and peroral Se intake with different time and duration of administration.

Results: Research results showed a decrease of antioxidant SOD and GPx in the administration of exposure to e-cigarette smoke, and they gradually increased after Se administration ($p=0.000$). Meanwhile, the malondialdehyde level was inversely proportional compared to antioxidant SOD and GPx.

Conclusion: Se is a micronutrient that can reduce free radicals due to exposure to e-cigarette smoke through enhancement of antioxidant enzymes such as SOD and GPx.

Keywords: Selenium, Malondialdehyde, E-cigarette, Glutathione peroxidase, Superoxide dismutase.

© 2019 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) DOI: <http://dx.doi.org/10.22159/ajpcr.2019.v12i8.34454>

INTRODUCTION

The use of electronic cigarettes keeps on increasing annually. It is caused by the absence of specific regulations targeting the use of e-cigarettes leading to their use at any place [1]. Besides, their great shapes and models compared to tobacco cigarettes along with their massive promotion attract young people [2]. Electronic cigarettes are also claimed as tools to reduce or stop the use of tobacco cigarettes. They are also considered to be safer and have less risk because they only produce water vapor claimed not to endanger health [3]. Various studies have shown the dangerous risks posed by exposure to electronic cigarettes. The danger of electronic cigarettes has been estimated to be greater compared to the one in tobacco cigarettes. Even in passive smokers inhaling the smoke of e-cigarettes, nicotine is also found in blood [4]. The inhaled smoke of e-cigarettes entering airways can trigger the existence of free radicals. The amount enhancement of excessive free radicals will cause amount imbalance of free radicals and antioxidants in the body so that oxidative stress occurs [5].

Free radicals in the body are molecules having one or more electrons which are not in pairs, but very reactive, and unstable. Free radicals can react fast with other molecules by catching electrons to stabilize themselves. Free radicals will be balanced by taking electrons, whereas the molecules under attacked will change into other free radicals causing damage to cells [6]. Oxidative stress will enhance lipid peroxidase on the attacked cell membrane [7]. One of lipid peroxidase results due to free radicals is malondialdehyde (MDA), which can be used as a parameter of free radicals enhancement in the body [8].

Reduction of free radicals can be done by administering antioxidants. Antioxidants mechanism works by giving electrons to the free radicals; thereby, they can stop cell damage process [9]. Besides, antioxidants

will neutralize free radicals so that they will not have any ability to take electrons from the cells [10]. Physiologically, human body can neutralize free radicals when their amount is not excessive [11]. This is done by means of endogenous antioxidant defense mechanism [12,13]. However, when there is a lack of endogenous antioxidants, the body needs exogenous antioxidants [14]. Various ways to enhance antioxidants in the body include adding intake of exogenous antioxidants from food, supplements, and doing sports [15]. Some of the endogenous antioxidants needed to neutralize free radicals free are glutathione peroxidase (GPx) and superoxide dismutase (SOD). Antioxidant GPx is an intracellular antioxidant enzyme which enzymatically changes hydrogen peroxidase (H_2O_2) to water (H_2O) [16]. Meanwhile, antioxidant SOD plays a role in changing superoxide radical into H_2O_2 [17].

Selenium (Se) has antioxidant and immunomodulation functions. In the body, Se is used to synthesize amino acid, called selenocysteine, which is very important for selenoprotein formation. Selenoprotein is composed of 25 types in human body and classified based on its function as various enzymes which one of them is antioxidant GPx [18]. Thus, Se is often called as an enzymatic component of GPx [19]. The main form of Se is selenomethionine, commonly found in food [20]. Main sources of Se in food are selenomethionine amino acid (in the form of cereals) and selenocysteine (from animal products). Some plants such as beet leaves, cabbage, and garlic contain a high level of Se, reaching up to 50% in the form of selenate. Se can also be obtained from supplements in the forms of selenate, selenite, selenomethionine, or other forms of Se [21]. Moreover, Se has a quite significant function for human's health because it is able to join into protein to be selenocysteine. Se effects have been proven, ranging from antioxidant to anti-inflammatory effects [22].

Various studies have proven that SE suppresses reactive oxygen species formation by improving GPx activity [23]. Antioxidant SOD is more influenced by copper and zinc micronutrients. Hence, the role of Se against SOD enhancement still needs further research [24].

METHODS

This research is an experimental study using a post-test control group design (Ethical Approval No:103/EA/KEPK/2018). The research was conducted for 5 weeks utilizing male Wistar rats (*Rattus norvegicus*). The research was divided into six groups with different treatments in each group.

The first group was the negative control group where the experimental animals did not undergo any intervention for 4 weeks. The second group was the cigarette control group where the experimental animals were given intervention of exposure to e-cigarette smoke for 4 weeks. The third group was the Se control group where the experimental animals were given intervention of peroral Se intake for 4 weeks. The fourth group was treatment Group I where the experimental animals were given exposure to e-cigarette smoke on the 1st week. Next, on the second until the 5th week, they were given exposure to e-cigarette smoke and peroral Se. The fifth group was treatment Group II, where the experimental animals were given exposure to e-cigarette smoke and peroral Se for 4 weeks. The sixth group was treatment Group III where the experimental animals were given peroral Se in the 1st week, next during the second until the 5th week they were given exposure to e-cigarette smoke and peroral Se.

Mice

Male Wistar rats aged 2–3 months weighing 150–200 g were used as animal models. The Wistar rats were never been subjects research and did not have macroscopic abnormalities. The experimental animals are adapted first for 5 days before giving intervention. The research was conducted in the Laboratory of the Faculty of Medicine, Universitas Airlangga.

Se

Se per oral used is an active form of selenomethionine. The recommended dose for adults is 200 mcg/day. Using the conversion table Laurence and Bacharach, calculation obtained was 3.6 mcg. Thus, the dosage that can be given to male white rats of the Wistar strain is 3.6 mcg/day per each experimental animal.

E-cigarette

The solution of an e-cigarette used contains 6 mg of nicotine. The room where exposure to e-cigarette smoke measures 50 cm × 40 cm × 20 cm there is space a pipe will be passed which can expend e-cigarette smoke. The exposure to e-cigarette smoke exposure is adjusted for the length of time planned for the research.

Measurement of GPx level in the blood

GPx level in blood measured using an enzyme-linked immunosorbent assay (ELISA) Kit, which can be applied in the quantitative *in vitro* determination of Rat GPx1 concentration in serum, plasma, and other biological fluids. The ELISA kit uses the competitive-ELISA method. The kit had been coated with Rat GPx1 as a specific antibody. The rat GPx1 concentration in the sample was then determined by comparing the sample result to the standard curve.

Measurement of SOD level in blood

SOD level in blood measured using an ELISA kit which can be applied in the quantitative *in vitro* determination of rat SOD1 concentration in serum, plasma, and other biological fluids. The ELISA kit uses the competitive-ELISA method. The kit had been coated with rat SOD1 as a specific antibody. The rat SOD 1 concentration in the sample was then determined by comparing the sample OD to the standard curve.

Measurement of MDA level in blood

MDA levels in blood measured using thiobarbituric acid (TBA)

reactive substance (TBARS) assay. Measurements were carried out using the bioassay system based on the reaction of TBARS with TBA in forming pink-colored compounds. The color intensity was measured at 535 nm or with fluorescence intensity at 560 nm/585 nm which was directly proportional to the concentration of TBARS in the sample.

Statistika

The collected data will be statistically tested using SPSS version 20 to analyze the difference and influence of Se on SOD levels, GPx, and MDA in the blood in all groups.

RESULTS

SOD levels in blood as a result of exposure to e-cigarette smoke

Research results were obtained by comparing the levels of SOD in each group. Based on Fig. 1, the average value and difference of levels of SOD in each group can be known. The results showed that the cigarette control group had the lowest average value reaching 2.29±0.12. Meanwhile, in the Se control group, it had the highest average value reaching 5.05±0.16. In the treatment groups, there was an increase in the average value of SOD sequentially.

To fulfill the parametric test requirements, the research results were subjected to a normality test ($p>0.05$) and homogeneity test ($p=0.008$). Based on Table 1, Kruskal-Wallis test was conducted to know the difference of SOD levels in the groups ($p=0.000$).

GPx levels in blood as a result of exposure to e-cigarette smoke

Research results were obtained by comparing levels of GPx in each group. Based on Fig. 2, the average value and difference of levels of GPx in each group can be known. The results showed that the cigarette control group had the lowest average value reaching 85.6±8.96. Meanwhile, in the Se control group, it had the highest average value reaching 330.6±13.22. In the treatment groups, there was an increase in the average value of GPx sequentially.

To fulfill the parametric test requirements, the research results were subjected to normality test ($p>0.05$) and homogeneity test ($p=0.385$). Based on Table 2, analysis of variance test was conducted to know the difference of GPx levels in the groups ($p=0.000$).

Table 1: Kruskal-Wallis test for superoxide dismutase in each group

Groups	Definition	Mean±SD	p-value
I	Negative control group	4.64±0.27	0.000
II	Cigarette control group	2.29±0.12	
III	Selenium control group	5.05±0.16	
IV	Treatment Group I	3.00±0.04	
V	Treatment Group II	3.33±0.1	
VI	Treatment Group III	4.67±0.07	




Fig. 1: The mean value for superoxide dismutase in each group




Fig. 2: The mean value for glutathione peroxidase in each group




Fig. 3: The mean value for malondialdehyde in each group

MDA levels in blood as a result of exposure to e-cigarette smoke

Research results were obtained by comparing levels of MDA in each group. Based on Fig. 3, the average value and difference of levels of MDA in each group can be known. The results showed that the negative control group had the lowest average value reaching 0.0144 ± 0.0018 . Meanwhile, in the cigarette control group, it had the highest average value reaching 0.2416 ± 0.0174 . In the treatment groups, there was an decrease in the average value of MDA sequentially.

To fulfill the parametric test requirements, the research results were subjected to normality test ($p > 0.05$) and homogeneity test ($p = 0.000$). Based on Table 3, Kruskal-Wallis test was conducted to know the difference of MDA levels in the groups ($p = 0.000$).

DISCUSSION

E-cigarette smoke entering the airways will increase free radicals in the body. The enhancement of free radicals can trigger an increase of MDA as well as a decrease of the body's enzymatic antioxidants in the forms of SOD and GPx [25]. The free radicals are also actually needed in the body because they play an important role in the body's defense system [26]. However, the imbalance between amount of free radicals and antioxidants will cause the occurrence of cell damage [27,28]. Additional intake of selenium per oral is needed to stimulate antioxidant increase in the body [29,30].

Administering Se in this research showed enhancement existence of SOD and GPx. The group with the highest increase of antioxidants was the Se control group. It was proven by higher enhancement of antioxidants compared to the negative control groups. Meanwhile, the cigarette control group had the lowest level of antioxidants. This was caused by an increase of the amount of free radicals entering the airways because of exposure to e-cigarette smoke. On the three treatment groups, there was antioxidant enhancement gradually. The first treatment group had a higher level of antioxidant due to the administration of Se done earlier than exposure to e-cigarette smoke. As a result, the body's enzymatic

Table 2: Analysis of variance test for glutathione peroxidase in each group

Groups	Definition	Mean \pm SD	p-value
I	Negative control group	236.2 \pm 7.3	0.000
II	Cigarette control group	85.6 \pm 8.96	
III	Selenium control group	330.6 \pm 13.22	
IV	Treatment Group I	118.2 \pm 6.46	
V	Treatment Group II	154.0 \pm 5.66	
VI	Treatment Group III	256.4 \pm 6.69	

Table 3: Kruskal-Wallis test for malondialdehyde in each group

Groups	Definition	Mean \pm SD	p-value
I	Negative control group	0.0144 \pm 0.0018	0.000
II	Cigarette control group	0.2416 \pm 0.0174	
III	Selenium control group	0.0204 \pm 0.0028	
IV	Treatment Group I	0.2104 \pm 0.0051	
V	Treatment Group II	0.1802 \pm 0.0043	
VI	Treatment Group III	0.1410 \pm 0.0032	

antioxidants were still able to reduce the free radicals entering through the airways. In the second treatment group, the administration of both exposures to e-cigarette and Se was done at the same time. Therefore, there was a decrease in the antioxidant level. The third treatment group indicated the lowest level of antioxidants of all other treatment groups. It happened because the free radicals had entered the airways earlier. Hence, the antioxidant enhancement did not show any significant increase.

Se is a supplement that plays an important role in human's health. The given Se intake is expected to be able to prevent the increase of antioxidants SOD and GPx as well as provide with potential hope as a therapy to prevent cell damage caused by exposure to e-cigarette [31].

CONCLUSION

Se is a micronutrient that can reduce free radicals due to exposure to e-cigarette smoke through enhancement of antioxidant enzymes such as SOD and GPx.

AUTHORS' CONTRIBUTIONS

All the authors have contributed equally in research, review, and finalization of the contents of the journal.

CONFLICTS OF INTEREST

Declare None.

REFERENCES

- Canistro D, Vivarelli F, Cirillo S, Marquillas CB, Buschini A, Lazzaretti M, et al. E-cigarettes induce toxicological effects that can raise the cancer risk. *Sci Rep* 2017;7:2028.
- Grana R, Benowitz N, Glantz SA. E-cigarettes: A scientific review. *Circulation* 2014;129:1972-86.
- Vardavas CI, Anagnostopoulos N, Kougias M, Evangelopoulou V, Connolly GN, Behrakis PK, et al. Short-term pulmonary effects of using an electronic cigarette: Impact on respiratory flow resistance, impedance, and exhaled nitric oxide. *Chest* 2012;141:1400-6.
- Polosa R, Caponnetto P, Morjaria JB, Papale G, Campagna D, Russo C, et al. Effect of an electronic nicotine delivery device (e-cigarette) on smoking reduction and cessation: A prospective 6-month pilot study. *BMC Public Health* 2011;11:786.
- Suryadinata RV, Wirjatmadi B, Adriani M. Effectiveness Decrease Combined with Supplements Malondialdehyde Antioxidant Superoxide Dismutase Gliadin Melon with Due to Exposure to Cigarette. *Glob Med Heal Commun* 2017;5:79-83.
- Suryadinata RV. Effect of free radicals on inflammatory process in chronic obstructive pulmonary disease (COPD). *Amerta Nutr*

- 2018;2:317-24.
7. Raghavendra U, Rao A, D'souza J, Pai VR, Nair S, Kumar V, et al. Comparative estimation of salivary total antioxidant capacity in periodontal health and chronic periodontitis a pilot study. *Asian J Pharm Clin Res* 2018;11:523.
 8. Ayala A, Muñoz MF, Argüelles S. Lipid peroxidation: Production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. *Oxid Med Cell Longev* 2014;2014:360438.
 9. Mapunda PE, Mligo C, Lyaruu HV. Evaluation of free radical scavenging ability and antiradical activities of *Ximenia caffra* fruit extracts at different ripening stages. *Int J Pharm Pharm Sci* 2019;11:55.
 10. Pratiwi SR, Lorensia A, Suryadinata RV. Asupan Vitamin C dan E dengan SQ-FFQ terhadap fungsi paru perokok dan non-perokok. *Media Kesehatan Masyarakat Indones* 2018;14:101.
 11. Subba A, Mandal P. Standardization of a traditional polyherbal formulation with pharmacognostic study; its phytochemical content, antioxidant, and antidiabetic activity. *Asian J Pharm Clin Res* 2019;12:5.
 12. Rahal A, Kumar A, Singh V, Yadav B, Tiwari R, Chakraborty S, et al. Oxidative stress, prooxidants, and antioxidants: The interplay. *Biomed Res Int* 2014;2014:761264.
 13. Pratima H. Antioxidant and antibacterial activity of alkaloid extract of *Cucumis trigonus* Roxb. *Int J Pharm Pharm Sci* 2019;11:44-8.
 14. Liguori I, Russo G, Curcio F, Bulli G, Aran L, Della-Morte D, et al. Oxidative stress, aging, and diseases. *Clin Interv Aging* 2018;13:757-72.
 15. Suryadinata RV, Wirjatmadi B, Adriani M. Effects of Changes in Goblet Cell Hyperplasia for 28 Days of Cigarette Smoke Exposure through the administration of Antioxidant Superoxide Dismutase. *Indones J Public Health* 2017;11:60.
 16. Lubos E, Loscalzo J, Handy DE. Glutathione peroxidase-1 in health and disease: From molecular mechanisms to therapeutic opportunities. *Antioxid Redox Signal* 2011;15:1957-97.
 17. Phaniendra A, Jestadi DB, Periyasamy L. Free radicals: Properties, sources, targets, and their implication in various diseases. *Indian J Clin Biochem* 2015;30:11-26.
 18. Kim Y, Kim DC, Cho ES, Ko SO, Kwon WY, Suh GJ, et al. Antioxidant and anti-inflammatory effects of selenium in oral buccal mucosa and small intestinal mucosa during intestinal ischemia-reperfusion injury. *J Inflamm (Lond)* 2014;11:36.
 19. Socha K, Kochanowicz J, Karpińska E, Soroczyńska J, Jakoniuk M, Mariak Z, et al. Dietary habits and selenium, glutathione peroxidase and total antioxidant status in the serum of patients with relapsing-remitting multiple sclerosis. *Nutr J* 2014;13:62.
 20. Bermingham EN, Hesketh JE, Sinclair BR, Koolaard JP, Roy NC. Selenium-enriched foods are more effective at increasing glutathione peroxidase (GPx) activity compared with selenomethionine: A meta-analysis. *Nutrients* 2014;6:4002-31.
 21. Mehdi Y, Hornick JL, Istasse L, Dufrasne I. Selenium in the environment, metabolism and involvement in body functions. *Molecules* 2013;18:3292-311.
 22. Cai X, Wang C, Yu W, Fan W, Wang S, Shen N, et al. Selenium exposure and cancer risk: An updated meta-analysis and meta-regression. *Sci Rep* 2016;6:19213.
 23. Kim KS, Suh GJ, Kwon WY, Kwak YH, Lee K, Lee HJ, et al. Antioxidant effects of selenium on lung injury in paraquat intoxicated rats. *Clin Toxicol (Phila)* 2012;50:749-53.
 24. Younus H. Therapeutic potentials of superoxide dismutase. *Int J Health Sci (Qassim)* 2018;12:88-93.
 25. Suryadinata RV, Wirjatmadi B, Adriani M, Sumarmi S. The effects of exposure duration to electronic cigarette smoke on differences in superoxide dismutase and malondialdehyde in blood of wistar rats. *Int J Curr Pharm Res* 2019;11:13-6.
 26. Chidambararajan P, Keerthana V, Priyadharshini K, Sakthivel B. *In vitro* antioxidant and anticancer activity of ulva lactuca l. Using MOLT-3 cell line. *Asian J Pharm Clin Res* 2019;12:3-6.
 27. Simioni C, Zauli G, Martelli AM, Vitale M, Sacchetti G, Gonelli A, et al. Oxidative stress: Role of physical exercise and antioxidant nutraceuticals in adulthood and aging. *Oncotarget* 2018;9:17181-98.
 28. Jing SL, Yen KP, Dash GK. *In vitro* antioxidant and photoprotective activities of *Carica papaya* fruits. *Asian J Pharm Clin Res* 2019;12:2-4.
 29. Lobo V, Patil A, Phatak A, Chandra N. Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacogn Rev* 2010;4:118-26.
 30. Pizzino G, Irrera N, Cucinotta M, Pallio G, Mannino F, Arcoraci V, et al. Oxidative stress: Harms and benefits for human health. *Oxid Med Cell Longev* 2017;2017:8416763.
 31. Wang N, Tan HY, Li S, Xu Y, Guo W, Feng Y, et al. Supplementation of micronutrient selenium in metabolic diseases: Its role as an antioxidant. *Oxid Med Cell Longev* 2017;2017:7478523.



IAS Services

- Book Publication
- Journal Publication
- Journal Subscription
- Special Thematic Issue
- Copy-editing & Typesetting
- Conference Proceedings & Abstracts
- Conference/Seminar/Workshop/Symposia
Event management

Why to publish in IAS ?

- Rapid publication
- Reprints (on request)
- Unlimited space for your data
- Quality, reputation and high standard of peer review.
- Wide range of article publication
- Indexing in major bibliographic citation
- Provide high visibility due to open access

INNOVARE ACADEMICS SCIENCES JOURNALS

Asian Journal of Pharmaceutical & Clinical Research
International Journal of Pharmacy & Pharmaceutical Sciences
International Journal of Chemistry Research
International Journal of Applied Pharmaceutics
International Journal of Current Pharmaceutical Research
Journal of Critical Reviews
Innovare Journal of Medical Sciences
Innovare Journal of Ayurvedic Sciences
Innovare Journal of Health Sciences
Innovare Journal of Engineering & Technology
Innovare Journal of Social Sciences
Innovare Journal of Business Management
Innovare Journal of Education
Innovare Journal of Agricultural Science
Innovare Journal of Food Science
Innovare Journal of Life Science
Innovare Journal of Sciences

Asian Journal of Pharmaceutical & Clinical Research
B-11, In front of Beema Hospital, Nayi Abadi
Mandsaur - 458001, MP, India
Contact No: +91 940 661 2909
Email: editor@ajpcr.com
www.innovareacademics.in