

Marine Sponge suberitidae 2021

by Sapto Andriyono

Submission date: 09-Jan-2023 05:07PM (UTC+0800)

Submission ID: 1990127319

File name: EJABF_Volume_25_Issue_5_Pages_409-418.pdf (478.65K)

Word count: 3725

Character count: 22457



Identification of Bacterial Symbionts from the Marine Sponge *Aptos suberitoides* (Demospongiae: Suberitidae)

Hafshoh Maisyaroh¹, Rahayu Kusdarwati², Heru Pramono³, Syafrina Sari Lubis⁴
Farikhah⁵, Supto Andriyono^{3,*}

¹Aquaculture Study Program, Fisheries and Marine Faculty, Universitas Airlangga, Campus C of Universitas Airlangga, Surabaya, East Java Indonesia

²Department of Aquaculture, Fisheries And Marine Faculty, Universitas Airlangga, Campus C of Universitas Airlangga, Surabaya, East Java Indonesia

³Department of Marine, Fisheries And Marine Faculty, Universitas Airlangga, Campus C of Universitas Airlangga, Surabaya, East Java Indonesia

⁴Department of Biology, Ar-Raniry Islamic State University of Banda Aceh, Indonesia

⁵Department of Aquaculture, Faculty of Agriculture, Muhammadiyah University of Gresik, East Java, Indonesia

*Corresponding Author: supto.andriyono@fpk.unair.ac.id

ARTICLE INFO

Article History:

Received: June 18, 2021

Accepted: Aug. 17, 2021

Online: Oct. 25, 2021

Keywords:

Sponges,
Symbiotic,
Bacteria,
Coastal water,
Diversity

ABSTRACT

The sponges are primitive multicellular animals (metazoans) that belong to the phylum of Porifera. They are symbiotic with bacteria, archaea, microalgae and fungi, and the common symbionts organisms. Symbionts bacteria are obtained vertically through gametes and horizontal through the filter feeder process. This study aimed to isolate and the species of symbionts bacteria in *Aptos suberitoides* sponge from Pasir Putih Beach waters Situbondo, East Java. Six symbionts bacterial were isolated from the sponge *Aptos suberitoides*; namely, *Corynebacterium hofmanni*, *Vibrio damsela*, *Oligella urethralis*, *Bacillus coagulan* and *Bordetella parapertusis*. Biochemical tests confirmed that the symbiotic bacteria can perform several processes: nitrification process, citrate, urease, Voges Proskauer, decarboxylase lysine, and decarboxylase ornithine and hydrolysis aesculin. In conclusion, excluding *Bacillus coagulant* bacteria, symbionts bacterial were proved mostly pathogenic to humans and fish.

INTRODUCTION

Sponges are filter feeders known to be symbiotic with bacteria, archaea, microalgae and fungi. These microorganisms can contribute significantly to spongy metabolism (Taylor *et al.*, 2007). Bacteria found to be symbiotic with sponges *Aptos* sp. dominated by bacteria of the class Actinobacteria, Flavobacterium, Alphaproteobacteria, Deltaproteobacteria and Gammaproteobacteria (Chasanah *et al.*, 2013). Sponges can produce chemical compounds as a form of self-defence response to predators and competitors (Taylor *et al.*, 2007). The resulting chemical compounds can induce

symbiotic microorganisms to produce specific secondary metabolites (Nofiani *et al.*, 2009). The sponge *Aaptos suberitoides* produces aaptamine and demethylaaptamine compounds with cytotoxic activity (Hanif *et al.* 2019). Bacteria *Rhodobacteraceae bacterium* (Murniasih *et al.* 2014), *Halomonas aquamarina*, *Alphaproteobacterium* and *Pseudoalteromonas luteviolaceae* symbiotic with sponges of genus *Aaptos* sp. are known to produce antibacterial compounds (Radjasa *et al.*, 2007). Identification of symbiotic bacterial species with sponges is essential for assessing the composition and function of the microbial community (Martin, 2002).

This study aimed to identify bacteria species that are symbiotic with marine sponges. The symbiotic bacteria are expected to be developed and become bioremediation candidates in cultivation activities such as shrimp pond processing and waste fish processing industry, which can decrease the water quality around the activity area.

MATERIALS AND METHODS

Samples Collection

Samples were taken from Pasir Putih beach water Situbondo at a depth of 12-14 meters. After the sponge was taken and cut, the samples were put in a cool box and seawater with fixed aeration. The samples were then taken to the laboratory to be identified and analyzed (Abubakar *et al.*, 2011). Sponge identification refers to the world sponge database accessed through online database (www.marinespecies.org/porifera) and based on sponge morphology (Subagio & Aunurohim, 2013).

Isolation Bacterial Symbiont Sponge

The sterilisation of the sponges surface was done with ethanol (70%) and rinsed thrice with sterile seawater. The rinsed sponges were taken as much as one gram, smoothed using a mortar, and five (5) mL of sterile seawater was added on each sampled gram. The homogeneous solution was diluted by adding one (1) mL of the solution into nine (9) mL of sterile seawater, then the outcome was diluted to 10^{-3} of the initial concentration (Rua *et al.*, 2014). The sample planting was done by surface/spread plate method [11] on Tryptic Soy Agar Sea Water (TSA-SW) media, and the TSA medium diluted with sterile seawater (Whitman *et al.*, 2012). The cultured sample was an outcome of 10^{-1} to 10^{-3} dilution and was incubated at 30°C for one to three days (Cicirelli, 2007). The bacterial colonies were collected using a sterile ose. They were scraped on TSA-SW medium and incubated for 48 hours at 30°C (Setyati & Subagiyo, 2012) until a single colony was obtained.

Bacterial Isolate Identification

Identification of spongy symbiont bacteria includes morphological and biochemical tests. Morphological tests include colony shape, colony colour, colony edge, elevation or

colony surface, colony and Gram staining structures. Biochemical tests included catalase, oxidase, O/F, TSIA, nitrate reduction, gelatin, MR-VP, phenylalanine deaminase, urease, citrate utilisation, glucose, lactose, sucrose, maltose, mannitol, dulcitol, inositol, sorbitol, arabinose, raffinose and xylose, lysine, malonate, MIO and aesculine. The results of further identification tests were compared to Cowan and Steel's Manual for the Identification of Medical Bacteria (Cowan and Steel, 1965) and Bergey's Manual of Determinative Bacteriology (Whitman *et al.*, 2012), which was used for the analysis and determination species from isolates of symbiotic bacteria.

RESULTS

Isolation of Bacterial Symbiont

Bacterial isolation from the *Aptos suberitoides* spongy tissue was grown on Tryptic Soy Agar-Sea Water (TSA-SW) medium, selected by colour, size and colony shape. Based on the colony morphology, though seven (7) bacterial isolates were planted on TSA-SW media, only six (6) isolates were grown (Table 1). The six bacterial isolates were purified with the same medium for the identification process.

Identification of Bacterial Symbiont

The six isolate symbiont bacteria were then conventionally identified, including morphological and biochemical tests (Table 2). The identification results of each isolate of the symbiotic bacteria were compared with identification manuals, Cowan and Steel's Manual for the Identification of Medical Bacteria (Cowan & Steel, 1965) and Bergey's Manual of Determinative Bacteriology (Whitman *et al.*, 2012) to determine the species of bacteria.

Table 1. Colony morphology of bacterial symbionts on *Aptos suberitoides*

No.	isolate code	color	elevate	Edge	form	size (mm)	isolate growth (+/-)
1.	SC1	white	Convex	Flat	rounded	1.5	+
2.	SC2	cream	Convex	Flat	rounded	1.0	-
3.	SC3	cream	Convex	Flat	amorf	6.0	+
4.	SC4	cream	Convex	Flat	amorf	2.0	+
5.	SC5	white	Convex	Flat	rounded	1.0	+
6.	SC6	white	Convex	Flat	rounded	2.0	+
7.	SC7	cream	Convex	Flat	rounded	1.0	+

DISCUSSION

Bacteria that are found and symbiotic with sponges have different characteristics. The *Corynebacterium hofmanni* bacterium has another name; *Corynebacterium pseudodiphtheriticum* (Cowan & Steel, 1965; Whitman *et al.*, 2012). *Corynebacterium hofmanni* bacteria are normal bacteria in the human nasopharynx (Whitehouse *et al.*, 2018). The bacteria of this genus are found to live in seawater (Mudryk, 1998), and one of the species of this genus; *Corynebacterium maris*, is found in *Fungia granulosa* coral

mucus (Ben-Dov *et al.*, 2009). *Corynebacterium hofmanni* bacteria rarely cause infections, but they can cause endocarditis, urinary tract infection (UTIs) and may infect skin lesions (Whitehouse *et al.*, 2018).

Table 2. Morphological and biochemical tests of symbiotic sponge bacteria

Parameter	Isolate					
	SC1	SC3	SC4	SC5	SC6	SC7
Gram Characteristic						
Bacterial form	coccus ^{SA}	coccus ^{SA}	coccus ^{SA}	coccus ^{SA}	coccus ^{SA}	coccus ^{SA}
Colour/Gram	+ ^{SA}	- ^{SA}	- ^{SA}	- ^{SA}	+ ^{SA}	- ^{SA}
Biochemistry Test						
TSI Agar	K/K	A/A	A/A	K/K	A/A	K/K
Gas	-	-	-	-	-	-
H ₂ S	-	-	-	-	-	-
Catalase	+ ^S	+ ^S	+ ^S	+ ^{SA}	+ ^{SA}	+ ^S
Oxidase	- ^S	+ ^{SA}	+ ^{SA}	+ ^{SA}	-	- ^{SA}
O/F	NR ^S	F ^S	F	NR ^S	NR	NR ^S
Nitrate Reduction	+ ^{SA}	- ^{SA}	- ^{SA}	- ^{SA}	+ ^d	W ^{SA}
Gelatin	- ^S	- ^A	- ^A	- ^{SA}	- ^{Ad}	- ^S
Motility	- ^S	+ ^{SA}	+ ^{SA}	- ^{SA}	+ ^{SA}	- ^{SA}
Indole	-	- ^{SA}	- ^{SA}	- ^A	- ^S	-
Simmons Citrate	+ ^S	+ ^A	+ ^A	- ^d	- ^d	- ^{SA}
Malonate	-	- ^A	- ^A	-	-	-
Christine's Urease	+ ^{SA}	- ^{SA}	- ^{SA}	- ^{SA}	+ ^{SA}	- ^{SA}
Methyl Red (MR)	- ^A	-	-	-	-	-
Voges Proskauer (VP)	- ^S	+ ^{SA}	+ ^{SA}	-	- ^{dd}	-
Lysine Decarboxylase	-	- ^{SA}	- ^{SA}	+ ^{SA}	- ^{SA}	+ ^A
Ornithine Decarboxylase	+ ^S	-	- ^{SA}	+ ^S	+ ^A	+ ^S
Phenylalanine Deaminase	-	- ^A	- ^A	-	-	-
Aesculin Hydrolysis	- ^{SA}	+ ^A	+ ^A	-	- ^{Ad}	-
Carbohydrate Fermentation						
Glucose	- ^S	+ ^{SA}	+ ^{SA}	- ^S	+ ^{SA}	- ^S
Lactose	- ^S	- ^A	- ^A	-	+ ^{Ad}	-
Sucrose	- ^S	+ ^{SA}	+ ^{SA}	-	+ ^{Ad}	-
Maltose	- ^S	+ ^A	+ ^A	- ^S	+ ^A	- ^S
Mannitol	- ^S	+ ^A	+ ^A	-	- ^A	-
Dulcitol	-	- ^A	- ^A	-	- ^A	-
Inositol	-	- ^A	- ^A	-	-	-
Sorbitol	-	- ^A	- ^A	-	- ^{Ad}	-
Arabinose	-	- ^{SA}	- ^{SA}	-	- ^{Ad}	-
Raffinose	-	- ^A	- ^A	-	- ^{S+Ad}	-
Xylose	-	- ^{SA}	- ^{SA}	-	- ^{SA}	-
Genus	<i>Corynebacterium</i>	<i>Vibrio</i>	<i>Vibrio</i>	<i>Oligella</i>	<i>Bacillus</i>	<i>Bordetella</i>
Species	<i>C. hofmannii</i>	<i>V. damsela</i>	<i>V. damsela</i>	<i>O. uretralis</i>	<i>B. coagulans</i>	<i>B. Parapertussis</i>

Remark: (+) = Positive; (-) = Negative; A = Acid; K = Alkaline; NR = Non Reaction; F = Fermentative; W = weak;

d = 16-84% strain positive

14

The bacterium *Vibrio damsela* or now known as *Photobacterium damsela* subsp. *damsela* (Rivas *et al.*, 2011) can be found in some damselfish sea fish, yellowtail, seabream, and brown shark. Another report explained that these bacteria are also common in lemon sharks, dolphins, turtles, octopus, uninfected fish and wounds in

humans. These bacteria can cause ulcers in *Chromis punctipinnis* fish and cause infection in human wounds (Brenner *et al.*, 2005). Bacteria *Photobacterium damsela* has been isolated from the marine sponges, family Geodiidae and Halichondriidae (Sfanos *et al.*, 2005). Other photobacterium bacterial species known to be symbiotic with marine sponges are known as *Photobacterium rosenbergii* (Thompson *et al.*, 2005), *Photobacterium jeanii* (Chimetto *et al.*, 2010) and *Photobacterium phosphoreum* (Sfanos *et al.*, 2005). In addition, *Vibrio* sp. bacteria are present in the water of white shrimp (*Litopenaeus vannamei*). If the abundance of bacteria *Vibrio* sp. in white shrimp exceed the minimum threshold of bacteria in the waters of 10^4 CFU/ml, it becomes susceptible to the attack of Vibriosis (Kharisma & Manan, 2012).

Oligella urethralis, formerly known as *Moraxella urethralis* (Brenner *et al.*, 2005), is a bacterium that is generally susceptible to most antibiotics. It was observed that this bacterium is susceptible to penicillin (Welch *et al.*, 1983). *Oligella urethralis* is widespread (Rossau *et al.*, 1987) but it is primarily isolated from the human urogenital tract and has been reported to cause urosepsis (Pugliese *et al.*, 1993). Some *Oligella urethralis* are also isolated from ear, blood and leg wounds (Rossau *et al.*, 1987; Brenner *et al.*, 2005). These bacteria have also been isolated from rabbit conjunctiva (Marini *et al.*, 1996). This genus, *Oligella ureolytica*, has been isolated from Gokceada-Turkey island in the coastal and offshore areas (Türetken & Altuğ, 2016).

The *Bacillus coagulans* has been found in corneal infections, bacteremia and cow abortions. These bacteria are also found in compost, milk, paper, cardboard and silage (De Vos & Garrity, 2009). They are also found in gemstones (Khan *et al.*, 2001). In addition, they were isolated from Gokceada-Turkey Island in the coastal and offshore areas (Türetken & Altuğ, 2016). *Bacillus coagulans* bacteria have an essential role in food spoilage as a producer of commercially valuable products such as lactic acid, thermostable enzymes, and antimicrobial peptides coagulin, and are used as probiotics (De Vos & Garrity, 2009). This bacterium was known to play a role in degrading chitin (Clements *et al.*, 2002). Several *Bacillus* species have been isolated from marine sponges; namely, *Bacillus benzoevorans*, *Bacillus cereus*, *Bacillus firmus*, *Bacillus gibsonii*, *Bacillus methanolicus*, *Bacillus niacini*, *Bacillus pumilus*, *Bacillus* sp., *Bacillus anthracis* and *Bacillus vietnamensis* (Sfanos *et al.*, 2005). Furthermore, *Bordetella parapertusis* bacteria have been found in the human respiratory tract (Wolfe *et al.*, 2005) and sheep (Martin, 1996). One species of this genus *Bordetella petrii* isolated from the sea sponges of the family Coelosphaeridae and Pseudoceratinidae (Sfanos *et al.*, 2005). *Bordetella parapertusis* bacteria can cause mild pertussis in humans and cause chronic progressive pneumonia in sheep (Brenner *et al.*, 2005).

The acquired bacterial symbionts can perform several processes in the body of the sponge. Collectively the microbes in the sponge can perform many processes, such as photosynthesis, methane oxidation, nitrification, nitrogen fixation, sulfate reduction, and dehalogenation (Taylor *et al.*, 2007). Bacterial symbionts are also involved in various

processes of sponge metabolism, such as vitamin production, nutrient transport, utilisation, redox sensing and response (Thomas *et al.*, 2010).

The data obtained showed that the mechanisms of these bacteria in the sponges body are vertical. Vertical transmission is transmitted through sponge gametes through oocytes or sponge larvae (Aknin *et al.*, 2010). The vertical transmission process of the sponge begins with the inclusion of the symbiotic bacteria present in mesohyl into the oocyte by phagocytosis. During the process of the division of the embryo, bacteria exist between blastomeres or are found in vacuoles. In the blastula, all bacteria are removed in the blastocoele (Ereskovsky *et al.*, 2005). The presence of bacterial cells in the central cavity of the embryo (blastocoele) is consistent with the description of bacteria present in the larval cavity (Sharp *et al.*, 2007).

Based on the sponge properties of the filter feeder, there is a possibility of horizontal transmission. Horizontal transmission is the bacterial selective absorption process of bacterial diversity in the surrounding water column that passes through the sponge during the filter feeder process (Taylor *et al.*, 2007). Known bacteria found in coastal waters of Pasir Putih Situbondo are *Pseudomonas cepacia*, *Bacillus subtilis* and *Vibrio alginolyticus* (Andriyono *et al.*, 2015). *Pseudomonas cepacia* or *Burkholderia cepacia* was found in *Aplysina fulva* marine sponges in coastal waters of Brazil (Hardoim *et al.*, 2009). *Bacillus subtilis* bacteria have also been isolated from marine sponges *Haliclona simulans* (Phelan *et al.*, 2013) and *Fasciospongia cavernosa* (Pandey *et al.*, 2014). The *Vibrio alginolitycus* is known as symbiotic bacteria with a sea sponge *Algelas* sp. and *Spongia* sp. (Hassanzadeh *et al.*, 2014). However, the current study did not find the three species of bacteria on the body of the *Aaptos suberitoides* sponge as a symbiont. This condition is possible due to the selection process in the body of the sponge during the filter feeder process. Sponges have an effective defence system against microbes and parasites involving bacterial engulfment into specialised cells, beside the fact that sponges also use signal pathways that actively kill bacteria transduction. The archaeocyte cell in the sponge body can be thought of as the macrophage of the sponge, where it is used for self-defense against foreign microorganisms (Müller & Müller, 2003).

CONCLUSION

This study has obtained isolates that are symbiotic with *Aaptos suberitoides*. Symbiotic bacteria with *Aaptos suberitoides* spongewere isolated and identified as *Corynebacterium hofmanni*, *Vibrio damsela*, *Oligella urethralis*, *Bacillus coagulant* and *Bordetella parapertusis*. The bacteria can not be used as bioremediation agents because most bacteria are pathogenic to humans and fish, except *Bacillus coagulant*. However, further research is required to determine the effect of bacteria as a probiotic agent candidate on bioremediation.

REFERENCES

- Abubakar, H.; Wahyudi, A. T. and Yuhana, M.** (2011). Skrining bakteri yang berasosiasi dengan spons *Jaspis* sp. sebagai penghasil senyawa antimikroba. *ILMU KELAUTAN: Indonesian Journal of Marine Sciences*, 16(1) : 35-40.
- Aknin, M.; Gros, E.; Vacelet, J.; Kashman, Y. and Gauvin-Bialecki, A.** (2010). Sterols from the Madagascar Sponge *Fascaplysinopsis* sp. *Marine drugs*, 8(12) :2961-2975.
- Andriyono, S.; Jalasena, B.; Tjahtjaningsih, W. and Pramono, H.** (2015). Characterisation of Symbiotic Bacteria Isolated from Sponge *Haliclona* sp. *Exploration and Conservation of Biodiversity*, The First International Conference On Life Sciences And Biotechnology 2015 :110-115.
- Ben-Dov, E.; Yosef, D. Z. B.; Pavlov, V. and Kushmaro, A.** (2009). *Corynebacterium* maris sp. nov., a marine bacterium isolated from the mucus of the coral *Fungia granulosa*. *International journal of systematic and evolutionary microbiology*, 59(10) :2458-2463.
- Brenner, D. J.; Krieg, N.; Staley, J. and Garrity, G.** (2005). *Bergey's manual of systematic bacteriology*, Vol 2: The Proteobacteria. NY: Springer.
- Chasanah, E.; Patantis, G.; Dewi, A. S.; Marraskuranto, E.; Januar, H. I.; Stella, S.; Soka, S. and Yogiara, Y** (2013). Analysis of Bacterial Community Associated with *Aaptos* sp. from Rote and Seribu Islands. *Microbiology Indonesia*, 7(1) :5-5.
- Chimetto, L. A.; Cleenwerck, I.; Thompson, C. C.; Brocchi, M.; Willems, A.; De Vos, P. and Thompson, F. L.** (2010). *Photobacterium jeanii* sp. nov., isolated from corals and zoanthids. *International journal of systematic and evolutionary microbiology*, 60(12) :2843-2848.
- Cicirelli, E. M.** (2007). *Bacterial quorum-sensing in the marine sponge environment: Implications on motility and flagellar biosynthesis*. Indiana University.
- Clements, L. D.; Miller, B. S. and Streips, U. N.** (2002). Comparative growth analysis of the facultative anaerobes *Bacillus subtilis*, *Bacillus licheniformis*, and *Escherichia coli*. *Systematic and applied microbiology*, 25(2) :284-286.
- Cowan, S. T., and Steel, K. J.** (1965). Manual for the identification of medical bacteria. *Manual for the Identification of Medical Bacteria*.
- De Vos, P. and Garrity, G. M.** (2009). *Bergey's manual of systematic bacteriology*: Springer.
- Ereskovsky, A. V.; Gonoboleva, E. and Vishnyakov, A.** (2005). Morphological evidence for vertical transmission of symbiotic bacteria in the viviparous sponge *Halisarca dujardini* Johnston (Porifera, Demospongiae, Halisarcida). *Marine Biology*, 146(5) :869-875.
- Hanif, N.; Murni, A.; Tanaka, C. and Tanaka, J.** (2019). Marine natural products from Indonesian waters. *Marine drugs*, 17(364) :1-69.

- Hardoim, C.; Costa, R.; Araujo, F.; Hajdu, E.; Peixoto, R.; Lins, U.; Rosado A, S. and Van Elsas, D.J.** (2009). Diversity of bacteria in the marine sponge *Aplysina fulva* in Brazilian coastal waters. *Applied and environmental microbiology*, 75(10): 3331-3343.
- Hassanzadeh, Y.; Bahador, N. and Baseri Salehi, M.** (2014). Isolation and Identification of symbiotic *Vibrio alginolyticus* from Persian Gulf sponges with API 20NE kit. *Journal of Marine Biology*, 6(3) :61-68.
- Khan, M.; Saha, M. and Afroz, H.** (2001). Microorganisms associated with gemstones. *Bangladesh Journal of Botany*, 30(2) :93-96.
- Kharisma, A. and Manan, A.** (2012). Kelimpahan Bakteri *Vibrio* sp. pada air pembesaran udang vannamei (*Litopenaeus vannamei*) sebagai deteksi dini serangan penyakit vibriosis. *Jurnal Ilmiah Perikanan dan Kelautan*, 4(2) :128-134.
- Marini, R. P.; Foltz, C. J.; Kersten, D.; Batchelder, M.; Kaser, W. and Li, X.** (1996). Microbiologic, radiographic, and anatomic study of the nasolacrimal duct apparatus in the rabbit (*Oryctolagus cuniculus*). *Laboratory animal science*, 46(6) :656-662.
- Martin, A.P.** (2002). Phylogenetic approaches for describing and comparing the diversity of microbial communities. *Applied and Environmental Microbiology* 68 (8) :3673-3682.
- Martin, W.** (1996). Respiratory infections of sheep. *Comparative immunology, microbiology and infectious diseases*, 19(3) :171-179.
- Mudryk, Z.** (1998). Generic composition and metabolic activity of bacteria inhabiting surface seawater layers. *Oceanological Studies*, 27(3) :57-70.
- Müller, W. E. and Müller, I. M.** (2003). Origin of the metazoan immune system: identification of the molecules and their functions in sponges. *Integrative and comparative biology*, 43(2) :281-292.
- Murniasih, T.; Kosela, S.; Kardono, L. B.; Hanafi, M. and Suwarso, W. P.** (2014). An antibacterial compound isolated from sponge-associated bacteria *Rhodobacter cicea* bacterium. *Journal of Medical Sciences (Faisalabad)*, 14(2) :75-80.
- Nofiani, R.; Nurbetty, S., and Sapar, A.** (2009). Aktivitas Antimikroba Ekstrak Metanol Bakteri Berasosiasi Spons Dari Pulau Lemukutan, Kalimantan Barat Antimicrobial Activities Of Methanol Extract From Unidentified Sponge Associated Bacteria In Lemukutan Island, Kalimantan Barat. *E-Jurnal Ilmu dan Teknologi Kelautan Tropis*, 1(2):33-41.
- Pandey, S.; Sree, A.; Sethi, D. P.; Kumar, C. G.; Kakollu, S.; Chowdhury, L. and Dash, S. S.** (2014). A marine sponge associated strain of *Bacillus subtilis* and other marine bacteria can produce anticholinesterase compounds. *Microbial cell factories*, 13(1) :1-9.

-
- Phelan, R. W.; Barret, M.; Cotter, P. D.; O'Connor, P. M.; Chen, R.; Morrissey, J. P.; Dobson, A.D.W.; O'Gara, F. and Barbosa, T. M.** (2013). Subtilomycin: a new lantibiotic from *Bacillus subtilis* strain MMA7 isolated from the marine sponge *Haliclona simulans*. *Marine drugs*, 11(6) :1878-1898.
- Pugliese, A.; Pacris, B.; Schoch, P. E. and Cunha, B. A.** (1993). *Oligella urethralis* urosepsis. *Clinical infectious diseases*, 17(6) :1069-1070.
- Radjasa, O. K.; Kencana, D. S.; Sabdono, A.; Hutagalung, R. A. and Lestari, E. S.** (2007). Antibacterial activity of marine bacteria associated with sponge *Aaptos* sp. against Multi Drugs Resistant (MDR) strains. *Jurnal Matematika dan sains*, 12(4):147-152.
- Rivas, A. J.; Balado, M.; Lemos, M. L. and Osorio, C. R.** (2011). The *Photobacterium damsela* subsp. *damsela* hemolysins *damselysin* and *HlyA* are encoded within a new virulence plasmid. *Infection and Immunity*, 79(11) :4617-4627.
- Rossau, R.; Kersters, K.; Falsen, E.; Jantzen, E.; Segers, P.; Union, A.; Nehls, L. and De Ley, J.** (1987). *Oligella*, a new genus including *Oligella urethralis* comb. nov.(formerly *Moraxella urethralis*) and *Oligella ureolytica* sp. nov.(formerly CDC group IVe): relationship to *Taylorella equigenitalis* and related taxa. *International journal of systematic and evolutionary microbiology*, 37(3) :198-210.
- Rua, C. P.; Trindade-Silva, A. E.; Appolinario, L. R.; Venas, T. M.; Garcia, G. D.; Carvalho, L. S.; Lima, A.; Kruger, R.; Pereira, R.C.; Berlinck, R.G.S.; Valle, R, A.B.; Thompson, C.C. and Thompson, F** (2014). Diversity and antimicrobial potential of culturable heterotrophic bacteria associated with the endemic marine sponge *Arenosclera brasiliensis*. *PeerJ*, 2, e419.
- Setyati, W. A. and Subagiyo, S.** (2012). Isolasi dan Seleksi Bakteri Penghasil Enzim Ekstraseluler (proteolitik, amilolitik, lipolitik dan selulolitik) yang Berasal dari Sedimen Kawasan Mangrove (Isolation and Selection of Extracellular Enzyme Producing Bacteria Originating from Mangrove Sedimen. *ILMU KELAUTAN: Indonesian Journal of Marine Sciences*, 17(3) :164-169.
- Sfanos, K.; Harmody, D.; Dang, P.; Ledger, A.; Pomponi, S.; McCarthy, P. and Lopez, J.** (2005). A molecular systematic survey of cultured microbial associates of deep-water marine invertebrates. *Systematic and applied microbiology*, 28(3):242-264.
- Sharp, K. H.; Eam, B.; Faulkner, D. J. and Haygood, M. G.** (2007). Vertical transmission of diverse microbes in the tropical sponge *Corticium* sp. *Applied and environmental microbiology*, 73(2):622-629.
- Subagio, I. B. and Aunurohim, A.** (2013). Struktur Komunitas Spons Laut (Porifera) di Pantai Pasir Putih, Situbondo. *Jurnal Sains dan Seni ITS*, 2(2) :159-165.

- Taylor, M. W.; Radax, R.; Steger, D. and Wagner, M.** (2007). Sponge-associated microorganisms: evolution, ecology, and biotechnological potential. *Microbiology and molecular biology reviews*, 71(2) :295-347.
- Thomas, T., Rusch, D., DeMaere, M. Z., Yung, P. Y., Lewis, M., Halpern, A., Heidelberg, K.B., Egan, S., Steinberg, P. D., Kjelleberg, S.** (2010). Functional genomic signatures of sponge bacteria reveal unique and shared features of symbiosis. *The ISME journal*, 4(12) :1557-1567.
- Thompson, F.L; Thompson, C.C; Naser, S.; Hoste, B.; Vandemeulebroecke, K.; Munn, C.; Bourne, D. and Swings, J.** (2005). *Photobacterium rosenbergii* sp. nov. and *Enterovibrio corallii* sp. nov., vibrios associated with coral bleaching. *International journal of systematic and evolutionary microbiology*, 55(2), :913-917.
- Türetken, P. S. Ç. and Altuğ, G.** (2016). Bacterial pollution, activity and heterotrophic diversity of the northern part of the Aegean Sea, Turkey. *Environmental monitoring and assessment*, 188(122) :1-12.
- Welch, W.; Porschen, R. and Luttrell, B.** (1983). Minimal inhibitory concentrations of 19 antimicrobial agents for 96 clinical isolates of group IVe bacteria. *Antimicrobial agents and chemotherapy*, 24(3) :432-433.
- Whitehouse, C. A.; Zhao, S., and Tate, H.** (2018). Antimicrobial resistance in *Campylobacter* species: mechanisms and genomic epidemiology. *Advances in applied microbiology*, 103 :1-47.
- Whitman, W. B.; Goodfellow, M.; Kämpfer, P.; Busse, H.-J.; Trujillo, M. E.; Ludwig, W. and Suzuki, K.-i.** (2012). *Bergey's manual of systematic bacteriology: Volume 5: The Actinobacteria*: Springer New York.
- Wolfe, D. N.; Kirimanjesswara, G. S. and Harvill, E. T.** (2005). Clearance of *Bordetella parapertussis* from the lower respiratory tract requires humoral and cellular immunity. *Infection and Immunity*, 73(10) :6508-6513.

Marine Sponge suberitidae 2021

ORIGINALITY REPORT

6%

SIMILARITY INDEX

0%

INTERNET SOURCES

6%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Rosa Amalia, Diah Ayuningrum, Agus Sabdono, Ocky Karna Radjasa. "Antipathogenic Activity of Acroporid Bacterial Symbionts Against Brown Band Disease-Associated Bacteria", Squalen Bulletin of Marine and Fisheries Postharvest and Biotechnology, 2021
Publication 1%
- 2** "Encyclopedia of Marine Biotechnology", Wiley, 2020
Publication <1%
- 3** "Natural Products from Actinomycetes", Springer Science and Business Media LLC, 2022
Publication <1%
- 4** "Symbiotic Microbiomes of Coral Reefs Sponges and Corals", Springer Science and Business Media LLC, 2019
Publication <1%
- 5** A.M. Labella, J.J. Rosado, M. Balado, M.L. Lemos, J.J. Borrego. " Virulence properties of <1%

three new species affecting cultured fish ",
Journal of Applied Microbiology, 2019

Publication

6

Ana Rita Moura Polónia, Daniel Francis
Richard Cleary, Francisco José Riso da Costa
Coelho, Leontine E. Becking et al.

"Compositional analysis of archaeal
communities in high and low microbial
abundance sponges in the Misool coral reef
system, Indonesia", Marine Biology Research,
2018

Publication

<1 %

7

Avi Bittan. "Asymptotic performance of a
buffer model in a data sharing environment",
ACM SIGMETRICS Performance Evaluation
Review, 5/1/1994

Publication

<1 %

8

Awanis Azizan, Andrea C Alfaro, Diana
Jaramillo, Leonie Venter et al. " Pathogenicity
and virulence of bacterial strains associated
with summer mortality in marine mussels () ",
FEMS Microbiology Ecology, 2022

Publication

<1 %

9

Detmer Sipkema, Sònia de Caralt, Jose A.
Morillo, Waleed Abu Al-Soud, Søren J.
Sørensen, Hauke Smidt, María J. Uriz. "Similar
sponge-associated bacteria can be acquired

<1 %

via both vertical and horizontal transmission",
Environmental Microbiology, 2015

Publication

10

Goodfellow, Michael. "The Family
Nocardiaceae", The Prokaryotes, 2014.

Publication

11

Hardoim, Cristiane C. P., and Rodrigo Costa.
"Temporal dynamics of prokaryotic
communities in the marine sponge
Sarcotragus spinosulus", Molecular Ecology,
2014.

Publication

12

Sulav Indra Paul, Md. Mahbubur Rahman,
Mohammad Abdus Salam, Md. Arifur Rahman
Khan, Md. Tofazzal Islam. "Identification of
marine sponge-associated bacteria of the
Saint Martin's island of the Bay of Bengal
emphasizing on the prevention of motile
Aeromonas septicemia in Labeo rohita",
Aquaculture, 2021

Publication

13

Alexander V. Ereskovsky, Elizaveta
Gonobobleva, Andrey Vishnyakov.
"Morphological evidence for vertical
transmission of symbiotic bacteria in the
viviparous sponge Halisarca dujardini
Johnston (Porifera, Demospongiae,
Halisarcida)", Marine Biology, 2004

Publication

<1 %

<1 %

<1 %

<1 %

14

M.L. Lemos, M. Balado. "Iron uptake mechanisms as key virulence factors in bacterial fish pathogens", *Journal of Applied Microbiology*, 2020

Publication

<1 %

15

Tian, Ren-Mao, Jin Sun, Lin Cai, Wei-Peng Zhang, Guo-Wei Zhou, Jian-Wen Qiu, and Pei-Yuan Qian. "The deep-sea glass sponge *Lophophysema eversa* harbors potential symbionts responsible for the nutrient conversions of carbon, nitrogen and sulfur : Potential symbionts in a glass sponge", *Environmental Microbiology*, 2015.

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Marine Sponge suberitidae 2021

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10
