

ISSN-0976-0245 (Print) • ISSN-0976-5506 (Electronic)

**Volume 11 / Number 03 / March 2020**



# Indian Journal of Public Health Research & Development

An International Journal

Website:  
[www.ijphrd.com](http://www.ijphrd.com)

## **XVIII**

231. Development of Doctors Attire According to Patients Preferences ..... 1238  
*Rizki Fadillah, Rizma Adlia Syakurah, Riana Sari Puspita Rasyid*
232. Lower Abdominal Pain During Office Hysteroscopy as a Guide in the Assessment of Tubal Patency in Infertile Women..... 1243  
*Sally Elattar, Ahmed Maged, Moutaz Elsherbini*
233. Overview of Smoking History in Lung Cancer Patients at RSUD Dr. Soetomo, Surabaya ..... 1249  
*Santi Martini, Kurnia Dwi Artanti, Sri Widati, Dessy Arumsari*
234. Radiographic Evaluation of Two Different Designs for Treatment of Mandibular Bilateral Free End Saddle with Osseo-integrated Implants ..... 1255  
*Seham Sabra, GehanF ekry, Emad Agamy*
235. Adaptive Physical Education in Rehabilitation after Injuries Maxillofacial Area ..... 1261  
*Sergey Petrovich Evseev, Olga Eduardovna Evseeva, Andrey Vladimirovich Aksenov*
236. Role of Autophagy in the Myelodysplastic Syndrome and Myeloproliferative Disorders ..... 1266  
*Shaimaa F. Kamel, Mohamed E. Abd El-Fattah, Hisham A. Aly, Hisham M. Tawfik, Elwy M. Soliman, Khaled M. Salah*
237. The Influence of Reward and Communication to Cohesion Team of Tuberculosis Prevention Team in Community Health Centers ..... 1274  
*Sinta Dewi Lestyoningrum*
238. The Effect of Burnout on Employee Satisfaction in 9 Local Health Centers in Surabaya City..... 1279  
*Siti Fatimah Hariyati, Ernawaty*
239. The Influence of Social and Cultural Factors on the Decision of Selecting a Place for Preeclampsia Mother Giving Birth ..... 1285  
*Siti Nur Azizah, Ernawaty*



# Indian Journal of Public Health Research & Development

An International Journal

Home

Editorial Team

Article Submission

Subscriptions Informations

Publication Ethics

Issues

Contact Us



## Indian Journal of Public Health Research & Development

An International Journal

### EDITORIAL PANEL



#### Editor

Vidya Surwade

Associate Professor, Dr Baba Saheb Ambedkar, Medical College & Hospital Rohinee, Delhi

#### Associate Editor

Prof. (Dr.) Anil Gupta

Vice principal, Professor and Head Department of Biochemistry, Eklavya Dental College and Hospital, Kotputli, Affiliated with Rajasthan University of Health Sciences, Jaipur.

[editor.ijphrd@gmail.com](mailto:editor.ijphrd@gmail.com)

#### Editorial (International Board )

- Dr Abdul Rashid Khan B. Md Jagar Din, Associate Professor, Department of Public Health Medicine, Penang Medical College, Penang , Malaysia
- Dr Luca Testarelli, Associate Professor at Sapienza University of Rome. Associate Professor of Endodontics and Restorative Dentistry
- Kamal Kant Sahu Internal Medicine Residency Saint Vincent Hospital, 123 Summer street, Worcester, 01608, Massachusetts, United States
- Dr V Kumar, Consulting Physician, Mount View Hospital, Las Vegas , USA
- Basheer A. Al-Sum , Botany and Microbiology Deptt , College of Science, King Saud University, Riyadh, Saudi Arabia
- Dr Ch Vijay Kumar, Associate Professor, Public Health and Community Medicine, University of Buraimi, Oman
- Dr VMC Ramaswamy, Senior Lecturer, Department of Pathology, International Medical University, Bukit Jalil, Kuala Lumpur
- Kartavya J. Vyas , Clinical Researcher, Department of Deployment Health Research, Naval Health Research Center, San Diego, CA (USA)
- Prof PK Pokharel, Community Medicine, BP Koirala Institute of Health Sciences, Nepal
- Sajjad Salim Issa AL-Musawi supervisor for student of the Arab board in family medicine

- Prof. Dr. Ayad F. Alkaim Professor of surface chemistry and applications of nano-materials, Babylon University, Iraq
- Prof. Dr. Imad Hadi Hameed Biomedical Science Department, College of Nursing, University of Babylon, Hillah city, Iraq
- Prof. Dr. Amean A Yasir. Department of Community , College of Nursing, University of Babylon, Hillah city, Iraq
- Dr. Sadeq Al-Murshedi, Lecturer. University of AL Zahraa for Women, Karbala, Iraq.
- Dr. Rajaa Ahmed Mahmoud University of Basrah, Al Zahraa College of Medicine, Iraq

#### **Editorial (National Board )**

- Prof. Sushanta Kumar Mishra, Community Medicine, GSL Medical College – Rajahmundry, Karnataka
- Prof D.K.Srivastava, Medical Biochemistry, Jamia Hamdard Medical College, New Delhi
- Dr Sunil Mehra, Paediatrician & Executive Director, MAMTA Health Institute of Mother & Child, New Delhi
- Prof M.Sriharibabu , General Medicine , GSL Medical College , Rajahmundry , Andhra Pradesh
- Prof Pankaj Datta, Principal & Prosthodontist, Indraprastha Dental College , Ghaziabad
- Prof Samarendra Mahapatro, Pediatrician, Hi-Tech Medical College, Bhubaneswar, Orissa
- Dr Abhiruchi Galhotra, Additional Professor, Community and Family Medicine, AII India Institute of Medical Sciences, Raipur
- Prof Deepti Pruthvi, Pathologist, SS Institute of Medical Sciences & Research Center. Davangere, Karnataka
- Prof G S Meena, Director Professor, Maulana Azad Medical College, New Delhi
- Prof Pradeep Khanna, Community Medicine, Post Graduate Institute of Medical Sciences, Rohtak , Haryana
- Dr Shailendra Handu Associate professor, Phrma , DM (Pharma, PGI Chandigarh)
- Dr. A.C. Dhariwal Directorate of National Vector Borne Disease Control Programme, Dte. DGHS, Ministry of Health Services, Govt. of India, Delhi.

#### **Scientific Committee**

- Dr Anju Ade, Professor, Community Medicine Department, SVIMS, Sri Padamavati Medical College, Tirupati. Andhra Pradesh.
- Dr.E.Venkata Rao, Associate Professor, Community Medicine, Institute of Medical Sciences & SUM Hospital, Bhubaneswar, Orissa.
- Dr. Amit K. Singh, Associate Professor, Community Medicine, VCSG Govt. Medical College, Srinagar – Garhwal, Uttarakhand
- Dr R G Viveki, Prof n head, Dept of Community Medicine, BIMS Belgavi
- Dr. Santosh Kumar Mulage, Assistant Professor, Anatomy, Raichur Institute of Medical Sciences Raichur(RIMS) , Karnataka
- Dr Gouri Ku. Padhy, Associate Professor, Community and Family Medicine, AII India Institute of Medical Sciences, Raipur
- Dr Ritu Goyal, Associate Professor, Anaesthesia , Sarswathi Institute of Medical Sciences, Panchsheel Nagar
- Dr. Anand Kalaskar, Associate Professor, Microbiology Prathima Institute of Medical Sciences , AP
- Dr.Md.Amirul Hassan , Associate Professor, Community Medicine Government Medical College, Ambedkar nagar, UP
- Dr.N.Girish, Associate Professor, Microbiology, VIMS&RC, Bangalore
- Dr BR Hungund, Associate Professor Pathology, JNMC ,Belgaum.
- Dr Sartaj Ahmad PhD Medical Sociology Associate Professor Swami Vivekananda Subharti University Meerut UP India

- Dr Manoj Kumar Tripathi (Associate Professor), Political Science and public administration, Swami Vivekananda Subharti University, Meerut, Uttar Pradesh, India.
  - Dr Sumeeta Soni, Associate Professor, Microbiology Department, B.J. Medical College, Ahmedabad ,Gujarat,India
- 

**OUR CONTACT INFO:****Institute of Medico-Legal Publications**

Logix Office Tower, Unit No. 1704, Logix City Centre Mall

Sector- 32, Noida - 201 301 (Uttar Pradesh)

Email: [editor.ijphrd@gmail.com](mailto:editor.ijphrd@gmail.com)

Website: [www.ijphrd.com](http://www.ijphrd.com)

**Visit our Website:** [www.imlp.in](http://www.imlp.in)

---

# Overview of Smoking History in Lung Cancer Patients at RSUD Dr. Soetomo, Surabaya

Santi Martini<sup>1</sup>, Kurnia Dwi Artanti<sup>1</sup>, Sri Widati<sup>2</sup>, Dessy Arumsari<sup>1</sup>

<sup>1</sup>Department of Epidemiology, <sup>2</sup>Department of Health Promotion and Behavior,  
Public Health Faculty, Universitas Airlangga

## Abstract

**Introduction:** Lung cancer belongs to the top ten diseases which caused death in the world. World Health Organization (WHO) estimated that there is about 2,09 million cases of lung cancer and 1,76 million death accident due to this disease. Smoking is defined as the most factor that can promote the lung cancer about 80% cases in male and 50% cases in female.

**Aims:** To identify the history of smoking activity in the lung cancer patients in the RSUD Dr. Soetomo, Surabaya.

**Method:** Research was determined as descriptive research with cross-sectional design. Variable in this research smoking status, age on the first time of smoking, duration of smoking, kind of smoke, and smoking level. Lung cancer patients in the RSUD Dr. Soetomo, Surabaya was addressed as subject in this research. Accidental sampling was performed as sampling method to the 53 respondents. Data was collected by distribution of questionnaire which contained risk factor of lung cancer. Data was analyzed by univariate analysis.

**Result:** Result saw that about 31 out of 53 respondents had history of smoking. Most of them had experienced as active smoker (22 respondents, 41,5%), had experience as first-time smoker on the age less than 15 years old (25 respondents, 47,20%), had been smoked for more than 20 years (29 respondents, 54,70%), have preferred to consume smoke with filter (21 responden, 39,60%), and considered as heavy level of smoker (15 respondents, 28,30%).

**Conclusion:** Most of lung cancer patient in the RSUD Dr. Soetomo, Surabaya had history as smoker.

**Keywords:** *Epidemiology, history of smoking, lung cancer, tobacco.*

## Introduction

World Health Organization (WHO) estimates 9.6 million people worldwide deaths caused cancer<sup>(1)</sup>. Lung cancer is the highest cause of death in men in 2018. The

incident Lung cancer amount 22,440 cases Or 14% of all cancer cases<sup>(2)</sup>. The main risk factor for lung cancer is smoking. In general, smoking causes 80% of lung cancer cases in men and 50% of lung cancer cases in women<sup>(3)</sup>.

The Result from Minister of Health Survey 2018 showed that proportion of tobacco consumption in the population aged 15 years and over were 62,9% in men and 33,8% for all. The prevalence of smoking in aged 10-18 years increased from 8.8 in 2016 to 9.1 in 2018<sup>(4)</sup>.

According to Sudoyo<sup>(5)</sup> the definition of smokers is people who have smoked 1 stick or more every day for at least a year. If people were not smoking for a

---

### Corresponding Author:

**Santi Martini**

Public Health Faculty, Universitas Airlangga, Ir. H. Soekarno Street, Mulyorejo, Surabaya, East Java, 60115, Indonesia  
e-mail: santi-m@fkm.unair.ac.id

month, they called people with a history of smoking. The increased risk factors are related to the duration of smoking in a year and the factors when starting smoking (the younger the individual starts smoking, the higher risk of lung cancer). Other factors that can be considered include the types of cigarettes smoked (tar content, filter, and *kretek* cigarettes) (6).

The various research studies found that Lung cancer not only associate with active and passive smoking, but some occupational agents, and indoor or outdoor air pollution(7,8).Lung cancer has most commonly been associated with occupational exposures. Occupational exposures such as Asbestos, Vinyl chloride, Arsenic, Beryllium, Chromium and Nickel(8).

Lung cancer cases in East Java Province calculated about 34,706 cases(9). Based on the recapitulation data for the last 10 months at Dr. Soetomo in 2018, lung cancer morbidity was 733 cases. Lung cancer morbidity is still high in Dr. Hospital Soetomo, it is necessary to conduct research on the description of smoking history as the main risk factor. This research aims to identify the history of smoking activity in the lung cancer patients.

### Methodology

The research was conducted in RSUD Dr. Soetomo, Surabaya on August 2018-January 2019. The

descriptive method obtained from secondary data of research by Dr. Santi Martini, dr., M. Kes about “Risk of diseases caused by smoking”. Cross-sectional was performed in this research. The research sample was about 53 respondents by accidental sampling technic. Moreover, variable in this research were smoking status, age on the first time of smoking, duration of smoking, kind of smoke, and smoking level. However, another risk factor was also included in this research such, for instance, family history, exposure of asbestos, pollutant exposure at home, and pollutant exposure at workplace. The questionnaire of lung cancer risk was used in this research. Then, research was analyzed by univariate analysis and presented frequency distribution by epidemiological approach.

### Results

Distribution of respondents by place and time.East Java consists of 38 districts/cities which consist of 29 districts and 9 cities. Based on the results of the study, it found that the highest lung cancer cases were in Surabaya with 14 respondents (26.41%). It can be due to the location of Dr. Soetomo located in Surabaya. Figure 1. shows that the distribution of respondents based on the year diagnosed was mostly in 2018 as many as 46 respondents (86.79%).




Figure 1. Distribution frequency respondent base on year when diagnose

Distribution of respondents based on people. Characteristics of lung cancer patients based on age showed that the majority of respondents belong to the age of more than 40 years as many as 47 respondents

(88.70%). The characteristics of lung cancer patients based on sex are mostly male about 35 respondents (66.00%) like at Table 1. The characteristics of lung cancer patients based on education are mostly classified

into the basic education category as many as 29 respondents (54.70%) are elementary or equal. More than 50% of lung cancer patients have stage IV as many as 46 respondents (86.8%).

**Table 1. Distribution frequency respondent base on person**

Characteristic	Frequency	Porcentage (%)
Age >40 year	47	88,70
Men	35	66,00
Work	36	67,90
Stage IV	46	86,80
<b>Type of Education</b>		
Basic	29	54,70
Intermediate	18	34,00
High	6	11,30
<b>Total</b>	<b>53</b>	<b>100,0</b>

Distribution of respondents based on smoking history. Based on table 3, the data showed that most lung cancer patients have a history of active smoking as many as 22 respondents (41.50%). Most respondents have a history of smoking by consuming filter cigarettes as many as 21 respondents (39.6%). Most of respondents first time smoking at the age of 15-55 years as many as 25 respondents (47.2%), most respondents had a long history of smoking more than 20 years as many as 29 respondents (54.7%), and most smoking level of respondents is belong to heavy category namely 15 respondents (28.3%).

**Table 2. Distribution Frequency base on history of smoking**

Variable	Frequency	Porcentage (%)
<b>History of Smoking</b>		
Never Smoke	12	22,60
Active Smoker	22	41,50
Passive Smoker	10	18,90
Active and passive smoker	9	17,00
<b>Type of Cigarette</b>		
No Type (Never & Passive smoker)	22	41,50
Filter	21	39,60
Non Filter	10	18,90
<b>Initial Smoking Age</b>		
Never & Passive smoker	22	41,50
1-14 years	6	11,30
15-55 years	25	47,20

Variable	Frequency	Porcentage (%)
<b>Duration Smoking</b>		
Never & Passive smoker	22	41,50
≤20 years	2	3,80
>20 years	29	54,70
<b>Level of Smoking</b>		
Never & Passive smoker	22	41,50
Mild	4	7,50
Middle	12	22,60
Heavy	15	28,3
<b>Total</b>	<b>53</b>	<b>100,0</b>

Distribution of respondents based on other risk factors. Based on Table 2, most respondents did not have a family history of lung cancer about 51 respondents (96.2%), most of the respondents were not exposed to asbestos as many as 40 respondents (75.5%), most respondents were not exposed to pollutants in the residence as many as 49 respondents (92.5%), and the majority of respondents were not exposed to pollutants in the workplace about 44 respondents (83.0%).

**Table 3. Distribution frequency of respondent base on others factor**

Other risk factor	Frequency	Porcentage (%)
<b>History of Family</b>		
Yes	2	3,8
No	51	96,2
<b>Asbestos Exposure</b>		
Exposure	13	24,5
Not Exposure	40	75,5
<b>Pollutant exposure at home</b>		
Exposure	4	7,5
Not Exposure	49	92,5
<b>Pollutant exposure at work place</b>		
Exposure	9	17,0
Not Exposure	44	83,0
<b>Total</b>	<b>53</b>	<b>100,0</b>

### Discussion

Distribution of respondents based on people. Based on the results, it exhibited that the majority of respondents was on the age more than 40 years old. About 95% of lung cancer patients at the age more than 40 years old which conducted by Malhotra as a trend of lung cancer in Delhi, India. Moreover, data showed that most of the respondents were male<sup>(3)</sup>. This study was in line with the research conducted by Neupane<sup>(10)</sup>. It stated that most



lung cancer is suffered by men. Then, based on the results, the respondents had the type of basic education<sup>(10)</sup>. This research is in agreement with the research conducted by Ernawati et al<sup>(11)</sup> which stated that most lung cancer patients have a history of junior high school level. Based on the results, most of the respondents are worker. Furthermore, other data showed that the majority of respondents belonging to stage IV. Some Research performed<sup>(12)</sup>,<sup>(13)</sup> that the stage of lung cancer patients is found in the late stages. Lung detection is generally slow in process. This also can promote patients to have less awareness to see themselves. Most patients are referrals from regional hospitals and health centers, including some services which do not have specialist doctors or supporting tools for diagnosing lung cancer. Another thing considered as a factor is the policy to go to the health center before going to the hospital. There are many reasons for improving the quality of care, evaluation of the cost, and reluctant to do the examination.

Distribution of respondents based on history of smoking. Based on the results, it found that most respondents had a history of active smoking. This research is in line with the research conducted by Hulma, Basyar, and Mulyani<sup>(14)</sup> found that smoking history in most detected in lung cancer patients. Based on the results of the study, it was found that the majority of respondents consumed filter cigarettes. This research is supported by research conducted by Herlina, Rahmalia, and Dewi<sup>(15)</sup>. They stated that lung cancer patients who have a history of smoking more often consume filter cigarettes. Other study found that Smoker have Risk more than 5 times death rate higher than No Smoker for Asian people<sup>(16)</sup>.

Based on the results of the study it was found that the majority of respondents who had a history of smoking, first smoked at the age of less than 15 years with 25 respondents (47.20%). This research is in line with the research conducted by Putera et al<sup>(17)</sup> states that most of lung cancer patients smoke at the age of 17-25 years old. This study is in line with the research conducted by Setyanda et al<sup>(18)</sup> that the average person has a long history of smoking with more than 20 years. The British Doctors' Study found that the relative risk (RR) of smoking for lung cancer have increased over the 20-year, 40-year, and 50-year follow up<sup>(16)</sup>. Based on the results of the study, it was found that most of the respondents were classified as heavy smoking level. This study is similar with Naser, Medison, and Erly<sup>(19)</sup>, that

the highest number of smoking is in the level of heavy smoking. The number of lung cancer patients who have a history of heavy smoking consuming large quantities of cigarettes per day has a greater risk of lung cancer.

Distribution of respondents based on other risk factors. Based on the results of the study, it found that most respondents did not have a family history of lung cancer. This research is in line with the research conducted by Ernawati et al<sup>(11)</sup> stated that most of lung cancer patients do not have a family with a history of cancer. The research by Yoshida et al<sup>(20)</sup> stated that a family history of lung cancer in the close relatives was associated with an increased risk of lung cancer between the sexes. Families who have cancer will increase the risk of lung cancer by 13.8%. First sisters allow a greater risk than brothers<sup>(21)</sup>.

This study was found that the majority of respondents were not exposed to asbestos. This can be seen from the top use (platonik) of occupied houses made of abscesses. Continuous exposure to asbestos and increased amount of exposure will increase the risk of lung cancer. Smokers exposed to asbestos have a higher risk of lung cancer<sup>(22)</sup>. Based on the results of the study, it was found that more than 80% of respondents were not exposed to pollutants in their homes. The pollutants came from the use of wood fuel stoves, the use of kerosene fuel stoves, and mosquito coils. Continuous exposure to the smoke will increase the risk of lung cancer<sup>(23)</sup>.

## Conclusion

Most of lung cancer patient in the RSUD Dr. Soetomo, Surabaya had history as smoker.

**Acknowledgments:** We would like to thank all participants in this study including the patient, patient's family, nurse and doctor. We would like to a big thank you to Dr. Soetomo General Hospital for allowing this research to be carried out.

**Conflict of Interest:** The authors have no conflicts.

**Source of Funding:** It is research and publication of this article funded by Ministry of Research and technology and higher education through Universitas Airlangga 2018.

**Ethical Clearance:** Received from Health Research Ethics Committee, Faculty of Public Health, Universitas Airlangga.

## References

1. World Health Organization. Latest global cancer data: Cancer burden rises to 18.1 million new cases and 9.6 million cancer deaths in 2018 [Internet]. Geneva: World Health Organization; 2018 [cited 2019 Jan 2]. Available from: <https://www.who.int/cancer/PRGlobocanFinal.pdf>
2. Globocan. Indonesia [Internet]. Geneva; 2018. Available from: <https://gco.iarc.fr/today/data/factsheets/populations/360-indonesia-fact-sheets.pdf>
3. Malhotra RK, Manoharan N, Nair O, Deo S, Rath GK. Trends in lung cancer incidence in Delhi, India 1988-2012: age-period-cohort and joinpoint analyses. *Asian Pacific J Cancer Prev.* 2012;19(6):1647–54.
4. Ministry of Health RI. Basic health research 2018. Jakarta; 2018.
5. Sudoyo A., Setiyohadi B, Alwi I, Simadibarata M, Setiati S. Internal medicine textbook jilid II. 5th ed. Jakarta: Internal Publishing; 2009. 1973-1982 p.
6. Muttaqin A. Client nursing care with immunological system disorder textbook. Jakarta: Salemba Medika; 2008.
7. Samet JM. Environmental causes of lung cancer: what do we know in 2003? *Chest.* 2004;125(5): 80-3.
8. Shankar A, Dubey A, Saini D, Singh M, Prasad CP, Roy S, et al. Environmental and occupational determinants of lung cancer. *Transl Lung Cancer Res* [Internet]. 2019;8(1):31–49. Available from: <http://tldr.amegroups.com/article/view/28014/21200>
9. Ministry of Health RI. Basic health research 2013. Jakarta; 2013. K. Available from: <http://www.depkes.go.id/resources/download/general/HasilRiskasdas2013.pdf>
10. Neupane PR, Poudel KK, Huang ZB, Steel R, Poudel JK. Distribution of cancer by sex and site in Nepal. *Asian Pacific J Cancer Prev.* 2017;18(6):1611–5.
11. Ernawati Y, Ermayanti S, Herman D, Russilawati. Risk factors for lung cancer in women treated in the lung section of RSUP Dr. M. Djamil Padang and RSUD Solok: penelitian case control. *J Kesehatan Andalas* [Internet]. 2019;8(2):1–8. Available from: <http://jurnal.fk.unand.ac.id/index.php/jka/article/view/951>
12. Tatun R, Icksan AG, Syahrudin E, Kekalih A. Thoracic CT scan according to cytology/histology type in lung cancer patients smoker. *eJKI.* 2015;3(3):177–85.
13. Siregar YS, Widyaningsih W, Syahrudin E. Body mass index, percentage of skeletal muscle and albumin in non small cell carcinoma lung cancer patients before and after chemotherapy. *J Respirologi Indones.* 2016;36(2):73–82.
14. Hulma MA, Basyar M, Mulyani H. Relationship between characteristics of patients with cytopathology in cases of pulmonary carcinoma treated at RSUP Dr. M. Djamil Padang. *J Kesehatan Andalas.* 2014;3(2):196–201.
15. Herlina, Rahmalisa S, Dewi YI. Relationship to smoking history with stage lung Ca . *J Online Mhs Bid Ilmu Keperawatan.* 2014;1(1):1–6.
16. Jung KJ, Jeon C, Jee SH. The effect of smoking on lung cancer: ethnic differences and the smoking paradox. *Epidemiol Health.* 2016;38(e2016060):1–6.
17. Putera PJC, Kalesaran AFC, Sekeon SAS. The relationship between smoking status and exposure to cigarette smoke with quality of life in residents in Kinilow village, north Tomohon subdistrict, Tomohon city.. *Media Kesehatan.* 2017;9(3):1–10.
18. Setyanda YOG, Sulastri D, Lestari Y. Relationship between smoking and incidence of hypertension in men aged 35 - 65 years in Padang city. *J Kesehatan Andalas.* 2015;4(2):434–40.
19. Naser F, Medison I, Erly. Description of the degree of smoking in patient with COPD in the lung at RSUP Dr. M. Djamil. *J Kesehatan Andalas.* 2016;5(2):306–11.
20. Yoshida K, Takizawa Y, Nishino Y, Takahashi S, Kanemura S, Omori J, et al. Association between family history of cancer and lung cancer risk among Japanese Men and Women. *Tohoku J Exp Med.* 2019;247(2):99–110.
21. Lin H, Huang Y-S, Yan H, Yang X-N, Zhong W-Z, Ye H-W, et al. A family history of cancer and lung cancer risk in never - smokers : a clinic - based case-control study. *Lung Cancer.* 2015;89(2):94-8.

22. Gilham C, Rake C, Burdett G, Nicholson AG, Davison L, Franchini A, et al. Pleural mesothelioma and lung cancer risks in relation to occupational history and asbestos lung burden. *Occup Environ Med.* 2016;73(5):290–9.
23. Raspanti GA, Hashibe M, Siwakoti B, Wei M, Thakur BK, Pun CB, et al. Household air pollution and lung cancer risk among never-smokers in Nepal. *Environ Res [Internet].* 2016;147(6):141–5. Available from: <http://dx.doi.org/10.1016/j.envres.2016.02.008>