

Effect of L-Carnosine in Patients with Age-Related Diseases: A Systematic Review and Meta- Analysis

by Chrismawan Ardianto

Submission date: 21-Feb-2023 04:41PM (UTC+0800)

Submission ID: 2019507684

File name: h_Age-Related_Diseases_A_Systematic_Review_and_Meta-Analysis.pdf (4.02M)

Word count: 13981

Character count: 77758

Systematic Review

Effect of L-Carnosine in Patients with Age-Related Diseases: A Systematic Review and Meta-Analysis

Kaoshik Sureshkumar¹, Mahesh Durairaj¹, Kaviya Srinivasan¹, Khang Wen Goh^{2,*}, Krishna Undela³, Vijayakumar Thangavel Mahalingam^{4,*}, Chrismawan Ardianto^{5,*}, Long Chiau Ming^{5,6,7}, Rajanandh Muhasaparur Ganesan^{8,*}

¹ Department of Pharmacy Practice, Sri Ramachandra Faculty of Pharmacy, Sri Ramachandra Institute of Higher Education and Research, 600116 Chennai, Tamil Nadu, India

² Faculty of Data Science and Information Technology, INTI International University, 71800 Nilai, Malaysia

³ Department of Pharmacy Practice, National Institute of Pharmaceutical Education and Research (NIPER) Guwahati, Changsari, 781101 Kamrup (R), Assam, India

⁴ Department of Pharmacy Practice, SRM College of Pharmacy, SRM Institute of Science and Technology, 603203 Chennai, Tamil Nadu, India

⁵ Department of Pharmacy Practice, Faculty of Pharmacy, Universitas Airlangga, 60115 Surabaya, Indonesia

⁶ School of Medical and Life Sciences, Sunway University, Bandar Sunway, 47500 Selangor, Malaysia

⁷ PAP Rashidah Sa'adatul Bolkliah Institute of Health Sciences, Universiti Brunei Darussalam, Jalan Tungku Link, Gadong BE1410, Brunei

⁸ Saveetha Medical College and Hospital, Saveetha Institute of Medical and Technical Sciences, Thandalam, 602105 Chennai, Tamil Nadu, India

*Correspondence: khangwen.goh@newinti.edu.my (Khang Wen Goh); vijaypractice@yahoo.com (Vijayakumar Thangavel Mahalingam);

chrismawan-a@ff.unair.ac.id (Chrismawan Ardianto); rajanandh.phd@gmail.com (Rajanandh Muhasaparur Ganesan)

Academic Editors: Leonid A. Gavrilov and Thomas Heinbockel

Submitted: 8 August 2022 Revised: 5 December 2022 Accepted: 8 December 2022 Published: 18 January 2023

Abstract

Introduction: L-carnosine has been found to have multimodal activity. **Aim:** The aim of this review was to find out the efficacy of L-carnosine in patients with age-related diseases. **Methods:** Clinical studies evaluated the effect of L-carnosine on cancer, cardiovascular disease, diabetes, and neurodegenerative disorders were searched in electronic bibliographic databases. The protocol has been registered with PROSPERO (CRD42022314033). The revised Cochrane risk of bias tool for randomized trials was used to assess all of the reports for risk of bias. RevMan 5.4 was used to conduct the meta-analysis. **Results:** Following the screening process, 14 papers were selected for systematic review, with 9 of them being qualified for meta-analysis. Many of the included studies showed that L-carnosine has potential therapeutic activity in age related diseases. Results from the meta-analysis showed that in diabetes mellitus, HbA1c [mean difference (MD) 95% CI = -1.25 (-2.49, -0.022); $p = 0.05$; $p = 0.001$; $I^2 = 85\%$] and fasting blood sugar (FBS) [MD 95% CI = -12.44 (-22.44, -2.44); $p = 0.01$; $p = 0.40$; $I^2 = 0\%$] and in neurodegenerative disorder, Wechsler Memory Scale Logical Memory 2 (WMS-LM2) [MD 95% CI = 1.34 (0.83, 1.85); $p < 0.00001$; $p = 0.43$; $I^2 = 0\%$], showed statistically significant difference, favoring the L-carnosine group over the control group. While in neurodegenerative disorder, Alzheimer's Disease Assessment Scale (ADAS) [MD 95% CI = 0.98 (-1.55, -0.42); $p = 0.0007$; $p = 0.86$; $I^2 = 0\%$] and Back Depression Inventory (BDI) [MD 95% CI = -1.12 (-1.87, -0.37); $p = 0.003$; $p = 0.73$; $I^2 = 0\%$] showed statistically significant difference, favoring the control group over L-carnosine group. **Conclusions:** Clinical studies were conducted to manage chemotherapy induced toxicities and there are no clinical studies available for its anti-cancer use, and the current evidence does not support its use in the treatment of cardiovascular disease.

Keywords: Alzheimer's disease; amino acid; cancer; diabetes; cardiovascular diseases; neurological diseases; 2-(3-aminopropanoylamino)-3-(3H-imidazol-4-yl) propanoic acid; polaprezinc

1. Introduction

Gulevitch discovered carnosine (β -alanine-L-histidine) as an endogenous and water-soluble dipeptide in Russia in 1900 [1]. Carnosine is produced by the enzyme carnosine synthase from L-histidine and beta-alanine, and it is degraded by the enzyme carnosinase. Carnosine levels in the human body are affected by gender (males have higher levels), age (carnosine concentration decreases with age), and food (vegetarian diet have a lower concentration of carnosine in the skeletal muscles) [2,3]. It has various health properties, including anti-oxidant, anti-glycating, ion-chelating, pH buffering, anti-inflammatory, and

anti-aging properties [4].

Aging is defined as the irreversible loss of physiological function and ageing processes are described as those that enhance an individual's susceptibility to conditions that eventually lead to death as they grow older. It's a complicated multi-factorial process in which numerous components interact at the same time and work at several levels of functional organisation. Aging leads to age-related diseases like cancer, cardiovascular disease, diabetes, and neurological disorders.



1.1 Role of L-Carnosine in Cancer

Toxicity from chemotherapy is a common and unfortunate side effect of treatment that can occur even at standard doses. Oral mucositis is a common consequence in patients who have undergone high-dose chemotherapy and after hematopoietic stem cell transplantation (HSCT). Oral mucositis is frequently accompanied by pain, odynophagia, xerostomia, and dysgeusia, and it can lead to malnutrition, dehydration, and lowers the patients' quality of life significantly (QoL) [5]. Polaprezinc (PZ), a chelated form of zinc and L-carnosine, inhibits the production of inflammatory cytokines such as tumour necrosis factor- α (TNF- α), interleukin-1 (IL-1), interleukin-6 (IL-6) and matrix metalloproteinase-2 (MMP-2) and induces an anti-inflammatory effect and immune responses via inhibition of nuclear factor-kappa-light-chain-enhancer of activated B cells (NF- κ B) [5,6].

Chemotherapy-induced peripheral neuropathy (CIPN) is a typical side effect of neurotoxic chemotherapeutic drugs used to treat cancer patients. It is characterized by inflammation, damage, or degeneration of the peripheral nerve fibres. As a commonly used third-generation platinum analogue, oxaliplatin is responsible for more than 70% of symptomatic neurotoxicity of any severity, which frequently leads to treatment discontinuation [7,8].

The oxidative stress, inflammatory, and apoptotic pathways are the responsible pathophysiological factors for Oxaliplatin-induced peripheral neuropathy (OPIN). The inflammatory cascade includes the up-regulation of the transcriptional factor NF- κ B and its downstream molecule TNF- α , while the redox pathway is associated with the elevation of malondialdehyde and the suppression of nuclear factor-2 erythroid related factor-2 (Nrf-2). Furthermore, elevated apoptotic markers such as caspase-3 can be used to assess the apoptotic signalling pathway [9,10]. By scavenging free radicals and suppressing the inflammatory cascade, L-carnosine has an antioxidant and anti-inflammation effect on OPIN pathways that have been widely studied [11].

1.2 Role of L-Carnosine in Cardiovascular Diseases

Pro-inflammatory cytokines, such as IL-6, and acute-phase proteins, such as C-reactive protein (CRP) are elevated in individuals with myocardial infarction (MI). The role of inflammation in the onset of MI is regarded to be significant [12]. L-carnosine has been shown to reduce the production of IL-6, and thereby it helps in the management of MI [13,14].

Despite breakthroughs in pharmacological therapy and the introduction of novel devices, patients with heart failure continue to have a poor QoL and a lower exercise tolerance. Several factors contribute to the decline in fractional capacity and exercise tolerance, including changes in endothelial and vasodilatory function, abnormalities in skeletal muscle metabolism, and reduction in muscle blood flow during exercise [15]. L-carnosine is a key antioxidant

that protects against oxidative damage caused by anaerobic exercise. Lactic acid production increases during exercise, followed by dissociation into lactate and H⁺, lowering pH levels [16]. L-carnosine has an excellent buffering activity because the imidazole group of histidine binds a proton and lowers the pH value. As carnosine's pKa is closer to that of physiological pH, it could be used as a buffer more quickly during exercise [17,18]. Increased L-carnosine levels in the muscles may help to improve contractility and reduce fatigue [19]. Other mechanisms include regulating myocardial calcium levels, increasing the sensitivity of calcium-release channels and the contractile machinery, and improving cardiac contractility and function [20,21].

1.3 Role of L-Carnosine in Diabetes Mellitus

Type 2 diabetes mellitus (T2DM) accounts for 90% of all diabetes cases. Insulin resistance is a condition in which the body does not fully respond to insulin, causing blood glucose levels to rise. Obesity, hypertension, and dyslipidemia are all common co-morbidities of T2DM [22]. Advanced glycation end products (AGEs) are a group of heterogeneous compounds that form in greater quantities in hyperglycemic conditions. The formation of covalent cross-links with proteins, increased oxidative stress, and up-regulation of inflammation are biological processes for AGEs' detrimental effects. Increased reactive oxygen species formation and NF- κ B targets-gene transcription is caused by AGEs interacting with the receptor for advanced glycation end products (RAGE), resulting in low-grade inflammation, endothelial dysfunction, platelet activation, and insulin resistance [23]. The identification of AGEs and their binding to RAGE contribute to diabetes' micro- and macro-vascular complications [24]. L-carnosine has antioxidant properties and the ability to protect against the production of AGEs, making it useful in the treatment of T2DM [25–27].

Hyperglycemia causes aberrant glucose-dependent pathway activation, i.e., the polyol pathway, hexosamine pathway and protein kinase C pathway. Multiple components, such as transforming growth factor-beta (TGF- β), vascular endothelial growth factor (VEGF), IL-1, IL-6, and TNF, are produced by these pathways [28,29]. Increased urine TGF- β activates both the canonical (ALK 5, Smad 2/3) and alternative (ALK 1, Smad 1/5) pathways in diabetic individuals. Extracellular matrix accumulates at the glomerular basement membrane (GBM) and mesangium when the canonical pathway is activated. In addition, activation of the alternative pathway induces podocyte damage, which results in the effacement of the foot process. As a result, TGF- β and metabolic pathway activation are essential determinants in the development of diabetic nephropathy [30,31]. One of the proposed mechanisms is that L-carnosine inhibits the synthesis of TGF- β [32].

Skin complications are common in diabetic patients, with over 30% of individuals developing some form of skin

Table 1. Selection criteria of articles for systematic review and meta-analysis.

Category	Inclusion criteria	Exclusion criteria
Population	Patients with age-related diseases such as cancer, cardiovascular disease, diabetes, and neurodegenerative disorders, any gender, ethnicity, geographic area, comorbidities, or medication use	Studies involving pediatrics or animal sciences
Intervention	L-carnosine supplemented in any dose, any dosage form, either alone or in combination with other intervention/s, of any dosage and for any duration	Studies without L-carnosine
Comparator	Placebo, usual care, before-after/pre-post comparison, or other pharmacological or non-pharmacological interventions	The comparison of only one group without considering before and after
Outcome	Effect of L-carnosine for the management of age-related diseases such as cancer, cardiovascular disease, diabetes, and neurodegenerative disorders	Studies without well-defined outcome parameter
Study design	Randomized Controlled Trials	Non-randomized controlled trials

involvement during the course of their condition [33]. Xerosis of the skin is found in up to 40% of diabetic patients [34]. In T2DM, topical use of urea 5% combined with arginine and carnosine increases skin hydration and alleviates skin dryness [33].

1.4 Role of L-Carnosine in Neurodegenerative Disorders

Individuals aged 55 and up will soon make up a quarter or more of the total population of schizophrenia patients globally. Glutamate dysfunction and dysregulation of the *N*-methyl-d-aspartate (NMDA) receptor (glutamatergic receptor) are thought to be linked to the pathogenesis of cognitive deficits of schizophrenia [35,36]. L-carnosine is a known antioxidant that is co-localized at glutamatergic synapses, and it helps people with schizophrenia with their deficient cellular antioxidant defences [37–39]. Furthermore, L-carnosine has been shown to modulate NMDA receptor activity, lower glutamate levels, and limit glutamate release [40,41].

Alzheimer's disease (AD) is the most common type of dementia, accounting for 60 to 70% of all cases. According to Hipkiss's [42] study, certain areas of the brain are the most sensitive to damage associated with AD when carnosine levels decline with age. Furthermore, heavy metal ions can be found in the brains of people with AD, which is defined by the presence of a pathogenic form of amyloid beta protein and intracellular lesions made up primarily of the cytoskeletal protein tau [43]. L-carnosine binds to metal ions and inhibits amyloid beta formation while also acting as a "scavenger" of free radicals [44].

Given this background, we aimed to find out the efficacy of L-carnosine versus placebo, or usual treatment, or other pharmacological/non-pharmacological interventions in patients with age-related diseases such as cancer, cardiovascular disease, diabetes, and neurodegenerative disorders. We, therefore, conducted a systematic review and meta-analysis of clinical studies, which will raise awareness of the clinical usage of L-carnosine for the aforementioned disorders among healthcare professionals, researchers, patients, and the general public.

2. Methods

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines were followed in this systematic review [45]. The protocol for this

systematic review was registered into PROSPERO, an international database of prospectively registered systematic reviews in health and social care, with the registration number CRD42022314033.

2.1 Selection Criteria

The population, intervention, comparison, outcomes, and study design (PICOS) approach was used to rate studies for eligibility [46]. Table 1 shows the PICOS criteria for inclusion and exclusion. Papers published before 2011 and articles written in languages other than English were excluded from the study.

2.2 Search Strategy

Before defining our research question, we ran a preliminary search in PROSPERO to see if there were any current systematic reviews or meta-analyses on the same topic. PubMed, Science Direct, Proquest, Scopus, Cochrane Library, Google Scholar, Clinical Trials Registry India (CTRI), and clinical trials.org were used by two investigators (KSur and KSri) to perform an independent literature search. Various combinations and phrases of major Medical Subject Heading (MeSH) terms and non-MeSH terms, such as L-carnosine, Carnosine, β -alanine-l-histidine, L-carnosine, randomized controlled trial (RCT), ageing, cancer, cardiovascular diseases, diabetes, and neurodegenerative disorders were used in search strategies.

The EndNote reference manager was used to remove duplicates, and a final duplicate removal was done. Disagreements over the study selection were settled through discussion among the reviewers. To validate eligibility, the full texts of the selected publications were extracted and independently examined by the three reviewers (KSri, KSur, MD). In the literature search, lists of references to important articles were also checked for intelligibility. Any discrepancies were handled through consensus; with the help of a third author (KU).

2.3 Study Selection

One reviewer (KSur) conducted the primary literature search. The identified papers' titles, abstracts, keywords, authors, author affiliations, year of publication, volume, issue, and DOI were exported to Microsoft Office Excel ver. 2019 (Redmond, Washington, United States). Furthermore, two reviewers (RMG and VTM) independently examined




Fig. 1. PRISMA flow chart illustrating the selection process for all included and excluded studies.

the titles and abstracts of the retrieved records to find studies that would fit the aforementioned inclusion criteria. Full-text screening was performed on an article with confusing qualifying criteria. The authors of the papers that might have an impact on the review but whose full text couldn't be found were contacted via email and asked for the full-text article. Following that, two review team members (RMG and VTM) independently reviewed the complete texts of these possibly eligible papers for eligibility. Any differences between the RMG and VTM over the eligibility of certain studies were resolved by discussion and consensus with a third reviewer (KU). After a month, the searches were conducted again.

2.4 Data Extraction

In a specially designed data extraction form, two independent investigators (KSri and KSur) extracted key data from all of the included publications. A discussion with the third independent investigator (MD) addressed the discrepancy amongst the reviewers, and data accuracy was validated. For each article, we extracted data on authors, year of publication, a number of participants, age, gender, history of diseases, comorbidities, clinical parameters, route of administration, dose and therapeutic regimen of the intervention and control groups, study follow-up time, ran-

domization, concealment of allocation, blindness scheme, presence of an intention-to-treat analysis, loss of follow-up, early termination, expected results and results.

2.5 Assessment of Methodological Quality

Two independent reviewers (MD and KSur) assessed the risk of bias (RoB) in the included papers using the revised Cochrane tool to assess the risk of bias, consisting of five domains: randomization procedure, deviation from intended intervention, missing outcome data, measurement of the outcome and selection of the reported result [47]. Finally, any discrepancies were resolved through discussion until an agreement was reached or by consulting a third reviewer (KS).

2.6 Statistical Analysis

Among the studies that reported similar outcomes, the meta-analysis was conducted using RevMan 5.4 software (Cochrane Training, New York, NY, USA). If continuous data had baseline and endpoint scores, we looked at the change from baseline to endpoint and estimated mean difference and standard error for continuous exposure. The mean difference (MD) and 95% CI were provided as the final pooled result. A forest plot was used to show the results and study level estimations. A *p*-value of 0.05 was used to

determine statistical significance. In each of the included studies' analyses, the I^2 statistic was utilised to assess heterogeneity. Due to the insufficient number of papers, we were unable to perform a funnel plot, hence publication bias could not be detected. When the study's heterogeneity (I^2) was less than 50%, a fixed-effects model was used for all analyses, and when the study's heterogeneity was greater than 50%, a random-effects model was employed.

3. Results

Cohen's kappa statistics were used to verify the inter-rater reliability of the reviewers. The Kappa test was used to determine the level of agreement and likelihood of inconsistency among the reviewers. Between MD and KSur, a Kappa test was conducted, yielding a value of kappa = 0.654 (standard error (SE) = 0.171; 95% CI: 0.321–1.000; weighted kappa = 0.774), indicating substantial agreement. The third reviewer (KSri) did not participate in the title, abstract, or full-text screening, but did analyses and explained inconsistencies between MD and KSur in the quality assessment. However, kappa tests were conducted across all four reviewers to confirm their consistency, revealing high agreement. RMG and KSri had a Kappa score of 0.641 (SE of kappa = 0.148; 95% CI: 0.421–0.994; weighted kappa = 0.499), while RMG and KSri had a Kappa test score of 0.632 (SE of kappa = 0.157; 95% CI: 0.263–0.969; weighted kappa = 0.625).

3.1 Study Search and Selection

An electronic database search yielded a total of 116 studies. The titles and abstracts of 105 potentially relevant articles were evaluated after duplicate studies were removed. Based on title and abstract screening, 86 articles were eliminated. The full-text screening included the remaining 21 studies. As shown in the PRISMA flow diagram (Fig. 1), 5 studies were excluded for the following reasons: unavailable data, inappropriate study design, and unrelated outcomes. At the end of the selection process, 14 studies [5,22,33,48–58] meeting the inclusion criteria were eventually selected for systematic review and meta-analysis. The PRISMA checklist is presented as a **Supplementary File**.

3.2 Study Characteristics

Table 2 (Ref. [5,22,33,48–58]) and **Supplementary Table 1** list the basic characteristics of the studies that were included. These studies were published between 2011 and 2022. Studies in neurodegenerative disorders were found in 43% (6 out of 14) of the included studies [48–51,53,56]. Diabetes mellitus was found in 29% (4 out of 14) of the studies [22,33,52,57]. Two of the remaining four studies were on cancer [5,55], while the other two were on cardiovascular diseases [54,58]. Both genders were included in the majority of the studies. In 8 of the 14 studies, researchers looked at safety measures and reported on adverse drug reactions [5,33,49,50,52,54,56,57]. Five of the studies included were

from Japan [5,51,53,56,58], two from Iran [22,50] and Italy [33,54] and one from Poland [48], USA [49], Greece [52], Egypt [55], and Thailand [57].

3.3 Intervention Characteristics

All of the studies are RCT that provide clinical evidence for the efficacy of L-carnosine as an individual supplement or in combination with other supplements. Of 14 studies, 4 studies used anserine as an add-on supplement with L-carnosine in neurodegenerative disorders [48,51,53,56], 2 studies used zinc as an additional supplement with L-carnosine [5,58], and 6 studies did not use any supplements with L-carnosine [22,49,50,54,55,57]. Federici *et al.* [33], study used urea and arginine as supplements along with L-carnosine. Karkabounas *et al.* [52], study used α -Linolenic acid (ALA) and thiamine as a supplement to L-carnosine for diabetes mellitus. As an intervention group regimen for L-carnosine supplementation, one of the included studies used chicken meat extract (CME) containing 40% carnosine related compounds (CRC) [48]. The dose and mode of administration, on the other hand, differed significantly between the included studies.

3.4 Outcome

Across the studies that were considered, a wide range of outcome measures were observed. L-carnosine has been shown in cancer studies to minimize treatment-related toxicities [5,55]. In addition, Magdy *et al.* [55], examined neuropathy grade and rated it using the National Cancer Institute Common Terminology Criteria for Adverse Events-version 4 (NCI-CTCAE-v4) system, revealing that L-carnosine improved OIPN by reducing the pathogenic triad of inflammation, oxidative stress, and apoptosis. The Kitagawa *et al.* [5], study looked at the incidence of Grade ≥ 3 oral mucositis and concluded that the PZ lozenge could be helpful in preventing oral mucositis associated with chemotherapy in HSCT patients.

Lombardi *et al.* [54], study had the primary outcome measure to evaluate the exercise parameters [distance at 6-minute walk test (6-MWT)] and cardiopulmonary exercise test and the effects on QoL [EuroQoL-5D (EQ-5D) and visual analogue scale (VAS)] among congestive heart failure (CHF) patients and concluded that L-carnosine added to conventional therapy may improve exercise performance and QoL. The zinc concentration, cardiac enzymes, and the level of inflammation markers in blood and urine samples were analysed in the Yoshikawa *et al.* [58], study, and it was found that it improves cardiac function post-acute myocardial infarction (AMI), and that this effect may be due to anti-inflammatory processes through the suppression of IL-6. Ejection fraction was a common outcome in cardiovascular diseases between these 2 studies [54,58].

Federici *et al.* [33], study examined the evolution of a 9-point grading xerosis assessment scale (XAS) score and a 4-point grading overall cutaneous score (OCS) score in

Table 2. Characteristics of the included studies.

Author, year	Age		Gender		Drug regimen		L-carnosine Dose & route		Duration		Other supplements		Outcome measures		Result	Conclusion	ROB
	Control	Intervention	Control (M/FM)	Intervention (M/FM)	Control	Intervention	Control	Intervention	Primary	Secondary	ADR						
Studies in cancer patients																	
Maibly R. <i>et al.</i> (2018) [53]	45.9 ± 8.6	45.6 ± 10.5	13/18	16/14	FLOFOX-6	Control drug regimen plus L-carnosine	500 mg/day, orally	Patients were assessed on baseline and 3 months after treatment	Nil	Neuropathy grade assessment according to NCI-CTCAE-v4 and analysis of oxidative (Nrf-2, MDA, NF-sB), /inflammation (TNF-α) /apoptotic markers (caspase-3)	Bio-chemical markers (BUN, AST, ALT, TBL, Hb, PLT and TLC)	Not mentioned	In both arms a significant correlation was only evident between TNF-α and the neuropathy grading score ($p < 0.03$); the correlation analysis was significantly positive between NF-sB and both Nrf-2 and caspase 3	L-carnosine proved to improve OPN by amelioration of the pathophysiological oxidative stress and apoptosis. These results led to the recommendation of the safe add-on therapy of carnosine to chemotherapeutic agents, especially those associated with CIPN	Low risk		
Kitagawa J <i>et al.</i> (2020) [5]	58 (29-72)*	48 (18-67)*	24/23	24/17	Ara-C/CPA ± Flu or ETP/MCNU; Ara-C/MCNU/L-PAM ± ETP; CPA, CPA/L-PAM ± ETP or Flu, CPA/BU ± ATG, Flu ± BU ± L-PAM, L-PAM/MCNU, L-PAM or BU	Control drug regimen plus PZ lozenges	18.75 mg/ four times a day, orally	Patients were assessed daily from beginning of chemotherapy until 35 days after transplantation	Zinc	Incidence of Grade ≥ 3 oral mucositis, Grade ≥ 3 oral mucositis, taste disturbance, The incidence of other non-haematological AEs, the median time of engraftment and rate of engraftment	Incidence and severity of oral mucositis, xerostomia and taste disturbance. The incidence of other non-haematological AEs, the median time of engraftment and rate of engraftment	Nausea, Vomiting, Constipation, Diarrhoea, Perioral neuropathy, Skin disorder, Febrile neutropenia, WBC decreased, Anaemia, PLT decreased	The incidence of Grade ≥ 2 but not Grade ≥ 3 oral mucositis was significantly reduced by preventive treatment with PZ lozenges. The PZ lozenge preparation had no influence on engraftment. The lozenge preparation was highly safe without any serious side effects. Therefore, PZ lozenge may be useful for prevention of oral mucositis associated with chemotherapy in patients undergoing HSCT	Low risk			
Studies in cardiovascular disease patients																	
Lombardi C <i>et al.</i> (2014) [54]	62.3 ± 9.9	61.2 ± 9.3	22/3	22/3	ACE inhibitors, ARB, β-blocker	Control drug regimen plus L-carnosine	500 mg/day orally before food at morning	Patients were evaluated at baseline and after 6 months	Nil	The changes in exercise parameters at 6-MWT and cardiopulmonary exercise test) and the effects on QoL (EQ-5D and VAS)	Echocardiographic parameters: haemalogical and biochemical assays	Nil	After 6 months of treatment with orodispersible L-carnosine was associated with an improvement in 6MWT distance (419 ± 85 to 478 ± 80 meters, $p = 0.014$). QoL significantly improved in the intervention group, compared to the control group, as shown by EQ-5D test (-0.56 ± 1.00 in L-carnosine group vs 0.40 ± 1.7 in control group; $p = 0.018$) and VAS score (11.0 ± 17.7 in L-carnosine group vs -5.8 ± 12.3 control group; $p < 0.0001$)	L-carnosine, added to conventional therapy, may improve exercise performance and QoL in ambulatory patients with stable CHF	Low risk		

Table 2. Continued.

Author, year	Age		Gender		Drug regimen		L-carnosine		Duration		Other supplements		Outcome measures		Result	Conclusion	ROB
	Control	Intervention	Control	Intervention	Control	Intervention	Control	Intervention	Control	Intervention	Primary	Secondary	ADR				
Yoshikawa F <i>et al.</i> (2019) [58]	60.2 ± 14.6	61 ± 13.1	18/8	23/1	300 mg clopidogrel and 200 mg aspirin pre-PCI. Heparin was administered in the catheterization laboratory. After PCI patients continued to take 100 mg aspirin and 75 mg clopidogrel daily until the follow-up of the study	Control: 75 mg L-carnosine Intervention: 75 mg PZ	75 mg twice a day	Patients were evaluated at baseline and after 9 months	Zinc	Zinc	Cardiac function in terms of the EF, LVDD, and LVDS, and echocardiography was performed upon admission to the hospital and at 9 months post-AMI of PZ	Not mentioned	The mean IL-6-maximal CPK level was significantly reduced in the PZ group (0.024 [0.003–0.066] vs. 0.076 [0.015–0.212]; $p = 0.045$). A significant increase was detected in the ejection fraction of the PZ group at the 2 time points (54 [51–57] vs. 62 [55–71], respectively; $p < 0.01$)	PZ was reported to improve cardiac function post-AMI. This effect may be due to anti-inflammatory processes through the suppression of IL-6.	Low risk		
Studies in diabetes mellitus patients																	
Federici A <i>et al.</i> (2015) [33]	61 ± 10	62 ± 9	12/13	8/17	SEC	UC	4.5 (0.01%) finger-tip units (2.5-g of cream). Both treatments were applied on the feet (dorsal and plantar regions) and on the distal third of the legs, twice daily	Patients were evaluated at baseline and after 4, 12, and 32 weeks for primary end point and at baseline and after 32 weeks for secondary end point	Urea and (0.5%)	9-point XAS score from 0 to 8) and of a 4-point grading OCS (from 0 to 3)	skin hydration and desquamation	Nil	UC induced greater hydration than SEC ($p = 0.001$) with a 91% reduction at week 32 in XAS score vs. baseline. After 4 weeks, compared with the SEC treated group, the XAS score in the UC treated group was significantly lower. OCS was reduced by 27% from baseline to end of the study in the UC group, and increased by 8% in the SEC group ($p = 0.02$; between groups)	UC based cream increases skin hydration and alleviates the condition of severe skin dryness in T2DM patients in comparison with a control SEC product, with a greater efficacy observed as early as 4 weeks into treatment. In addition, this approach is effective in improving significantly severe foot xerosis and fissuring	Low risk		
Hojjehghani S <i>et al.</i> (2017) [22]	40.4 ± 5.1	43.0 ± 7.6	10/13	12/9	Microcrystalline cellulose with anti-diabetic medications	Control: 500 mg L-carnosine Intervention: 500 mg L-carnosine	Two capsules of 500 mg (1 g/day) after each meal	Patients were evaluated at baseline and after 12 weeks	Nil	FBS, TC, HDL, LDL, HbA1c, fasting insulin, HOMA-IR, and TG, serum levels of penicillamine, CML, sRAGE, TNF- α , IL-18 and IL-6	Not mentioned	L-carnosine supplementation resulted in significant decrease in fat mass and an increase in fat free mass, better control of FBS, HbA1c, and decreased levels of TG, CML and TNF- α levels with almost no meaningful effect on HOMA-IR, TC, LDL, HDL, serum sRAGE, IL-6 and IL-18	L-carnosine resulted in a decrease of fat mass and an increase in fat free mass, better control of FBS, HbA1c, and decreased levels of TG, CML and TNF- α levels with almost no meaningful effect on HOMA-IR, TC, LDL, HDL, serum sRAGE, IL-6 and IL-18	Low risk			

Table 2. Continued.

Author, year	Age		Gender		Drug regimen		L-carnosine		Duration	Other supplements		Outcome measures		ROB	
	Control	Intervention	Control	Intervention	Control	Intervention	Control	Intervention		Primary	Secondary	ADR	Result		Conclusion
Karkabounas S <i>et al.</i> (2018) [52]	57 ± 5	57 ± 5	35/47	(M/FM)	Placebo	ALA, carnosine, thiamine a day orally	6 mg/kg three times a day orally	2 × 500 mg taken after breakfast and dinner	Patients were evaluated after overnight fast before and after the 8 weeks supplementation	ALA (7 mg/kg) and thiamine (1 mg/kg)	Glycaemic profile, Lipid profile, Kidney function, Liver damage, Thyroid function, Cancer markers, Electrolytes, Minerals and Oxidative Stress, ADP, PAF, Arachidonic acid, Epinephrine, Thrombin and Collagen	Nil	Glucose and HbA1c levels were significantly reduced after supplementation (135.7 ± 9.5 mg/dL vs. 126.5 ± 16.8 mg/dL and 8.3% ± 0.3% vs. 6.03% ± 0.58%, respectively, $p < 0.05$); however, insulin was significantly increased from the pancreas. The supplementation had only a minor effect on insulin sensitivity since the steep increase in insulin levels was accompanied by a small but significant drop in glucose level	Low risk	
Sirivatamasit N <i>et al.</i> (2021) [57]	57.0 ± 6.9	55.6 ± 4.8	4/16	11/9	Anti-hypertensive drugs, anti-glycaemic drugs with Placebo (starch)	Control drug regimen plus L-carnosine	2 g/day orally, The dose was split into 2 × 500 mg taken after breakfast and dinner	Patients were evaluated after overnight fast before and after the 8 weeks supplementation	Nil	The change of urinary TGF-β level	The improving level of UACR	Nil	Urinary TGF-β significantly decreased with carnosine supplementation (-17.8% of the baseline values), whereas it tended to increase with placebo (+16.9% of the baseline values) (between-group difference $p < 0.05$). However, BUN, sCr, GFR and other biochemical parameters remained unchanged during the study period including urinary albuminuria	The study showed that oral carnosine supplementation could reduce urinary TGF-β level in T2DM with diabetic nephropathy, but without significant effect on urine albumin, indicating an additional reno-protective effect from conventional therapy	Low risk
Changappa KN <i>et al.</i> (2012) [49]	46.5 ± 9.0	46.6 ± 8.5	23/14	21/12	Placebo (rice powder and magnesium stearate)	L-carnosine	500 mg of L-carnosine (or placebo) 500 mg/day began during the first week, increasing to 500 mg/p.o. twice daily in the second week, 1500 mg/day in the third week and finally to 1000 mg per oral twice daily during the fourth week	Patients were assessed for cognitive domains at baseline, 4 weeks, and 12 weeks along with psychopathology and safety parameters	Nil	Psychopathology, QoL and SF-36 scales, QLI, and the SF-36 survey including the PCS and MCS summary score	(i) Cardiovascular: Increased blood pressure; (ii) Digestive: Dry mouth, Diarrhoea, Constipation, Dyspepsia and Nausea; (iii) Metabolic and nutritional: Decreased appetite, Weight loss, Weight gain and Hyperglycaemia; (iv) Dermatological: Skin rash, Dry skin and Pruritus; (v) Musculo-skeletal: Foot pain, Joint pain; (vi) Nervous: Dizziness, Sleepiness, Paraesthesia; (vii) Psychiatric: Worsening psychosis Vivid dreams; (viii) Respiratory: Pneumonia	On the strategic target detection test, the L-carnosine group displayed significantly improved strategic efficiency and made fewer perseverative errors compared with placebo. Other cognitive tests showed no significant differences between treatments.	These preliminary findings suggest that L-carnosine further consideration as adjunctive treatment to improve executive dysfunction in persons with schizophrenia	Low risk	

Table 2. Continued.

Author, year	Age		Gender		Drug regimen		L-carnosine		Duration		Other supplements		Outcome measures		Result	Conclusion	ROB
	Control	Intervention	Control	Intervention	Control	Intervention	Control	Intervention	Control	Intervention	Primary	Secondary	ADR				
Budzen S <i>et al.</i> (2014) [48]	80.5 ± 7.5	81 ± 7.0	13/13 (M/FM)	12/13 (M/FM)	Placebo (ly-40% of chicken components enzymatic treatment)	CME containing 40% of CRC meat to carnosine (2:1 ratio prepared following breast meat extraction with water (cold and at 80 °C) and spray drying)	Control: 2.5 g per day	Intervention: 2.5 g per day	Patients were assessed for cognitive and physical capacity at baseline and 13 week follow-up supplementation	Anserine	(i) The cognitive function, presence of depressive symptoms, physical capacity, body measurements, BP and HR; (ii) Behavioural measurements: The MMSE and STMS to assess cognitive functions, GDS to assess the presence of depressive symptoms, CDR to evaluate the severity of dementia	Nil	Not mentioned	The groups differed significantly at baseline according to certain clinical and physical parameters, such as HR before exercise ($p < 0.001$). The perceived exertion differed significantly ($p < 0.05$) in the CRC group between baseline and follow up results, suggesting that exercises were seen as less comprehensive. The mean value of the MMSE scores increased significantly in the CRC group ($p < 0.001$), the mean value of STMS scores showed significant increase in CRC group ($p < 0.001$). The value of MMSE correlated negatively with age. The Spearman rank correlation coefficient was ($p = -0.25$)	Anserine and carnosine supplementation in the elderly may bring promising effects on cognitive functioning and physical capacity	Low risk	
Hisatsune T <i>et al.</i> (2016) [51]	70.6 ± 5.1	67.8 ± 5.6	8/10 (M/FM)	8/11 (M/FM)	Placebo (43 mg/day L-lysine and L-histidine) (150 mg/day)	Anserine and carnosine formula (dry)	Control: 1.0 g of anserine-incarnosine (3:1) formula daily	Intervention: 1.0 g of anserine-incarnosine (3:1) formula daily	Patients were evaluated at baseline and after 3 months	Anserine	(i) A dietary survey was conducted using a semi-quantitative method as reported. Inventory of food intake during the 3-month test period; (ii) Cognitive testing: WMS-LM1, WMS-LM2, ADAScog, BDI, SF-36 MCS and SF-36 PCS; (iii) Blood samples and immunoassays: Concentrations of 27 cytokines in serum samples collected from the volunteers at the baseline and during the follow-up. (iv) Microarray analysis; (v) PBMCs from volunteers were used for total RNA extraction. The total RNA quality was assessed; (v) Acquisition of human MRI data and MRI data analysis	Nil	Not mentioned	Among the tests, delayed recall verbal memory assessed by the WMS-LM1 showed significant preservation in the ACS group compared to the placebo group ($p = 0.0128$). Blood analysis revealed a decreased secretion of inflammatory cytokines, including CCL-2 and IL-8, in the ACS group. MRI analysis using arterial spin labeling showed a suppression in the age-related decline in brain blood flow in the posterior cingulate cortex area in the ACS group, compared to the placebo group ($p = 0.0248$). In another RCT, delayed recall verbal memory showed significant preservation in the ACS group, compared to the placebo group ($p = 0.0202$)	The survey results indicated that the anserine and carnosine consumption from the diet was equivalent between the ACS and placebo groups in this study, a better method is needed to estimate their levels more accurately	Low risk	

Table 2. Continued.

Author, year	Age		Gender		Drug regimen		L-carnosine		Duration	Other supplements		Outcome measures		Result	Conclusion	ROB
	Control	Intervention	Control	Intervention	Control	Intervention	Control	Intervention		Primary	Secondary	ADR				
Katakura Y <i>et al.</i> (2017) [53]	65.3 ± 1.6	60.4 ± 2.1	10/20 (M/FM)	10/20 (M/FM)	Placebo (43 mg/day L-lysine and L-histidine (150 mg/day))	43 mg/day L-carnosine	1.0 g anserine/carnosine (3:1) formula daily	Patients were anserine evaluated at baseline and 3 months	Anserine	(i) A dietary survey was conducted using a semi-quantitative method as reported. Inventory of food intake during the 3-month test period. (ii) Cognitive testing: WMS-LMI, WMS-LM2, ADAS-cog, BDI, SF-36, MCS and SF-36 PCS; (iii) Microarray analysis: PBMCs from volunteers were used for total RNA extraction. The total RNA quality was assessed; (iv) qRT-PCR: Gene expression levels were normalized to the corresponding β -actin level.	Nil	Not mentioned	A microarray analysis of PBMCs showed decreased expression of CCL24, an inflammatory chemokine ($p < 0.05$). Verbal memory assessed using the WMS-LM, was preserved in the ACS group. An age-restricted sub-analysis showed significant verbal memory preservation by ACS in participants who were in their 60s (active, $n = 12$; placebo, $n = 9$; $p = 0.048$) and 70s (active, $n = 7$; placebo, $n = 11$; $p < 0.017$). The suppression of CCL24 expression was greatest in people who were in their 70s ($p < 0.01$). There was a significant correlation between the preservation of verbal memory and suppression of CCL24 expression in the group that was in the 70s (Poisson correlation, $r = 0.46$, $p < 0.05$).	ACS may preserve verbal episodic memory, probably owing to CCL24 suppression in the blood, especially in elderly participants.	Low risk	
Ghajar A <i>et al.</i> (2018) [50]	24.93 ± 6.78	22.00 ± 7.76	28/2 (M/FM)	26/4 (M/FM)	Risperidone (up to 6 mg/day) + Placebo. Patients with sleep problems received 1 mg lorazepam every night for the first week	Control drug regimen plus L-carnosine	2 g/day, participants received 500 mg/day of L-carnosine for 5 days, on day 6, the dose of L-carnosine was increased to 1000 mg/day in two divided doses for 5 days. On day 11, patients received the maximum fixed dose of 2 g/day in 2 divided dose	Patients were evaluated at baseline and 2, 4, 6 and 8 week	Nil	(i) The difference of change in the PANSS subscale scores and the PANSS and ESRs total scores (at each visit); (ii) HDRS to assess changes in depressive symptoms; (iii) ADRs	Headache, Dry mouth, Nausea, Daytime drowsiness and sweating	The difference in PANSS negative subscale score reduction	L-carnosine resulted in greater improvement of negative scores as well as total PANSS scores but not positive subscale scores compared to placebo. HDRS scores and its changes did not differ between the two groups. Both groups demonstrated a constant ESRs score during the trial course. Frequency of other side effects was not significantly different between the two groups. In a multiple regression analysis model (controlled for positive, general psychopathology; depressive and extrapyramidal symptoms, as well as other variables), the treatment group significantly predicted changes in primary negative symptoms	L-carnosine as an add-on to risperidone showed good tolerability and significant beneficial effects on negative symptoms of patients with chronic stable schizophrenia	Low risk	

Table 2. Continued.

Author, year	Age		Gender		Drug regimen		L-carnosine		Duration	Other supplements	Outcome measures		ADR	Result	Conclusion	ROB
	Control	Intervention	Control (M/F)	Intervention (M/F)	Control	Intervention	Dose & route	Primary			Secondary					
Masaoka N <i>et al.</i> (2019) [56]	73.6 ± 6.1	72.9 ± 8.8	12/13	12/13	Placebo	Anserine and carnosine	1 g of ACS (250 mg of carnosine)	Anserine were evaluated at baseline and 4, 8 and 12 week	Patients were evaluated at baseline and 4, 8 and 12 week	Anserine (750 mg)	Nil	Nil	Nil	The score improvement in the glnCDR was superior in the active group than placebo ($p = 0.023$). No beneficial effect in MCI subjects but might suggest that beneficial effect in the active group was detected in the other psychometric tests including the MMSE, WAIS and ADAS. When APOE4 help to prevent a transition from MCI to AD as a secondary (APOE4 (-)) subjects were separately analyzed, beneficial change in the APOE4 (+) subjects was observed in MMSE ($p = 0.025$) as well as in glnCDR ($p = 0.026$) assessing these benefits	Low risk	

Age-data expressed as mean ± SD, * median (range).

M/F, Male/Female; ADRs, Adverse drug reactions; ROB, Risk of Bias; FLOFOX-6, oxaliplatin, 5 fluorouracil and leucovorin; NCI-CTCAE-v4, National Cancer Institute Common Terminology Criteria for Adverse Events; NF- κ B, human nuclear factor erythroid2-related factor; MDA, malondialdehyde; NF- κ B, human nuclear factor- κ B; TNF- α , Tumor necrosis factor- α ; BUN, Blood Urea Nitrogen; AST, Aspartate Amino Transferase; ALT, Alanine transaminase; T.Bil, Total bilirubin; Hb, Hemoglobin; PLT, Platelets; T.LC, total leukocyte count; OPN, Oxaliplatin induced peripheral neuropathy; Ara-C, cytarabine; CPA, cyclophosphamide; Flu, fludabine; ETP, etoposide; MCNU, ramustine; L-PAM, melphalan; BU, busulfan; ATG, thymoglobulin; AEs, Adverse events; CIPN, Chemotherapy-induced peripheral neuropathy; PZ, Polaprezinc; AE, Adverse event; sCr, Serum creatinine; WBC, White blood cells; HsCT, Hematopoietic stem cell transplantation; ACEinhibitors, Angiotensin Converting Enzyme inhibitors; ARB, Angiotensin Receptor Blockers; 6-MWT, 6-minute walk test; Quality of life; EQ-5D, EuroQol-5D; VAS, Visual Analogue Scale; LVEF, Left ventricular ejection fraction; CHF, Congestive Heart Failure; PCI, Percutaneous coronary intervention; CRP, C-reactive protein; IL-6, Interleukin-6; EF, Ejection fraction; LVDd, Left ventricular diastolic dimension; LVDs, Left ventricular end-systolic dimension; AMI, Acute myocardial infarction; CPK, Creatine phosphokinase; SEC, Glycerin-based emollient cream; UC, Urea, arginine and carnosine cream; XAS, Xerosis assessment scale; OCS, Overall cutaneous score; T2DM, Type 2 diabetes mellitus; FBS, Fasting blood sugar; TC, Total Cholesterol; HDL-C, High-density lipoprotein cholesterol; LDL-C, Low-density lipoprotein cholesterol; HbA1c, Glycated hemoglobin; HOMA-IR, Homeostatic model assessment of insulin resistance; HOMA- β , Homeostatic model assessment of insulin secretion; TG, Triglycerides; CML, Carboxymethyl lysine; sRAGE, Soluble receptor of advanced glycation end products; IL-1 β , Interleukin-1 β ; ALA, α -lipoic acid; ADP, Adenosine diphosphate; PAF, Platelet activating factor; TGF- β , Transforming growth factor beta; UACR, Urine albumin-creatinine ratio; GFR, Glomerular filtration rate; p.o. per oral; QoL, Quality of life interview; SF-36MCS, 36-item short form survey for mental health component summary scores; SF-36PCS, 36-item short form survey for physical health component summary scores; CME, Chicken meat extract; CRC, carnosine related compounds; MMSE, Mini mental state examination; STMS, Short Test of mental status; GDS, Geriatric Depression Scale; CDR, Clinical Dementia Rating; BP, Blood pressure; HR, Heart rate; WMS-LMI, Wechsler memory scale-revised logical memory immediate recall; WMS-LM2, Wechsler memory scale-revised logical memory delayed recall; ADAS-cog, Alzheimer's disease Assessment Scale; BDI, Beck Depression Inventory; FBMCs, peripheral blood mononuclear cells; RNA, Ribonucleic acid; MRI, Magnetic Resonance Imaging; ACS, Anserine and Carnosine Supplementation; RCT, Randomized controlled trial; IL-8, Interleukin-8; qRT-PCR, Real-time quantitative reverse transcription polymerase chain reaction; FBMCs, peripheral blood mononuclear cells PANSS, Positive and negative syndrome scale; ESRs, Erythrocyt sedimentation rate; HDRS, Hamilton depression rating scale; APOE4, Apolipoprotein E4; ELISA, Enzyme-linked immunosorbent assay; EEG, Electroencephalogram; MCI, Mild cognitive impairment; AD, Alzheimer's disease; NAI, Neuronal activity topography.

T2DM patients, which enhances skin hydration and alleviates the state of severe skin dryness. Fasting blood sugar (FBS), total cholesterol (TC), high-density lipoprotein cholesterol (HDL-c), low-density lipoprotein cholesterol (LDL-c), Glycated hemoglobin (HbA1c), fasting insulin, homeostatic model assessment of insulin resistance (HOMA-IR), homeostatic model assessment of insulin secretion (HOMA- β), triglycerides (TG) and serum levels of pentosidine Carboxymethyl lysine (CML), soluble receptor of advanced glycation end products (sRAGE), TNF- α , interleukin-1 β (IL-1 β), IL-6 were examined in Houjehani *et al.* [22], study. They found that L-carnosine improved glycemic control, lipid profile, AGEs, and low-grade inflammation without changing soluble receptor of AGEs, IL-6 and IL-1 β .

The glycemic profile, lipid profile, kidney function, liver function, tissue damage, thyroid function, cancer markers, electrolytes, minerals, oxidative stress, platelet activating factor (PAF), arachidonic acid, epinephrine, thrombin, and collagen were all examined in the Karkabounas *et al.* [52], study, which found that L-carnosine as an added supplement reduced glucose concentration in T2DM patients by potentially increasing insulin production from the pancreas.

Siriwattanasit *et al.* [57], looked at the change in urine TGF- β levels in T2DM patients with diabetic nephropathy and showed lower urinary TGF- β levels. HbA1c, FBS, LDL-c, HDL-c, TG, and TC are typical laboratory variables examined in the included diabetic studies [22,52,57]. A comparable analysis of serum creatinine (serum Cr.) was found in [52,57] study.

Budzen *et al.* [48], study investigated cognitive function, the existence of depressive symptoms, physical capacity, body measures, blood pressure (BP), and heart rate (HR) in neurodegenerative disorder patients. They also looked at the mini mental state examination (MMSE) and short test of mental status (STMS) to test cognitive functions, the geriatric depression scale (GDS) to examine the presence of depressive symptoms, and the clinical dementia rating (CDR) to assess dementia severity, and found that L-carnosine supplementation in the elderly could have positive effects on cognitive functioning and physical capacity.

Based on the results of a neurocognition set-shifting test and a strategic target detection test, L-carnosine was found to be an additional treatment to ameliorate executive dysfunction in people with schizophrenia [49]. The difference in positive and negative syndrome scale (PANSS) negative subscale score reduction and hamilton depression rating scale (HDRS) was used to assess changes in depressive symptoms, and L-carnosine as an add-on to risperidone was found to have good tolerability and considerable favourable effects on negative symptoms in individuals with chronic stable schizophrenia [50].

The outcome measured in the Hisatsune *et al.* [51], study included a semi-quantitative dietary survey; cog-

nitive testing: Wechsler memory scale-revised logical memory immediate recall (WMS-LM1), Wechsler memory scale-revised logical memory delayed recall (WMS-LM2), Alzheimer's disease assessment scale (ADAScog), beck depression inventory (BDI), study, 36-item d physical health component summary (SF-36 MCS), and study, 36-item d physical health component summary (SF-36 PCS); blood samples and immunoassays; microarray analysis and acquisition of human Magnetic Resonance Imaging (MRI) data and MRI data analysis; and microarray analysis and acquisition of human MRI data. The Anserine and Carnosine (ACS) and placebo groups in this trial consumed the same amount of anserine and carnosine from their diets, according to the survey results.

Katakura *et al.* [53], evaluated dietary survey outcome measures using a semi-quantitative method; cognitive testing: WMS-LM1, WMS-LM2, ADAScog, BDI, SF-36 MCS and SF-36 PCS; microarray analysis and real-time quantitative reverse transcription polymerase chain reaction (qRT-PCR), which found that ACS may preserve verbal episodic memory, likely due to CCL24 suppression in the blood, especially in elderly participants.

Inventory of anserine and carnosine in the Normal Diet; Blood Analysis; Cognitive Testing: MMSE, gloCDR, CDRsob, WMS-1, WMS-2, ADAS and GDS; and electroencephalogram (EEG) recording were used in the Masuoka *et al.* [56], study, which showed that the results did not indicate that ACS has significant efficacy on cognitive change in mild cognitive impairment (MCI) subjects but might suggest that ACS has a beneficial effect in APOE4 (+) MCI subjects exists, and thereby, ACS might help to prevent a transition from MCI to AD as a secondary effect in APOE4 (+) MCI subjects.

WMS-1, WMS-2, and ADAS [51,53,56] were the most commonly used outcome measures in neurodegenerative disorder studies, followed by BDI, SF-36 MCS, and SF-36 PCS [51,53], GDS [48,51,56] and MMSE [48,56].

3.5 Risk of Bias and Quality of Reporting

Supplementary Table 2 shows the RoB assessment for each of the studies. All reports were critically examined using the standard checklist for the revised Cochrane RoB tool. The method of randomization is rarely described, despite the fact that randomization is an important tool to reduce bias. The method of randomization was not reported in any of the included trials, or there was insufficient information. Fourteen RCTs were analyzed for the effect of L-carnosine in various age-related diseases such as cancer [5,55], cardiovascular disease [54,58], diabetes mellitus [22,33,52,57] and neurodegenerative disorder [48–51,53,56].

The bias arising from randomization technique was “low risk” for all 14 included studies in a quality assessment using the Revised Cochrane Risk of Bias Tool for RCT, whereas bias due to deviation from intended intervention

had “some concerns” for 5 studies [5,33,54,55,58] and low risk for the other 9 [22,48–53,56,57].

Bias for missing outcome data was found to be “low risk” in all of the included studies. Only one study [33] found a “low risk” of bias in outcome measurement, whereas the remaining 13 studies had “some concerns” about outcome measuring. All of the studies that were considered demonstrated a “low risk” of selecting the reported result. All of the studies were evaluated separately, and the aggregate assessment had a “low risk” of bias (Supplementary Table 2).

Only four studies [5,33,55,57] identified a “high risk” of allocation concealment in the selection bias (Fig. 2), and one study identified an “unclear risk” of allocation concealment in the selection bias [54]. Only six studies had a “high risk” of performance bias due to participant and personnel blinding [5,33,54,55,57,58]. However, in one trial, the outcome assessors were blinded [33]. The random sequence generation in selection bias was addressed in all of the studies. All of the papers were evaluated to see if the research question, study population, and L-carnosine indices were well described. Other causes of bias, such as medication pooling, dropouts, unit-of-analysis mistakes, design-specific bias, and conflicts of interest, were not mentioned in these studies.

3.6 Meta-Analysis

Primarily, we analysed whether or not L-carnosine supplementation had a positive impact on age-related diseases such as cancer, cardiovascular disease, diabetes mellitus, and neurodegenerative disorders. We used 14 studies in our qualitative analysis [5,22,33,48–58]. Five studies were excluded from the quantitative synthesis [5,33,49,50,55] because they did not correlate with the outcome variables and did not report any data with the other included studies under each age-related disease.

3.6.1 Effect of L-Carnosine in Cardiovascular Disease Patients

Two studies [54,58] looked at the effects of L-carnosine on patients with cardiovascular disease. On the ejection fraction, there was no statistically significant difference between the L-carnosine and control groups [MD 95% CI = 0.82 (–2.38, 4.01); $p = 0.62$; $p = 0.56$; $I^2 = 0\%$] (Fig. 3).

3.6.2. Effect of L-Carnosine in Diabetes Mellitus Patients

HbA1c, FBS, LDL-c, HDL-c, TGs, TCs, and serum Cr. were used to study the effect of L-carnosine in diabetes mellitus patients. Three studies (n = 248) [22,52,57] showed a statistically significant difference between the L-carnosine group and the control group on HbA1c [MD 95% CI = –1.25 (–2.49, –0.022); $p = 0.05$; $p = 0.001$; $I^2 = 85\%$] (Fig. 4A) and FBS [MD 95% CI = –12.44 (–22.44, –2.44); $p = 0.01$; $p = 0.40$; $I^2 = 0\%$] (Fig. 4B), favouring the L-

carnosine group.

On the other hand, there was no statistically significant difference between the L-carnosine and control groups on the LDL-c [MD 95% CI = –4.70 (–13.33, 3.94); $p = 0.29$; $p = 0.36$; $I^2 = 2\%$] (Fig. 4C), HDL-c [MD 95% CI = –0.50 (–3.82, 2.82); $p = 0.77$; $p = 0.77$; $I^2 = 0\%$] (Fig. 4D), TGs [MD 95% CI = –0.42 (–6.02, 5.19); $p = 0.88$; $p = 0.44$; $I^2 = 0\%$] (Fig. 4E) and TCs [MD 95% CI = –5.09 (–11.81, 1.63); $p = 0.14$; $p = 0.60$; $I^2 = 0\%$] (Fig. 4F) in three studies (n = 248) [22,52,57]. L-carnosine does not alter the levels of serum creatinine [MD 95% CI = 0.06 (–0.12, 0.24); $p = 0.51$; $p = 0.27$; $I^2 = 19\%$] in two studies (n = 204) [52,57] (Fig. 4G).

3.6.3. Effect of L-Carnosine in Neurodegenerative Disorder Patients

Variables such as WMS-1, WMS-2, ADAS, BDI, SF-36 MCS, SF-36 PCS, MMSE, and GDS were used to find the effect of L-carnosine in patients with neurodegenerative disorder. On WMS-1, the association between the three studies (n = 149) [51,53,56] revealed no statistically significant difference between the L-carnosine and control groups [MD 95% CI = 0.26 (–0.79, 0.26); $p = 0.33$; $p = 0.75$; $I^2 = 0\%$] (Fig. 5A), however the same studies showed the statistically significant difference between the groups on WMS-2 [MD 95% CI = 1.34 (0.83, 1.85); $p < 0.00001$; $p = 0.43$; $I^2 = 0\%$] (Fig. 5B).

On the ADAS, a statistically significant difference was observed between the L-carnosine and control groups of the three studies (n = 149) [51,53,56] [MD 95% CI = 0.98 (–1.55, –0.42); $p = 0.0007$; $p = 0.86$; $I^2 = 0\%$] favouring the control group over the L-carnosine group (Fig. 5C). Similarly, on BDI, a statistically significant difference was observed between the L-carnosine and control groups of the two studies (n = 99) [51,53], favouring the control group over the L-carnosine group (Fig. 5D).

Pooling analysis of the SF-36 MCS [MD 95% CI = –0.25 (–1.29, 0.79); $p = 0.64$; $p = 0.55$; $I^2 = 0\%$] and SF-36 PCS [MD 95% CI = 0.53 (–0.47, 1.52); $p = 0.30$; $p = 0.71$; $I^2 = 0\%$], as shown in Fig. 5E,F, revealed that there was no statistically significant difference between the L-carnosine group and the control group in the two studies (n = 99) [51, 53].

Meta-analysis of data from three studies (n = 140) [48,51,53] showed no statistically significant difference in MMSE score between the L-carnosine and control groups [MD 95% CI = 0.50 (–0.60, 1.60); $p = 0.37$; $p = 0.97$; $I^2 = 0\%$] (Fig. 5G). Similarly, in GDI also no statistically significant difference between the L-carnosine and control groups in two studies (n = 101) [48,53] [MD 95% CI = 0.01 (–1.73, 1.76); $p = 0.99$; $p = 0.97$; $I^2 = 0\%$] (Fig. 5H).

4. Discussion

The purpose of this review is to assess the effect of L-carnosine in the context of age-related illnesses and




Fig. 2. Methodological quality of the included studies based on the revised Cochrane risk of bias tool for randomized trials. (A) Overall risk of bias. Each item is presented as a percentage based on all included studies. (B) Individual risk of bias. Each item is scored as 'weak', 'moderate', or 'strong'.




Fig. 3. Effect of L-carnosine in cardiovascular disease patients. Effect of L-carnosine treated versus control on measures of ejection fraction in patients with cardiovascular disorders. CI, confidence interval; df, degrees of freedom; IV, inverse variance; SE, standard error.

makeweight in the literature, as well as to convey the relevance of existing data as evidence for clinical practise and future research. Clinical studies on the efficacy of L-carnosine have been widely published in the last decade, indicating a growing interest among health care professionals in the use of L-carnosine for the treatment of age-related illnesses. Egypt, Japan, Italy, Iran, Greece, Thailand, the United States, and Poland have all conducted studies that

look at the preference for dietary/therapeutic intervention in L-carnosine, particularly in oriental medicine, in these countries.

L-carnosine has been found to have multimodal activity. Its impact on age-related diseases such cancer, cardiovascular disease, diabetes, and neurological problems is comprehensively reviewed in this paper. Fig. 6 depicts the mode of action of L-carnosine in the aforementioned




Fig. 4. Effect of L-carnosine in diabetes mellitus patients. (A) Effect of L-carnosine treated versus control on measures of HbA1c in patients with diabetes mellitus. (B) Effect of L-carnosine treated versus control on measures of FBS in patients with diabetes mellitus. (C) Effect of L-carnosine treated versus control on measures of LDL-c in patients with diabetes mellitus. (D) Effect of L-carnosine treated versus control on measures of HDL-c in patients with diabetes mellitus. (E) Effect of L-carnosine treated versus control on measures of TGs in patients with diabetes mellitus. (F) Effect of L-carnosine treated versus control on measures of TCs in patients with diabetes mellitus. (G) Effect of L-carnosine treated versus control on measures of serum Cr. in patients with diabetes mellitus. HbA1c, glycated hemoglobin; FBS, fasting blood sugar; LDL-c, low-density lipoprotein cholesterol; HDL-c, high-density lipoprotein cholesterol; TGs, triglycerides; TCs, total cholesterol; Cr, creatinine; CI, confidence interval; df, degrees of freedom; IV, inverse variance; SE standard error.

illnesses. Apart from the selected 14 RCTs, the therapeutic potential of L-carnosine was studied in pre-clinical and other clinical (non-RCTs) studies. The anti-cancer effect of L-carnosine has been investigated in animals for various

cancer types such as ovarian cancer [59], HER2/neu cancer [60], bladder cancer [61], gastric cancer [62,63], hepatic cancer [64] and colon cancer [65]. L-carnosine has also been demonstrated to help with chemotherapy-induced




Fig. 5. Effect of L-carnosine in neurodegenerative disorder patients. (A) Effect of L-carnosine treated versus control on measures of WMS-1 in patients with neurodegenerative disorder. (B) Effect of L-carnosine treated versus control on measures of WMS-2 in patients with neurodegenerative disorder. (C) Effect of L-carnosine treated versus control on measures of ADAS in patients with neurodegenerative disorder. (D) Effect of L-carnosine treated versus control on measures of BDI in patients with neurodegenerative disorder. (E) Effect of L-carnosine treated versus control on measures of SF-36 MCS in patients with neurodegenerative disorder. (F) Effect of L-carnosine treated versus control on measures of SF-36 PCS in patients with neurodegenerative disorder. (G) Effect of L-carnosine treated versus control on measures of MMSE in patients with neurodegenerative disorder. (H) Effect of L-carnosine treated versus control on measures of GDS in patients with neurodegenerative disorder. GDS, geriatric depression scale; WMS-1, wechsler memory scale-revised logical memory immediate recall; WMS-2, wechsler memory scale-revised logical memory delayed recall; ADAS, Alzheimer's disease assessment scale; BDI, beck depression inventory; SF-36 MCS, 36-item short form survey for mental-health component summary scores; SF-36 PCS, 36-item short form survey for physical-health component summary scores; MMSE, mini mental state examination; CI, confidence interval; df, degrees of freedom; IV, inverse variance; SE, standard error.

toxicities [66–68]. L-carnosine has clinical evidence for its action against chemotherapy-induced toxicities [69,70],

but no research has been conducted on its anticancer activity. Pre-clinical evidence from cardiovascular research




Fig. 6. Mechanism of L-carnosine in age-related disorders (cancer, cardiovascular disease, diabetes, and neurodegenerative disorders).

suggests that L-carnosine improves functional capability in ischemia circumstances [71–78]. Similarly, pre-clinical evidence from diabetes studies showed that L-carnosine had an effect on diabetes [79,80] and its complications [81–83]. Pre-clinical and clinical (other than RCTs) studies from neurodegenerative disorders explored the effect of L-carnosine on cognitive decline and neurophysiological effects [84–88].

This systematic review and meta-analysis of the literature presents data using the PRISMA guidelines and the PICOS approach. The 14 RCTs included in the study recruited a total of 794 individuals. Neurodegenerative studies had 329 (41%) participants, diabetes studies had 216 (27%) participants, cancer studies had 149 (18%) participants, and cardiovascular studies had 100 (12.5%) participants.

When we looked at all of the studies on the effects of L-carnosine supplementation on age-related disorders, we found that just 9 of them provided adequate data to make

a quantitative comparison of variables. In the remaining 5 studies, 2 of the cancer studies [5,55] investigated the mechanism of dissimilar toxicities induced by chemotherapy, and 1 of the diabetes studies [33] investigated L-carnosine on severe xerosis, which is one of a skin complication of diabetes, where they measured skin hydration, so it wasn't eligible for quantitative comparison, and the other 2 studies [49,50] looked at neurodegenerative diseases and weren't eligible because of their dissimilar assessments.

The majority of the studies included in this meta-analysis used a combination of anserine and L-carnosine. Anserine is a natural derivative of carnosine that is commonly used because human carnosinase does not cleave it. Anserine and carnosine have been shown to have similar physiological activities [89]. Just a few of the studies employed PZ, a synthetic carnosine derivative in which zinc and carnosine are linked in a one-to-one ratio to form a polymeric structure. Several companies are currently marketing this product as a zinc dietary supplement with "additional

value for gastric health” [90]. For a better understanding of L-carnosine’s therapeutic potential against age-related disorders, future clinical research focusing on L-carnosine should attempt L-carnosine alone.

Our meta-analysis of diabetes studies [22,52,57] suggests that the intervention group (L-carnosine alone or in combination with the control group regimen, or with or without add-on supplements) can improve FBS and HbA1c levels in diabetes patients. Despite the fact that individual studies showed increases in exercise capacity and functional capacity, statistical significance was not found in the intervention group compared to the control group in cardiovascular studies [54,58]. Neurodegenerative studies [51,53,56] suggest that L-carnosine improves memory deficiencies among elderly populations accordance with WMS-2 scale but other than that failed to show significant results for other tools. It’s likely that L-carnosine affects specific cognitive functions, which may be observed in the clinical setting using more specialised techniques rather than a broad assessment of cognitive status [91].

According to the Caruso *et al.* [91], study T2DM is known to be a risk factor for cognitive impairment and AD, and different neurobiological links have been identified between T2DM and AD, such as insulin resistance, low-grade inflammation, increased oxidative stress, and AGE accumulation. In accordance with Caruso *et al.* [91], study and our meta-analysis, L-carnosine has anti-diabetic and cognitive properties. To validate the therapeutic potential of L-carnosine, more double-blind, randomised, placebo-controlled trials on age-related illnesses are needed.

The diversity in measures (for example, different scales for cognitive function) used across studies made meta-analysis challenging, and as a result, not all studies included in the systematic review could be included in the meta-analysis, resulting in lower statistical power and potential overestimation of the results.

5. Conclusions and Future Perspectives

The findings of the review indisputably demonstrated a significant beneficial effect of L-carnosine only for diabetes mellitus and cognitive impairment, but not for cardiovascular illnesses. L-carnosine research in cancer was solely focused on chemotherapy-induced toxicity. Its anti-cancer abilities have not been investigated. Future research should use preclinical evidence to look at anticancer potential of L-carnosine in a variety of cancer patients to fill this gap in the literature. To establish the therapeutic potential of L-carnosine in various age-related disorders, more RCTs with larger sample size are needed.

Abbreviations

PROSPERO, The International Prospective Register of Systematic Reviews; PICO, population, intervention, comparison, outcomes, and study design; PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-

Analysis; RevMan, Revenue manager; HbA1c, Glycated haemoglobin; FBS, Fasting blood sugar; WMS-LM2, Beck Depression Inventory; ADAS, Alzheimer’s disease Assessment Scale; BDI, Beck Depression Inventory.

Author Contributions

KSur, MD, KSri, KU, RMG contributed to the database search, data collection; KSur, MD, KWG, KU, VTM, and RMG contributed to data extraction, data analysis; KSur, KWG, KU, VTM, CA, LCM, RMG contributed to writing and revising of the manuscript. The topic was conceptualized by VTM, CA and RMG. All authors read and approved the final manuscript.

Ethics Approval and Consent to Participate

Not applicable.

Acknowledgment

Not applicable.

Funding

This research received no external funding.

Conflict of Interest

The authors declare no conflict of interest.

Supplementary Material

Supplementary material associated with this article can be found, in the online version, at <https://doi.org/10.31083/j.fbl2801018>.

References

- [1] Gariballa SE, Sinclair AJ. Carnosine: physiological properties and therapeutic potential. *Age and Ageing*. 2000; 29: 207–210.
- [2] Jukić I, Kolobarić N, Stupin A, Matić A, Kozina N, Mihaljević Z, *et al.* Carnosine, small but mighty-prospect of use as functional ingredient for functional food formulation. *Antioxidants*. 2021; 10: 1037.
- [3] Everaert I, Mooyaart A, Baguet A, Zutinic A, Baelde H, Achten E, *et al.* Vegetarianism, female gender and increasing age, but not CNDP1 genotype, are associated with reduced muscle carnosine levels in humans. *Amino Acids*. 2011; 40: 1221–1229.
- [4] Bellia F, Vecchio G, Rizzarelli E. Carnosine derivatives: new multifunctional drug-like molecules. *Amino Acids*. 2012; 43: 153–163.
- [5] Kitagawa J, Kobayashi R, Nagata Y, Kasahara S, Ono T, Sawada M, *et al.* Polaprezinc for prevention of oral mucositis in patients receiving chemotherapy followed by hematopoietic stem cell transplantation: A multi-institutional randomized controlled trial. *International Journal of Cancer*. 2021; 148: 1462–1469.
- [6] Tang W, Liu H, Ooi TC, Rajab NF, Cao H, Sharif R. Zinc carnosine: Frontiers advances of supplement for cancer therapy. *Biomedicine & Pharmacotherapy*. 2022; 151: 113157.
- [7] Grothey A, Goldberg RM. A review of oxaliplatin and its clinical use in colorectal cancer. *Expert Opinion on Pharmacotherapy*. 2004; 5: 2159–2170.
- [8] McWhinney SR, Goldberg RM, McLeod HL. Platinum neu-

- rotoxicity pharmacogenetics. *Molecular Cancer Therapeutics*. 2009; 8: 10–16.
- [9] Li W, Khor TO, Xu C, Shen G, Jeong W, Yu S, *et al.* Activation of Nrf2-antioxidant signaling attenuates NfκB-inflammatory response and elicits apoptosis. *Biochemical Pharmacology*. 2008; 76: 1485–1489.
- [10] Negi G, Kumar A, Sharma S. Adopting Nrf2 and NF-κB from cancer: Is there any role of the duo in diabetes. *Nature Precedings*. 2011; 8: 1.
- [11] Peng S, Ying AF, Chan NJH, Sundar R, Soon YY, Bandla A. Prevention of Oxaliplatin-Induced Peripheral Neuropathy: A Systematic Review and Meta-Analysis. *Frontiers in Oncology*. 2022; 12: 731223.
- [12] Liebetrau C, Hoffmann J, Dörr O, Gaede L, Blumenstein J, Biermann H, *et al.* Release kinetics of inflammatory biomarkers in a clinical model of acute myocardial infarction. *Circulation Research*. 2015; 116: 867–875.
- [13] Liu W, Liu T, Yin M. Beneficial effects of histidine and carnosine on ethanol-induced chronic liver injury. *Food and Chemical Toxicology: an International Journal Published for the British Industrial Biological Research Association*. 2008; 46: 1503–1509.
- [14] Guney Y, Turku UO, Hicsonmez A, Andrieu MN, Guney HZ, Bilgihan A, *et al.* Carnosine may reduce lung injury caused by radiation therapy. *Medical Hypotheses*. 2006; 66: 957–959.
- [15] Piña IL, Apstein CS, Balady GJ, Belardinelli R, Chaitman BR, Duscha BD, *et al.* Exercise and heart failure: A statement from the American Heart Association Committee on exercise, rehabilitation, and prevention. *Circulation*. 2003; 107: 1210–1225.
- [16] Hultman E, Sahlin K. Acid-base balance during exercise. *Exercise and Sport Sciences Reviews*. 1980; 8: 41–128.
- [17] Caruso J, Charles J, Unruh K, Giebel R, Learmonth L, Potter W. Ergogenic effects of β-alanine and carnosine: proposed future research to quantify their efficacy. *Nutrients*. 2012; 4: 585–601.
- [18] Culbertson JY, Kreider RB, Greenwood M, Cooke M. Effects of beta-alanine on muscle carnosine and exercise performance: a review of the current literature. *Nutrients*. 2010; 2: 75–98.
- [19] Derave W, Everaert I, Beekman S, Baguet A. Muscle carnosine metabolism and beta-alanine supplementation in relation to exercise and training. *Sports Medicine*. 2010; 40: 247–263.
- [20] Batrukova MA, Rubtsov AM. Histidine-containing dipeptides as endogenous regulators of the activity of sarcoplasmic reticulum Ca-release channels. *Biochimica Et Biophysica Acta*. 1997; 1324: 142–150.
- [21] Ririe DG, Roberts PR, Shouse MN, Zaloga GP. Vasodilatory actions of the dietary peptide carnosine. *Nutrition*. 2000; 16: 168–172.
- [22] Houjehani S, Kheirouri S, Faraji E, Jafarabadi MA. L-Carnosine supplementation attenuated fasting glucose, triglycerides, advanced glycation end products, and tumor necrosis factor-α levels in patients with type 2 diabetes: a double-blind placebo-controlled randomized clinical trial. *Nutrition Research*. 2018; 49: 96–106.
- [23] Semba RD, Nicklett EJ, Ferrucci L. Does accumulation of advanced glycation end products contribute to the aging phenotype? *The Journals of Gerontology Series A, Biological Sciences and Medical Sciences*. 2010; 65: 963–975.
- [24] Goldin A, Beckman JA, Schmidt AM, Creager MA. Advanced glycation end products: sparking the development of diabetic vascular injury. *Circulation*. 2006; 114: 597–605.
- [25] Alhamedani MS, Al-Azzawie HF, Abbas FKH. Decreased formation of advanced glycation end-products in peritoneal fluid by carnosine and related peptides. *Peritoneal Dialysis International*. 2007; 27: 86–89.
- [26] Baye E, Menon K, de Courten MP, Earnest A, Cameron J, de Courten B. Does supplementation with carnosine improve cardiometabolic health and cognitive function in patients with pre-diabetes and type 2 diabetes? Study protocol for a randomised, double-blind, placebo-controlled trial. *BMJ Open*. 2017; 7: e017691.
- [27] Pfister F, Riedl E, Wang Q, vom Hagen F, Deinzer M, Harnsen MC, *et al.* Oral carnosine supplementation prevents vascular damage in experimental diabetic retinopathy. *Cellular Physiology and Biochemistry*. 2011; 28: 125–136.
- [28] Satirapoj B, Adler SG. Comprehensive approach to diabetic nephropathy. *Kidney Research and Clinical Practice*. 2014; 33: 121–131.
- [29] Schena FP, Gesualdo L. Pathogenetic mechanisms of diabetic nephropathy. *Journal of the American Society of Nephrology*. 2005; 16 Suppl 1: S30–S33.
- [30] Chang AS, Hathaway CK, Smithies O, Kakoki M. Transforming growth factor-β1 and diabetic nephropathy. *American Journal of Physiology Renal Physiology*. 2016; 310: F689–F696.
- [31] Zheng X, Bhalla V. The missing link: studying the alternative TGF-β pathway provides a unifying theory for different components of diabetic nephropathy. *Diabetes*. 2015; 64: 1898–1900.
- [32] Hobart LJ, Seibel I, Yeagans GS, Seidler NW. Anti-crosslinking properties of carnosine: significance of histidine. *Life Sciences*. 2004; 75: 1379–1389.
- [33] Federici A, Federici G, Milani M. Use of a urea, arginine and carnosine cream versus a standard emollient glycerol cream for treatment of severe xerosis of the feet in patients with type 2 diabetes: a randomized, 8 month, assessor-blinded, controlled trial. *Current Medical Research and Opinion*. 2015; 31: 1063–1069.
- [34] Uhoda E, Debatisse B, Paquet P, Piérard-Franchimont C, Piérard GE. The so-called dry skin of the diabetic patient. *Revue Médicale De Liege*. 2005; 60: 560–563.
- [35] Beck K, Javitt DC, Howes OD. Targeting glutamate to treat schizophrenia: lessons from recent clinical studies. *Psychopharmacology*. 2016; 233: 2425–2428.
- [36] Field JR, Walker AG, Conn PJ. Targeting glutamate synapses in schizophrenia. *Trends in Molecular Medicine*. 2011; 17: 689–698.
- [37] Solana-Manrique C, Sanz FJ, Martínez-Carrión G, Paricio N. Antioxidant and Neuroprotective Effects of Carnosine: Therapeutic Implications in Neurodegenerative Diseases. *Antioxidants*. 2022; 11: 848.
- [38] Karton A, O'Reilly RJ, Pattison DI, Davies MJ, Radom L. Computational design of effective, bioinspired HOCl antioxidants: the role of intramolecular Cl⁺ and H⁺ shifts. *Journal of the American Chemical Society*. 2012; 134: 19240–19245.
- [39] Kulebyakin K, Karpova L, Lakonsteva E, Krasavin M, Boldyrev A. Carnosine protects neurons against oxidative stress and modulates the time profile of MAPK cascade signaling. *Amino Acids*. 2012; 43: 91–96.
- [40] Shen Y, He P, Fan Y, Zhang J, Yan H, Hu W, *et al.* Carnosine protects against permanent cerebral ischemia in histidine decarboxylase knockout mice by reducing glutamate excitotoxicity. *Free Radical Biology & Medicine*. 2010; 48: 727–735.
- [41] Shen Y, Hu W, Fan Y, Dai H, Fu Q, Wei E, *et al.* Carnosine protects against NMDA-induced neurotoxicity in differentiated rat PC12 cells through carnosine-histidine-histamine pathway and H(1)/H(3) receptors. *Biochemical Pharmacology*. 2007; 73: 709–717.
- [42] Hipkiss AR. Could carnosine or related structures suppress Alzheimer's disease? *Journal of Alzheimer's Disease*. 2007; 11: 229–240.
- [43] Taylor JP, Hardy J, Fischbeck KH. Toxic proteins in neurodegenerative disease. *Science*. 2002; 296: 1991–1995.
- [44] Corona C, Frazzini V, Silvestri E, Lattanzio R, La Sorda R, Piantelli M, *et al.* Effects of dietary supplementation of carnosine

- on mitochondrial dysfunction, amyloid pathology, and cognitive deficits in 3xTg-AD mice. *PLoS ONE*. 2011; 6: e17971.
- [45] Park HY, Suh CH, Woo S, Kim PH, Kim KW. Quality Reporting of Systematic Review and Meta-Analysis According to PRISMA 2020 Guidelines: Results from Recently Published Papers in the *Korean Journal of Radiology*. *Korean Journal of Radiology*. 2022; 23: 355–369.
- [46] Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JPA, *et al.* The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *Journal of Clinical Epidemiology*. 2009; 62: e1–34.
- [47] Higgins JPT, Altman DG, Gotzsche PC, Juni P, Moher D, Oxman AD, *et al.* The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *British Medical Journal*. 2011; 343: d5928.
- [48] Szczeniaki D, Budzeń S, Kopeć W, Rymaszewska J. Anserine and carnosine supplementation in the elderly: Effects on cognitive functioning and physical capacity. *Archives of Gerontology and Geriatrics*. 2014; 59: 485–490.
- [49] Chengappa KNR, Turkin SR, DeSanti S, Bowie CR, Brar JS, Schlicht PJ, *et al.* A preliminary, randomized, double-blind, placebo-controlled trial of L-carnosine to improve cognition in schizophrenia. *Schizophrenia Research*. 2012; 142: 145–152.
- [50] Ghajar A, Khoae-Ardakani M, Shahmoradi Z, Alavi A, Afarideh M, Shalbfan M, *et al.* L-carnosine as an add-on to risperidone for treatment of negative symptoms in patients with stable schizophrenia: A double-blind, randomized placebo-controlled trial. *Psychiatry Research*. 2018; 262: 94–101.
- [51] Hisatsune T, Kaneko J, Kurashige H, Cao Y, Satsu H, Totsuka M, *et al.* Effect of Anserine/Carnosine Supplementation on Verbal Episodic Memory in Elderly People. *Journal of Alzheimer's Disease*. 2016; 50: 149–159.
- [52] Karkabounas S, Papadopoulos N, Anastasiadou C, Gubili C, Peschos D, Daskalou T, *et al.* Effects of α -Lipoic Acid, Carnosine, and Thiamine Supplementation in Obese Patients with Type 2 Diabetes Mellitus: A Randomized, Double-Blind Study. *Journal of Medicinal Food*. 2018; 21: 1197–1203.
- [53] Katakura Y, Totsuka M, Imabayashi E, Matsuda H, Hisatsune T. Anserine/Carnosine Supplementation Suppresses the Expression of the Inflammatory Chemokine CCL24 in Peripheral Blood Mononuclear Cells from Elderly People. *Nutrients*. 2017; 9: 1199.
- [54] Lombardi C, Carubelli V, Lazzarini V, Vizzardini E, Bordonali T, Ciccarese C, *et al.* Effects of oral administration of orodispersible levo-carnosine on quality of life and exercise performance in patients with chronic heart failure. *Nutrition*. 2015; 31: 72–78.
- [55] Yehia R, Saleh S, El Abhar H, Saad AS, Schaalan M. L-Carnosine protects against Oxaliplatin-induced peripheral neuropathy in colorectal cancer patients: A perspective on targeting Nrf-2 and NF- κ B pathways. *Toxicology and Applied Pharmacology*. 2019; 365: 41–50.
- [56] Masuoka N, Yoshimine C, Hori M, Tanaka M, Asada T, Abe K, *et al.* Effects of Anserine/Carnosine Supplementation on Mild Cognitive Impairment with APOE4. *Nutrients*. 2019; 11: 1626.
- [57] Siriwardhanasit N, Satirapoj B, Supasyndh O. Effect of Oral carnosine supplementation on urinary TGF- β in diabetic nephropathy: a randomized controlled trial. *BMC Nephrology*. 2021; 22: 236.
- [58] Yoshikawa F, Nakajima T, Hanada M, Hirata K, Masuyama T, Aikawa R. Beneficial effect of polaprezinc on cardiac function post-myocardial infarction: A prospective and randomized clinical trial. *Medicine*. 2019; 98: e14637.
- [59] Mikula-Pietrasik J, Książek K. L-Carnosine Prevents the Pro-carcinogenic Activity of Senescent Peritoneal Mesothelium Towards Ovarian Cancer Cells. *Anticancer Research*. 2016; 36: 665–671.
- [60] Renner C, Zemitzsch N, Fuchs B, Geiger KD, Hermes M, Hengstler J, *et al.* Carnosine retards tumor growth in vivo in an NIH3T3-HER2/neu mouse model. *Molecular Cancer*. 2010; 9: 2.
- [61] Hwang B, Shin S, Song J, Choi YH, Kim W, Moon S. Carnosine exerts antitumor activity against bladder cancers in vitro and in vivo via suppression of angiogenesis. *The Journal of Nutritional Biochemistry*. 2019; 74: 108230.
- [62] Cheng J, Yang J, Liu Y, Xu M, Huang Y, Zhang J, *et al.* Profiling and targeting of cellular mitochondrial bioenergetics: inhibition of human gastric cancer cell growth by carnosine. *Acta Pharmacologica Sinica*. 2019; 40: 938–948.
- [63] Zhang Z, Miao L, Wu X, Liu G, Peng Y, Xin X, *et al.* Carnosine Inhibits the Proliferation of Human Gastric Carcinoma Cells by Retarding Akt/mTOR/p70S6K Signaling. *Journal of Cancer*. 2014; 5: 382–389.
- [64] Ding M, Jiao G, Shi H, Chen Y. Investigations on in vitro anticarcinogenic potential of L-carnosine in liver cancer cells. *Cytotechnology*. 2018; 70: 163–167.
- [65] Horii Y, Shen J, Fujisaki Y, Yoshida K, Nagai K. Effects of L-carnosine on splenic sympathetic nerve activity and tumor proliferation. *Neuroscience Letters*. 2012; 510: 1–5.
- [66] Kumral A, Giriş M, Soluk-Tekkeşin M, Olgaç V, Doğru-Abbasoğlu S, Türkoğlu Ü, *et al.* Beneficial effects of carnosine and carnosine plus vitamin E treatments on doxorubicin-induced oxidative stress and cardiac, hepatic, and renal toxicity in rats. *Human & Experimental Toxicology*. 2016; 35: 635–643.
- [67] Mohamed MA. The protective effect of carnosine on kidneys of albino rats in methotrexate induced oxidative injury. *Egyptian Journal of Forensic Sciences*. 2018; 18: 69–81.
- [68] Xu M, He R, Zhai Y, Abe K, Kurihara H. Effects of carnosine on cyclophosphamide-induced hematopoietic suppression in mice. *The American Journal of Chinese Medicine*. 2014; 42: 131–142.
- [69] Fujii H, Hirose C, Ishihara M, Iihara H, Imai H, Tanaka Y, *et al.* Improvement of Dysgeusia by Polaprezinc, a Zinc-L-carnosine, in Outpatients Receiving Cancer Chemotherapy. *Anticancer Research*. 2018; 38: 6367–6373.
- [70] Funato M, Ozeki M, Suzuki A, Ishihara M, Kobayashi R, Nozawa A, *et al.* Prophylactic Effect of Polaprezinc, a Zinc-L-carnosine, Against Chemotherapy-induced Oral Mucositis in Pediatric Patients Undergoing Autologous Stem Cell Transplantation. *Anticancer Research*. 2018; 38: 4691–4697.
- [71] Bae O, Serfozo K, Baek S, Lee KY, Dorrance A, Rumbelha W, *et al.* Safety and efficacy evaluation of carnosine, an endogenous neuroprotective agent for ischemic stroke. *Stroke*. 2013; 44: 205–212.
- [72] Fedorova TN, Devyatov AA, Berezhnoi DS, Stvolinskii SL, Morozova MP, Gavrilova SA, *et al.* Oxidative Status in Different Areas of the Cerebral Cortex of Wistar Rats during Focal Ischemia and Its Modulation with Carnosine. *Bulletin of Experimental Biology and Medicine*. 2018; 165: 746–750.
- [73] Jain S, Kim E, Kim D, Burrows D, De Felice M, Kim M, *et al.* Comparative Cerebroprotective Potential of d- and l-Carnosine Following Ischemic Stroke in Mice. *International Journal of Molecular Sciences*. 2020; 21: 3053.
- [74] Keskin A, Kanar BG, Kanar RG, Kepez A, Tunerir B. An investigation on the impact of carnosine on the myocardium in lower extremity ischemia-reperfusion injury in rats. *International Journal of the Cardiovascular Academy*. 2017; 3: 109–113.
- [75] Menini S, Iacobini C, Ricci C, Scipioni A, Blasetti Fantauzzi C, Giaccari A, *et al.* D-Carnosine octylester attenuates atherosclerosis and renal disease in ApoE null mice fed a Western diet through reduction of carbonyl stress and inflammation. *British Journal of Pharmacology*. 2012; 166: 1344–1356.

- [76] Stefani GP, Capalunga L, da Silva LR, Heck TG, Frizzo MN, Sulzbacher LM, *et al.* Effects of aerobic and resistance exercise training associated with carnosine precursor supplementation on maximal strength and $\dot{V}O_{2max}$ in rats with heart failure. *Life Sciences*. 2021; 282: 119816.
- [77] Zeb A, Cha JH, Noh AR, Qureshi OS, Kim KW, Choe YH, *et al.* Neuroprotective effects of carnosine-loaded elastic liposomes in cerebral ischemia rat model. *Journal of Pharmaceutical Investigation*. 2020; 50: 373–381.
- [78] Zhang X, Xia L, Huang Z, Xu F. The effects of carnosine pretreatment on the inflammatory response and the PI3K/Akt signaling pathway following hypoxia-ischemia in neonatal rats. *bioRxiv*. 2019: 766063.
- [79] Ahshin-Majid S, Zamani S, Kiamari T, Kiasalari Z, Baluchnejadmojarad T, Roghani M. Carnosine ameliorates cognitive deficits in streptozotocin-induced diabetic rats: Possible involved mechanisms. *Peptides*. 2016; 86: 102–111.
- [80] Peters V, Riedl E, Braunagel M, Höger S, Hauske S, Pfister F, *et al.* Carnosine treatment in combination with ACE inhibition in diabetic rats. *Regulatory Peptides*. 2014; 194–195: 36–40.
- [81] Abdelkader H, Longman M, Alany RG, Pierscionek B. On the Anticataractogenic Effects of L-Carnosine: Is It Best Described as an Antioxidant, Metal-Chelating Agent or Glycation Inhibitor? *Oxidative Medicine and Cellular Longevity*. 2016; 2016: 3240261.
- [82] Albrecht T, Schilperoort M, Zhang S, Braun JD, Qiu J, Rodriguez A, *et al.* Carnosine Attenuates the Development of both Type 2 Diabetes and Diabetic Nephropathy in BTBR ob/ob Mice. *Scientific Reports*. 2017; 7: 44492.
- [83] Ansurudeen I, Sunkari VG, Grünler J, Peters V, Schmitt CP, Catrina S, *et al.* Carnosine enhances diabetic wound healing in the db/db mouse model of type 2 diabetes. *Amino Acids*. 2012; 43: 127–134.
- [84] Brown JM, Baker LS, Seroogy KB, Genter MB. Intranasal Carnosine Mitigates α -Synuclein Pathology and Motor Dysfunction in the Thy1-aSyn Mouse Model of Parkinson's Disease. *ACS Chemical Neuroscience*. 2021; 12: 2347–2359.
- [85] Colín-Barenque L, Bizarro-Nevarés P, González Villalva A, Pedraza-Chaverri J, Medina-Campos ON, Jimenez-Martínez R, *et al.* Neuroprotective effect of carnosine in the olfactory bulb after vanadium inhalation in a mouse model. *International Journal of Experimental Pathology*. 2018; 99: 180–188.
- [86] Ding Q, Tanigawa K, Kaneko J, Totsuka M, Katakura Y, Imabayashi E, *et al.* Anserine/Carnosine Supplementation Preserves Blood Flow in the Prefrontal Brain of Elderly People Carrying APOE e4. *Aging and Disease*. 2018; 9: 334–345.
- [87] Herculano B, Tamura M, Ohba A, Shimatani M, Kutsuna N, Hisatsune T. β -alanyl-L-histidine rescues cognitive deficits caused by feeding a high fat diet in a transgenic mouse model of Alzheimer's disease. *Journal of Alzheimer's Disease*. 2013; 33: 983–997.
- [88] Tsuji T, Furuhashi K, Gerasimenko M, Shabalova A, Cherepanov SM, Minami K, *et al.* Oral Supplementation with L-Carnosine Attenuates Social Recognition Deficits in CD157KO Mice via Oxytocin Release. *Nutrients*. 2022; 14: 803.
- [89] Chmielewska K, Dzierzbicka K, Inkielewicz-Stepniak I, Przybyłowska M. Therapeutic Potential of Carnosine and Its Derivatives in the Treatment of Human Diseases. *Chemical Research in Toxicology*. 2020; 33: 1561–1578.
- [90] Mahmood A, FitzGerald AJ, Marchbank T, Ntatsaki E, Murray D, Ghosh S, *et al.* Zinc carnosine, a health food supplement that stabilises small bowel integrity and stimulates gut repair processes. *Gut*. 2007; 56: 168–175.
- [91] Caruso G, Godos J, Castellano S, Micek A, Murabito P, Galvano F, *et al.* The Therapeutic Potential of Carnosine/Anserine Supplementation against Cognitive Decline: A Systematic Review with Meta-Analysis. *Biomedicines*. 2021; 9: 253.

Effect of L-Carnosine in Patients with Age-Related Diseases: A Systematic Review and Meta-Analysis

ORIGINALITY REPORT

16%

SIMILARITY INDEX

14%

INTERNET SOURCES

10%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	www.ncbi.nlm.nih.gov Internet Source	2%
2	forum.schizophrenia.com Internet Source	1%
3	www.sciencedirect.com Internet Source	1%
4	Alireza Ghajar, Faezeh Gholamian, Maryam Tabatabaei-Motlagh, Mohsen Afarideh et al. "Citicoline (CDP-choline) add-on therapy to risperidone for treatment of negative symptoms in patients with stable schizophrenia: A double-blind, randomized placebo-controlled trial", Human Psychopharmacology: Clinical and Experimental, 2018 Publication	1%
5	www.j-alz.com Internet Source	1%
6	www.researchsquare.com Internet Source	

1 %

7 Yuko Horii, Jiao Shen, Yoshiyuki Fujisaki, Kokichi Yoshida, Katsuya Nagai. "Effects of l-carnosine on splenic sympathetic nerve activity and tumor proliferation", Neuroscience Letters, 2012
Publication

1 %

8 www.nice.org.uk
Internet Source

<1 %

9 Kevin Peters. "Neuropsychological Subgroups of Cognitively-Impaired-Not-Demented (CIND) Individuals: Delineation, Reliability, and Predictive Validity", Journal of Clinical and Experimental Neuropsychology, 2/23/2005
Publication

<1 %

10 theses.gla.ac.uk
Internet Source

<1 %

11 pure.rug.nl
Internet Source

<1 %

12 spinesection.org
Internet Source

<1 %

13 Verena Peters, Johannes Zschocke, Claus P. Schmitt. "Carnosinase, diabetes mellitus and the potential relevance of carnosinase

<1 %

deficiency", Journal of Inherited Metabolic Disease, 2017

Publication

14

info.elitelearning.com

Internet Source

<1 %

15

malariajournal.biomedcentral.com

Internet Source

<1 %

16

A Mahmood. "Zinc carnosine, a health food supplement that stabilises small bowel integrity and stimulates gut repair processes", *Gut*, 2/1/2007

Publication

<1 %

17

"Accepted Abstracts from the International Brain Injury Association's Ninth World Congress on Brain Injury", *Brain Injury*, 2012

Publication

<1 %

18

"Exercise-based cardiac rehabilitation for coronary heart disease", *Cochrane Database of Systematic Reviews*, 2016.

Publication

<1 %

19

daneshyari.com

Internet Source

<1 %

20

www.naturalhealthresearch.org

Internet Source

<1 %

21

doaj.org

Internet Source

<1 %

22	onlinelibrary.wiley.com Internet Source	<1 %
23	"Posters", <i>Annals of Oncology</i> , 06/01/2011 Publication	<1 %
24	discovery.ucl.ac.uk Internet Source	<1 %
25	dalspace.library.dal.ca Internet Source	<1 %
26	etheses.bham.ac.uk Internet Source	<1 %
27	www.emjreviews.com Internet Source	<1 %
28	Anne Sicard, Jean-Louis Zeddou, Michel Yvon, Yannis Michalakis, Serafin Gutiérrez, Stéphane Blanc. "Circulative Nonpropagative Aphid Transmission of Nanoviruses: an Oversimplified View", <i>Journal of Virology</i> , 2015 Publication	<1 %
29	Sarah Jeffrey, Punitha Isaac Samraj, Behin Sundara Raj. "Therapeutic Benefits of Alpha-Lipoic Acid Supplementation in Diabetes Mellitus: A Narrative Review", <i>Journal of Dietary Supplements</i> , 2021 Publication	<1 %
30	doi.org Internet Source	<1 %

31

www.isrctn.com

Internet Source

<1 %

32

Abolfazl Avan, Tjeerd J. Postma, Cecilia Ceresa, Amir Avan, Guido Cavaletti, Elisa Giovannetti, Godefridus J. Peters. "Platinum - Induced Neurotoxicity and Preventive Strategies: Past, Present, and Future", *The Oncologist*, 2015

Publication

<1 %

33

Christian Keinki, Annika Momberg, Katrin Clauß, Gültekin Bozkurt et al. "Effect of question prompt lists for cancer patients on communication and mental health outcomes —A systematic review", *Patient Education and Counseling*, 2021

Publication

<1 %

34

Hideyuki Hayashi, Yuming Ning, Paul Kurlansky, Anna Vaynrub, Matthew Bacchetta, Erika B. Rosenzweig, Koji Takeda. "Characteristics and prognostic significance of right heart remodeling and tricuspid regurgitation after pulmonary endarterectomy", *The Journal of Thoracic and Cardiovascular Surgery*, 2022

Publication

<1 %

35

William E. Mansbach, Elizabeth E. MacDougall. "Development and validation of the short

<1 %

form of the Brief Cognitive Assessment Tool (BCAT-SF)", Aging & Mental Health, 2012

Publication

36	psychiatriapolska.pl Internet Source	<1 %
37	www.bioinfopublication.org Internet Source	<1 %
38	www.mif-ua.com Internet Source	<1 %
39	www.science.gov Internet Source	<1 %
40	K.N. Roy Chengappa, Scott R. Turkin, Susan DeSanti, Christopher R. Bowie et al. "A preliminary, randomized, double-blind, placebo-controlled trial of l-carnosine to improve cognition in schizophrenia", Schizophrenia Research, 2012 Publication	<1 %
41	bmcmusculoskeletdisord.biomedcentral.com Internet Source	<1 %
42	core.ac.uk Internet Source	<1 %
43	ecommons.aku.edu Internet Source	<1 %
44	silo.pub Internet Source	<1 %

45	www.mdpi.com Internet Source	<1 %
46	www.medrxiv.org Internet Source	<1 %
47	Hu, J.. "Health-related quality of life in individuals with chronic obstructive pulmonary disease", Heart & Lung - The Journal of Acute and Critical Care, 200511/12 Publication	<1 %
48	Mohamad Fawzi Mahomoodally, Sanaa Dilmar A., S. Khatoon Khadaroo. "Carnosine", Elsevier BV, 2022 Publication	<1 %
49	docplayer.fi Internet Source	<1 %
50	forskning.ruc.dk Internet Source	<1 %
51	pubag.nal.usda.gov Internet Source	<1 %
52	Estifanos Baye, Kirthi Menon, Maximilian PJ de Courten, Arul Earnest, James Cameron, Barbora de Courten. "Does supplementation with carnosine improve cardiometabolic health and cognitive function in patients with pre-diabetes and type 2 diabetes? study	<1 %

protocol for a randomised, double-blind,
placebo-controlled trial", BMJ Open, 2017

Publication

53

api.bmrb.io

Internet Source

<1 %

54

minerva.usc.es

Internet Source

<1 %

55

worldwidescience.org

Internet Source

<1 %

56

Nam Y. Lee, Gerard C. Blobe. "The Interaction of Endoglin with β -Arrestin2 Regulates Transforming Growth Factor- β -mediated ERK Activation and Migration in Endothelial Cells", Journal of Biological Chemistry, 2007

Publication

<1 %

57

bsdwebstorage.blob.core.windows.net

Internet Source

<1 %

58

clinicaltrials.gov

Internet Source

<1 %

59

coek.info

Internet Source

<1 %

60

hdl.handle.net

Internet Source

<1 %

61

journals.viamedica.pl

Internet Source

<1 %

62	publications.aap.org Internet Source	<1 %
63	www.annalsmedres.org Internet Source	<1 %
64	www.lifeextension.com Internet Source	<1 %
65	www.omicsonline.org Internet Source	<1 %
66	"27th Annual Meeting Chicago, Illinois May 12-15, 2004", <i>Journal of General Internal Medicine</i> , 01/2004 Publication	<1 %
67	"Therapist-supported Internet cognitive behavioural therapy for anxiety disorders in adults", <i>Cochrane Database of Systematic Reviews</i> , 2016. Publication	<1 %
68	Audrey Baguet, Jan Bourgois, Lander Vanhee, Eric Achten, Wim Derave. "Important role of muscle carnosine in rowing performance", <i>Journal of Applied Physiology</i> , 2010 Publication	<1 %
69	cdn-links.lww.com Internet Source	<1 %
70	cris.maastrichtuniversity.nl Internet Source	<1 %

71	eprints.lancs.ac.uk Internet Source	<1 %
72	hrcak.srce.hr Internet Source	<1 %
73	old.viamedica.pl Internet Source	<1 %
74	www.atsjournals.org Internet Source	<1 %
75	www.jonbarron.org Internet Source	<1 %
76	www.thieme-connect.com Internet Source	<1 %
77	Ai Bo, Brenna Durand, Yi Wang. "A scoping review of parent-involved ethnic and racial socialization programs", Children and Youth Services Review, 2023 Publication	<1 %
78	Maryam Noroozian, Sina Ghasemi, Seyed-Mohammad-Reza Hosseini, Amirhossein Modabbernia et al. "A placebo-controlled study of tropisetron added to risperidone for the treatment of negative symptoms in chronic and stable schizophrenia", Psychopharmacology, 2013 Publication	<1 %

79 Peter P. Toth. "When high is low: Raising low levels of high-density lipoprotein cholesterol", *Current Cardiology Reports*, 11/2008
Publication <1 %

80 academic.oup.com
Internet Source <1 %

81 bmcgastroenterol.biomedcentral.com
Internet Source <1 %

82 content.iospress.com
Internet Source <1 %

83 de Courten, Barbora, Michaela Jakubova, Maximilian PJ de Courten, Ivica Just Kukurova, Silvia Vallova, Patrik Krumpolec, Ladislav Valkovic, Timea Kurdiova, Davide Garzon, Silvia Barbaresi, Helena J. Teede, Wim Derave, Martin Krssak, Giancarlo Aldini, Jozef Ukropec, and Barbara Ukropcova. "Effects of carnosine supplementation on glucose metabolism: Pilot clinical trial : Carnosine Supplementation and Insulin Resistance", *Obesity*, 2016.
Publication <1 %

84 espace.library.uq.edu.au
Internet Source <1 %

85 link.springer.com
Internet Source <1 %

86 lup.lub.lu.se

Internet Source

<1 %

87 pubmed.ncbi.nlm.nih.gov
Internet Source

<1 %

88 www.dovepress.com
Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Effect of L-Carnosine in Patients with Age-Related Diseases: A Systematic Review and Meta-Analysis

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14

PAGE 15

PAGE 16

PAGE 17

PAGE 18

PAGE 19

