

Illuminating the Molecular Intricacies of Exosomes and ncRNAs in Cardiovascular Diseases: Prospective Therapeutic and Biomarker Potential

by Chrismawan Ardianto

Submission date: 21-Feb-2023 04:28PM (UTC+0800)

Submission ID: 2019503473

File name: lar_Diseases_Prospective_Therapeutic_and_Biomarker_Potential.pdf (1.66M)

Word count: 12804

Character count: 76007

Review

Illuminating the Molecular Intricacies of Exosomes and ncRNAs in Cardiovascular Diseases: Prospective Therapeutic and Biomarker Potential

Farheen Badrealam Khan ^{1,*}, Shahab Uddin ^{2,3}, Abozer Y. Elderderly ⁴, Khang Wen Goh ⁵, Long Chiau Ming ^{6,7}, Chrismawan Ardianto ^{7,*}, Abdul Rasheed Palakot ¹, Irfa Anwar ¹, Mohsina Khan ⁸, Mohammad Owais ⁹, Chih-Yang Huang ^{10,11,12,13,14}, Jayasimha Rayalu Daddam ¹⁵, Meraj Alam Khan ¹⁶, Shoaib Shoaib ¹⁷, Md Khursheed ¹⁸, Sara Reshadat ¹⁹, Hamid Reza Khayat Kashani ²⁰, Sameer Mirza ²¹, Abbas A. Khaleel ²¹ and Mohammed Akli Ayoub ^{1,22,23,*}



Citation: Khan, F.B.; Uddin, S.; Elderderly, A.Y.; Goh, K.W.; Ming, L.C.; Ardianto, C.; Palakot, A.R.; Anwar, I.; Khan, M.; Owais, M.; et al. Illuminating the Molecular Intricacies of Exosomes and ncRNAs in Cardiovascular Diseases: Prospective Therapeutic and Biomarker Potential. *Cells* **2022**, *11*, 3664. <https://doi.org/10.3390/cells11223664>

Academic Editors: Y-h. Taguchi and Hsiuying Wang

Received: 11 September 2022

Accepted: 1 November 2022

Published: 18 November 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

- ¹ Department of Biology, College of Science, The United Arab Emirates University, Al Ain 15551, United Arab Emirates
 - ² Translational Research Institute, Academic Health System, Hamad Medical Corporation, Doha 3050, Qatar
 - ³ Dermatology Institute, Academic Health System, Hamad Medical Corporation, Doha 3050, Qatar
 - ⁴ Department of Clinical Laboratory Sciences, College of Applied Medical Sciences, Jouf University, Sakaka 72388, Saudi Arabia
 - ⁵ Faculty of Data Sciences and Information Technology, INTI International University, Nilai 78100, Malaysia
 - ⁶ PAPRSB Institute of Health Sciences, Universiti Brunei Darussalam, Gadong BE1410, Brunei
 - ⁷ Department of Pharmacy Practice, Faculty of Pharmacy, Universitas Airlangga, Surabaya 60115, Indonesia
 - ⁸ Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA
 - ⁹ Interdisciplinary Biotechnology Unit, Aligarh Muslim University, Aligarh 202002, UP, India
 - ¹⁰ Department of Biotechnology, Asia University, Taichung 404, Taiwan
 - ¹¹ Graduate Institute of Biomedical Sciences, China Medical University, Taichung 404, Taiwan
 - ¹² Cardiovascular and Mitochondrial Related Disease Research Center, Hualien Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation, Hualien 970, Taiwan
 - ¹³ Centre of General Education, Buddhist Tzu Chi Medical Foundation, Tzu Chi University of Science and Technology, Hualien 970, Taiwan
 - ¹⁴ Department of Medical Research, China Medical University Hospital, China Medical University, Taichung 404, Taiwan
 - ¹⁵ Department of Ruminant Science, Institute of Animal Sciences, Agriculture Research Organization, Volcani Center, Rishon LeZion 7505101, Israel
 - ¹⁶ Program in Translational Medicine, Peter Gilgan Centre for Research and Learning, The Hospital for Sick Children & DigiBiomics Inc, Toronto, ON M51X8, Canada
 - ¹⁷ Department Biochemistry, Jawaharlal Nehru Medical College, Faculty of Medicine, Aligarh Muslim University, Aligarh 202002, UP, India
 - ¹⁸ College of Medicine, Mohammed Bin Rashid University of Medicine and Health Sciences, Dubai 505055, United Arab Emirates
 - ¹⁹ Department of Internal Medicine, Semnan University of Medical Sciences, Semnan 3513119111, Iran
 - ²⁰ Imam Hossein Hospital, Madani Street, Tehran 1617763141, Iran
 - ²¹ Department of Chemistry, United Arab Emirates University, Al Ain 15551, United Arab Emirates
 - ²² Zayed Center for Health Sciences, United Arab Emirates University, Al Ain 15551, United Arab Emirates
 - ²³ Department of Biology, College of Arts and Sciences, Khalifa University, Abu Dhabi 127788, United Arab Emirates
- * Correspondence: farheen_k@uaeu.ac.ae (F.B.K.); mayoub@uaeu.ac.ae (M.A.A.); chrisawan-a@ff.unair.ac.id (C.A.)

Abstract: Cardiovascular diseases (CVDs) are one of the leading causes of death worldwide. Accumulating evidences have highlighted the importance of exosomes and non-coding RNAs (ncRNAs) in cardiac physiology and pathology. It is in general consensus that exosomes and ncRNAs play a crucial role in the maintenance of normal cellular function; and interestingly it is envisaged that their potential as prospective therapeutic candidates and biomarkers are increasing rapidly. Considering all these aspects, this review provides a comprehensive overview of the recent understanding of exosomes and ncRNAs in CVDs. We provide a great deal of discussion regarding their role in the cardiovascular system, together with providing a glimpse of ideas regarding strategies exploited to harness their potential as a therapeutic intervention and prospective biomarker against CVDs.

Thus, it could be envisaged that a thorough understanding of the intricacies related to exosomes and ncRNA would seemingly allow their full exploration and may lead clinical settings to become a reality in near future.

Keywords: cardiovascular disease; exosomes; ncRNA; miRNA; lncRNA; circRNA

1. Introduction

Cardiovascular diseases (CVDs) represent one of the major causes of death annually and poses a serious burden to the healthcare sector of the society. The World Health Organization estimates that the number of people succumbing to CVDs may cross almost 25 million by 2030 [1]. With the advancements in healthcare systems and infrastructure, the quality of life of CVDs patients has improved substantially. Nevertheless, despite such interventions, the prevalence of heart failure (HF) still remains relatively high. As a matter of fact, cardiac tissues are composed of different types of cells which work in perfect harmony with each other owing to various delicate inter- and intra-cellular communication systems between these cells. This homeostasis is basically achieved through regulated orchestration of various signaling pathways involving autocrine, paracrine, and endocrine release of chemicals/mediators in a feedback loop system. Nevertheless, when this homeostasis is perturbed, pathological conditions are inevitable, and CVDs represent such a multifaceted phenomenon with wide range of pathologies. Accumulating evidences have highlighted the importance of exosomes and non-coding RNAs (ncRNAs) in cardiac physiology and pathology [2–4]. It is widely accepted that exosomes and ncRNAs play crucial role in maintenance of the normal cellular function and their potential as prospective biomarkers and therapeutic candidates are rapidly increasing. Considering all these aspects in mind, this review collates a comprehensive overview of the recent understanding of exosomes and ncRNAs in CVDs with special converge on hypertension induced cardiac complication. We provide a great deal of discussion regarding their role in cardiovascular system together with providing a glimpse of ideas regarding strategies exploited to harness their potential as therapeutic intervention and prospective biomarker against CVDs.

1.1. General Introduction of Exosomes

Extracellular vesicles (EVs) are membranous lipid assemblies, which carries a variety of cellular cargo including lipids, proteins, nucleic acids, metabolites, and so on [5]. Generally, these EVs are categorized based on their size and the nature of their biogenesis [6]; nevertheless, there is some overlap within this nomenclature leading to some contradiction [7]. As of yet, there are no set rules to fully categorize EVs. As a result, the International Society of Extracellular Vesicles has advocated the generic term “EVs” for the vesicles released from the cell [8]. Nevertheless, broadly speaking, there are two major classes namely microvesicles (MVs) and exosomes. MVs are also known as ectosomes, microparticles, or shedding vesicles, are vesicles having size ranging from ~100–1000 nm and are formed from the outward budding of the plasma membrane [9,10]; whereas, exosomes are the vesicles ranging from ~40 to 120 nm and are formed through a complex process that involves inward budding of endosomes [10–12]. Since the discovery of EVs, intensive research has been on-going; nevertheless, as of yet the biology of these EVs especially exosomes are not completely understood. It has been envisaged that exosomes are virtually being released from almost every cell type and they basically facilitate transport of various molecular entities, including nucleic acids, proteins, lipids, and metabolites, both locally and systemically [5,13–17]. Research in the frontiers of exosomes are rapidly increasing; basically a PubMed search with the keyword “exosomes” shows more than thousands of literature been published on the subject, highlighting their importance in the present scenario. Accumulating evidences have ascertained their imperative role in the context of cardiovascular physiology and pathology [18–20]. The origin and evolutionary

perspective of exosomes and their primordial origin remains enigmatic and understanding of its plausible relation with single celled organism also remains relatively obscure. Exosomes which were once thought to be merely associated with the recycling machinery of the cell, playing role in cellular homeostasis, have undergone pragmatic shift in the field of translational medicine. They are released from wide spectrum of cells, including immune cells such as B cells, T cells, dendritic cells and stem cells, and are present in various biological fluids, such as cerebrospinal fluid, serum, saliva, urine, etc. Evidence has shown that exosomes are mechanistically and functionally diverse from its canonical counterpart and are also more heterogeneous, depending upon its origin [21]. Persistent to its endosomal origin, studies have shown the presence of major lipid rafts components consisting of ceramide, cholesterol, sphingomyelin, phosphoglycerides, long and saturated fatty-acyl chains, etc., in the exosomes. Additionally, since exosomes and multivesicular bodies (MVBs) generally originate with the aid of endosomal sorting complex required for transport (ESCRT) pathway, the proteins related to ESCRT are very prevalent and, in fact, many of them, such as HSP70, HSP90, TSG101, Alix, and tetraspanin family proteins, are considered “signature proteins” of exosomes. This, however, does not imply the absence of any other proteins since exosomes can also arise independent of classical ESCRT pathway and also it is to be noted that they act as a carrier for various protein molecules; thus, their protein profile seems to be wide and varied depending on the conditions. Recent studies have highlighted the importance of membrane proteins in the exosomes which can be leveraged to understand their origin, their preferred cellular destination and pathology of diseased state [21]. In addition to lipids and proteins, exosomes also comprise nucleic acid molecules, including mRNA, miRNA, lncRNA, circRNA, etc., as discussed below. A representative figure highlighting the biogenesis of exosomes and the typical structure of exosomes are presented in Figure 1. As a matter of fact, it is in general consensus that once these exosomes are secreted from the parent cell, they interact with the recipient/responder cells through various mechanism including clathrin-mediated endocytosis, lipid-raft mediated, and/or caveolin-mediated endocytosis, receptor-ligand mediated internalization, phagocytosis or micropinocytosis, and/or direct fusion with the plasma membrane. Lately, it has been evident that these pathways are not mutually explicit and plausibly could co-exist for the internalization of a same population of exosomes [10,22]. For example, Isabella et al., 2009 showed exosome uptake by melanoma cells through the plasma membrane fusion [23]. Similarly, another study identified exosome uptake in neurosecretory PC12 through clathrin-mediated endocytosis [24]. Perhaps, through these mechanisms, these exosomal particles modulate the activity of the recipient cells. The mechanism of exosome uptake is shown in classical cellular cargo transport physiology [10,25]. Further, it has been envisaged that the mode and level of internalization of exosomes by different cells varies widely depending on the cell type and environmental conditions. Unfortunately, but not surprisingly, it has been highlighted that the uptake of exosomes is highest in fibroblast cells and least in cardiomyocytes. Nevertheless, the underlying intricacies regulating exosomal targeting/internalization by cardiomyocyte still remains incompletely understood. Interestingly, Eguchi et al., highlighted that stem cell-derived exosomes containing the anti-apoptotic miRNA-214 are up-taken by the cardiac cells through clathrin-mediated endocytosis [26]. With paucity in the literature underlying molecular intricacies in exosomal internalization and interaction in cardiac cells, not much could be ascertained in the present scenario. Albeit certain speculations could be made based on the understanding obtained from reports on exosomes cell interaction with other cell types. It is envisaged that alteration in its profile gives plethora of information in relation to perturbation in the physiological homeostasis of the body. Interestingly, multiple lines of studies have shown that exosomes with their signature molecules plausibly act as an excellent and minimally invasive biomarker for diagnosis and prognosis of various diseases in general and CVDs in particular. To this end, much literature reviews are available highlighting the potential of exosomal signatures molecules as intriguing biomarkers for variety of pathological conditions including CVDs [27–29].




Figure 1. Representative figure highlighting the biogenesis of exosomes (A) and the typical structure of exosomes (B). Basically, exosome biogenesis starts with the inward vagination of the cellular membrane to form early endosomes. Thereafter, the intraluminal vesicles (ILVs) are formed, and the endosomes mature to multivesicular bodies (MVBs). MVBs fuse with the cellular membrane to release ILVs into the extracellular space, where thereafter they are denoted as exosomes. On the other hand, these MVBs can fuse with lysosomes of the cell, resulting in the degradation of ILVs (A). Exosomes contain various molecular entities, including nucleic acids (DNA and/or RNA), membrane anchored-proteins, cytosolic proteins, and lipids (B). The figures are prepared with the BioRender Software ([biorender.com](https://www.biorender.com)).

1.1.1. Exosomes in Cardiac Physiology and Pathology

As a matter of fact, exosome-mediated crosstalk amongst various cell types in heart tissues have been highlighted to play crucial role in the maintenance of cardiac homeostasis, as well as in the pathogenesis of cardiac diseases [27,30]. It is well recognized that in response to various stresses, heart tissue undergoes cardiac remodeling and development of cardiac hypertrophy, apoptosis, and fibrotic responses, which eventually contribute to HF [31,32]. Albeit, understanding the molecular intricacies underlying cardiac remodeling is one of the main challenges in cardiovascular medicine. However, it has been highlighted that these responses, in part, involves vesicle-mediated cellular cross talk among cardiomyocytes and other cells in the myocardium [33,34]. Reports have shown that cardiac cells under stress have increased secretion of exosomes and the exosomal content/composition are also altered; all these aspects eventually activate or suppress various molecular signaling in the recipient cells [30,35]. Interestingly, Lyu and group have highlighted that cardiac fibroblast (CF)-derived exosomes enhanced Renin–Angiotensin System (RAS) signaling in cardiomyocytes; and it was found that attenuation of these exosome secretion considerably reversed Angiotensin II-induced cardiac injuries [36]. Similarly, researchers have highlighted that CF-derived exosomes, which were plausibly enriched with miRNAs, ensues in induction of hypertrophic responses[37]; whereas Yang et al., highlighted that exosomes derived from cardiomyocytes ensued in cardiac fibrosis through myocyte–fibroblast cross-talk [38]. Li and group has shown that plasma exosomal seemingly regulates inflammatory responses

during cardiopulmonary bypass surgery through plausible involvement of miR-223 [39]. These studies explicitly highlighted the importance of exosomes in cardiac homeostasis and disease biology. In addition to playing an imperative role in maintaining cardiac homeostasis and pathophysiology; they have been highlighted to endow with potentials to revolutionize cell based therapeutic intervention against CVDs by being a potential means of cell free therapeutic strategy [40]. Accordingly, in the subsequent section, newer area into the exploration of exosomes as cell free therapeutic intervention, intriguing drug delivery platform, and novel biomarkers for CVDs had been discussed.

1.1.2. Exosomes-Based Therapeutic Interventions against CVDs

Over the years, efforts have continuously been laid down to develop effective therapeutic strategies that would certainly improve the quality of the CVDs clientele. Newer therapeutic strategies are being developed, focusing not only to protect the heart tissue but also to regenerate the myocardium. To this end, accumulating evidence has highlighted the potential of stem cell therapies against CVDs; nevertheless, as of yet, these therapies refrain from showing promising results in clinical trials. Meanwhile, it has been envisaged that most of the favorable outcomes of the transplanted cells were usually indirect. Reports have highlighted that when mesenchymal stem cells (MSCs) were injected in animal model, only 6% of the injected cells were finally being retained in the infarct site [41]. It has been argued that the transplanted cells may secrete various factors/mediators, including extracellular vesicles (EVs), exosomes, growth factors, etc., that might actually play important role in mediating the beneficial effects of cell therapy. This has reinforced the holistic and emerging view of exosomes as an alternative and viable therapy. Nevertheless, despite many promising studies, the precise mechanism of exosome induced perturbations in the recipient cell still remains poorly understood. Meanwhile, taking note of other aspects, a forward leap in the arena of exosomes-based therapeutic interventions has been development of synthetic exosomes with drug delivery potentials, especially the bio-engineered targeted exosomes as detailed in the subsequent sections. Interestingly, many studies clearly indicated that exosomes in general and engineered exosomes in particular have opened newer frontier in arena of intriguing drug delivery platform and there is a high probably that these strategies may find a prosperous status in biomedical sciences in near future.

(A) Exosomes as Cell-Free Therapeutic Strategies against CVDs

Owing to their various intriguing characteristics, they are increasingly being employed as a means of cell-free therapeutic interventions for myriads of obstinate diseases, including CVDs [40]. Accumulating evidence has reported that exosomes from cardiosphere-derived stem cells (CDCs) have been shown to simulate the therapeutic effects of CDCs to a large extend in animal models of heart disease [42–45]. They have been underscored to modulate cardiomyocyte hypertrophic and apoptotic responses, induce angiogenesis, and stimulate endogenous cardiomyocyte proliferation [46]. Interestingly, Zhu and the group have reported the application of human umbilical cord mesenchymal stem cell (UMSC) derived exosomes against aging related cardiac complications. In their study, the authors have ascertained that UMSC derived exosomes through the release of novel metastasis-associated lung adenocarcinoma transcript 1 (MALAT1) lncRNA suppressed aging-related cardiac complications through subsequent attenuation of NF- κ B/TNF- α signaling cascade [47]. Further, it has been highlighted that exosomes produced by CDCs have been demonstrated to stimulate myocardial regeneration via transportation of miRNA to the cardiac cells [42,44,48]. In addition, Limana and group have demonstrated that exosomal from pericardial fluid considerably improved myocardial performance following myocardial infarction (MI) and has ascertained that exosomal protein clustering, an important mediator of TGF- β signaling, was plausibly responsible for the underlying cardiac protective effects [49]. Interestingly, these discoveries rationalize the use of exosomes as intriguing therapeutic intervention against CVDs.

(B) Bio-Engineered Exosomes as Next-Generation Therapeutic Intervention

As a matter of fact, exosomes have been comprehended as an important cellular communication agent embodying potentials to transport diverse range of molecular entities within the biological system [50,51]. Because of their intrinsic ability to deliver molecular entities, they are considered as a promising drug delivery system (DDS) for various bioactive compounds and small molecular drugs and has been demonstrated to considerably improve their pharmacological properties against various diseases in general, and CVDs in particular. Compared with conventional drug delivery platforms, such as micelles, microemulsion, nanospheres, liposomes, and metallic nano-particulate system; exosomes offer many desirable advantages, such as lower toxicity, lower immunogenicity, high stability in circulation, better biocompatibility, and biological barrier permeability, which makes them attractive platforms for efficient delivery of therapeutic agents. Interestingly, exosomes have been used to deliver therapeutic drug and small molecules to many tissues, including the heart [52–58]. In fact, in recent years, engineered exosomes has been harnessed for targeted co-delivery of chemotherapeutics drug and RNA in fight against various diseases [59]. Nevertheless, exosomes in analogy with other drug delivery platforms also suffer from the drawback of endocytosis by the mononuclear phagocyte system (MPS). It has been highlighted that, when unmodified/neat exosomes were administrated systemically in animal model, they were found preferentially accumulated in the MPS organs such as liver, kidney, and spleen, which, thereafter, were rapidly cleared by bile excretion, renal filtration, and/or were phagocytized, leading to minimal accumulation of the therapeutics in the intended tissues or organs and undue delivery to un-intended tissues [60]. This bio-distribution profile and off-target effects limited the clinical acceptability of the unmodified exosomes [60–62]. Therefore, attempts have been made to modify exosomes for effective targeting to desired tissue. One method that has been harnessed is modification of exosomes with homing ligands or peptides, which confers them targeting capability to tissues or organs carrying the corresponding receptors. In cardiovascular system, several homing ligands/peptides are been explored for targeted therapy [52,63–65]. Moreover, many peptides endowed with homing potential to different cardiovascular systems, such as normal cardiomyocyte, ischemia/reperfusion injured cardiomyocytes, the vascular system etc. offers exciting avenues for exosome targeting ligands [63,64,66–68]. Interestingly, exosomes can be derived from an individual differentiated hematopoietic stem cells (HSC) and used for tissue-targeted cargo delivery through the expression of tissue-specific peptides. Thereafter, by loading miRNA and/or siRNA of the targeted gene, these modified tissue targeted exosomes can selectively regulate gene expression in the specific tissue corresponding to the homing peptides. Interestingly, Vandergriff et al., developed an infarct-targeting exosomes, through the use of cardiac homing peptide (CHP: CSTSMLKAC (IMTP)) to increase the efficacy and decrease the effective dose of intravenously delivered exosomes [63,64]. They basically conjugated cardiac stem cell-derived exosomes with cardiac homing peptide IMTP through a click chemistry approach using dioleoylphosphatidyl ethanolamine N-hydroxy succinimide linker. Interestingly, increased retention of the IMTP-exosomes within the ischemia/reperfusion injured heart tissues were observed to a considerable extent and improvement in cardiac function was also achieved thereof [69]. Similarly, molecular cloning and lentivirus packaging techniques were employed to engineer exosomal enriched membrane protein, i.e., Lamp2b fused with ischemic myocardium-targeting peptide IMTP. Such a fusion resulted in peptides being displayed on the surface of exosomes. Interestingly, these IMTP-exosomes displayed efficient internalization by hypoxia-injured embryonic cardiomyocyte H9c2 cells compared to blank-exosomes and subsequent increased accumulation in ischemic heart tissue were also obtained [65]. Meanwhile, attenuation of the inflammatory, apoptotic, and fibrotic responses was observed and enhanced vasculogenesis, and improved cardiac function were detected following IMTP-exosome treatment in ischemic heart. Further, Mentkowski and Lang bio-engineered a cardiomyocyte targeted exosomes that demonstrated improved cardiac retention in in vivo system [52]. Further, Mentkowski and Lang bio-engineered

a cardiomyocyte targeted exosomes that demonstrated improved cardiac retention in an in vivo system [52]. To this end, the researcher selected a cardiomyocyte-specific peptide (Cardiomyocyte Peptide (CMP): WLSEAGPVVTVRALRGTGS) [63,70]; which has proven ability to specifically target cardiac tissues [53,69,71,72]. The researcher ligated this CMP to the extra-exosomal N-terminus of Lamp2b. Interestingly, these cardiac-targeted CDC exosomes showed improved uptake into cardiac cells in an in vitro model; thereby leading to improved cardiac retention in in vivo system and, eventually, reduced cardiac apoptosis [52]. It has been envisaged that decorating the surfaces of the exosomes with homing ligand/entities will certainly reduce the time exosomes require to reach the therapeutic concentration in targeted tissues, and will considerably reduce the off-target effect, thereby leading to enhanced therapeutic potential. For detailed outline for the generation and isolation of the engineered exosomes; readers are advised to go through various previously published articles [52,59,65,68,69,73]. An overview of procedures for generation of engineered exosomes for specific targeting of the therapeutic molecules to desired tissue along with the workflow of differential ultracentrifugation for the isolation of the exosome are represented in Figure 2.




Figure 2. Representative figure highlighting the procedures for generation of engineered/modified exosomes for specific targeting of the therapeutic molecules to desired tissue (A) along with the workflow of differential ultracentrifugation for exosome isolation (B). The figures are prepared with the BioRender Software ([biorender.com](https://www.biorender.com)).

1.1.3. Exosomes as Prospective Biomarkers for CVDs

Accumulating evidences have shown that exosomes contain diverse biological contents that plausibly is a reflection of a particular state of the system [74]. Along these lines, the vast repertoire of molecular entities that are packaged within exosomes, their versatile appearance in nearly all body fluids marks their potential candidature for prospective novel non-invasive biomarkers [75].

Amongst the exosomes content, exosomes proteins and RNA molecules especially miRNA are increasingly been reported as promising biomarkers [76]. In fact, exosomal miRNAs have been the most studied for their role as novel biomarkers for CVDs. A distinct miRNA profile has been reported by various workers in CVD patients compared to normal individuals. To this end, Matsumoto and group reported that p53-responsive circulating exosomes miRNAs viz. hsa-miR-192, hsa-miR-194 and hsa-miR-34a, were considerably upregulated in the serum of acute MI clientele that have experienced development of HF in short period. This study highlights the importance of these exo-miRNA as plausible prognostic biomarkers for acute MI [77]. Further, studies have also shown that serum exosomal miR-9 and miR-124 levels were significantly higher in stroke patients. Concomitantly, circulating exosomal miR-9 and miR-124 might be promising biomarkers for stroke diagnosis [78]. Further, Gidlof and colleagues have demonstrated that upregulation of plasma levels of hsa-miR-208b and hsa-miR-499-5p corresponded to increase in the risk of HF, highlighting their prognostic biomarker potential [79].

Further, studies have envisaged the importance of various other exosomal proteins for prospective biomarkers for CVDs. To this end, Pironti et al., have reported that circulating exosomes induced by cardiac pressure overload contain functional angiotensin II type 1 receptors (AT₁Rs); they have envisaged that the transfer of AT₁Rs plausibly deteriorates cardiac function during blood pressure (BP) overload, thus, could help in analyzing the prognosis of the pressure overload diseased patients [80]. Similarly, the adenosine 2A receptors and dopamine receptors have also been packed within EVs and transferred to other cells, leading to an increase in BP and cardiac remodeling thereof [81]. These findings seemingly highlight for usage of these exosomal proteins as prognostic biomarkers for hypertension clientele.

Collectively, it is reasonable to argue that many studies are being performed in basic and clinical research to understand the roles of exosomes in CVDs and to explore their prospective therapeutic, drug delivery, and biomarker potential. In parallel, it is also envisaged that albeit exosome therapy for CVDs looks promising and tempting; nevertheless, researchers in this field face numerous problems. Not only lack of thorough knowledge about cellular and molecular intricacies; but also purely technical issues as well. The issue related to low level of endocytosis in cardiomyocytes well describes the situation, and, therefore, it seems really challenging to treat CVDs or other diseases with exosomes that have not been sufficiently modified. Another problem of all the works investigating exosomes is the impossibility of isolating pure exosome preparation. Nonetheless, the research fraternities are highly optimistic and as more and more are gleaned about these aspects, it will be highly helpful in providing scope for improvisation.

1.2. General Introduction of Non-Coding RNAs

It is widely accepted notion that, albeit the human genomes are transcribed into RNA; nevertheless, approximately only 2% of these transcripts have protein coding functions. Reckoning with these, researchers have started to investigate the role of ncRNAs in regulation of various physiological and pathological conditions, including CVDs. In fact, in recent years, the role of ncRNAs in cardiovascular physiology and pathophysiology has become the focus of many research endeavors [82–84]. It is argued that a better understanding of the involvement of ncRNAs in CVDs will offer better comprehension of the underlying intricacies which will certainly aid in novel therapeutic insights [85]. In terms of classification, basically, these ncRNAs are broadly classified based on their size; usually transcripts with nucleotide lengths < 200 nucleotides are considered as small noncoding RNAs (sncRNA);

²¹ for, e.g., microRNA (miRNA), piwi-interacting RNA (piRNA), small nucleolar RNAs (snoRNAs) etc.; whereas transcripts with nucleotide lengths > 200 nucleotides are considered as long noncoding RNAs (lncRNA); for, e.g., lncRNA, which comprises of long intergenic RNA (lincRNA), enhancer RNAs (eRNAs), and sense or antisense transcripts (AS), as discussed below [86–88]. It is in general consensus that the cellular and temporal specificity drives the mechanism of action of ncRNAs. Basically, ncRNAs are found within the nucleus, nucleolus, cytoplasm, and even in the mitochondria. Nevertheless, extracellular ncRNAs were found outside of the cells as well. For example, ncRNA, specifically miRNA, was first observed in the plasma of the esophageal and melanoma cancer patients [89] and later on established as a potential blood biomarker for various cancer diagnoses [90]. Furthermore, ncRNAs can be transported from one cell to another through various means. For example, Valadi et al. showed that miRNA transport through exosome and they termed these RNA components as exosomal shuttle RNA [17]. Interestingly, another study found that miRNA-126 can also be transported between cells through apoptotic bodies [91]. In addition to these, ncRNAs could also be transported through carrier proteins. For example, Kasey et al., demonstrated the stable transfer of functional miRNA through high density lipoprotein (HDL) into the atherogenic mouse model [92]. Interestingly, ncRNAs can be sorted and packaged into the exosome and circulate into the plasma and transported to the recipient cells [93]. Another pioneering study of plasma derived exosomal RNA profiling revealed the presence of various forms of ncRNA, including miRNA as the most abundant form in the blood circulation [94]. Horizontal transfer of ncRNAs from one cell type to another cell type has been recently established as means of intercellular communication. For example, an interesting report showed the exosome mediated miRNA transport from T-cells to the antigen presenting cells (APCs), wherein they modulate the gene expression profile of APCs [95]. Furthermore, these exosomal RNAs are involved in many pathophysiological conditions. Recently, a group showed exosomal associated lncRNAs mediated modulation of the function of l-Lacto-dehydrogenase B (LDHB), high mobility group protein 17 (HMG17) and CSF2RB which causes changes in nucleosomal architecture, and thereof enhances the cell viability [96]. These studies are attracting much attention; nevertheless, at this moment of time, it is reasonable to argue that the detailed intricacies about the exact mode of function and biology of these ncRNAs is still a matter of great interest; and with much concerted efforts from different research fraternities; more and more would be gleaned about these intricacies in near future. A representative table providing the lists of ncRNAs highlighted in various cardiac pathophysiology are depicted in Table 1.

³² 1.2.1. General Introduction of Long Non-Coding RNAs (lncRNAs)

In the recent past, studies have envisaged the importance of lncRNAs in orchestration of various cardiovascular signaling cascades [97–99]. As already mentioned, lncRNAs comprise a subclass of ncRNAs broadly classified as transcripts > 200 nucleotides in length with limited coding functions. Further, depending on its location in the genome and its relative distance from protein encoding genes; they can be classified as sense lncRNAs, antisense lncRNAs, bidirectional lncRNAs, intronic lncRNAs, and intergenic lncRNAs, respectively [100]. Evidence has shown that most of these lncRNAs are nuclear; however, recent studies have shown that they are also present in the cytoplasmic compartment as well. The genetic loci of lncRNAs are quite similar to that of mRNAs, but they show less coherent co-transcriptional splicing and, in addition, they predominantly possess only one intronic region. Further, in general, the expression level of lncRNAs is relatively less but more specific than normal protein coding genes, although some discrepancies can be observed in this depending on the tissue type [101]. Since the expression of lncRNAs are finely regulated, they provide vital clues regarding the developmental stages of the cell and/or disease state. Additionally, they are increasingly becoming popular for their regulatory roles in gene expression, chromatin modification, cellular differentiation besides acting as scaffolds/guides with intriguing spatial control. These functional roles are not very exclusive and many lncRNAs seem to obfuscate this notion and perform more than

43

one function simultaneously. Interestingly but not surprisingly, the advent of newer high throughput technologies, such as RNA-seq, microarray, next-generation sequencing, and advanced transcriptomic technologies together with bioinformatic tools have heralded a new paradigm shift in our understanding of diverse functionalities of lncRNAs. Although the prevalence of such class of RNA has been known since the 1980s, there has been surge of studies showing its increasing novel regulatory functions and its role in disease progression over several decades. Interestingly, an intricate lncRNAs map was developed by Iyer et al., wherein they have characterized lncRNAs from different tissues, cancerous cells, and cell lines [102]. Similarly, Cabili et al., assembled a reference catalogue of lncRNAs from variety of body tissues and cell type [103]. This has also necessitated a comprehensive annotation resource for lncRNAs which would help researchers in better understanding of lncRNAs. These resources include GENCODE (<https://www.genencodegenes.org/>), LNCipedia (<https://lncipedia.org/info>), NONCODE (<http://www.noncode.org/>), TANRIC (<https://www.tanric.org/>), LNCat (<http://biocc.hrbmu.edu.cn/LNCat/>), etc. [104].

(A) lncRNAs in Cardiac Physiology and Pathology

As a matter of fact, many studies have highlighted the role of lncRNAs in regulation of CVDs; studies have envisaged the role of lncRNAs in cardiac remodeling, including cardiac hypertrophy, apoptosis, and fibrotic responses [47,105–113]. To this end, Zhang and group have characterized the intricacies of a lncRNA named cardiac hypertrophy-associated regulator (CHAR) in cardiac hypertrophy and delineated the underlying signaling cascade thereof [105]. Further, several studies have ascertained linkage between lncRNA and miRNA in cardiac injuries. For example, the lncRNA Plscr4 and lncRNA taurine upregulated gene 1 (TUG1) has been shown to regulate cardiac hypertrophy seemingly through regulation of miR-214 and miR-29b-3p, respectively [114,115]. Similarly, lncRNA H19/miR-675 axis has been ascertained to regulate cardiac apoptosis through suppression of VDAC1 in diabetic cardiomyopathy [116]; moreover, lncRNA myocardial infarction-regulatory factor (MIRF), i.e., lncRNA MIRF has been highlighted to promote cardiac apoptosis through regulation of the miR-26a–Bak1 axis [109]. Moreover, lncRNA ANRIL has been shown to regulate myocardial apoptosis through regulation of IL-33/ST2 pathway in acute MI animal model [117]. In addition, lncRNA NONMMUT022555, also known as pro-fibrotic lncRNA, has been reported to play intriguing role in fibrogenesis process plausibly by favoring proliferation of cardiac fibroblasts through modulation of let-7d level in MI mouse model [118]. Further, Micheletti et al., have shown that Wisp2 super enhancer associated RNA (Wisper) was associated with cardiac fibrosis and cardiac dysfunction in a murine model of MI and in aortic stenosis human patients [119]. Moreover, exosomal lncRNA AK139128 derived from cardiomyocytes under hypoxia condition has been reported to induce apoptosis and attenuate cellular proliferation in cardiac fibroblasts [120].

(B) lncRNA as Therapeutic Interventions and Biomarkers in CVDs

Meanwhile, lncRNAs have been attracting lots of attention as a potential therapeutic candidates, as well as a prospective biomarker for CVDs. A recent comprehensive study by Hu and group sheds light on the differential profile of exosomal lncRNA and mRNA in rheumatic heart disease (RHD). Interestingly, it was found that there were almost 231 lncRNA, which were differentially expressed in RHD patients in comparison to healthy clientele. This pioneering transcriptomic analysis of the exosomal lncRNA and mRNA has provided valuable information not only for plausible biomarker for prognosis but also provided insights into intriguing therapeutic targets [121]. Further, a study by Shao et al., 2017 showed that terminal differentiation-induced ncRNA (TINCR) considerably attenuated cardiac hypertrophy through epigenetic regulation of the protein kinase CAMKII in transverse aortic constriction mouse model [122]. Previously, Micheletti and group have shown that silencing Wisper lncRNA through antisense oligonucleotide technology resulted in attenuation of cardiac dysfunction and MI-induced fibrosis in an in vivo model [119].

As a matter of fact, evidences have shown that lncRNAs displays dynamic alteration under pathological conditions and have long term stability in the body fluids [123]. All these features make lncRNAs as a potential non-invasive prognostic and diagnostic biomarker. To this end, Kumarswamy and group have explored the potential of lncRNAs as a prognostic biomarker for HF. Basically, the group have identified mitochondria-derived lncRNA long intergenic non-coding RNA predicting cardiac remodeling (LIPCAR), as a novel biomarker of cardiac remodeling and could certainly predicts future death in patients with HF [123]. In addition, LIPCAR was also ascertained as an intriguing biomarker for diastolic dysfunction and remodeling in type 2 diabetic clientele [124]. Furthermore, Wang and group has revealed augmented plasma levels of LIPCAR and the paternally imprinted lncRNA H19 in clientele with coronary artery disease (CAD) [125]. Further, Xuan et al., 2017 have ascertained the role circulating lncRNAs, i.e., non-coding repressor of NFAT (NRON) and myosin heavy-chain-associated RNA transcripts (MHRT) as novel predictive biomarkers of HF [126].

1.2.2. General Introduction of Circular Non-Coding RNAs (circRNAs)

CircRNA represents another large class of ncRNAs which as the name suggests are circular covalently closed and show spatiotemporal expression pattern in tissues and cells. They carry out their function by acting as RNA-binding proteins, sequestering agents, transcriptional regulators, as well as miRNA sponges. In addition to these, it has been reported that some selected circRNAs are converted into functional proteins as well. Despite the absence of poly-adenylation site and capping region, circRNA localizes to the cytoplasmic compartment and forms a very stable circular structure resistant to exonuclease. Albeit studies related to circRNA are accelerating; nevertheless, information regarding the function of circRNA and the ability of it to regulate the physiological and pathological conditions are relatively in infancy. In addition, most of the studies on circRNA have been carried out with limited size of cohort which results in inconclusive interpretations. Additionally, the lack of standardized procedure for evaluating circRNA has resulted in data inconsistency between different groups. These factors have limited the scope of circRNA. Nevertheless, as with lncRNA, several annotation resources for circRNA have been created which the help of researchers in better comprehension of these molecules. A repertoire of tissue specific circRNA database was created recently by Liu et al. and named generically as circRNA database (<http://circnet.mbc.nctu.edu.tw/>). Several other databases to further accelerate research on circRNA has also been created, such as CircRNABase (<http://starbase.sysu.edu.cn/starbase2/mirCircRNA.php>), circBase (<http://www.circbase.org/>), Circ2Traits (<http://gyanxet-beta.com/circdb/>), CircInteractome (<https://circinteractome.nia.nih.gov/>), etc.

(A) CircRNA in Cardiac Physiology and Pathology

Evidence has shown that circRNA has emerged as regulatory molecule in CVDs. In addition, they are considered as novel biomarkers for CVDs, besides being considered as important therapeutic targets. With the help of high sequencing technologies, recent studies have found plenty of circRNA in heart tissues from human and mouse origin. Interestingly, heart-related circRNA (HRCR) was the pioneer circRNA, which was found to be considerably suppressed in hypertrophic heart and in HF model [127–129]. It has been reported that HRCR acts as a sponge for miR-223, which has been implicated in cardiac hypertrophic responses [128]. Intriguingly, it has also been shown that overexpression of HRCR could provide protection against hypertrophy plausibly through attenuation of miR-223 in a mouse model [127]. Further, whole transcriptome analysis revealed that five circRNA, namely circRNA26, circRNA261, circRNA1191, circRNA4251, and circRNA6913, were differentially expressed following cardiac hypertrophic induced by high glucose treatment. These circRNA was found to have around ~60 target miRNA for regulation [130]. Concomitantly, these differentially expressed circRNA ascertained the biologically relevant RNA markers and corresponding regulatory network in high glucose induced cardiomy-

opathies. Further, Wu et al., using circRNA microarray and in silico analysis ascertained that 59 plasma circRNA were differentially expressed (46 circRNAs were significantly upregulated and 13 were significantly downregulated) in human hypertensive plasma samples. Amongst these differentially expressed circRNA, *has_circ_0005870* was further validated to be considerably downregulated in hypertensive clientele [131]. In another study, researchers have highlighted the role of *circRNA_000203* to promote cardiac hypertrophy plausibly through inhibition of miR-26b-5p and miR-140-3p which regulate Gata4 expression levels [132]. Further, CircRNA microarray studies have ascertain that three circRNA namely chr8:71336875j71337745, chr5:90817794j90827570, and chr6:22033342j22038870 were overexpressed in case of rat coronary artery endothelial cells (CAEC) treated with TGF- β 1 [133]. This study interestingly highlighted the potential role of differentially expressed circRNAs during TGF- β 1-related CVDs.

(B) CircRNA as Therapeutic Interventions and Biomarkers against CVDs

Further, recent studies have exploited circRNA as therapeutic interventions against CVDs. Interestingly, in the case of atherosclerosis, circANRIL has been demonstrated to bestow athero-protection through modulation of ribosomal RNA (rRNA) maturation and governing pathways related to atherogenesis [134]. Similarly, *circRNA_010567* was shown to ameliorate MI through attenuation of TGF- β 1 [135]. Likewise, a promising study by Zeng et al. evaluated the potential of circRNA *circ_Amotl1*, which is highly expressed in neonatal cardiac tissue and manifests cardio-protective functions by binding to PDK1 and AKT1 [136].

As already mentioned, linear RNA molecules have been highlighted as potential biomarkers [137]; to this end, as circRNAs are more stable and resistant to exonucleases compared to linear RNAs, as a result it bestows more advantageous properties for its potential to act as a biomarker [138]. Thus, it is highly reasonable to argue that circRNAs are more superior to its analogous mRNAs and lncRNAs as prospective biomarker candidates in terms of abundance, stability, and specificity. In analogy with lncRNAs, Zhao and group have envisaged the potential of peripheral blood circular RNA *hsa_circ_0124644* as a diagnostic biomarker of CAD [139]. A recent meta-analysis of several databases have demonstrated that two circRNA namely *circCDKN2BAS* and *circMACF1* have prospective potentials to be used as circulating biomarker in CVDs [140]. Furthermore, a study in 2017 highlighted the usage of the circRNA, myocardial infarction associated circular RNA (MICRA) to predict the risk in MI clientele [141]. Further, the circRNA HRCR described above could also be potentially considered for biomarker repository [138,142]. Likewise, *hsa-circ-0005870* described above might represent a novel diagnostic biomarker for hypertension [131].

Collectively, the aspects that they are abundant, stable, as well as evolutionally conserved in tissues, saliva, exosomes, and blood offers enormous potential to extend the current landscape of prognostic and diagnostic biomarkers for CVDs, as well as for other diseases [123,143]. However, it is a matter of great interest that amongst exosomes, ncRNA, and exosomal ncRNA, which one would show better candidature as prospective biomarkers for CVDs is still not known.

Taken together, although each molecular entities viz. exosomes and ncRNA have distinctive role in cardiovascular system; nevertheless, the importance of cross talks between these molecular entities as regulator of various events in cardiovascular system should not be overlooked at the same time [144].

Table 1. Representative table providing the lists of ncRNAs highlighted in various cardiac pathophysiology.

RNA	Disease	Mechanism/Functions	References
<i>LucRNA</i>			
TINCR	Cardiac Hypertrophy	Silencing of CaMKII protein kinase	[122]
Plscr4	Cardiac Hypertrophy/ Heart Failure	Down-regulation of miR-214 expression	[114]
TUG1	Cardiac Hypertrophy	Down-regulation of miR-29b-3p	[115]
CHRF	Cardiac Hypertrophy	Down-regulation of miR-489	[106]
MIAT	Cardiac Hypertrophy	Regulation of TLR4 expression by sponging miR-93 in cardiomyocytes	[107]
FTX	Cardiac Apoptosis	Modulation of Bcl212 expression; Inhibition of miR-29b-1-5p	[108]
MIRF	Cardiac Apoptosis	Inhibition of miR-26a	[109]
ANRIL	Cardiac Apoptosis	Regulation of IL33/ST2	[117]
H19	Cardiac Apoptosis/ Cardiac Fibrosis	Reduction of VDAC-1; Increment in collagen and TGF- β levels, reduction of Dus5 expression	[116,125]
NONMMUT022555	Cardiac Fibrosis	Reduction in level of let-7d	[118]
Wisper	Cardiac Fibrosis	Regulation of expression of a profibrotic form of lysyl hydroxylase 2	[119]
(GAS5)	Cardiac Fibrosis	Inhibition of miR-21; Modulation of endothelial cells via exosomes and macrophage apoptosis	[110,111]
Mhrt	Cardiac Fibrosis	Interaction with the chromatin-remodeling factor Brg1	[126]
SRA1	Cardiac Fibrosis	Inhibition of miR-148b	[112]
MEG3	Myocardial infarction	Regulation miR-183 level	[113]
AK139128	Myocardial infarction	Regulation of cellular activities of cardiac fibroblasts in vitro and in vivo	[120]
MALTA1	Aging induced cardiac dysfunction	Inhibition of NF- κ B/TNF- α signaling pathway	[47]
<i>CircRNA</i>			
HRCR	Cardiac Hypertrophy/ Heart Failure	Acts as sponge for miR-223	[127–129]
CircRNA_0005870	Hypertension	-	[131]
CircRNA_000203	Cardiac Hypertrophy	Inhibition of miR-26b-5p and miR-140-3p which regulate Gata4 expression levels	[132]
CircANRIL	Artherosclerosis	Regulation of ribosomal RNA (rRNA) maturation and modulation of pathways related to atherogenesis	[134]
CircRNA_010567	Myocardial Infarction	Attenuation of TGF- β 1	[135]
CircRNA_Amot1	Doxorubicin induced cardiomyopathy	Binding to PDK1 and AKT1	[136]

2. Conclusions

Since the discovery of exosomes and ncRNAs, they have garnered much attention across the research fraternities; nevertheless, their intricacies, especially in relation with CVDs, are not completely understood. Nonetheless, in recent years, research in these fields

has expanded greatly. It is argued that as the challenges in the field are gradually addressed, it will be highly instrumental to better understand the underlying intricacies regarding their biology and function, especially in CVDs. However, there are still various daunting challenges that are important stumbling blocks to truly harness their potential in clinical settings. These include establishment of optimal dose and route of administration, better understanding of the immunogenicity of these molecular entities upon administration to the model animals, improved understanding of their pharmacokinetics and pharmacodynamic parameters, development/optimization of tools to comprehensively characterize them, etc. At this moment of time, it is reasonable to argue that these challenges need to be addressed on an urgent basis. Accordingly, a better understanding of these intricacies, along with addressing the underlying challenges will provide a fundamental basis for improving their efficacy for improved therapeutic intervention to efficiently deal with not only CVDs but also other debilitating diseases as well with equal potency.

Author Contributions: Conceptualization: F.B.K.; writing—original draft preparation: F.B.K., S.U., A.Y.E., K.W.G., L.C.M., C.A., A.R.P., I.A., M.K. (Mohsina Khan) M.O., C.-Y.H., J.R.D., M.A.K., S.S., M.K. (Md Khursheed), S.R., H.R.K.K., S.M., A.A.K. and M.A.A.; writing—review and editing: F.B.K. All authors have read and agreed to the published version of the manuscript.

Funding: Zayed Center for Health Sciences, United Arab Emirates University (UAEU-ZCHS) (Grant number #31R235) and the UAEU postdoc grant (Grant number #31R241).

Acknowledgments: We acknowledge the funding support by a center-based grant from Zayed Center for Health Sciences, United Arab Emirates University.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Timmis, A.; Mimosza, L.; Vardas, P.; Townsend, N.; Torbica, A.; Katus, H.; De Smedt, D.; Gale, C.P.; Maggioni, A.P.; Petersen, S.E.; et al. European Society of Cardiology: Cardiovascular disease statistics 2021. *Eur. Heart J.* **2022**, *43*, 716–799. [[CrossRef](#)]
2. Li, C.; Ni, Y.-Q.; Xu, H.; Xiang, Q.-Y.; Zhao, Y.; Zhan, J.-K.; He, J.-Y.; Li, S.; Liu, Y.-S. Roles and mechanisms of exosomal non-coding RNAs in human health and diseases. *Signal Transduct. Target. Ther.* **2021**, *6*, 383. [[CrossRef](#)] [[PubMed](#)]
3. Yuan, Z.; Huang, W. New developments in exosomal lncRNAs in cardiovascular diseases. *Front. Cardiovasc. Med.* **2021**, *8*, 709169. [[CrossRef](#)] [[PubMed](#)]
4. Xiao, Y.C.; Wang, W.; Gao, Y.; Li, W.Y.; Tan, X.; Wang, Y.K.; Wang, W.Z. The Peripheral Circulating Exosomal microRNAs Related to Central Inflammation in Chronic Heart Failure. *J. Cardiovasc. Transl. Res.* **2022**, *15*, 500–513. [[CrossRef](#)] [[PubMed](#)]
5. Yokoi, A.; Ochiya, T. Exosomes and extracellular vesicles: Rethinking the essential values in cancer biology. *Semin. Cancer Biol.* **2021**, *74*, 79–91. [[CrossRef](#)] [[PubMed](#)]
6. Yáñez-Mó, M.a.; Siljander, P.R.M.; Andreu, Z.; Zavec, A.B.; Borràs, F.E.; Buzas, E.I.; Buzas, K.; Casal, E.; Cappello, F.; Carvalho, J.; et al. Biological properties of extracellular vesicles and their physiological functions. *J. Extracell. Vesicles* **2015**, *4*, 27066. [[CrossRef](#)]
7. Gould, S.J.; Raposo, G.A. As we wait: Coping with an imperfect nomenclature for extracellular vesicles. *J. Extracell. Vesicles* **2013**, *2*, 20389. [[CrossRef](#)]
8. Théry, C.; Witwer, K.W.; Aikawa, E.; Alcaraz, M.J.; Anderson, J.D.; Andriantsitohaina, R.; Antoniou, A.; Arab, T.; Archer, F.; Atkin-Smith, G.K.; et al. Minimal information for studies of extracellular vesicles 2018 (MISEV2018): A position statement of the International Society for Extracellular Vesicles and update of the MISEV2014 guidelines. *J. Extracell. Vesicles* **2018**, *7*, 1535750. [[CrossRef](#)]
9. Cocucci, E.; Racchetti, G.; Meldolesi, J. Shedding microvesicles: Artefacts no more. *Trends Cell Biol.* **2009**, *19*, 43–51. [[CrossRef](#)]
10. Gurung, S.; Perocheau, D.; Touramanidou, L.; Baruteau, J. The exosome journey: From biogenesis to uptake and intracellular signalling. *Cell Commun. Signal.* **2021**, *19*, 1–19. [[CrossRef](#)]
11. French, K.C.; Antonyak, M.A.; Cerione, R.A. Extracellular vesicle docking at the cellular port: Extracellular vesicle binding and uptake. *Semin. Cell Dev. Biol.* **2017**, *67*, 48–55. [[CrossRef](#)] [[PubMed](#)]
12. Johnstone, R.M.; Adam, M.; Hammond, J.R.; Orr, L.; Turbide, C. Vesicle formation during reticulocyte maturation. Association of plasma membrane activities with released vesicles (exosomes). *J. Biol. Chem.* **1987**, *262*, 9412–9420. [[CrossRef](#)]
13. Safdar, A.; Saleem, A.; Tarnopolsky, M.A. The potential of endurance exercise-derived exosomes to treat metabolic diseases. *Nat. Rev. Endocrinol.* **2016**, *12*, 504–517. [[CrossRef](#)] [[PubMed](#)]
14. Martínez, M.C.; Andriantsitohaina, R. Extracellular vesicles in metabolic syndrome. *Circ. Res.* **2017**, *120*, 1674–1686. [[CrossRef](#)]
15. Théry, C.; Zitvogel, L.; Amigorena, S. Exosomes: Composition, biogenesis and function. *Nat. Rev. Immunol.* **2002**, *2*, 569–579. [[CrossRef](#)]

16. Huang-Doran, I.; Zhang, C.-Y.; Vidal-Puig, A. Extracellular vesicles: Novel mediators of cell communication in metabolic disease. *Trends Endocrinol. Metab.* **2017**, *28*, 3–18. [[CrossRef](#)]
17. Valadi, H.; Ekström, K.; Bossios, A.; Sjöstrand, M.; Lee, J.J.; Lötvall, J.O. Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat. Cell Biol.* **2007**, *9*, 654–659. [[CrossRef](#)]
18. Ibrahim, A.; Marbán, E. Exosomes: Fundamental biology and roles in cardiovascular physiology. *Annu. Rev. Physiol.* **2016**, *78*, 67. [[CrossRef](#)]
19. Patil, M.; Henderson, J.; Luong, H.; Annamalai, D.; Sreejit, G.; Krishnamurthy, P. The art of intercellular wireless communications: Exosomes in heart disease and therapy. *Front. Cell Dev. Biol.* **2019**, *7*, 315. [[CrossRef](#)]
20. Kishore, R.; Garikipati, V.N.S.; Gumpert, A. Tiny shuttles for information transfer: Exosomes in cardiac health and disease. *J. Cardiovasc. Transl. Res.* **2016**, *9*, 169–175. [[CrossRef](#)]
21. van Niel, G.; Carter, D.R.F.; Clayton, A.; Lambert, D.W.; Raposo, G.; Vader, P. Challenges and directions in studying cell–cell communication by extracellular vesicles. *Nat. Rev. Mol. Cell Biol.* **2022**, *23*, 369–382. [[CrossRef](#)] [[PubMed](#)]
22. Garikipati, V.N.S.; Shojja-Taheri, F.; Davis, M.E.; Kishore, R. Extracellular vesicles and the application of system biology and computational modeling in cardiac repair. *Circ. Res.* **2018**, *123*, 188–204. [[CrossRef](#)] [[PubMed](#)]
23. Parolini, I.; Federici, C.; Raggi, C.; Lugini, L.; Palleschi, S.; De Milito, A.; Coscia, C.; Iessi, E.; Logozzi, M.; Molinari, A. Microenvironmental pH is a key factor for exosome traffic in tumor cells. *J. Biol. Chem.* **2009**, *284*, 34211–34222. [[CrossRef](#)] [[PubMed](#)]
24. Tian, T.; Zhu, Y.-L.; Zhou, Y.-Y.; Liang, G.-F.; Wang, Y.-Y.; Hu, F.-H.; Xiao, Z.-D. Exosome uptake through clathrin-mediated endocytosis and macropinocytosis and mediating miR-21 delivery. *J. Biol. Chem.* **2014**, *289*, 22258–22267. [[CrossRef](#)] [[PubMed](#)]
25. Mathieu, M.; Martin-Jaular, L.; Lavieau, G.; Théry, C. Specificities of secretion and uptake of exosomes and other extracellular vesicles for cell-to-cell communication. *Nat. Cell Biol.* **2019**, *21*, 9–17. [[CrossRef](#)] [[PubMed](#)]
26. Eguchi, S.; Takefuji, M.; Sakaguchi, T.; Ishihama, S.; Mori, Y.; Tsuda, T.; Takikawa, T.; Yoshida, T.; Ohashi, K.; Shimizu, Y. Cardiomyocytes capture stem cell-derived, anti-apoptotic microRNA-214 via clathrin-mediated endocytosis in acute myocardial infarction. *J. Biol. Chem.* **2019**, *294*, 11665–11674. [[CrossRef](#)] [[PubMed](#)]
27. Zhang, Y.; Hu, Y.-W.; Zheng, L.; Wang, Q. Characteristics and roles of exosomes in cardiovascular disease. *DNA Cell Biol.* **2017**, *36*, 202–211. [[CrossRef](#)]
28. Otero-Ortega, L.; Laso-García, F.; Gómez-de Frutos, M.; Fuentes, B.; Diekhorst, L.; Díez-Tejedor, E.; Gutiérrez-Fernández, M. Role of exosomes as a treatment and potential biomarker for stroke. *Transl. Stroke Res.* **2019**, *10*, 241–249. [[CrossRef](#)]
29. Street, J.M.; Koritzinsky, E.H.; Glispié, D.M.; Star, R.A.; Yuen, P.S.T. Urine exosomes: An emerging trove of biomarkers. *Adv. Clin. Chem.* **2017**, *78*, 103–122.
30. Bang, C.; Batkai, S.; Dangwal, S.; Gupta, S.K.; Foinquinos, A.; Holzmann, A.; Just, A.; Remke, J.; Zimmer, K.; Zeug, A. Cardiac fibroblast-derived microRNA passenger strand-enriched exosomes mediate cardiomyocyte hypertrophy. *J. Clin. Investig.* **2014**, *124*, 2136–2146. [[CrossRef](#)]
31. Chiang, J.-T.; Badrealam, K.F.; Shibu, M.A.; Cheng, S.-F.; Shen, C.-Y.; Chang, C.-F.; Lin, Y.-M.; Viswanadha, V.P.; Liao, S.-C.; Huang, C.-Y. Anti-apoptosis and anti-fibrosis effects of *eriobotrya japonica* in spontaneously hypertensive rat hearts. *Int. J. Mol. Sci.* **2018**, *19*, 1638. [[CrossRef](#)] [[PubMed](#)]
32. Chiang, J.T.; Badrealam, K.F.; Shibu, M.A.; Kuo, C.H.; Huang, C.Y.; Chen, B.C.; Lin, Y.M.; Viswanadha, V.P.; Kuo, W.W. *Eriobotrya japonica* ameliorates cardiac hypertrophy in H9c2 cardiomyoblast and in spontaneously hypertensive rats. *Environ. Toxicol.* **2018**, *33*, 1113–1122. [[CrossRef](#)] [[PubMed](#)]
33. Burnap, S.A.; Mayr, M. Extracellular vesicle crosstalk between the myocardium and immune system upon infarction. *Circ. Res.* **2018**, *123*, 15–17. [[CrossRef](#)] [[PubMed](#)]
34. Sluijter, J.P.G.; Verhage, V.; Deddens, J.C.; van den Akker, F.; Doevendans, P.A. Microvesicles and exosomes for intracardiac communication. *Cardiovasc. Res.* **2014**, *102*, 302–311. [[CrossRef](#)] [[PubMed](#)]
35. Tian, J.; Guo, X.; Liu, X.-M.; Liu, L.; Weng, Q.-F.; Dong, S.-J.; Knowlton, A.A.; Yuan, W.-J.; Lin, L. Extracellular HSP60 induces inflammation through activating and up-regulating TLRs in cardiomyocytes. *Cardiovasc. Res.* **2013**, *98*, 391–401. [[CrossRef](#)] [[PubMed](#)]
36. Lyu, L.; Wang, H.; Li, B.; Qin, Q.; Qi, L.; Nagarkatti, M.; Nagarkatti, P.; Janicki, J.S.; Wang, X.L.; Cui, T. A critical role of cardiac fibroblast-derived exosomes in activating renin angiotensin system in cardiomyocytes. *J. Mol. Cell. Cardiol.* **2015**, *89*, 268–279. [[CrossRef](#)]
37. Ong, S.-G.; Lee, W.H.; Huang, M.; Dey, D.; Kodo, K.; Sanchez-Freire, V.; Gold, J.D.; Wu, J.C. Cross talk of combined gene and cell therapy in ischemic heart disease: Role of exosomal microRNA transfer. *Circulation* **2014**, *130*, S60–S69. [[CrossRef](#)]
38. Yang, J.; Yu, X.; Xue, F.; Li, Y.; Liu, W.; Zhang, S. Exosomes derived from cardiomyocytes promote cardiac fibrosis via myocyte-fibroblast cross-talk. *Am. J. Transl. Res.* **2018**, *10*, 4350.
39. Poon, K.-S.; Palanisamy, K.; Chang, S.-S.; Sun, K.-T.; Chen, K.-B.; Li, P.-C.; Lin, T.-C.; Li, C.-Y. Plasma exosomal miR-223 expression regulates inflammatory responses during cardiac surgery with cardiopulmonary bypass. *Sci. Rep.* **2017**, *7*, 10807. [[CrossRef](#)]
40. Davidson, S.M.; Yellon, D.M. Exosomes and cardioprotection—a critical analysis. *Mol. Asp. Med.* **2018**, *60*, 104–114. [[CrossRef](#)]
41. Tang, Y.; Zhou, Y.; Li, H. Advances in mesenchymal stem cell exosomes: A review. *Stem Cell Res. Ther.* **2021**, *12*, 71. [[CrossRef](#)] [[PubMed](#)]

42. Ibrahim, A.G.-E.; Cheng, K.; Marbán, E. Exosomes as critical agents of cardiac regeneration triggered by cell therapy. *Stem Cell Rep.* **2014**, *2*, 606–619. [[CrossRef](#)] [[PubMed](#)]
43. Gallet, R.; Dawkins, J.; Valle, J.; Simsolo, E.; De Couto, G.; Middleton, R.; Tseliou, E.; Luthringer, D.; Kreke, M.; Smith, R.R. Exosomes secreted by cardiosphere-derived cells reduce scarring, attenuate adverse remodelling, and improve function in acute and chronic porcine myocardial infarction. *Eur. Heart J.* **2017**, *38*, 201–211. [[CrossRef](#)] [[PubMed](#)]
44. Gray, W.D.; French, K.M.; Ghosh-Choudhary, S.; Maxwell, J.T.; Brown, M.E.; Platt, M.O.; Searles, C.D.; Davis, M.E. Identification of therapeutic covariant microRNA clusters in hypoxia-treated cardiac progenitor cell exosomes using systems biology. *Circ. Res.* **2015**, *116*, 255–263. [[CrossRef](#)]
45. Mao, L.; Li, Y.-D.; Chen, R.-L.; Li, G.; Zhou, X.-X.; Song, F.; Wu, C.; Hu, Y.; Hong, Y.-X.; Dang, X. Heart-targeting exosomes from human cardiosphere-derived cells improve the therapeutic effect on cardiac hypertrophy. *J. Nanobiotechnol.* **2022**, *20*, 435. [[CrossRef](#)]
46. Barile, L.; Moccetti, T.; Marbán, E.; Vassalli, G. Roles of exosomes in cardioprotection. *Eur. Heart J.* **2017**, *38*, 1372–1379. [[CrossRef](#)]
47. Zhu, B.; Zhang, L.; Liang, C.; Liu, B.; Pan, X.; Wang, Y.; Zhang, Y.; Xie, W.; Yan, B.; et al. Stem cell-derived exosomes prevent aging-induced cardiac dysfunction through a novel exosome/lncRNA MALAT1/NF- κ B/TNF- α signaling pathway. *Oxid. Med. Cell Longev.* **2019**, *2019*, 9739258. [[CrossRef](#)]
48. Chen, L.; Wang, Y.; Pan, Y.; Zhang, L.; Shen, C.; Qin, G.; Ashraf, M.; Weintraub, N.; Ma, G.; Tang, Y. Cardiac progenitor-derived exosomes protect ischemic myocardium from acute ischemia/reperfusion injury. *Biochem. Biophys. Res. Commun.* **2013**, *431*, 566–571. [[CrossRef](#)]
49. Foglio, E.; Puddighinu, G.; Fasanaro, P.; D'Arcangelo, D.; Perrone, G.A.; Mocini, D.; Campanella, C.; Coppola, L.; Logozzi, M.; Azzarito, T. Exosomal clusterin, identified in the pericardial fluid, improves myocardial performance following MI through epicardial activation, enhanced arteriogenesis and reduced apoptosis. *Int. J. Cardiol.* **2015**, *197*, 333–347. [[CrossRef](#)]
50. Zhang, Z.; Xu, Y.; Cao, C.; Wang, B.; Guo, J.; Qin, Z.; Lu, Y.; Zhang, J.; Zhang, L.; Wang, W. Exosomes as a messenger to regulate the crosstalk between macrophages and cardiomyocytes under hypoxia conditions. *J. Cell. Mol. Med.* **2022**, *26*, 1486–1500. [[CrossRef](#)]
51. Gabisonia, K.; Khan, M.; Recchia, F.A. Extracellular vesicle-mediated bidirectional communication between heart and other organs. *Am. J. Physiol. Heart Circ. Physiol.* **2022**, *322*, H769–H784. [[CrossRef](#)] [[PubMed](#)]
52. Mentkowski, K.I.; Lang, J.K. Exosomes engineered to express a cardiomyocyte binding peptide demonstrate improved cardiac retention in vivo. *Sci. Rep.* **2019**, *9*, 10041. [[CrossRef](#)] [[PubMed](#)]
53. Kim, H.; Yun, N.; Mun, D.; Kang, J.-Y.; Lee, S.-H.; Park, H.; Park, H.; Joung, B. Cardiac-specific delivery by cardiac tissue-targeting peptide-expressing exosomes. *Biochem. Biophys. Res. Commun.* **2018**, *499*, 803–808. [[CrossRef](#)] [[PubMed](#)]
54. Zheng, M.; Huang, M.; Ma, X.; Chen, H.; Gao, X. Harnessing exosomes for the development of brain drug delivery systems. *Bioconjugate Chem.* **2019**, *30*, 994–1005. [[CrossRef](#)]
55. Alvarez-Erviti, L.; Seow, Y.; Yin, H.; Betts, C.; Likhacheva, S.; Wood, M.J.A. Delivery of siRNA to the mouse brain by systemic injection of targeted exosomes. *Nat. Biotechnol.* **2011**, *29*, 341–345. [[CrossRef](#)]
56. Yang, T.; Fogarty, B.; LaForge, B.; Aziz, S.; Pham, T.; Lai, L.; Bai, S. Delivery of small interfering RNA to inhibit vascular endothelial growth factor in zebrafish using natural brain endothelia cell-secreted exosome nanovesicles for the treatment of brain cancer. *AAPS J.* **2017**, *19*, 475–486. [[CrossRef](#)]
57. Ohno, S.-i.; Takanashi, M.; Sudo, K.; Ueda, S.; Ishikawa, A.; Matsuyama, N.; Fujita, K.; Mizutani, T.; Ohgi, T.; Ochiya, T. Systemically injected exosomes targeted to EGFR deliver antitumor microRNA to breast cancer cells. *Mol. Ther.* **2013**, *21*, 185–191. [[CrossRef](#)]
58. Wu, C.Y.; Du, S.L.; Zhang, J.; Liang, A.L.; Liu, Y.J. Exosomes and breast cancer: A comprehensive review of novel therapeutic strategies from diagnosis to treatment. *Cancer Gene Ther.* **2017**, *24*, 6–12. [[CrossRef](#)]
59. Liang, G.; Zhu, Y.; Ali, D.J.; Tian, T.; Xu, H.; Si, K.; Sun, B.; Chen, B.; Xiao, Z. Engineered exosomes for targeted co-delivery of miR-21 inhibitor and chemotherapeutics to reverse drug resistance in colon cancer. *J. Nanobiotechnol.* **2020**, *18*, 10. [[CrossRef](#)]
60. Tseliou, E.; Fouad, J.; Reich, H.; Slipczuk, L.; De Couto, G.; Aminzadeh, M.; Middleton, R.; Valle, J.; Weixin, L.; Marbán, E. Fibroblasts rendered antifibrotic, antiapoptotic, and angiogenic by priming with cardiosphere-derived extracellular membrane vesicles. *J. Am. Coll. Cardiol.* **2015**, *66*, 599–611. [[CrossRef](#)]
61. Qiu, X.; Li, Z.; Han, X.; Zhen, L.; Luo, C.; Liu, M.; Yu, K.; Ren, Y. Tumor-derived nanovesicles promote lung distribution of the therapeutic nanovector through repression of Kupffer cell-mediated phagocytosis. *Theranostics* **2019**, *9*, 2618. [[CrossRef](#)] [[PubMed](#)]
62. Gustafson, H.H.; Holt-Casper, D.; Grainger, D.W.; Ghandehari, H. Nanoparticle uptake: The phagocyte problem. *Nano Today* **2015**, *10*, 487–510. [[CrossRef](#)] [[PubMed](#)]
63. Kanki, S.; Jaalouk, D.E.; Lee, S.; Alvin, Y.C.; Gannon, J.; Lee, R.T. Identification of targeting peptides for ischemic myocardium by in vivo phage display. *J. Mol. Cell. Cardiol.* **2011**, *50*, 841–848. [[CrossRef](#)] [[PubMed](#)]
64. Won, Y.-W.; McGinn, A.N.; Lee, M.; Bull, D.A.; Kim, S.W. Targeted gene delivery to ischemic myocardium by homing peptide-guided polymeric carrier. *Mol. Pharm.* **2013**, *10*, 378–385. [[CrossRef](#)]
65. Wang, X.; Chen, Y.; Zhao, Z.; Meng, Q.; Yu, Y.; Sun, J.; Yang, Z.; Chen, Y.; Li, J.; Ma, T. Engineered exosomes with ischemic myocardium-targeting peptide for targeted therapy in myocardial infarction. *J. Am. Heart Assoc.* **2018**, *7*, e008737. [[CrossRef](#)]
66. Lee, K.; Yu, P.; Lingampalli, N.; Kim, H.J.; Tang, R.; Murthy, N. Peptide-enhanced mRNA transfection in cultured mouse cardiac fibroblasts and direct reprogramming towards cardiomyocyte-like cells. *Int. J. Nanomed.* **2015**, *10*, 1841.

67. Toba, M.; Alzoubi, A.; O'Neill, K.; Abe, K.; Urakami, T.; Komatsu, M.; Alvarez, D.; Järvinen, T.A.H.; Mann, D.; Ruoslahti, E. A novel vascular homing peptide strategy to selectively enhance pulmonary drug efficacy in pulmonary arterial hypertension. *Am. J. Pathol.* **2014**, *184*, 369–375. [[CrossRef](#)]
68. Xitong, D.; Xiaorong, Z. Targeted therapeutic delivery using engineered exosomes and its applications in cardiovascular diseases. *Gene* **2016**, *575*, 377–384. [[CrossRef](#)]
69. Vandergriff, A.; Huang, K.E.; Shen, D.; Hu, S.; Hensley, M.T.; Caranasos, T.G.; Qian, L.; Cheng, K. Targeting regenerative exosomes to myocardial infarction using cardiac homing peptide. *Theranostics* **2018**, *8*, 1869. [[CrossRef](#)]
70. McGuire, M.J.; Samli, K.N.; Johnston, S.A.; Brown, K.C. In vitro selection of a peptide with high selectivity for cardiomyocytes in vivo. *J. Mol. Biol.* **2004**, *342*, 171–182. [[CrossRef](#)]
71. Wang, X.; Huang, H.; Zhang, L.; Bai, Y.; Chen, H. PCM and TAT co-modified liposome with improved myocardium delivery: In vitro and in vivo evaluations. *Drug Deliv.* **2017**, *24*, 339–345. [[CrossRef](#)] [[PubMed](#)]
72. Nam, H.Y.; McGinn, A.; Kim, P.-H.; Kim, S.W.; Bull, D.A. Primary cardiomyocyte-targeted bioreducible polymer for efficient gene delivery to the myocardium. *Biomaterials* **2010**, *31*, 8081–8087. [[CrossRef](#)] [[PubMed](#)]
73. Lai, J.J.; Chau, Z.L.; Chen, S.Y.; Hill, J.J.; Korpany, K.V.; Liang, N.W.; Lin, L.H.; Lin, Y.H.; Liu, J.K.; Liu, Y.C. Exosome Processing and Characterization Approaches for Research and Technology Development. *Adv. Sci.* **2022**, *9*, 2103222. [[CrossRef](#)] [[PubMed](#)]
74. Kalluri, R.; LeBleu, V.S. The biology, function, and biomedical applications of exosomes. *Science* **2020**, *367*, eaau6977. [[CrossRef](#)] [[PubMed](#)]
75. Tickner, J.A.; Urquhart, A.J.; Stephenson, S.-A.; Richard, D.J.; O'Byrne, K.J. Functions and therapeutic roles of exosomes in cancer. *Front. Oncol.* **2014**, *4*, 127. [[CrossRef](#)]
76. Bellin, G.; Gardin, C.; Ferroni, L.; Chachques, J.C.; Rogante, M.; Mitrečić, D.; Ferrari, R.; Zavan, B. Exosome in cardiovascular diseases: A complex world full of hope. *Cells* **2019**, *8*, 166. [[CrossRef](#)]
77. Matsumoto, S.; Sakata, Y.; Suna, S.; Nakatani, D.; Usami, M.; Hara, M.; Kitamura, T.; Hamasaki, T.; Nanto, S.; Kawahara, Y.; et al. Circulating p53-responsive microRNAs are predictive indicators of heart failure after acute myocardial infarction. *Circ. Res.* **2013**, *113*, 322–326. [[CrossRef](#)]
78. Ji, Q.; Ji, Y.; Peng, J.; Zhou, X.; Chen, X.; Zhao, H.; Xu, T.; Chen, L.; Xu, Y. Increased brain-specific MiR-9 and MiR-124 in the serum exosomes of acute ischemic stroke patients. *PLoS ONE* **2016**, *11*, e0163645. [[CrossRef](#)]
79. Gidlöf, O.; Smith, J.G.; Miyazu, K.; Gilje, P.; Spencer, A.; Blomquist, S.; Erlinge, D. Circulating cardio-enriched microRNAs are associated with long-term prognosis following myocardial infarction. *BMC Cardiovasc. Disord.* **2013**, *13*, 12. [[CrossRef](#)]
80. Pironi, G.; Strachan, R.T.; Abraham, D.; Mon-Wei Yu, S.; Chen, M.; Chen, W.; Hanada, K.; Mao, L.; Watson, L.J.; Rockman, H.A. Circulating exosomes induced by cardiac pressure overload contain functional angiotensin II type 1 receptors. *Circulation* **2015**, *131*, 2120–2130. [[CrossRef](#)]
81. Guescini, M.; Leo, G.; Genedani, S.; Carone, C.; Pederzoli, F.; Ciruela, F.; Guidolin, D.; Stocchi, V.; Mantuano, M.; Borroto-Escuela, D.O. Microvesicle and tunneling nanotube mediated intercellular transfer of g-protein coupled receptors in cell cultures. *Exp. Cell Res.* **2012**, *318*, 603–613. [[CrossRef](#)] [[PubMed](#)]
82. Bonnet, S.; Boucherat, O.; Paulin, R.; Wu, D.; Hindmarch, C.C.T.; Archer, S.L.; Song, R.; Moore Iv, J.B.; Provencher, S.; Zhang, L. Clinical value of non-coding RNAs in cardiovascular, pulmonary, and muscle diseases. *Am. J. Physiol. Cell Physiol.* **2020**, *318*, C1–C28. [[CrossRef](#)] [[PubMed](#)]
83. Zhang, Y.; Du, W.; Yang, B. Long non-coding RNAs as new regulators of cardiac electrophysiology and arrhythmias: Molecular mechanisms, therapeutic implications and challenges. *Pharmacol. Ther.* **2019**, *203*, 107389. [[CrossRef](#)] [[PubMed](#)]
84. Lim, T.B.; Aliwarga, E.; Luu, T.D.A.; Li, Y.P.; Ng, S.L.; Annadoray, L.; Sian, S.; Ackers-Johnson, M.A.; Foo, R.S.-Y. Targeting the highly abundant circular RNA circSlc8a1 in cardiomyocytes attenuates pressure overload induced hypertrophy. *Cardiovasc. Res.* **2019**, *115*, 1998–2007. [[CrossRef](#)]
85. Chen, C.; Tang, Y.; Sun, H.; Lin, X.; Jiang, B. The roles of long noncoding RNAs in myocardial pathophysiology. *Biosci. Rep.* **2019**, *39*, BSR20190966. [[CrossRef](#)]
86. Lorenzen, J.M.; Thum, T. Long noncoding RNAs in kidney and cardiovascular diseases. *Nat. Rev. Nephrol.* **2016**, *12*, 360–373. [[CrossRef](#)]
87. Gurha, P.; Marian, A.J. Noncoding RNAs in cardiovascular biology and disease. *Circ. Res.* **2013**, *113*, e115–e120. [[CrossRef](#)]
88. Djebali, S.; Davis, C.A.; Merkel, A.; Dobin, A.; Lassmann, T.; Mortazavi, A.; Tanzer, A.; Lagarde, J.; Lin, W.; Schlesinger, F.; et al. Landscape of transcription in human cells. *Nature* **2012**, *489*, 101–108. [[CrossRef](#)]
89. El-Hefnawy, T.; Raja, S.; Kelly, L.; Bigbee, W.L.; Kirkwood, J.M.; Luketich, J.D.; Godfrey, T.E. Characterization of amplifiable, circulating RNA in plasma and its potential as a tool for cancer diagnostics. *Clin. Chem.* **2004**, *50*, 564–573. [[CrossRef](#)]
90. Mitchell, P.S.; Parkin, R.K.; Kroh, E.M.; Fritz, B.R.; Wyman, S.K.; Pogosova-Agadjanyan, E.L.; Peterson, A.; Noteboom, J.; O'Briant, K.C.; Allen, A. Circulating microRNAs as stable blood-based markers for cancer detection. *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 10513–10518. [[CrossRef](#)]
91. Zernecke, A.; Bidzhekov, K.; Noels, H.; Shagdarsuren, E.; Gan, L.; Denecke, B.; Hristov, M.; Köppel, T.; Jahantigh, M.N.; Lutgens, E. Delivery of microRNA-126 by apoptotic bodies induces CXCL12-dependent vascular protection. *Sci. Signal.* **2009**, *2*, ra81. [[CrossRef](#)] [[PubMed](#)]
92. Vickers, K.C.; Palmisano, B.T.; Shoucri, B.M.; Shamburek, R.D.; Remaley, A.T. MicroRNAs are transported in plasma and delivered to recipient cells by high-density lipoproteins. *Nat. Cell Biol.* **2011**, *13*, 423–433. [[CrossRef](#)] [[PubMed](#)]

93. Zhang, J.; Li, S.; Li, L.; Li, M.; Guo, C.; Yao, J.; Mi, S. Exosome and exosomal microRNA: Trafficking, sorting, and function. *Genom. Proteom. Bioinform.* **2015**, *13*, 17–24. [[CrossRef](#)] [[PubMed](#)]
94. Huang, X.; Yuan, T.; Tschannen, M.; Sun, Z.; Jacob, H.; Du, M.; Liang, M.; Dittmar, R.L.; Liu, Y.; Liang, M. Characterization of human plasma-derived exosomal RNAs by deep sequencing. *BMC Genom.* **2013**, *14*, 319. [[CrossRef](#)]
95. Mittelbrunn, M.; Gutiérrez-Vázquez, C.; Villarroya-Beltrí, C.; González, S.; Sánchez-Cabo, F.; González, M.Á.; Bernad, A.; Sánchez-Madrid, F. Unidirectional transfer of microRNA-loaded exosomes from T cells to antigen-presenting cells. *Nat. Commun.* **2011**, *2*, 282. [[CrossRef](#)]
96. Hewson, C.; Capraro, D.; Burdach, J.; Whitaker, N.; Morris, K.V. Extracellular vesicle associated long non-coding RNAs functionally enhance cell viability. *Non-Coding RNA Res.* **2016**, *1*, 3–11. [[CrossRef](#)]
97. Hobuß, L.; Bär, C.; Thum, T. Long non-coding RNAs: At the heart of cardiac dysfunction? *Front. Physiol.* **2019**, *10*, 30. [[CrossRef](#)]
98. Zhang, Y.; Sun, L.; Xuan, L.; Pan, Z.; Hu, X.; Liu, H.; Bai, Y.; Jiao, L.; Li, Z.; Cui, L. Long non-coding RNA CCRR controls cardiac conduction via regulating intercellular coupling. *Nat. Commun.* **2018**, *9*, 4176. [[CrossRef](#)]
99. Li, M.; Duan, L.; Li, Y.; Liu, B. Long noncoding RNA/circular noncoding RNA–miRNA–mRNA axes in cardiovascular diseases. *Life Sci.* **2019**, *233*, 116440. [[CrossRef](#)]
100. Wu, Z.; Liu, X.; Liu, L.; Deng, H.; Zhang, J.; Xu, Q.; Cen, B.; Ji, A. Regulation of lncRNA expression. *Cell. Mol. Biol. Lett.* **2014**, *19*, 561–575. [[CrossRef](#)]
101. Chen, L.; Zhang, Y.-H.; Pan, X.; Liu, M.; Wang, S.; Huang, T.; Cai, Y.-D. Tissue expression difference between mRNAs and lncRNAs. *Int. J. Mol. Sci.* **2018**, *19*, 3416. [[CrossRef](#)] [[PubMed](#)]
102. Iyer, M.K.; Niknafs, Y.S.; Malik, R.; Singhal, U.; Sahu, A.; Hosono, Y.; Barrette, T.R.; Prensner, J.R.; Evans, J.R.; Zhao, S. The landscape of long noncoding RNAs in the human transcriptome. *Nat. Genet.* **2015**, *47*, 199–208. [[CrossRef](#)] [[PubMed](#)]
103. Cabili, M.N.; Trapnell, C.; Goff, L.; Koziol, M.; Tazon-Vega, B.; Regev, A.; Rinn, J.L. Integrative annotation of human large intergenic noncoding RNAs reveals global properties and specific subclasses. *Genes Dev.* **2011**, *25*, 1915–1927. [[CrossRef](#)] [[PubMed](#)]
104. Xu, J.; Bai, J.; Zhang, X.; Lv, Y.; Gong, Y.; Liu, L.; Zhao, H.; Yu, F.; Ping, Y.; Zhang, G. A comprehensive overview of lncRNA annotation resources. *Brief. Bioinform.* **2017**, *18*, 236–249. [[CrossRef](#)]
105. Zhang, M.; Jiang, Y.; Guo, X.; Zhang, B.; Wu, J.; Sun, J.; Liang, H.; Shan, H.; Zhang, Y.; Liu, J. Long non-coding RNA cardiac hypertrophy-associated regulator governs cardiac hypertrophy via regulating miR-20b and the downstream PTEN/AKT pathway. *J. Cell. Mol. Med.* **2019**, *23*, 7685–7698. [[CrossRef](#)]
106. Wang, K.; Liu, F.; Zhou, L.-Y.; Long, B.; Yuan, S.-M.; Wang, Y.; Liu, C.-Y.; Sun, T.; Zhang, X.-J.; Li, P.-F. The long noncoding RNA CHRF regulates cardiac hypertrophy by targeting miR-489. *Circ. Res.* **2014**, *114*, 1377–1388. [[CrossRef](#)]
107. Li, Y.; Wang, J.; Sun, L.; Zhu, S. LncRNA myocardial infarction-associated transcript (MIAT) contributed to cardiac hypertrophy by regulating TLR4 via miR-93. *Eur. J. Pharmacol.* **2018**, *818*, 508–517. [[CrossRef](#)]
108. Long, B.; Li, N.; Xu, X.-X.; Li, X.-X.; Xu, X.-J.; Guo, D.; Zhang, D.; Wu, Z.-H.; Zhang, S.-Y. Long noncoding RNA FTX regulates cardiomyocyte apoptosis by targeting miR-29b-1-5p and Bcl2L2. *Biochem. Biophys. Res. Commun.* **2018**, *495*, 312–318. [[CrossRef](#)]
109. Su, X.; Lv, L.; Li, Y.; Fang, R.; Yang, R.; Li, C.; Li, T.; Zhu, D.; Li, X.; Zhou, Y.; et al. lncRNA MIRF promotes cardiac apoptosis through the miR-26a-Bak1 Axis. *Mol. Nucleic Acids.* **2020**, *20*, 841–850. [[CrossRef](#)]
110. Wang, Y.-N.-Z.; Shan, K.; Yao, M.-D.; Yao, J.; Wang, J.-J.; Li, X.; Liu, B.; Zhang, Y.-Y.; Ji, Y.; Jiang, Q.J.H. Long noncoding RNA-GAS5: A novel regulator of hypertension-induced vascular remodeling. *Hypertension* **2016**, *68*, 736–748. [[CrossRef](#)]
111. Tao, H.; Zhang, J.-G.; Qin, R.-H.; Dai, C.; Shi, P.; Yang, J.-J.; Deng, Z.-Y.; Shi, K.-H.J.T. LncRNA GAS5 controls cardiac fibroblast activation and fibrosis by targeting miR-21 via PTEN/MMP-2 signaling pathway. *Toxicology* **2017**, *386*, 11–18. [[CrossRef](#)] [[PubMed](#)]
112. Zhang, S.; Gao, S.; Wang, Y.; Jin, P.; Lu, F.J.D.; Biology, C. lncRNA SRA1 promotes the activation of cardiac myofibroblasts through negative regulation of miR-148b. *DNA Cell Biol.* **2019**, *38*, 385–394. [[CrossRef](#)] [[PubMed](#)]
113. Zou, L.; Ma, X.; Lin, S.; Wu, B.; Chen, Y.; Peng, C. Long noncoding RNA-MEG3 contributes to myocardial ischemia–reperfusion injury through suppression of miR-7-5p expression. *Biosci. Rep.* **2019**, *39*, BSR20190210. [[CrossRef](#)] [[PubMed](#)]
114. Lv, L.; Li, T.; Li, X.; Xu, C.; Liu, Q.; Jiang, H.; Li, Y.; Liu, Y.; Yan, H.; Huang, Q. The lncRNA Plscr4 controls cardiac hypertrophy by regulating miR-214. *Molecular Ther.-Nucleic Acids* **2018**, *10*, 387–397. [[CrossRef](#)]
115. Zou, X.; Wang, J.; Tang, L.; Wen, Q. LncRNA TUG1 contributes to cardiac hypertrophy via regulating miR-29b-3p. *Vitr. Cell. Dev. Biol. Anim.* **2019**, *55*, 482–490. [[CrossRef](#)]
116. Li, X.; Wang, H.; Yao, B.; Xu, W.; Chen, J.; Zhou, X. lncRNA H19/miR-675 axis regulates cardiomyocyte apoptosis by targeting VDAC1 in diabetic cardiomyopathy. *Sci. Rep.* **2016**, *6*, 36340. [[CrossRef](#)]
117. Yang, J.; Huang, X.; Hu, F.; Fu, X.; Jiang, Z.; Chen, K. LncRNA ANRIL knockdown relieves myocardial cell apoptosis in acute myocardial infarction by regulating IL-33/ST2. *Cell Cycle* **2019**, *18*, 3393–3403. [[CrossRef](#)]
118. Liang, H.; Pan, Z.; Zhao, X.; Liu, L.; Sun, J.; Su, X.; Xu, C.; Zhou, Y.; Zhao, D.; Xu, B. LncRNA PFL contributes to cardiac fibrosis by acting as a competing endogenous RNA of let-7d. *Theranostics* **2018**, *8*, 1180. [[CrossRef](#)]
119. Micheletti, R.; Plaisance, L.; Abraham, B.J.; Sarre, A.; Ting, C.-C.; Alexanian, M.; Maric, D.; Maison, D.; Nemir, M.; Young, R.A. The long noncoding RNA Wisper controls cardiac fibrosis and remodeling. *Sci. Transl. Med.* **2017**, *9*, eaai9118. [[CrossRef](#)]
120. Wang, L.; Zhang, J. Exosomal lncRNA AK139128 derived from hypoxic cardiomyocytes promotes apoptosis and inhibits cell proliferation in cardiac fibroblasts. *Int. J. Nanomed.* **2020**, *15*, 3363. [[CrossRef](#)]

121. Luo, Y.; Huang, L.; Luo, W.; Ye, S.; Hu, Q. Genomic analysis of lncRNA and mRNA profiles in circulating exosomes of patients with rheumatic heart disease. *Biol. Open* **2019**, *8*, bio045633. [[CrossRef](#)] [[PubMed](#)]
122. Shao, M.; Chen, G.; Lv, F.; Liu, Y.; Tian, H.; Tao, R.; Jiang, R.; Zhang, W.; Zhuo, C. LncRNA TINCR attenuates cardiac hypertrophy by epigenetically silencing CaMKII. *Oncotarget* **2017**, *8*, 47565. [[CrossRef](#)] [[PubMed](#)]
123. Kumarswamy, R.; Bauters, C.; Volkmann, I.; Maury, F.; Fetisch, J.; Holzmann, A.; Lemesle, G.; de Groote, P.; Pinet, F.; Thum, T. Circulating long noncoding RNA, LIPCAR, predicts survival in patients with heart failure. *Circ. Res.* **2014**, *114*, 1569–1575. [[CrossRef](#)] [[PubMed](#)]
124. de Gonzalo-Calvo, D.; Kenneweg, F.; Bang, C.; Toro, R.; Van Der Meer, R.W.; Rijzewijk, L.J.; Smit, J.W.; Lamb, H.J.; Llorente-Cortes, V.; Thum, T. Circulating long-non coding RNAs as biomarkers of left ventricular diastolic function and remodelling in patients with well-controlled type 2 diabetes. *Sci. Rep.* **2016**, *6*, 37354. [[CrossRef](#)]
125. Zhang, Z.; Gao, W.; Long, Q.-Q.; Zhang, J.; Li, Y.-F.; Yan, J.-J.; Yang, Z.-J.; Wang, L.-S. Increased plasma levels of lncRNA H19 and LIPCAR are associated with increased risk of coronary artery disease in a Chinese population. *Sci. Rep.* **2017**, *7*, 7491. [[CrossRef](#)]
126. Xuan, L.; Sun, L.; Zhang, Y.; Huang, Y.; Hou, Y.; Li, Q.; Guo, Y.; Feng, B.; Cui, L.; Wang, X. Circulating long non-coding RNA s NRON and MHRT as novel predictive biomarkers of heart failure. *J. Cell. Mol. Med.* **2017**, *21*, 1803–1814. [[CrossRef](#)]
127. Wang, K.; Long, B.O.; Liu, F.; Wang, J.-X.; Liu, C.-Y.; Zhao, B.; Zhou, L.-Y.; Sun, T.; Wang, M.; Yu, T. A circular RNA protects the heart from pathological hypertrophy and heart failure by targeting miR-223. *Eur. Heart J.* **2016**, *37*, 2602–2611. [[CrossRef](#)]
128. Taibi, F.; Metzinger-Le Meuth, V.; Massy, Z.A.; Metzinger, L. miR-223: An inflammatory oncomiR enters the cardiovascular field. *Biochim. Biophys. Acta (BBA)-Mol. Basis Dis.* **2014**, *1842*, 1001–1009. [[CrossRef](#)]
129. Zhao, G. Significance of non-coding circular RNAs and micro RNAs in the pathogenesis of cardiovascular diseases. *J. Med. Genet.* **2018**, *55*, 713–720. [[CrossRef](#)]
130. Meng, Z.; Chen, C.; Cao, H.; Wang, J.; Shen, E. Whole transcriptome sequencing reveals biologically significant RNA markers and related regulating biological pathways in cardiomyocyte hypertrophy induced by high glucose. *J. Cell. Biochem.* **2019**, *120*, 1018–1027. [[CrossRef](#)]
131. Wu, N.; Jin, L.; Cai, J. Profiling and bioinformatics analyses reveal differential circular RNA expression in hypertensive patients. *Clin. Exp. Hypertens.* **2017**, *39*, 454–459. [[CrossRef](#)] [[PubMed](#)]
132. Li, H.; Xu, J.-D.; Fang, X.-H.; Zhu, J.-N.; Yang, J.; Pan, R.; Yuan, S.-J.; Zeng, N.; Yang, Z.-Z.; Yang, H. Circular RNA circRNA_000203 aggravates cardiac hypertrophy via suppressing miR-26b-5p and miR-140-3p binding to Gata4. *Cardiovasc. Res.* **2020**, *116*, 1323–1334. [[CrossRef](#)] [[PubMed](#)]
133. Huang, X.; Chen, Y.; Xiao, J.; Huang, Z.; He, L.; Xu, D.; Peng, J. Identification of differentially expressed circular RNAs during TGF- β 1-induced endothelial-to-mesenchymal transition in rat coronary artery endothelial cells. *Anatol. J. Cardiol.* **2018**, *19*, 192. [[PubMed](#)]
134. Holdt, L.M.; Stahlinger, A.; Sass, K.; Pichler, G.; Kulak, N.A.; Willfert, W.; Kohlmaier, A.; Herbst, A.; Northoff, B.H.; Nicolaou, A. Circular non-coding RNA ANRIL modulates ribosomal RNA maturation and atherosclerosis in humans. *Nat. Commun.* **2016**, *7*, 1–14. [[CrossRef](#)] [[PubMed](#)]
135. Bai, M.; Pan, C.L.; Jiang, G.X.; Zhang, Y.M. CircRNA 010567 improves myocardial infarction rats through inhibiting TGF- β 1. *Eur. Rev. Med. Pharmacol. Sci.* **2020**, *24*, 369–375. [[CrossRef](#)]
136. Zeng, Y.; Du, W.W.; Wu, Y.; Yang, Z.; Awan, F.M.; Li, X.; Yang, W.; Zhang, C.; Yang, Q.; Yee, A. A circular RNA binds to and activates AKT phosphorylation and nuclear localization reducing apoptosis and enhancing cardiac repair. *Theranostics* **2017**, *7*, 3842. [[CrossRef](#)]
137. Viereck, J.; Thum, T. Circulating noncoding RNAs as biomarkers of cardiovascular disease and injury. *Circ. Res.* **2017**, *120*, 381–399. [[CrossRef](#)]
138. Wang, W.; Wang, Y.; Piao, H.; Li, B.; Huang, M.; Zhu, Z.; Li, D.; Wang, T.; Xu, R.; Liu, K. Circular RNAs as potential biomarkers and therapeutics for cardiovascular disease. *PeerJ* **2019**, *7*, e6831. [[CrossRef](#)]
139. Zhao, Z.; Li, X.; Gao, C.; Jian, D.; Hao, P.; Rao, L.; Li, M. Peripheral blood circular RNA hsa_circ_0124644 can be used as a diagnostic biomarker of coronary artery disease. *Sci. Rep.* **2017**, *7*, 39918. [[CrossRef](#)]
140. Li, J.J.H.; Wang, W.; Wang, X.Q.; He, Y.; Wang, S.S.; Yan, Y.X. A novel strategy of identifying circRNA biomarkers in cardiovascular disease by meta-analysis. *J. Cell. Physiol.* **2019**, *234*, 21601–21612. [[CrossRef](#)]
141. Salgado-Somoza, A.; Zhang, L.; Vausort, M.; Devaux, Y. The circular RNA MICRA for risk stratification after myocardial infarction. *IJC Heart Vasc.* **2017**, *17*, 33–36. [[CrossRef](#)] [[PubMed](#)]
142. Wang, J.; Chen, J.; Sen, S. MicroRNA as biomarkers and diagnostics. *J. Cell. Physiol.* **2016**, *231*, 25–30. [[CrossRef](#)] [[PubMed](#)]
143. Zhang, Z.; Yang, T.; Xiao, J. Circular RNAs: Promising biomarkers for human diseases. *EBioMedicine* **2018**, *34*, 267–274. [[CrossRef](#)]
144. Li, C.; Wang, Z.; Zhang, J.; Zhao, X.; Xu, P.; Liu, X.; Li, M.; Lv, C.; Song, X. Crosstalk of mRNA, miRNA, lncRNA, and circRNA and their regulatory pattern in pulmonary fibrosis. *Mol. Ther. Nucleic Acids* **2019**, *18*, 204–218. [[CrossRef](#)] [[PubMed](#)]

Illuminating the Molecular Intricacies of Exosomes and ncRNAs in Cardiovascular Diseases: Prospective Therapeutic and Biomarker Potential

ORIGINALITY REPORT

11 %
SIMILARITY INDEX

7 %
INTERNET SOURCES

9 %
PUBLICATIONS

0 %
STUDENT PAPERS

PRIMARY SOURCES

1 www.ncbi.nlm.nih.gov **2** %
Internet Source

2 link.springer.com **1** %
Internet Source

3 peerj.com **1** %
Internet Source

4 Dang Xitong, Zeng Xiaorong. "Targeted therapeutic delivery using engineered exosomes and its applications in cardiovascular diseases", Gene, 2016 **1** %
Publication

5 Priyatansh Gurha. "Noncoding RNAs in cardiovascular diseases", Current Opinion in Cardiology, 2019 **<1** %
Publication

6 Binh Yen Nguyen, Tayyiba Azam, Xin Wang. "Cellular signaling cross-talk between different cardiac cell populations: an insight **<1** %

into the role of exosomes in the heart diseases and therapy", American Journal of Physiology-Heart and Circulatory Physiology, 2021

Publication

7 theses.gla.ac.uk <1 %
Internet Source

8 www.cell.com <1 %
Internet Source

9 Yoosoo Yang, Yeonsun Hong, Eunji Cho, Gi Beom Kim, In-San Kim. "Extracellular vesicles as a platform for membrane-associated therapeutic protein delivery", Journal of Extracellular Vesicles, 2018 <1 %
Publication

10 www.biomedres.us <1 %
Internet Source

11 Horst Pick, Ana Catarina Alves, Horst Vogel. "Single-Vesicle Assays Using Liposomes and Cell-Derived Vesicles: From Modeling Complex Membrane Processes to Synthetic Biology and Biomedical Applications", Chemical Reviews, 2018 <1 %
Publication

12 Zandberga, Elīna, Viktors Kozirovskis, Artūrs Ābols, Diāna Andrējeva, Gunta Purkalne, and Aija Linē. "Cell-free microRNAs as diagnostic, <1 %

prognostic, and predictive biomarkers for lung cancer", Genes Chromosomes and Cancer, 2013.

Publication

13

www.science.gov

Internet Source

<1 %

14

Ruifang Wu, Yuwen Su, Haijing Wu, Yong Dai, Ming Zhao, Qianjin Lu. "Characters, functions and clinical perspectives of long non-coding RNAs", Molecular Genetics and Genomics, 2016

Publication

<1 %

15

V. Vlassov, P. Laktionov, E. Rykova. "Circulating Nucleic Acids as a Potential Source for Cancer Biomarkers", Current Molecular Medicine, 2010

Publication

<1 %

16

Weitie Wang, Yong Wang, Hulin Piao, Bo Li, Maoxun Huang, Zhicheng Zhu, Dan Li, Tiance Wang, Rihao Xu, Kexiang Liu. "Circular RNAs as potential biomarkers and therapeutics for cardiovascular disease", PeerJ, 2019

Publication

<1 %

17

Yihua Bei, Pujiao Yu, Dragos Cretoiu, Sanda Maria Cretoiu, Junjie Xiao. "Chapter 5 Exosomes-Based Biomarkers for the Prognosis of Cardiovascular Diseases",

<1 %

18

Carol W Chen, Leo L Wang, Samir Zaman, Jon Gordon et al. "Sustained release of endothelial progenitor cell-derived extracellular vesicles from shear-thinning hydrogels improves angiogenesis and promotes function after myocardial infarction", Cardiovascular Research, 2018

Publication

<1 %

19

Chia-Feng Liu, W.H. Wilson Tang. "Epigenetics in Cardiac Hypertrophy and Heart Failure", JACC: Basic to Translational Science, 2019

Publication

<1 %

20

Rashmi Tripathi, Pavan Chakraborty, Prithvi Kumar Varadwaj. "Unraveling long non-coding RNAs through analysis of high-throughput RNA-sequencing data", Non-coding RNA Research, 2017

Publication

<1 %

21

Griffiths, B.B., and R.G. Hunter. "Neuroepigenetics of stress", Neuroscience, 2014.

Publication

<1 %

22

Goretti, Emeline, Daniel R. Wagner, and Yvan Devaux. "miRNAs as biomarkers of myocardial infarction: a step forward towards

<1 %

personalized medicine?", Trends in Molecular Medicine, 2014.

Publication

23

Janas, Teresa, Maja M. Janas, Karolina Sapoń, and Tadeusz Janas. "Mechanisms of RNA loading into exosomes", FEBS Letters, 2015.

Publication

<1 %

24

Miaomiao Yang, Xinxing Wang, Tianhui Wang. "Regulation of Mitochondrial Function by Noncoding RNAs in Heart Failure and Its Application in Diagnosis and Treatment", Journal of Cardiovascular Pharmacology, 2021

Publication

<1 %

25

Nour Shaheen, Rehab Adel Diab, Mariam Tarek Desouki, Ahmed Shaheen, Mohamed Elmasry, Mayssa Rebei, Sarya Swed. "MicroRNAs (miRNAs) Role in Hypertension: Pathogenesis and Promising Therapeutics", Research Square Platform LLC, 2023

Publication

<1 %

26

journals.physiology.org

Internet Source

<1 %

27

translationalneurodegeneration.biomedcentral.com

Internet Source

<1 %

28

www.karger.com

Internet Source

<1 %

www.thno.org

29

Internet Source

<1 %

30

Erdogan Pekcan Erkan, Okay Saydam.
"Extracellular Vesicles as Novel Delivery Tools
for Cancer Treatment", Current Cancer Drug
Targets, 2015

Publication

<1 %

31

Iram Mushtaq, Ayesha Ishtiaq, Tahir Ali,
Muhammad Ishtiaq Jan, Iram Murtaza.
"Chapter 1 An Overview of Non-coding RNAs
and Cardiovascular System", Springer Science
and Business Media LLC, 2020

Publication

<1 %

32

Maria-Theodora Melissari, Phillip Grote.
"Roles for long non-coding RNAs in physiology
and disease", Pflügers Archiv - European
Journal of Physiology, 2016

Publication

<1 %

33

Pan, ZW, YJ Lu, and BF Yang. "Advances in
exploring the role of microRNAs in the
pathogenesis, diagnosis, and therapy of
cardiac diseases in China : miRNAs and
cardiac diseases research in China", British
Journal of Pharmacology, 2014.

Publication

<1 %

34

Sara Napoli. "Chapter 1 LncRNAs and
Available Databases", Springer Science and

<1 %

Business Media LLC, 2021

Publication

35

Sveva Bollini. "Veni, Vidi, Vici": how to arm your troops in the battlefield of cardiac repair", Cardiovascular Research, 2020

Publication

<1 %

36

Yingke Liu, Zhihe Zhao, Man Li. "Overcoming the cellular barriers and beyond: Recent progress on cell penetrating peptide modified nanomedicine in combating physiological and pathological barriers", Asian Journal of Pharmaceutical Sciences, 2022

Publication

<1 %

37

ashpublications.org

Internet Source

<1 %

38

fus-stag.literatumonline.com

Internet Source

<1 %

39

onlinelibrary.wiley.com

Internet Source

<1 %

40

pubmed.ncbi.nlm.nih.gov

Internet Source

<1 %

41

qmro.qmul.ac.uk

Internet Source

<1 %

42

van der Meel, R., M. Krawczyk-Durka, W. W. van Solinge, and R. M. Schiffelers. "Toward routine detection of extracellular vesicles in

<1 %

clinical samples", International Journal of Laboratory Hematology, 2014.

Publication

43

www.biorxiv.org

Internet Source

<1 %

44

Antonia Teona Deftu, Beatrice Mihaela Radu, Dragos Cretoiu, Alexandru Florian Deftu, Sanda Maria Cretoiu, Junjie Xiao. "Exosomes as intercellular communication messengers for cardiovascular and cerebrovascular diseases", Elsevier BV, 2020

Publication

<1 %

45

Jenny Y. Y. Ooi, Bianca C. Bernardo. "Chapter 21 Translational Potential of Non-coding RNAs for Cardiovascular Disease", Springer Science and Business Media LLC, 2020

Publication

<1 %

46

Tian Tian, Shuya Qiao, Bakhos A. Tannous. "Nanotechnology-Inspired Extracellular Vesicles Theranostics for Diagnosis and Therapy of Central Nervous System Diseases", ACS Applied Materials & Interfaces, 2022

Publication

<1 %

Exclude bibliography On

Illuminating the Molecular Intricacies of Exosomes and ncRNAs in Cardiovascular Diseases: Prospective Therapeutic and Biomarker Potential

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14

PAGE 15

PAGE 16

PAGE 17

PAGE 18
