

Resubmission of submitted manuscript after minor correction.

---

Dari: anvpublications@anvpublication.org

Kepada: sharylita@gmail.com; dwimurtiastutik@yahoo.co.id

Tanggal: Rabu, 17 Maret 2021 pukul 10.10 WIB

---

## **Research Journal of Pharmacy and Technology**

Paper ID: **21313130622496372** Date of Submission: **13-Mar-2021**

Paper Title: **Evaluation of the antifungal effects of rosemary oil and comparison with nystatin on the growth of isolate Candida Species from HIV/AIDS patients with Candidiasis oral**

Authors: **Dwi Murtiastutik; Afif Nurul Hidayati; Septiana Widyantari; Astindari; Bernadya Yogatri Anjuwita Saputri; Evy Ervianti; Damayanti; Maylita Sari**

Dear Author(s),

Thanks for submission of paper at enlisted journals of 'A & V Publications'. In the manuscript, it is compulsory to cite references of the related articles from journals of A&V Publications. Therefore, we are requested you to kindly cite / refers at least ten published manuscript from the enlisted journals. The link of published articles is embedded at the bottom of this mail, which you would to click and select any ten manuscripts as applicable to the submitted manuscript.

After citing the related articles, please resubmit the revised manuscript at article submission portal within 7 days.

[Click here to view the simalar works as yours.](#)

Please Login to account from which you have submitted the article, for more details. ([Click Here to Login Now.](#))

Thanks for your kind submission of manuscript.

**Editor**

**This is auto generated mail. Please do not reply.**

You have new comments for your submitted article.

---

Dari: anvpublications@anvpublication.org

Kepada: sharylita@gmail.com; dwimurtiastutik@yahoo.co.id

Tanggal: Selasa, 31 Agustus 2021 pukul 11.09 WIB

---

## **Research Journal of Pharmacy and Technology**

Paper ID: **21313130622496372** Date of Submission: **13-Mar-2021**

Paper Title: **Evaluation of the antifungal effects of rosemary oil and comparison with nystatin on the growth of isolate Candida Species from HIV/AIDS patients with Candidiasis oral**

Authors: **Dwi Murtiastutik; Afif Nurul Hidayati; Septiana Widyantari; Astindari; Bernadya Yogatri Anjuwita Saputri; Evy Ervianti; Damayanti; Maylita Sari**

Dear Author(s),

Thanks for submission of your article. You have following comments from Resubmission of Article for your submitted article titled "**Evaluation of the antifungal effects of rosemary oil and comparison with nystatin on the growth of isolate Candida Species from HIV/AIDS patients with Candidiasis oral**"

Date: DD-Aug-YYYY, 09:39 AM

### **COMMENTS:**

Dear Author, PI Resubmit the article after making following corrections.. The Ref no. are not as per instruction to author. (Model Journal reference) Ballesteros B. Tobias G. Shao L. Pellicer E. Nogués J. Mendoza E. Green ML et al Steam Purification for the Removal of Graphitic Shells Coating Catalytic Particles and the Shortening of Single-Walled Carbon Nanotubes. Small. 2008; 4(9):1501-6.doi.org/10.1021/ja061680u Honmane S. Hajare A. More H. Osmani RAM. Salunkhe S. Lung delivery of nanoliposomal salbutamol sulfate dry powder inhalation for facilitated asthma therapy. Journal of liposome research 2019; 29(4):332-342. doi.org/10.1080/08982104.2018.1531022 Thanks Editor

Please Login to account from which you have submitted the article for more details. ([Click Here to Login Now.](#))

Thank you.

**Editor**

**This is auto generated mail. Please do not reply.**

Regarding submission of final version of article.

---

Dari: anvpublications@anvpublication.org

Kepada: sharylita@gmail.com; dwimurtiastutik@yahoo.co.id

Tanggal: Rabu, 26 Januari 2022 pukul 17.13 WIB

---

## **Research Journal of Pharmacy and Technology**

Paper ID: **21313130622496372** Date of Submission: **13-Mar-2021**

Paper Title: **Evaluation of the antifungal effects of rosemary oil and comparison with nystatin on the growth of isolate Candida Species from HIV/AIDS patients with Candidiasis oral**

Authors: **Dwi Murtiastutik; Afif Nurul Hidayati; Septiana Widyantari; Astindari; Bernadya Yogatri Anjuwita Saputri; Evy Ervianti; Damayanti; Maylita Sari**

Dear Author(s),

Editorial board has considered your article titled "**Evaluation of the antifungal effects of rosemary oil and comparison with nystatin on the growth of isolate Candida Species from HIV/AIDS patients with Candidiasis oral**". for publication in Year : 2022, Vol: 15, Issue: 9. You are requested to send the final version of your article by clearly mentioning the *Paper Title* and *Author(s) name, affiliation and email address* in proper sequence.

**Note:- Please submit the final version of article with in 7 Days.**

Please Login to account from which you have submitted the article for more details. (To login now click : [http://www.anvpublication.org/Subscribed\\_User\\_Login.aspx](http://www.anvpublication.org/Subscribed_User_Login.aspx))

Thank you.

**Editor**

**This is auto generated mail. Please do not reply.**

Date: January 30, 2022

## Research Journal of Pharmacy and Technology

Paper ID:21313130622496372

Author's Name:Dwi Murtiastutik, Afif Nurul Hidayati, Septiana Widyantari, Astindari, Bernadya Yogatri Anjuwita Saputri, Evy Ervianti, Damayanti, Maylita Sari

Paper Title: Evaluation of the antifungal effects of rosemary oil and comparison with nystatin on the growth of isolate Candida Species from HIV/AIDS patients with Candidiasis oral



### Acceptance of Manuscript

With reference to your article titled '**Evaluation of the antifungal effects of rosemary oil and comparison with nystatin on the growth of isolate Candida Species from HIV/AIDS patients with Candidiasis oral**' Author by **Dwi Murtiastutik, Afif Nurul Hidayati, Septiana Widyantari, Astindari, Bernadya Yogatri Anjuwita Saputri, Evy Ervianti, Damayanti, Maylita Sari**. We wish to bring to your kind notice the following:

- We acknowledge the receipt of the above mentioned article.
- The above mentioned article has been sent to the reviewer of expert comments.
- The above mentioned article have been accepted for publication in Year : 2022, Vol: 15, Issue: 9 of the journal.

Thanking you for submission of manuscript.

Yours sincerely,



*Sr. Baharawal*  
Editor in Chief

**Note: -**

1. To verify the originality of this certificate please visit the following link:

[http://www.anvpublication.org/Download\\_Acceptance.aspx?PID=21313130622496372](http://www.anvpublication.org/Download_Acceptance.aspx?PID=21313130622496372)

PRINT

Payment details for your submitted article is verified successfully.

---

Dari: anvpublications@anvpublication.org

Kepada: dwimurtiastutik@yahoo.co.id

Tanggal: Senin, 31 Januari 2022 pukul 17.17 WIB

---

Dear Sir/Madam,  
Regards,

Thanks for submitting your article. It is our great pleasure to inform you that, payment proof for your submitted article (PID: 21313130622496372 ) is verified by the publisher.

Please Login to your account from which you have submitted the article for more details. ([Click Here to Login Now.](#))

Thanking you.  
**Editor-in-Chief**

**This is auto generated mail. Please do not reply.**

RESEARCH ARTICLE

## Evaluation of the Antifungal effects of Rosemary Oil and comparison with Nystatin on the Growth of *Candida* species isolates from HIV/AIDS patients with Oral Candidiasis

Dwi Murtiastutik<sup>1\*</sup>, Afif Nurul Hidayati<sup>1,2</sup>, Septiana Widyantari<sup>1</sup>, Astindari<sup>1</sup>, Bernadya Yogatri A. Saputri<sup>1</sup>, Lunardi Bintanjoyo<sup>1</sup>, Evy Ervianti<sup>1</sup>, Damayanti<sup>1</sup>, Maylita Sari<sup>1</sup>

<sup>1</sup>Department of Dermatology and Venereology, Faculty of Medicine, Universitas Airlangga, Dr. Soetomo General Academic Hospital, Surabaya 60285, Indonesia.

<sup>2</sup>Universitas Airlangga Teaching Hospital, Surabaya 60115, Indonesia.

\*Corresponding Author E-mail: [dwimurtiastutik@yahoo.co.id](mailto:dwimurtiastutik@yahoo.co.id)

### ABSTRACT:

**Background:** Oral candidiasis is an infection due to the activity of *Candida albicans* in the oral cavity. Oral candidiasis is one of the most common opportunistic infections occurring among Human Immunodeficiency Virus (HIV)/Acquired immune deficiency syndrome (AIDS) patients. Due to increasing resistance and adverse effects to commonly used antifungal drugs, many recent studies have examined the use of herbal essential oils as antifungal agents. In this study, essential oil of *Rosmarinus officinalis* (*Lamiaceae*) and nystatin were examined for in vitro antifungal activity against *Candida* species. **Aim:** To evaluate antifungal activity of essential oil of *Rosmarinus officinalis* (*Lamiaceae*) and nystatin by comparing inhibition zone diameters. **Methods:** This study was an experimental laboratory study with a posttest only design conducted in Dr. Soetomo General Academic Hospital, Surabaya. Forty isolates consisted of 20 isolates of *Candida albicans* and *Candida non-albicans* were subjected to test for antifungal activity using the diffusion disk method using paper discs or blank discs and inhibitory zones were recorded. **Results:** Diffusion test results revealed stronger antifungal effect of nystatin against all analyzed *Candida* strains. This study showed the mean diameter of the inhibitory zone for *Candida albicans* formed by rosemary essential oil is 2.25 mm and the average inhibition zone formed by rosemary essential oil for *Candida non-albicans* is 1.5 mm. **Conclusion:** The antifungal activity of nystatin is stronger when compared to rosemary essential oil as seen from a greater inhibition zone than rosemary essential oil in the diffusion method.

**KEYWORDS:** Antifungal activity, nystatin, *Rosmarinus officinalis*, HIV, *Candida albicans*, *Candida non-albicans*.

### INTRODUCTION:

*Candida* species are opportunistic yeasts and the etiologic agent of candidiasis, the most frequent fungal infection in humans. In the oral cavity, *Candida albicans* (*C. albicans*) is the most abundant *Candida* species in health and disease state. As an opportunistic pathogen, it usually causes infection in immunosuppressed or immunodeficient individuals such as HIV/AIDS patients.<sup>1</sup>

Various strategies are being used to manage candida infections, including the use of topical or systemic fungistatic or fungicidal agents. Antifungal agents including polyenes (nystatin, amphotericin B), azoles (fluconazole, itraconazole and miconazole), and antimetabolites agents (5-fluorocytosine) are most common medications used to treat candidiasis. However, occasionally consumption of the effective dose cannot be tolerated due to side effects.<sup>2</sup>

antimicrobial compounds of botanical extracts, in the form of crude extracts, essential oils and molecules, including flavonoids and terpenes. Recent studies have also addressed the association between conventional medications and natural products to achieve better treatment efficacy.<sup>3</sup>

Essential oils are defined as volatile and hydrophobic aromatic natural substances extracted from various parts, such as flowers, leaves, peels, barks, roots, seeds, resins, of aromatic plants, containing highly concentrated effective compounds. Rosemary (*Rosmarinus officinalis L.*), of the *Lamiaceae* family, is a perennial shrub with pleasant scent that grows in several regions globally. Most of these are used in the local folk tradition for many purposes.<sup>4</sup> Historically, rosemary has been used as a medicinal agent to treat renal colic and dysmenorrhea. It has also been applied for relief of respiratory symptoms, stimulation of hair growth, as aromatherapy for anxiety treatment and increasing alertness.

Rosemary essential oils have varying biological activities according to their chemical compositions. Around 13 chemotypes of rosemary essential oil have been recognized, which differ in their relative amounts of camphor, borneol, verbenone, pinene, 1, 8-cineole, and bornyl acetate. Nature-derived compounds which are effective against resistant *Candida* species are necessary options to control oral candidiasis. Fungistatic properties of *Rosmarinus officinalis* essential oil has been studied in several researches.<sup>5</sup> Fontenelle et al. stated that at low concentrations rosemary essential oil produced antifungal activity against *Candida*, but the anti-adherent activity of this oil has not been studied. Thus, further investigation of the antifungal activity of these oils is necessary to substantiate the use of essential oils clinically. The mechanism of antifungal action of rosemary essential oils has not been fully elucidated. Hammer et al. proposed that interaction of essential oils with lipid components results in changes in the cell membranes.<sup>6</sup> Other studies found that *Rosmarinus officinalis L.* possesses several antifungal mechanisms. It has been shown to alter cellular structure and fungal membrane permeability thus inhibiting the adhesion of *Candida albicans*. A study even showed that bionanosystem consisting of nanoparticles coated with rosemary essential oil was able to reduce the adhesiveness and to prevent the development of highly resistant biofilms of *Candida*.<sup>7</sup>

Various in vitro studies have examined the antifungal activity of rosemary essential oil against *C. albicans*. Hosein Nejati et al. studied topical rosemary for wound caused by *C. albicans* in mice, and found that topical rosemary has antifungal and anti-inflammatory properties in the wound due to its high terpenoids, limonene, 1,8

cineol content.<sup>8</sup> Other studies examined the effect of rosemary essential oil against *C. albicans* isolated from vaginal smears, and found that rosemary essential oil has a minimum inhibition concentration (MIC) of 0.1 mg/ml.<sup>4</sup> Cavalcanti et al. compared rosemary essential oil with nystatin against *Candida* species isolates and showed that rosemary essential oil has a MIC value of 2.25mg/ml which was better than 100,000 IU/ml nystatin.<sup>6</sup> This study evaluated the antifungal activity of rosemary essential oil compared with synthetic nystatin against isolates of *Candida* species.

## MATERIAL AND METHODS:

This is an experimental laboratory study with a posttest only design conducted in Dr. Soetomo General Academic Hospital, Surabaya. Forty isolates used in this study, which consisted of 20 isolates of *Candida albicans* and 20 isolates of *Candida non-albicans*, were obtained from forty HIV/AIDS patients with oral candidiasis. This research has been reviewed and approved by the Ethics Committee of the Dr. Soetomo General Academic Hospital (1614/KEPK/XI/2019). Analyzed essential of rosemary (*Rosmarinus officinalis L.*), were commercial product produced by Young Living USA, according to the producer's notification, characterized as 100% natural essential oil. This essential oil were analyzed by GC containing 45.12% 1,8-cineol, 12.78% alpha-pinene, 10.79% camphor, 3.50% borneol. Nystatin used as susceptibility test. The isolates were tested for antifungal activity using the disc diffusion method. The colonies were mixed in 0.85% sterile normal saline (5ml volume). Mueller Hinton agar with 2% glucose and 0.5µg/ml of methylene blue was used. Strain suspensions (10<sup>6</sup> CFU/ml) were swabbed on Saboraud Dextrose Agar. Filter paper disc (6 mm diameter) were placed on the agar surface and added with 10µl of 100% concentrate rosemary essential oil and 100% concentrate of nystatin. Plates were incubated at 37°C for 48 hours. The zone diameters were interpreted according to Clinical Laboratory Standards Institute (CLSI) guidelines. The obtained results (mean and standard deviation) were analyzed using SPSS. The data for diffusion methods were analyzed by Mann-Whitney test. Statistical significance was determined at p≤0.05.

## RESULTS:

The isolates used in this study consisted of 20 isolates of *Candida albicans* and 20 isolates of *Candida non-albicans* taken from HIV/AIDS patients with oral candidiasis. The investigation on antimycotic activity of rosemary essential oils and nystatin against *Candida* species was done by agar diffusion technique. The zone of inhibition in 100% concentrations of essential oils and nystatin against *Candida albicans* is recorded and tabulated in Table 1.

**Table 1: Inhibitory zones (mm±sd) of rosemary essential oils and nystatin in disc-diffusion assay**

Species	Concentration	n	Means±SD (mm)	p value
<i>Candida albicans</i>	Nystatin 100%	20	16.50 ±4.7	< 0.001*
	Rosemary 100%	20	2.25 ± 4.1	
<i>Candida non-albicans</i>	Nystatin 100%	20	12.45±4.1	< 0.001*
	Rosemary 100%	20	1.5±3.1	

\*Significant differences (p<0.05), SD: Standard deviation

Table 1 shows the inhibition zone formed by rosemary essential oil is smaller when compared to the inhibition zone formed by nystatin. The mean value of inhibition zone formed by nystatin is 16.5mm which indicates that nystatin is still sensitive against *Candida albicans*. Statistically, the p value obtained from this study is p <0.001, showing a significant difference between inhibition zones formed by nystatin and inhibitory zones formed by rosemary essential oil. This study also showed inhibition zones formed by rosemary essential oil on the growth of *Candida non-albicans* were smaller when compared to inhibition zones formed by nystatin. The average value of inhibition zone formed by nystatin is 12.45mm, where this value indicates that nystatin is not sensitive but is included in the intermediate classification against *Candida non-albicans*. The statistical value obtained from this study is the value of p<0.001 where there is a significant difference between the inhibition zone formed by nystatin and the inhibition zone formed by rosemary essential oil.

## DISCUSSION:

This study was conducted to examine the in vitro antifungal activity of rosemary essential oils and nystatin against *Candida albicans* and *Candida non-albicans*. When fungal isolate is given substances with antifungal properties, its growth will be inhibited. In disc diffusion method, inhibition of fungal growth can be observed as inhibitory clear zones found around paper discs on media that have been inoculated with *Candida albicans* and *Candida non-albicans*.<sup>9</sup> The formation of inhibition zones around disc paper depends on the presence or absence of active compounds in the extract, while the size of the inhibition zone correspond to the level of activity of the substances in the extract. Hermawan et al. in 2007 stated that interpretation of inhibition zones refers to the general standard issued by the Ministry of Health in 1988 in which microbes are defined to be sensitive to active compounds of a plant if the measurement of power diameter resistance is 12 to 24mm.<sup>10</sup> Disc diffusion method is easier to perform, can be interpreted within 24hours and can be used for diagnostic purposes on daily basis whereas broth microdilution method is more cumbersome and requires more practical skills.<sup>9</sup>

In this study, the mean diameters of the inhibitory zone formed by nystatin and rosemary essential oil against *Candida albicans* are 16.5mm and 2.25mm respectively.

*Candida non-albicans* were also included in this study which consisted of 9 isolates of *Candida krusei*, 4 isolates of *Candida glabrata*, 2 isolates of *Candida dubliniensis*, 2 isolates of *Candida parapsilosis*, 2 isolates of *Candida tropicalis* and 1 isolate of *Candida lipolytica*. The mean diameters of the inhibitory zone formed by nystatin and rosemary essential oil against *Candida non-albicans* are 12.45mm and 1.5mm respectively. The largest inhibitory zone formed by nystatin was observed against two isolates of *Candida dubliniensis* which was 20mm and 18mm respectively. The largest inhibitory zone formed by rosemary essential oils were observed against *Candida krusei* with a diameter of 8mm.

Diameter of inhibitory zone more than 15mm indicates sensitivity to nystatin.<sup>11</sup> This study showed that *Candida albicans* is still sensitive to nystatin. Carvalho et al. in 2012 examined the sensitivity of nystatin and essential oils, one of which was rosemary, showing that all 40 isolates of *Candida albicans* species were still sensitive to nystatin and rosemary essential oils with respective inhibitory zone's diameters of more than 15 mm and 12mm.<sup>5</sup> Murtiastutik et al. in 2019 studied the sensitivity of *Candida albicans* and *non-albicans* isolated from HIV/AIDS patients at Dr. Soetomo General Academic Hospital to nystatin by the diffusion disk method, and showed that all isolates was still sensitive to nystatin.<sup>12</sup> This study also showed also that *Candida non-albicans* has intermediate sensitivity to nystatin, except *Candida krusei* which showed to be sensitive to nystatin. The result of this study slightly differs from study by Kengne et al. in 2017 on sensitivity of *Candida* species isolated from vulvovaginal candidiasis patients which showed that nystatin was the most sensitive antifungal against *Candida non-albicans*.<sup>11</sup> The mechanism of antifungals that causes inhibition of fungal growth is damage to cell membrane. This will disrupt the integrity of cellular components and impair the process of fungal cellular respiration. The end result is insufficient energy to transport active substances so that fungal growth is disrupted.<sup>13,14</sup>

In this study, the formation of inhibitory zone by rosemary essential oil although smaller than those formed by nystatin indicated that rosemary essential oil has antifungal activity against both *Candida albicans* and *Candida non-albicans*, also that the antifungal



activity of nystatin was better than rosemary essential oil. This is possibly attributed to nystatin which could diffuse well on agar media and which was the first choice drug for infections by *Candida*. This also could be due other factors including the content of antifungal compound within rosemary essential oil that is not too high, diffusion power, types of fungi and the concentration of the extract.<sup>15</sup>

Periodic antifungal resistance surveillance protocols by susceptibility testing of all *Candida* isolates is essential to provide crucial information about the resistance pattern in a particular region. This also will guide in choosing the empiric or prophylactic drug before the antifungal resistance test result is available.<sup>16-20</sup> In conclusion, plants with medicinal properties are considered to be important source of novel potentially therapeutic compounds. The present study has shown the antimycotic activity of rosemary oil on *Candida* species. However, the antifungal activity of nystatin is stronger when compared to rosemary essential oil as seen from a greater inhibition zone than rosemary essential oil in the diffusion method.

#### ACKNOWLEDGEMENT:

We would like to thank Pepy Dwi Endraswari for helping us in carrying out the culture of *Candida* species in the Department of Microbiology Dr. Soetomo General Academic Hospital, Surabaya, Indonesia and Alvian Arifin Saiboo for helping us in the publication process of this journal.

#### CONFLICT OF INTEREST:

The authors declare no conflict of interest.

#### REFERENCES:

1. Abdulaziz SM, Shaswary IA, Muhammad AA. In vitro antifungal activity of essential oils from local plants against fluconazole-resistant oral *Candida albicans* isolates. *Zanco J Med Sci*. 2015; 19(2):965–71.
2. da Silva ICG, de Pontes Santos HB, Cavalcanti YW, Nonaka CFW, de Sousa SA, de Castro RD. Antifungal Activity of Eugenol and its Association with Nystatin on *Candida albicans*. *Pesqui Bras Odontopediatria Clin Integr*. 2017; 17(1):e3235.
3. Sakkas H, Gousia P, Economou V, Petsios S, Papadopoulou C. Antifungal activity of four essential oils against *Candida* clinical isolates. *Asian J Ethnopharmacol Med Foods*. 2016; 2(1):22–5.
4. Jahani S, Bazi S, Shahi Z, Asadi MS, Mosavi F, Baigi GS. Antifungal Effect of the Extract of the Plants Against *Candida albicans*. *Int J Infect*. 2017; 4(2):e36807.
5. Carvalhinho S, Costa AM, Coelho AC, Martins E, Sampaio A. Susceptibilities of *Candida albicans* Mouth Isolates to Antifungal Agents, Essentials Oils and Mouth Rinses. *Mycopathologia*. 2012; 174:69–76.
6. Cavalcanti YW, de Almeida L de FD, Padilha WWN. Anti-adherent activity of *Rosmarinus officinalis* essential oil on *Candida albicans*: an SEM analysis. *Rev Odonto Cienc*. 2011; 26(2):139–44.
7. Patturaja K, Geetha R V. Evaluation of Antimycotic Activity of Three Essential Oils on *Candida Albicans* -An Invitro Study. *J Pharm Sci Res*. 2017; 9(4):480–2.
8. Nejati H, Farahpour MR, Nagadehi MN. Topical Rosemary officinalis essential oil improves wound healing against disseminated *Candida albicans* infection in rat model. *Comp Clin Path*. 2015; 24:1377–83.
9. Moghtader M, Salari H, Farahmand A. Evaluation of the antifungal effects of rosemary oil and comparison with synthetic borneol and fungicide on the growth of *Aspergillus flavus*. *J Ecol Nat Environ*. 2011; 3(6):210–4.
10. Andayani N, Susilowati A, Pangastuti A. Anti candida minyak atsiri lengkuas putih (*Alpinia galanga*) terhadap *Candida albicans* penyebab candidiasis secara invitro. *EL-Vivo*. 2014; 2(2):1–9.
11. Kengne M, Shu SV, Nwobegahay JM, Achonduh O. Antifungals susceptibility pattern of *Candida* spp. isolated from female genital tract at the Yaoundé Bethesda Hospital in Cameroon. *Pan Afr Med J*. 2017; 28:294.
12. Murtiastutik D, Maharani CS, Rahmadewi, Listiawan MY. Nystatin Profile on *Candida* Species in HIV / AIDS Patients with Oral Candidiasis: A Phenomenology Study. *J Pure Appl Microbiol*. 2019; 13(4):2013–9.
13. Lely N, Pratiwi RI, Imanda YL. Efektivitas Antijamur Kombinasi Ketokonazol dengan Minyak Atsiri Sereh Wangi (*Cymbopogon nardus* (L.) Rendle). *Indones J Appl Sci*. 2017; 7(2):10–5.
14. Nenoff P, Krüger C, Neumeister C, Schwantes U, Koch D. In vitro susceptibility testing of yeasts to nystatin – low minimum inhibitory concentrations suggest no indication of in vitro resistance of *Candida albicans*, *Candida* species or non-*Candida* yeast species to nystatin. *Clin Med Investig*. 2016; 1(3).
15. Khorram Z, Hakimaneh SM, Naeini A, Rafeinezhad R, Salari AM, Shayegh SS. The Antifungal Effects of Two Herbal Essences in Comparison with Nystatin on the *Candida* Strains Isolated from the Edentulous Patients. *J Contemp Dent Pract*. 2019; 20(6):716–9.
16. Ćosić J, Vrandečić K, Postić J, Jurković D, Ravlić M. In vitro antifungal activity of essential oils on growth of phytopathogenic fungi. *Poljoprivreda*. 2010; 16(2):25–8.
17. Bozin B, Mimica-Dukić N, Samojlik I, Jovin E. Antimicrobial and Antioxidant Properties of Rosemary and Sage (*Rosmarinus officinalis* L. and *Salvia officinalis* L., Lamiaceae) Essential Oils. *J Agric Food Chem*. 2007; 55:7879–85.
18. Murtiastutik D, Listiawan MY, Bintanjoyo L, Hidayati AN, Widiantari S, Astindari, et al. Ketoconazole: A Re-emerging choice for Oral candidiasis in patients with human immunodeficiency virus infection/acquired Immunodeficiency Syndrome. *Res J Pharm Technol*. 2022;15(3):1071–6.
19. Rezeki S, Pradono SA, Subita GP, Rosana Y, Sunnati S, Gani BA. The antifungal susceptibility of *Candida albicans* isolated from HIV/AIDS patients. *Dent J (Majalah Kedokt Gigi)*. 2021;54(2):82–6.
20. Reza NR, Tantari SHW, Basuki S. Uji Kepekaan In Vitro Flukonazol Terhadap Spesies *Candida* penyebab Kandidiasis Oral Pada Pasien HIV/AIDS dengan Vitek II (In Vitro Susceptibility Test of Fluconazole to *Candida* spp in Patients with Oropharyngeal Candidiasis and HIV/AIDS with Vitek II). *Berk Ilmu Kesehat Kulit dan Kelamin - Period Dermatology Venereol*. 2017;29(3):234–42.