



Dr. Yi-Ping Hsieh (<https://sciprofiles.com/profile/684275>)

E-Mail () **Website** (<https://und.edu/directory/yiping.hsieh>)

Guest Editor

Department of Social Work, University of North Dakota, 225 Centennial Drive, Stop 7135, Grand Forks, ND 58202-7135, USA

Interests: Internet addiction; bullying and cyberbullying; online victimization; family processes; adverse childhood experiences; children's and adolescents' mental health and behavioral problems

Dr. Chung-Ying Lin (<https://sciprofiles.com/profile/664254>)

E-Mail () **Website** (<http://ahs.ncku.edu.tw/p/412-1129-84.php?Lang=en>)

Guest Editor

Institute of Allied Health Sciences, National Cheng Kung University Hospital, College of Medicine, National Cheng Kung University, Tainan 701, Taiwan

Interests: patient-reported outcomes; childhood adversity and mental health in children/adolescents; therapeutic effects on people with mental illness; psychometric testing

Special Issues, Collections and Topics in MDPI journals

Special Issue Information

Dear Colleagues,

With technological advancement and the development of convenient application functions, modern society is living in a digital era. Although digital technology has been developed to improve human health and life satisfaction, some people are involved in technology-based addictive behaviors, which have negative impacts on their wellbeing. Common types of technology addiction include internet addiction, smartphone addiction, gaming addiction, social media addiction, pornography addiction (online sexual addiction), and online shopping addiction. The American Society of Addiction Medicine has acknowledged that addictions are not limited to drugs and alcohol, but can also be behavioral in nature. In addition to the symptoms of addiction or dependency (e.g., loss of control, tolerance, withdrawal, changes in mood, and continuation of use despite adverse consequences), a technology-based addiction is apparent if and when digital usage interferes with one's daily life and general wellbeing, and when the individual shows a preoccupation with using their digital medium of choice. It is important to examine how technology-based addictive behaviors influence various aspects of wellbeing (e.g., physical wellbeing, social wellbeing, emotional wellbeing, psychological wellbeing, economic wellbeing, life satisfaction, and engagement in activities and work), and what the risk and protective factors are which can reduce or increase the likelihood of digital addiction, which can and buffer or amplify the negative effects of digital addiction on wellbeing.

This Special Issue aims to examine and advance understanding surrounding technology-based addictive behaviors and their correlates and impacts on wellbeing through rigorous and empirical research. It is also important to address policy implementation that involves all stakeholders in screening and preventing technology-based addictive behavior; such legislation has the potential to reduce its harm on human wellbeing, capture the broad lessons emerging from global cross-country scholars, inform practices, and enhance capacity for tackling such addictions.

We welcome original research articles and reviews for submission to this Special Issue. We look forward to receiving your contributions.

Dr. Yi-Ping Hsieh ([View PDF](chrome-extension://dagcmkpagjllhakfdhnbomgmjdpkdklff/enhanced-reader.html?openApp&pdf=https%3A%2F%2Fmdpi-view-pdf) (chrome-extension://dagcmkpagjllhakfdhnbomgmjdpkdklff/enhanced-reader.html?openApp&pdf=https%3A%2F%2Fmdpi-view-pdf))

Dr. Chung-Ying Lin ([View PDF](http://ahs.ncku.edu.tw/p/412-1129-84.php?Lang=en&attachment%2Fhealthcare%2Fhealthcare-11-00700%2Farticle_deploy%2Fhealthcare-11-00700.pdf%3Fversion%3D1677478556) (http://ahs.ncku.edu.tw/p/412-1129-84.php?Lang=en&attachment%2Fhealthcare%2Fhealthcare-11-00700%2Farticle_deploy%2Fhealthcare-11-00700.pdf%3Fversion%3D1677478556))

Guest Editors

Manuscript Submission Information

Manuscripts should be submitted online at www.mdpi.com (<https://www.mdpi.com/>) by **registering** (<https://www.mdpi.com/user/register/>) and **logging in to this website** (<https://www.mdpi.com/user/login/>). Once you are registered, **click here to go to the submission form** (<https://susy.mdpi.com/user/manuscripts/upload/?journal=healthcare>). Manuscripts can be submitted until the deadline. All submissions that pass pre-check are peer-reviewed. Accepted papers will be published continuously in the journal (as soon as accepted) and will be listed together on the special issue website. Research articles, review articles as well as short communications are invited. For planned papers, a title and short abstract (about 100 words) can be sent to the Editorial Office for announcement on this website.

Submitted manuscripts should not have been published previously, nor be under consideration for publication elsewhere (except conference proceedings papers). All manuscripts are thoroughly refereed through a single-blind peer-review process. A guide for authors and other relevant information for submission of manuscripts is available on the **Instructions for Authors** (<https://www.mdpi.com/journal/healthcare/instructions>) page. **Healthcare** (<https://www.mdpi.com/journal/healthcare/>) is an international peer-reviewed open access semimonthly journal published by MDPI.

Please visit the **Instructions for Authors** (<https://www.mdpi.com/journal/healthcare/instructions>) page before submitting a manuscript. The **Article Processing Charge (APC)** (<https://www.mdpi.com/about/apc/>) for publication in this **open access** (<https://www.mdpi.com/about/openaccess/>) journal is 2000 CHF (Swiss Francs). Submitted papers should be well formatted and use good English. Authors may use MDPI's **English editing service** (<https://www.mdpi.com/authors/english>) prior to publication or during author revisions.

Keywords

- internet addiction
- smartphone addiction
- technology addiction
- pathological technology use
- online gaming
- social media use
- wellbeing
- mental health
- risk and protective factors
- prevention/intervention

Published Papers (6 papers)

[Download All Papers](#)

Order results

Content type

Result details

Normal

[Show export options](#) ▾

Research

Jump to: [Review](#)

Open Access Article

[The Psychometric Properties of the DASS-21 and Its Association with Problematic Internet Use among Chinese College Freshmen \(/2227-9032/11/5/700\)](#)

by [Cui-Hong Cao](#) (<https://sciprofiles.com/profile/2020792>),
[Chang-Yan Dang](#) (<https://sciprofiles.com/profile/author/LzB3dTRuNTdmdIVtZk8xQXJKV1luTS9BSnNnQWd3TFNqaU9Gc1NWW3VSVT0=>),
[Xia Zheng](#) (<https://sciprofiles.com/profile/author/Um41eJXemNDTnkwTUYvUUkrYy8xMFErQjhRWFgvcEI5SnRtRGireHFTdz0=>),
[Wang-Guang Chen](#) (<https://sciprofiles.com/profile/author/R2RVY1VXaUdiQy9ZNU5qRDF1Zy9Tb0U2ZmhsYm5hdC95bIV4YVWkdU5Rbz0=>),
[Li-Hua Chen](#) (<https://sciprofiles.com/profile/1537537>) and [Jeffrey H. Gamble](#) (<https://sciprofiles.com/profile/1058941>).

Healthcare 2023, 11(5), 700; <https://doi.org/10.3390/healthcare11050700> (<https://doi.org/10.3390/healthcare11050700>) - 27 Feb 2023

Viewed by 820

Abstract This study examines the psychometric properties of the DASS-21-item version (DASS-21) commonly used for mental health assessment in China. However, evidence is lacking regarding its applicability with freshmen as a demographic. Debates also exist regarding its use in this population. [Read more.](#)

(This article belongs to the Special Issue [The Correlates of Technology-Based Addictive Behaviors and Their Impacts on Wellbeing](#) ([/journal/healthcare/special_issues/Technology_Based_Addictive_Behaviors_Impacts](#)))

► [Show Figures](#)

(https://pub.mdpi-res.com/healthcare/healthcare-11-00700/article_deploy/html/images/healthcare-11-00700-g001-550.jpg?1677478623)

Open Access Article

[The Mediating Role of Selfitis in the Associations between Self-Esteem, Problematic Social Media Use, Problematic Smartphone Use, Body-Self Appearance, and Psychological Distress among Young Ghanaian Adults \(/2227-9032/10/12/2500\)](#)

by [Derek Oppong](#) (<https://sciprofiles.com/profile/2604252>),
[Emma Sethina Adjaottor](#) (<https://sciprofiles.com/profile/author/czZqbVdyWStRNGF0SzgrWHhNM3ZoMkJFSEZyaFUvUHirajM2dHZTaDEyOD0=>),

[Frimpong-Manso Addo](#) (<https://sciprofiles.com/profile/author/SWJ1Q1FVZXpzQldmUmJFOC9QM1czRzRGeGhNb2ErdW5MVCtTUk9BSHBjST0=>),

[Worlali Nyaledzigbor](#) (<https://sciprofiles.com/profile/author/ZjFFeHVkZ2d1QXBkZkdKUVJQR2YweTBDZS9yRmdBdDd1d0JzQktHQzgaz0=>),

[Amma Serwaa Ofori-Amanfo](#) (<https://sciprofiles.com/profile/2635409>), [Hsin-Pao Chen](#) (<https://sciprofiles.com/profile/1676934>) and

[Daniel Kwasi Ahorsu](#) (<https://sciprofiles.com/profile/1303687>)

Healthcare 2022, 10(12), 2500; <https://doi.org/10.3390/healthcare10122500> (<https://doi.org/10.3390/healthcare10122500>) - 10 Dec 2022

Viewed by 772

Abstract Selfie-related activities have become pervasive to the point that they may affect the mental health of people who engage in them. To ascertain this mechanism, this study examined the mediating role of selfitis in the associations between self-esteem, problematic social media use, problematic smartphone use, body-self appearance, and psychological distress among young Ghanaian adults. [Read more.](#)

Read more

(This article belongs to the Special Issue [The Correlates of Technology-Based Addictive Behaviors and Their Impacts on Wellbeing](#) ([/journal/healthcare/special_issues/Technology_Based_Addictive_Behaviors_Impacts](#)))

► Show Figures

(toggle desktop layout cookie)

(https://pub.mdpi-res.com/healthcare/healthcare-10-02500/article_deploy/html/images/healthcare-10-02500-g001-550.jpg?1671000259).

Open Access Article

([/2227-9032/10/9/1745/pdf?version=1662890957](#))

[The Mediating Roles of Anxiety, Depression, Sleepiness, Insomnia, and Sleep Quality in the Association between Problematic Social Media Use and Quality of Life among Patients with Cancer](#) ([/2227-9032/10/9/1745](#))

by [Vida Imani](#) (<https://sciprofiles.com/profile/author/dDBhcGt6RTNTT3lCa1ZBMXdjZTYvcVpSTehJa0VaelpRb0l5TjNSS3JLOD0=>), [Daniel Kwasi Ahorsu](#) (<https://sciprofiles.com/profile/1303687>), [Nasrin Taghizadeh](#) (<https://sciprofiles.com/profile/author/Z0xaY1ZoUnpCcVY0am95Vk1XTUJwcjcyTEIPMEhQVVgwT0NtTmFoeVFDZz0=>), [Zahra Parsapour](#) (<https://sciprofiles.com/profile/author/MDg5K1FWRVFSRHJtVXd6cm1SQXkxRlBcG5wbXJtT2lzRUVUt1RvbE1lbz0=>), [Babak Nejati](#) (<https://sciprofiles.com/profile/2394693>), [Hsin-Pao Chen](#) (<https://sciprofiles.com/profile/2394694>) and [Amir H. Pakpour](#) (<https://sciprofiles.com/profile/795065>)

Healthcare **2022**, *10*(9), 1745; <https://doi.org/10.3390/healthcare10091745> (<https://doi.org/10.3390/healthcare10091745>) - 11 Sep 2022

Cited by [3](#) ([/2227-9032/10/9/1745#metrics](#)) | Viewed by 1420

Abstract The present study examined the mediating role of anxiety, depression, sleepiness, insomnia, and sleep quality in the association between problematic social media use and quality of life (QoL) among patients with cancer. This cross-sectional survey study recruited 288 patients with cancer to respond [...] [Read more](#).

(This article belongs to the Special Issue [The Correlates of Technology-Based Addictive Behaviors and Their Impacts on Wellbeing](#) ([/journal/healthcare/special_issues/Technology_Based_Addictive_Behaviors_Impacts](#)))

► Show Figures

(https://pub.mdpi-res.com/healthcare/healthcare-10-01745/article_deploy/html/images/healthcare-10-01745-g001-550.jpg?1662891026).

Open Access Article

([/2227-9032/10/8/1448/pdf?version=1660809674](#))

[Internet-Related Instruments \(Bergen Social Media Addiction Scale, Smartphone Application-Based Addiction Scale, Internet Gaming Disorder Scale-Short Form, and Nomophobia Questionnaire\) and Their Associations with Distress among Malaysian University Students](#) ([/2227-9032/10/8/1448](#))

by [Serene En Hui Tung](#) (<https://sciprofiles.com/profile/author/U25JdndBODhsQ0hZbzVOekw0Z2czazVid3NDd2pHU1MreGhYdUE3ME04TT0=>), [Wan Ying Gan](#) (<https://sciprofiles.com/profile/327775>), [Jung-Sheng Chen](#) (<https://sciprofiles.com/profile/1690464>), [Ruckwongpat Kamolthip](#) (<https://sciprofiles.com/profile/author/ZWo1NEXqOVBI5FRVU5Ua2tlb2Fackhpb1FQcDlCS1U2aCtBbnJKRURZbz0=>), [Iqbal Pramukti](#) (<https://sciprofiles.com/profile/author/MkRHdFFTM91U0d6b0JyeTJwempPYTFpLzBjRFN5bVNZam9kR3FFTE5iWT0=>), [Siti R. Nadhiroh](#) (<https://sciprofiles.com/profile/2360229>), [Yen-Ling Chang](#) (<https://sciprofiles.com/profile/2292790>), [Chien-Chin Lin](#) (<https://sciprofiles.com/profile/author/S2J3K0NqaEFYb1RLUGUrUWl3ZGNXbGVnbGJZnJ1e1tQ1lqK3k0dDQ3Zz0=>), [Amir H. Pakpour](#) (<https://sciprofiles.com/profile/1747839>), [Chung-Ying Lin](#) (<https://sciprofiles.com/profile/664254>) and [Mark D. Griffiths](#) (<https://sciprofiles.com/profile/6341>)

Healthcare **2022**, *10*(8), 1448; <https://doi.org/10.3390/healthcare10081448> (<https://doi.org/10.3390/healthcare10081448>) - 02 Aug 2022

Cited by [8](#) ([/2227-9032/10/8/1448#metrics](#)) | Viewed by 2287

Abstract The present study evaluated the psychometric properties of four instruments related to internet use, namely the Bergen Social Media Addiction Scale (BSMAS), Smartphone Application-Based Addiction Scale (SABAS), (nine-item) Internet Gaming Disorder Scale Short Form (IGDS9-SF), and Nomophobia Questionnaire (NMPQ) as well as their [...] [Read more](#).

(This article belongs to the Special Issue [The Correlates of Technology-Based Addictive Behaviors and Their Impacts on Wellbeing](#) ([/journal/healthcare/special_issues/Technology_Based_Addictive_Behaviors_Impacts](#)))

Open Access Article

([/2227-9032/10/5/963/pdf?version=1653382910](#))

[Does Smartphone Addiction, Social Media Addiction, and/or Internet Game Addiction Affect Adolescents' Interpersonal Interactions?](#) ([/2227-9032/10/5/963](#))

by [Shang-Yu Yang](#) (<https://sciprofiles.com/profile/309404>), [Yu-Chi Wang](#) (<https://sciprofiles.com/profile/author/RUIVRHh4c0tQNVRRHvYVFptbHdhbTQwOEhIYtJcDNPMnk0N3YrVkrZaz0=>), [Ya-Chen Lee](#) (<https://sciprofiles.com/profile/author/aWR3WE9PRFA2TmxhWmY5aHI5FRUd2JISjZLly9kanZLd3BFYwP3Nm54Yz0=>), [Ying-Lien Lin](#) (<https://sciprofiles.com/profile/author/S2RPWWpveTNpbCtXTzFoUnVhYTJwRTR2cHpFOGIFdURQT0xnTFJzK2xMOD0=>), [Pei-Lun Hsieh](#) (<https://sciprofiles.com/profile/author/L0NuRtFVRXAYWINbDNmZi9VZjNMY05NRGJQbjdrSTNvcmlPbDdyemp2ND0=>) and [Pin-Hsuan Lin](#) (<https://sciprofiles.com/profile/1261709>)





Healthcare **2022**, *10*(5), 963; <https://doi.org/10.3390/healthcare10050963> (<https://doi.org/10.3390/healthcare10050963>) - 23 May 2022

Cited by [8](#) ([/2227-9032/10/5/963#metrics](#)) | Viewed by 4478

Abstract The purpose of this study is to investigate the correlations that levels of addiction to smartphones, social media, and online games have with levels of real-life and online interpersonal interactions among adolescents. In this cross-sectional study of adolescents in a college in Taiwan, [...] [Read more](#).

(This article belongs to the Special Issue [The Correlates of Technology-Based Addictive Behaviors and Their Impacts on Wellbeing](#) ([/journal/healthcare/special_issues/Technology_Based_Addictive_Behaviors_Impacts](#)))

Reliability Generalization Meta-Analysis of Internet Gaming Disorder Scale ((2227-9032/10/10/1992))

by  [Júlia Gisbert-Pérez \(https://sciprofiles.com/profile/2434225\)](https://sciprofiles.com/profile/2434225),  [Manuel Martí-Vilar \(https://sciprofiles.com/profile/1670116\)](https://sciprofiles.com/profile/1670116),
 [César Merino-Soto \(https://sciprofiles.com/profile/1823482\)](https://sciprofiles.com/profile/1823482) and  [Miguel Vallejos-Flores \(https://sciprofiles.com/profile/2440193\)](https://sciprofiles.com/profile/2440193)
Healthcare **2022**, *10*(10), 1992; <https://doi.org/10.3390/healthcare10101992> (<https://doi.org/10.3390/healthcare10101992>) - 11 Oct 2022
Viewed by 1071

Abstract The objective of this study was to carry out a reliability generalization meta-analysis of the Internet Gaming Disorder Scale (IGDS) to find out if it presents adequate values that justify its application in its four versions: original and abbreviated with dichotomous or ordinal [...] [Read more.](#)
(This article belongs to the Special Issue [The Correlates of Technology-Based Addictive Behaviors and Their Impacts on Wellbeing \(/journal/healthcare/special_issues/Technology_Based_Addictive_Behaviors_Impacts_\)](#))

► Show Figures

https://pub.mdpi-res.com/healthcare/healthcare-10-01992/article_deploy/html/images/healthcare-10-01992-g001-550.jpg?1665487247

Show export options ▾

Displaying articles 1-6

[Healthcare \(/journal/healthcare\)](#), EISSN 2227-9032, Published by MDPI

[RSS \(/rss/journal/healthcare\)](#) [Content Alert \(/journal/healthcare/toc-alert\)](#)

Further Information

[Article Processing Charges \(/apc\)](#)

[Pay an Invoice \(/about/payment\)](#)

[Open Access Policy \(/openaccess\)](#)

[Contact MDPI \(/about/contact\)](#)

[Jobs at MDPI \(https://careers.mdpi.com\)](https://careers.mdpi.com)

Guidelines

[For Authors \(/authors\)](#)

[For Reviewers \(/reviewers\)](#)

[For Editors \(/editors\)](#)

[For Librarians \(/librarians\)](#)

[For Publishers \(/publishing_services\)](#)

[For Societies \(/societies\)](#)

[For Conference Organizers \(/conference_organizers\)](#) [bomgmjdpkdklff/enhanced-reader.html?openApp&pdf=https%3A%2F%2Fmdpi-res.com%2Fdl-attachment%2Fhealthcare%2Fhealthcare-11-00700%2Farticle_deploy%2Fhealthcare-11-00700.pdf%3Fversion%3D1677478556](#)

[Sciforum \(https://sciforum.net\)](https://sciforum.net)

[MDPI Books \(https://www.mdpi.com/books\)](https://www.mdpi.com/books)

[Preprints.org \(https://www.preprints.org\)](https://www.preprints.org)

[Scilit \(https://www.scilit.net\)](https://www.scilit.net)

[SciProfiles \(https://sciprofiles.com\)](https://sciprofiles.com)

[Encyclopedia \(https://encyclopedia.pub\)](https://encyclopedia.pub)

[JAMS \(https://jams.pub\)](https://jams.pub)

[Proceedings Series \(/about/proceedings\)](#)

Follow MDPI

[LinkedIn \(https://www.linkedin.com/company/mdpi\)](https://www.linkedin.com/company/mdpi)

[Facebook \(https://www.facebook.com/MDPIOpenAccessPublishing\)](https://www.facebook.com/MDPIOpenAccessPublishing)




[Twitter \(https://twitter.com/MDPIOpenAccess\)](https://twitter.com/MDPIOpenAccess)

Subscribe to receive issue release notifications and newsletters from MDPI journals

Select options ▼

Enter your email address...

Subscribe

 (/toggle_desktop_layout_cookie)  

© 1996-2023 MDPI (Basel, Switzerland) unless otherwise stated

[Disclaimer](#) [Terms and Conditions \(/about/terms-and-conditions\)](#) [Privacy Policy \(/about/privacy\)](#)

View PDF (chrome-extension://dagcmkpagjlhakfdhnbomgmjdpkdklff/enhanced-reader.html?openApp&pdf=https%3A%2F%2Fmdpi-res.com%2Fd_attachment%2Fhealthcare%2Fhealthcare-11-00700%2Farticle_deploy%2Fhealthcare-11-00700.pdf%3Fversion%3D1677478556)

In addition to the main databases, we also focus on ensuring coverage of our journals within as many scope-specific databases as possible, in order to increase visibility for our authors, and further disseminate their impactful data. Moreover, we also work with various universities and government organizations, to ensure that our journals are listed within country-specific journal ranking lists that are often required by funders or institutions for authors to publish with a journal.

Furthermore, all MDPI journals are digitally preserved cover-to-cover, by the **Swiss National Library** (<https://www.nb.admin.ch/snl/en/home/information-professionals/e-helvetica/e-helvetica-uebersicht.html>) and **CLOCKSS** (<https://clockss.org/>), ensuring long-term archival of all of our publications.

All Content is Open Access and Free for Readers

Journals published by MDPI are fully open access: research articles, reviews or any other content on this platform is available to everyone free of charge. To be able to provide open access journals, we finance publication through **article processing charges (APC)** ([/about/apc](#)); these are usually covered by the authors' institutes or research funding bodies. Read more about our **open access policy** ([/about/openaccess](#)).

MDPI Publication Ethics Statement

MDPI is a member of the Committee on Publication Ethics (**COPE** (<http://publicationethics.org/>)). MDPI takes the responsibility to enforce a rigorous peer-review together with strict ethical policies and standards to ensure to add high quality scientific works to the field of scholarly publication. Unfortunately, cases of plagiarism, data falsification, inappropriate authorship credit, and the like, do arise. MDPI takes such publishing ethics issues very seriously and our editors are trained to proceed in such cases with a zero tolerance policy. To verify the originality of content submitted to our journals, we use **iThenticate** (<https://www.ithenticate.com/>) to check submissions against previous publications.

Memberships

MDPI is a member of



(<http://oaspa.org/membership/members/>)



(<http://sparceurope.org>)

DOAJ







(<http://doaj.org>)

COPE

(<http://publicationethics.org/members/publishers?name=MDPI>)

Article

Internet-Related Instruments (Bergen Social Media Addiction Scale, Smartphone Application-Based Addiction Scale, Internet Gaming Disorder Scale-Short Form, and Nomophobia Questionnaire) and Their Associations with Distress among Malaysian University Students

Serene En Hui Tung^{1,†}, Wan Ying Gan^{2,†}, Jung-Sheng Chen³, Ruckwongpatr Kamolthip⁴, Iqbal Pramukti⁵, Siti R. Nadhiroh⁶, Yen-Ling Chang^{7,*}, Chien-Chin Lin^{8,9,10}, Amir H. Pakpour¹¹, Chung-Ying Lin^{4,12,13,14,*}, and Mark D. Griffiths¹⁵



Citation: Tung, S.E.H.; Gan, W.Y.; Chen, J.-S.; Kamolthip, R.; Pramukti, I.; Nadhiroh, S.R.; Chang, Y.-L.; Lin, C.-C.; Pakpour, A.H.; Lin, C.-Y.; et al. Internet-Related Instruments (Bergen Social Media Addiction Scale, Smartphone Application-Based Addiction Scale, Internet Gaming Disorder Scale-Short Form, and Nomophobia Questionnaire) and Their Associations with Distress among Malaysian University Students. *Healthcare* **2022**, *10*, 1448. <https://doi.org/10.3390/healthcare10081448>

Academic Editor: Daniele Giansanti

Received: 23 June 2022

Accepted: 29 July 2022

Published: 2 August 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

- ¹ Division of Nutrition and Dietetics, School of Health Sciences, International Medical University, Kuala Lumpur 57000, Malaysia; serenetung84@gmail.com
 - ² Department of Nutrition, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, Serdang 43400, Malaysia; wanying@upm.edu.my
 - ³ Department of Medical Research, E-Da Hospital, Kaohsiung 82445, Taiwan; ed113187@edah.org.tw
 - ⁴ Institute of Allied Health Sciences, College of Medicine, National Cheng Kung University, Tainan 701401, Taiwan; kamolthip.681@gmail.com
 - ⁵ Department of Community Health Nursing, Faculty of Nursing, Universitas Padjadjaran, Bandung 45363, Indonesia; iqbalrevi@gmail.com
 - ⁶ Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya 60115, Indonesia; sitinadhiroh@fkm.unair.ac.id
 - ⁷ Department of Family Medicine, Cardinal Tien Hospital, New Taipei 23148, Taiwan
 - ⁸ Department of Laboratory Medicine, National Taiwan University Hospital, Taipei 100225, Taiwan; jake9665@gmail.com
 - ⁹ Division of Hematology and Internal Medicine, National Taiwan University Hospital, Taipei 100225, Taiwan
 - ¹⁰ Graduate Institute of Clinical Medicine, National Taiwan University, Taipei 106319, Taiwan
 - ¹¹ Department of Nursing, School of Health and Welfare, Jönköping University, SE-551 11 Jönköping, Sweden; amir.pakpour@ju.se
 - ¹² Biostatistics Consulting Center, National Cheng Kung University Hospital, College of Medicine, National Cheng Kung University, Tainan 701401, Taiwan
 - ¹³ Department of Occupational Therapy, College of Medicine, National Cheng Kung University, Tainan 701401, Taiwan
 - ¹⁴ Department of Public Health, College of Medicine, National Cheng Kung University, Tainan 701401, Taiwan
 - ¹⁵ International Gaming Research Unit, Psychology Department, Nottingham Trent University, Nottingham NG1 4FQ, UK; mark.griffiths@ntu.ac.uk
- * Correspondence: th.yenlingchang@gmail.com (Y.-L.C.); cylin36933@gmail.com (C.-Y.L.)
† These authors contributed equally to this work.

Abstract: The present study evaluated the psychometric properties of four instruments related to internet use, namely the Bergen Social Media Addiction Scale (BSMAS), Smartphone Application-Based Addiction Scale (SABAS), (nine-item) Internet Gaming Disorder Scale Short Form (IGDS9-SF), and Nomophobia Questionnaire (NMPQ) as well as their associations with psychological distress among Malaysian university students. A total of 380 Malaysian university students (71.6% females, mean age 24.0 years) were recruited through convenience sampling and completed an online survey including questions concerning socio-demographic background, social media addiction, smartphone addiction, internet gaming disorder, and nomophobia. Confirmatory factor analysis and Rasch analysis were applied to evaluate the psychometric properties of the instruments and Cronbach's alpha value and McDonald's omega value were used to confirm the internal consistency reliability of the instruments. The unidimensional structure was confirmed for the BSMAS, SABAS, and IGDS9-SF while the four-factor structure was confirmed for NMPQ. All instruments showed good internal consistency reliability. Promising validity and reliability were confirmed for BSMAS, SABAS, IGDS9-SF, and NMPQ. Therefore, they are useful to assess different types of problematic internet use among

university students in Malaysia. Furthermore, a significant association was observed between internet use and psychological distress. The present study is the first to investigate the validity and reliability of BSMAS, SABAS, IGDS9-SF, and NMPQ among Malaysian university students using rigorous psychometric testing methods (i.e., Rasch analysis).

Keywords: smartphone addiction; social media addiction; problematic internet use; psychometrics; university students; Rasch analysis

1. Introduction

The advancement in technology has led to great use of the internet and young adults are major consumers of internet-based applications and platforms [1]. Because of its convenience, the internet has become an important part of individuals' lives, and this was especially prevalent during the COVID-19 pandemic given the various restrictions forcing individuals to rely on digital technology more than ever before [2,3]. It is estimated that approximately 60% of the world's population uses the internet and that Asia has the largest number of internet users followed by Europe [4].

In Malaysia (where the present study was carried out), the Department of Statistics Malaysia (in 2020) reported that 91.7% of Malaysian households had access to the internet via smartphones (98.2%) and computers (80.0%) and that the prevalence of internet use was 89.6%. Malaysian internet users were reported to use the internet for social networking use (98.0%) and playing or downloading games (87.9%) [5]. One of the reasons that internet use has become so prevalent is the global use of smartphones. The constant availability of smartphones with potentially addictive applications such as social networking sites and online video games can be problematic for a minority [6].

While many benefits have been proposed and confirmed for the use of the internet, such as communication, teaching and learning, information gathering, staying connected with friends and family, and entertainment, the lack of control regarding internet use has also led to an increase in detrimental health-related outcomes both physically and mentally [7]. Some of these health outcomes include poor sleep, musculoskeletal discomfort [8], and psychological distress, such as depression, anxiety, and stress [2,3]. These problems have been reported among social media users [9,10], those having problems with online gaming [11], and problematic smartphone users [12]. Moreover, a phenomenon of fear of not having a smartphone on hand has also been observed and has been termed 'nomophobia' (no mobile phobia). Similar to the aforementioned potentially addictive behaviors, nomophobia is related to negative mental health-related outcomes [13].

Although internet addiction is not classified as a mental disorder in the latest (fifth) edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-5), one online activity has been proposed as a tentative addictive disorder in the DSM-5 (i.e., internet gaming disorder) [14]. Given the importance of assessing various online addictive behavior uses among youth due to the various detrimental health outcomes, psychometric instruments to assess these potentially addictive behaviors are important to identify the severity. Although problematic internet use has been investigated in the past in Malaysia [15,16], these studies were conducted among children, medical students, and patients with depression. Therefore, there is still a lack of research conducted in understanding the addictive behaviors related to internet and smartphone use, especially among university students in Malaysia. As overdependence on the internet is prevalent among young people in Malaysia [16], there is a need to validate instruments that assess problematic use for prevalence profiles and prevent problematic internet use among Malaysian university students.

To date, there are few Malaysian studies to support the use of the Bergen Social Media Addiction Scale (BSMAS), Smartphone Application-Based Addiction Scale (SABAS), (nine-item) Internet Gaming Disorder Scale Short Form (IGDS9-SF), and Nomophobia Questionnaire (NMPQ) in Malaysia. More specifically, the psychometric evidence of these

instruments used in Malaysia has not previously been examined through the application of Rasch analysis, a powerful statistical method that more robustly evaluates the validity and reliability of an instrument in comparison to the classical test theory [17]. Therefore, the purpose of the present study was to examine the psychometric properties of BSMAS, SABAS, IGDS9-SF, and NMPQ among Malaysian university students. After validating the BSMAS, SABAS, IGDS9-SF, and NMPQ, their associations with psychological distress were further investigated.

2. Materials and Methods

2.1. Participants and Data Collection

The participants of the study were recruited between 19 August 2021 and 30 September 2021 utilizing a convenience sampling method. The recruitment was conducted using online social media platforms including *Facebook*, *WhatsApp*, *Instagram*, and *Telegram* and hosted on *Google Forms*. The inclusion criteria of the present study were (i) being a university student and (ii) residing in Malaysia during the survey period. The exclusion criteria were university students who (i) were pregnant or lactating females; (ii) self-reported as having chronic diseases such as cardiovascular diseases and cancer; and (iii) were physically disabled. These exclusion criteria were used because they may influence an individual's internet use and subsequently impact the answers to the internet-related instruments used in the present study. All participants were required to provide informed consent (provided on the first page of the online survey) by clicking the option "*I agree to participate in the study*" before completing the survey. The questions relating to the exclusion criteria were in the initial sociodemographic questions in the online survey and were answered after the participants had agreed to complete the survey. However, no participants were removed from the analysis as none of the participants said they were physically disabled, had a chronic illness, and/or were pregnant/lactating. Study approval was obtained from the Ethics Committee for Research Involving Human Subjects in Universiti Putra Malaysia (JKEUPM; Reference Number: JKEUPM-2021-455) before the commencement of the study.

2.2. Measures

The measures were all in English because English is the main medium of instruction in universities and colleges in Malaysia. In addition, Malaysia is a multicultural country where different languages are used by different ethnic populations (e.g., Malays, Chinese, and Indians) making English a more appropriate language to be used in the present setting. However, cultural bias was likely to exist, and all measures were checked for their cultural adaptations by the present authors using the international guidelines on patient-reported outcomes [18,19]. More specifically, the Malaysian authors checked that all the content descriptions in the measures could be fully understood by the Malaysian participants.

2.3. Socio-Demographic Information

The background information obtained from the participants included age, gender (male or female), ethnicity (Malay, Chinese, or other), study program level (undergraduate or postgraduate), and marital status (single, married, or other). In addition, information on the concerning number of daily hours spent on social media and gaming was also obtained.

2.4. Bergen Social Media Addiction Scale (BSMAS)

The Bergen Social Media Addiction Scale (BSMAS) was used to assess social media addiction and experience of using social media within the past week [20]. It contains six items rated on a five-point Likert scale ranging from 1 (*very rarely*) to 5 (*very often*), yielding a total score ranging from 6 to 30. A higher score on the BSMAS indicates a greater risk of addiction to social media. The BSMAS has been translated into several languages and has shown acceptable psychometric properties across studies. The scale has shown good internal consistency reliability with a Cronbach's α of 0.88 [20].

2.5. Smartphone Application-Based Addiction Scale (SABAS)

The Smartphone Application-Based Addiction Scale (SABAS) was used to assess the risk of developing smartphone addiction within the past week [21]. It contains six items rated on a 6-point Likert scale from 1 (*strongly disagree*) to 6 (*strongly agree*), yielding a total score ranging from 6 to 36. A higher score on the SABAS indicates that an individual is more at risk of addiction to a smartphone application. The scale has shown good internal consistency reliability ($\alpha = 0.81$) and promising concurrent validity [21].

2.6. Internet Gaming Disorder Scale-Short Form (IGDS9-SF)

The Internet Gaming Disorder Scale-Short form (IGDS9-SF) was used to assess the severity of internet gaming disorder in both online and/or offline gaming activities within the past week [22]. It contains nine items rated on a five-point Likert scale ranging from 1 (*never*) to 5 (*very often*), yielding a total score ranging from 9 to 45. A higher score on the IGDS9-SF indicates a higher risk of internet gaming disorder. The scale has shown high internal consistency reliability ($\alpha = 0.90$) and promising concurrent validity [23].

2.7. Nomophobia Questionnaire (NMPQ)

The Nomophobia Questionnaire (NMP-Q) was used to assess the risk of developing fear and being out of mobile phone contact within the past week [24]. The NMP-Q contains 20 items distributed across four factors (i.e., not being able to communicate, losing connectedness, not being able to access information, and giving up convenience). All the items are rated using a seven-point Likert scale from 1 (*strongly disagree*) to 7 (*strongly agree*), yielding a total score ranging from 20 to 140. A higher score on the NMP-Q indicates that an individual suffers more from the fear of being unable to use their mobile phone. The scale has shown high internal consistency reliability ($\alpha = 0.70$ – 0.91) and promising concurrent validity [25].

2.8. Depression, Anxiety, Stress Scale (DASS-21)

The Depression, Anxiety, Stress Scale (DASS-21) was used to assess the level of psychological distress [26]. It contains 21 items (seven items on each subscale of depression, anxiety, and stress) rated on a four-point Likert scale ranging from 0 (*not applied at all*) to 3 (*applied very much*), yielding a total score ranging from 0 to 21 for each subscale. A higher score on the DASS-21 indicates a higher level of psychological distress. The scale has shown high internal consistency reliability ($\alpha = 0.926$) and adequate construct validity among Malaysian university students [27].

2.9. Data Analysis

Descriptive statistics were first carried out to understand (i) the participants' characteristics and (ii) the item score distributions on all four psychometric instruments. Then, three sets of psychometric testing methods were applied to evaluate the properties of the four instruments: (i) internal consistency; (ii) confirmatory factor analysis (CFA); and (iii) Rasch analysis.

For internal consistency, Cronbach's α and McDonald's ω were used to assess how the items in each of the four instruments (and the subscales for the NMPQ) adhered to each other. An acceptable value for either Cronbach's α or McDonald's ω is 0.7 or above [28,29]. For CFA, a diagonally weighted least squares estimator was used to test the factor structures for the four instruments separately. More specifically, unidimensional structure was tested for the BSMAS, SABAS, and IGDS9-SF. A four-factor structure was tested for the NMPQ. Moreover, several fit indices, including a nonsignificant χ^2 test, a comparative fit index (CFI) > 0.9, a Tucker-Lewis index (TLI) > 0.9, a root mean square error of approximation (RMSEA) < 0.08, and a standardized root mean square residual (SRMR) < 0.08 were used together to evaluate whether the tested factor structures (i.e., unidimensional structure for BSMAS, SABAS, and IGDS9-SF; four-factor structure for NMPQ) were supported [30,31]. Factor loadings derived from the CFA were expected to be higher than 0.4 [32]. For Rasch

analysis, a partial credit model (PCM) was used to test whether the items for the BSMAS, SABAS, and IGDS9-SF; and the items for the subscales of the NMPQ fitted their latent underlying concepts. Again, a nonsignificant χ^2 test was used to support the fit between the item and its latent concept. Outfit and infit mean square (MnSq) were also used to test the fit. More specifically, both outfit and infit MnSq between 0.5 and 1.5 indicate good fit, while a value smaller than 0.5 indicates overfit (i.e., redundancy in the latent concept) and a value larger than 1.5 indicates misfit (i.e., out of the latent concept) [33,34]. All the statistical analyses were carried out using R software: the *psych* package was utilized for internal consistency [35]; the *lavaan* package was utilized for CFA [36]; and the *eRm* package was utilized for the Rasch analyses [37]. Moreover, a sample size of at least 300 was needed for the present study based on the participant-item ‘rule of thumb’ ratio. More specifically, a ratio of 15:1 was utilized (i.e., 15 participants for every item on the instrument with the most items). Given that the NMPQ has 20 items, a sample size of 300 was the minimal requirement (i.e., $15 \times 20 = 300$).

Lastly, Pearson correlations were used to examine the associations between the BSMAS, SABAS, IGDS9-SF, and NMPQ scores with the three DASS-21 subscale scores (i.e., depression, anxiety, and stress). IBM SPSS 20.0 (IBM Incorp.: Armond, NY, USA) was used for testing the Pearson correlation.

3. Results

The participants ($N = 380$) were generally young adults (mean age = 24.04 years; SD age = 5.07) with the majority being female ($n = 272$; 71.6%). Slightly over half of the participants were ethnic Chinese ($n = 200$; 52.6%), followed by Malay ($n = 117$; 30.8%). The participants spent an average spent 4.5 h daily on social media (SD = 3.21) and 1.36 h daily on gaming (SD = 2.01). Detailed information on the participants’ characteristics is shown in Table 1.

Table 1. Characteristics of the participants ($N = 380$).

	<i>n</i> (%) or <i>M</i> (SD)
Age in year	24.04 (5.07)
Gender	
<i>Male</i>	108 (28.4)
<i>Female</i>	272 (71.6)
Ethnicity group	
<i>Malay</i>	117 (30.8)
<i>Chinese</i>	200 (52.6)
<i>Indian</i>	39 (10.3)
<i>Others</i>	24 (6.3)
Study program	
<i>Undergraduate</i>	277 (72.9)
<i>Postgraduate</i>	103 (27.1)
Marital status	
<i>Single</i>	350 (92.1)
<i>Married</i>	28 (7.4)
<i>Other</i>	2 (0.5)
Daily hours on social media	4.50 (3.21)
Daily hours on gaming	1.36 (2.01)

The descriptive statistics and distributions of all the items (including BSMAS, SABAS, IGDS9-SF, and NMPQ) are presented in Table 2. All the items demonstrated normal distributions according to small values of skewness (ranging between -0.03 and 0.41 for BSMAS; between -0.47 and 0.27 for SABAS; between 0.40 and 1.61 for IGDS9-SF; between -0.33 and 0.43 for NMPQ) and kurtosis (ranging between -0.95 and -0.62 for BSMAS; between -0.90 and -0.38 for SABAS; between -1.07 and 1.59 for IGDS9-SF; between -1.24 and -0.85 for NMPQ).

Table 2. Score distributions for the items of the internet-related instruments.

	Mean	SD	Skewness	Kurtosis	Range
BSMAS	16.80	5.40	0.17	−0.45	6–30
Item B1	2.93	1.10	−0.03	−0.62	1–5
Item B2	3.03	1.14	0.01	−0.72	1–5
Item B3	2.93	1.25	−0.01	−0.95	1–5
Item B4	2.87	1.13	0.09	−0.67	1–5
Item B5	2.52	1.16	0.34	−0.75	1–5
Item B6	2.51	1.21	0.41	−0.75	1–5
SABAS	21.44	6.71	−0.06	−0.53	6–36
Item S1	3.96	1.41	−0.43	−0.73	1–6
Item S2	3.19	1.38	0.14	−0.76	1–6
Item S3	3.89	1.41	−0.46	−0.58	1–6
Item S4	4.00	1.34	−0.47	−0.38	1–6
Item S5	3.16	1.43	0.17	−0.85	1–6
Item S6	3.24	1.48	0.27	−0.90	1–6
IGDS9-SF	16.87	8.26	0.94	0.09	9–43
Item I1	2.05	1.12	0.76	−0.43	1–5
Item I2	1.77	1.04	1.27	0.86	1–5
Item I3	1.93	1.15	1.03	−0.01	1–5
Item I4	1.86	1.06	1.03	0.10	1–5
Item I5	1.74	1.06	1.33	0.88	1–5
Item I6	1.81	1.09	1.24	0.55	1–5
Item I7	1.68	1.04	1.46	1.23	1–5
Item I8	2.48	1.35	0.40	−1.07	1–5
Item I9	1.56	0.95	1.61	1.59	1–5
NMPQ	78.71	28.67	−0.003	−0.80	20–140
NMPQ_F1	17.05	6.28	−0.25	−0.82	4–28
Item N1	4.23	1.79	−0.19	−0.95	1–7
Item N2	4.53	1.74	−0.29	−0.89	1–7
Item N3	3.87	1.85	−0.08	−1.12	1–7
Item N4	4.42	1.75	−0.30	−0.85	1–7
NMPQ_F2	19.42	7.79	−0.04	−0.83	5–35
Item N5	3.88	1.94	0.07	−1.18	1–7
Item N6	3.47	1.88	0.26	−1.03	1–7
Item N7	4.29	1.93	−0.21	−1.12	1–7
Item N8	3.89	1.91	0.01	−1.13	1–7
Item N9	3.89	1.86	0.00	−1.09	1–7
NMPQ_F3	24.63	9.97	−0.12	−1.00	6–42
Item N10	3.81	1.93	0.08	−1.18	1–7
Item N11	4.47	1.87	−0.33	−1.00	1–7
Item N12	4.11	1.87	−0.14	−1.09	1–7
Item N13	4.29	1.85	−0.21	−1.05	1–7
Item N14	4.17	1.82	−0.18	−1.00	1–7
Item N15	3.79	1.86	0.05	−1.16	1–7
NMPQ_F4	17.60	8.11	0.21	−0.91	5–35
Item N16	3.22	1.88	0.43	−0.99	1–7
Item N17	3.43	1.83	0.26	−1.05	1–7
Item N18	3.58	1.89	0.15	−1.13	1–7
Item N19	3.62	1.87	0.13	−1.12	1–7
Item N20	3.75	1.97	0.09	−1.24	1–7

BSMAS = Bergen Social Media Addiction Scale; SABAS = Smartphone Application Based Addiction Scale; IGDS9-SF = [Nine-item] Internet Gaming Disorder Scale-Short Form; NMPQ = Nomophobia Questionnaire; NMPQ_F1 = Not being able to access information; NMPQ_F2 = Giving up convenience; NMPQ_F3 = Not being able to communicate; NMPQ_F4 = Losing connectedness.

The internal consistencies of all four instruments were very good to excellent (Cronbach's $\alpha = 0.86$ and McDonald's $\omega = 0.91$ for BSMAS; Cronbach's $\alpha = 0.88$ and McDonald's $\omega = 0.93$ for SABAS; Cronbach's $\alpha = 0.95$ and McDonald's $\omega = 0.96$ for IGDS9-SF; and Cronbach's $\alpha = 0.88$ to 0.95 and McDonald's $\omega = 0.90$ to 0.97 for NMPQ). Moreover, CFA

results supported the unidimensionality for the BSMAS (p -value of $\chi^2 = 0.11$; CFI = 0.996; TLI = 0.994; RMSEA = 0.039; and SRMR = 0.045), SABAS (p -value of $\chi^2 = 0.36$; CFI = 0.999; TLI = 0.999; RMSEA = 0.016; and SRMR = 0.038), and IGDS9-SF (p -value of $\chi^2 = 0.99$; CFI = 1.000; TLI = 1.005; RMSEA = 0.000; and SRMR = 0.037). The CFA results supported the four-factor structure of the NMPQ (p -value of $\chi^2 = 1.00$; CFI = 1.000; TLI = 1.003; RMSEA = 0.000; and SRMR = 0.037). Moreover, all the factor loadings in the CFA were larger than 0.4 (Table 3).

Table 3. Psychometric results derived from confirmatory factor analysis on the internet-related instruments.

	BSMAS	SABAS	IGDS9-SF	NMPQ			
Factor loading				NMPQ_F1 ^a	NMPQ_F2 ^a	NMPQ_F3 ^a	NMPQ_F4 ^a
Item 1	0.71	0.64	0.84	0.81	0.72	0.87	0.83
Item 2	0.76	0.75	0.87	0.82	0.73	0.84	0.86
Item 3	0.73	0.80	0.86	0.87	0.77	0.89	0.83
Item 4	0.70	0.81	0.88	0.84	0.83	0.86	0.77
Item 5	0.74	0.74	0.81	–	0.78	0.89	0.83
Item 6	0.68	0.74	0.89	–	–	0.85	–
Item 7	–	–	0.80	–	–	–	–
Item 8	–	–	0.66	–	–	–	–
Item 9	–	–	0.76	–	–	–	–
Fit indices							
χ^2 (df)	14.23 (9)	9.92 (9)	13.17 (27)			107.92 (164)	
p -value	0.11	0.36	0.99			1.00	
CFI	0.996	0.999	1.000			1.000	
TLI	0.994	0.999	1.005			1.003	
RMSEA	0.039	0.016	0.000			0.000	
SRMR	0.045	0.038	0.037			0.037	

BSMAS = Bergen Social Media Addiction Scale; SABAS = Smartphone Application Based Addiction Scale; IGDS9-SF = [Nine-item] Internet Gaming Disorder Scale-Short Form; NMPQ = Nomophobia Questionnaire; NMPQ_F1 = Not being able to access information; NMPQ_F2 = Giving up convenience; NMPQ_F3 = Not being able to communicate; NMPQ_F4 = Losing connectedness; CFI = comparative fit index; TLI = Tucker-Lewis index; RMSEA = root mean square error of approximation; SRMR = standardized root mean square residual. ^a The items depended on the item number for that specific factor; for example, Item 2 for NMPQ_F2 is the fifth item for the entire NMPQ.

Rasch analyses additionally showed the good properties of all the items for all four instruments. More specifically, nonsignificant χ^2 tests together with outfit MnSq and infit MnSq between 0.5 and 1.5 for all the items were found for the BSMAS (outfit MnSq = 0.78 to 0.96; infit MnSq = 0.79 to 0.96), the SABAS (outfit MnSq = 0.67 to 1.12; infit MnSq = 0.68 to 1.13), and the NMPQ (outfit MnSq = 0.59 to 1.13; infit MnSq = 0.62 to 1.05). Only one item in the IGDS9-SF showed misfit (i.e., Item 8 “Do you play in order to temporarily escape or relieve a negative mood (e.g., helplessness, guilt, anxiety)?”), with a significant χ^2 test ($p < 0.001$) and large MnSq (outfit MnSq = 1.87 and infit MnSq = 1.86). However, all other items in the IGDS9-SF fit with the Rasch model (Table 4).

Significant and moderate associations were observed between each psychological distress and the four instruments of BSMAS ($r = 0.393$ to 0.428 ; p -values < 0.001), SABAS ($r = 0.359$ to 0.373 ; p -values < 0.001), IGDS9-SF ($r = 0.331$ to 0.392 ; p -values < 0.001), and NMPQ ($r = 0.295$ to 0.339 ; p -values < 0.001) (Table 5).

Table 4. Psychometric results derived from Rasch analysis on the internet-related instruments.

	χ^2 (df)	<i>p</i> -Value	Outfit MnSq	Infit MnSq
BSMAS				
Item B1	318.94 (362)	0.95	0.88	0.88
Item B2	284.03 (362)	1.00	0.78	0.79
Item B3	315.25 (362)	0.96	0.87	0.83
Item B4	328.67 (362)	0.90	0.91	0.90
Item B5	299.58 (362)	0.99	0.83	0.81
Item B6	348.39 (362)	0.69	0.96	0.96
SABAS				
Item S1	412.55 (369)	0.06	1.12	1.13
Item S2	308.79 (369)	0.99	0.84	0.83
Item S3	269.52 (369)	1.00	0.73	0.73
Item S4	246.98 (369)	1.00	0.67	0.68
Item S5	331.30 (369)	0.92	0.90	0.88
Item S6	329.14 (369)	0.93	0.89	0.90
IGDS9-SF				
Item I1	260.66 (272)	0.68	0.96	0.91
Item I2	160.46 (272)	1.00	0.59	0.68
Item I3	201.62 (272)	1.00	0.74	0.80
Item I4	181.27 (272)	1.00	0.66	0.70
Item I5	233.58 (272)	0.96	0.86	0.90
Item I6	143.75 (272)	1.00	0.53	0.61
Item I7	209.63 (272)	1.00	0.77	0.84
Item I8	511.56 (272)	<0.001	1.87	1.86
Item I9	195.90 (272)	1.00	0.72	0.88
NMPQ				
NMPQ_F1				
Item N1	300.34 (358)	0.99	0.84	0.84
Item N2	226.97 (358)	1.00	0.63	0.62
Item N3	319.70 (358)	0.93	0.89	0.86
Item N4	264.56 (358)	1.00	0.74	0.76
NMPQ_F2				
Item N5	309.39 (355)	0.96	0.87	0.92
Item N6	273.10 (355)	1.00	0.77	0.81
Item N7	286.26 (355)	1.00	0.80	0.81
Item N8	262.99 (355)	1.00	0.74	0.75
Item N9	320.29 (355)	0.91	0.90	0.91
NMPQ_F3				
Item N10	372.04 (347)	0.17	1.07	1.05
Item N11	267.68 (347)	1.00	0.77	0.81
Item N12	275.19 (347)	1.00	0.79	0.79
Item N13	209.69 (347)	1.00	0.60	0.63
Item N14	296.00 (347)	0.98	0.85	0.83
Item N15	360.07 (347)	0.30	1.04	1.00
NMPQ_F4				
Item N16	294.66 (340)	0.96	0.86	0.87
Item N17	222.45 (340)	1.00	0.65	0.65
Item N18	201.80 (340)	1.00	0.59	0.62
Item N19	384.68 (340)	0.048	1.13	1.05
Item N20	329.46 (340)	0.65	0.97	0.95

BSMAS = Bergen Social Media Addiction Scale; SABAS = Smartphone Application Based Addiction Scale; IGDS9-SF = [Nine-item] Internet Gaming Disorder Scale-Short Form; NMPQ = Nomophobia Questionnaire; NMPQ_F1 = Not being able to access information; NMPQ_F2 = Giving up convenience; NMPQ_F3 = Not being able to communicate; NMPQ_F4 = Losing connectedness; Outfit MnSq = outfit mean square; Infit MnSq = infit mean square.

Table 5. Correlation between psychological distress and internet-related instruments.

	<i>r</i>		
	Depression	Anxiety	Stress
BSMAS	0.428	0.393	0.421
SABAS	0.368	0.359	0.373
IGDS9-SF	0.392	0.331	0.357
NMPQ	0.304	0.295	0.339

BSMAS = Bergen Social Media Addiction Scale; SABAS = Smartphone Application Based Addiction Scale; IGDS9-SF = (Nine-item) Internet Gaming Disorder Scale-Short Form; NMPQ = Nomophobia Questionnaire. Note: All *p*-values < 0.001.

4. Discussion

To the best of the present authors' knowledge, the present study is the first to examine the psychometric properties of four internet problem-related instruments (i.e., BSMAS, SABAS, IGDS9-SF, and NMPQ) among university students in Malaysia utilizing both classical and modern test theories. The psychometric results demonstrated good factor structures for all four instruments and confirmed the unidimensional structure of the BSMAS, SABAS, IGDS9-SF, and the four-factor structure of the NMPQ. Such factor structure findings were consistent with previous studies [38–43]. In short, all instruments demonstrated robust psychometric properties among Malaysian university students.

4.1. BSMAS

The results for BSMAS specifically showed that the scale had a unidimensional structure with acceptable internal consistency indicating that the instrument is able to assess the use of problematic social media use among Malaysia university students across different ethnic groups. The unidimensional structure of the instrument was found to be similar to recent Korean [38] and Spanish [41] studies. With the rise in the use of social media among youth for social interaction and information gathering [44], there is a need to ensure that the instrument used to assess problematic social media use is valid and tested for its psychometric properties. With the confirmed validity and the very good psychometric properties of BSMAS using the Rasch model analysis approach in the present study, the findings here echo the Rasch analysis findings of the Persian BSMAS, both showing that all BSMAS items embedded in the unidimensional structure [45].

4.2. SABAS

The findings for SABAS also indicated a unidimensional structure with very good internal consistency on this scale, confirming the reliability of the instrument in assessing problematic smartphone application use among university students. The findings related to the unidimensional structure and reliability were similar to a previous study [39]. Moreover, the findings of Rasch analysis in the present study concur with a prior Persian SABAS validation study [46] (i.e., all SABAS items embedded in the unidimensional structure). With the rise in the use of smartphones as a means of communication [47], there is a need to assess the dependency of users on smartphones due to the growing incidence of problematic smartphone use. Previous studies conducted in Malaysia have used the Smartphone Addiction Scale (SAS), [48] but a limitation of using the SAS is that it does not specifically assess the activity engagement of smartphone applications although the SAS has good psychometric properties. Therefore, the findings of the present study provide an alternative option with the focus on smartphone applications for healthcare providers and researchers to examine smartphone addiction among university students in Malaysia.

4.3. IGDS9-SF

Similar to BSMAS and SABAS, very good internal consistency and a unidimensional structure were found in the present study for IGDS9-SF, confirming findings from previous studies conducted among Turkish [40] and Sri Lankan [41] populations. The uni-

dimensional structure of IGDS9-SF was also confirmed in a recent systematic review by Poon et al. [49] which examined the psychometric properties of the instrument in 21 studies comprising 15 different language versions (i.e., English, European Portuguese, South American Portuguese, Spanish, Albanian, Italian, Turkish, Slovenian, Polish, Persian, Malay, Korean, and Chinese, with three sublanguages of traditional Chinese in Hong Kong, traditional Chinese in Taiwan, and simplified Chinese in mainland China). Similarly, the one-factor structure of the IGDS9-SF was also found for the Malay version of the instrument in a previous study [50]. However, the psychometric properties of the Malay version of the instrument in the previous study were only tested using classical test theory and did not incorporate any Rasch analysis.

Despite the unidimensional structure found for the IGDS9-SF, Item 8 (“Do you play in order to temporarily escape or relieve a negative mood [e.g., helplessness, guilt, anxiety]?”) which assesses behavior escape from negative mood through gaming was found to have psychometric problems with relatively high infit and outfit MnSq values. One of the possible explanations is that this item may be unable to differentiate between disordered and non-disordered gamers. One of the explanations is that relieving a negative mood could be a form of positive coping strategy rather than a loss of control [51]. Therefore, the behavior escape assessed by Item 8 might be practiced by both disordered and non-disordered gamers [52]. This was confirmed by the low accuracy and specificity in the ability of this item to differentiate between disordered and non-disordered gamers in a previous study [53].

4.4. NMPQ

Lastly, the findings for NMPQ confirmed the four-factor structure of the scale with an acceptable range of internal consistency for all four factors. These findings support previous findings among Persians [25], Greek [54], Portuguese [55], and Indonesian populations [56] where similar reliability and four-factor structure were observed in both populations. However, it should be noted that the previous Persian study was conducted among adolescents rather than university students. Nevertheless, findings from the present study confirm that the NMPQ is a valid and robust research instrument due to its good psychometric properties and internal consistencies. Therefore, the NMPQ is able to assess nomophobia among university students in Malaysia.

4.5. Associations between Psychological Distress and the Four Instruments

The findings from this study were consistent with previous studies [42,57–59], whereby BSMAS, SABAS, IGDS9-SF, and NMPQ were correlated with depression, anxiety, and stress. A systematic review looking into problematic social networking site use (PSNSU) assessed through BSMAS and comorbid psychiatric disorders in recent large-scale studies found positive associations between BSMAS and depression, anxiety, and stress [57]. Similarly, a study among a 964 sample of internet gamers recruited online found a positive relationship between anxiety, depression, and internet gaming assessed using the IGDS9-SF [58]. Another study among 11,014 primary and secondary schoolteachers examined the association between problematic internet use, psychological distress, and sleep problems during COVID-19 and found positive correlations between BSMAS and SABAS with psychological distress among both high sleep and low sleep problem groups [59]. As for nomophobia, similar results were found where a positive correlation was found between NMPQ with depression, anxiety, and stress among 500 European Portuguese adults [43]. The relationship between psychological distress and BSMAS was the strongest compared to SABAS, IGDS9-SF, and NMPQ which further confirms the conclusion that problematic social networking site use among adults is associated with comorbid psychiatric disorder outcomes [57].

4.6. Implications

Some of the practical implications of the present study include that these four instruments could be used to study problematic internet use behaviors among Malaysian university students by researchers and healthcare practitioners in the future. The risk of possible internet addiction could also be assessed through these instruments. Nevertheless, these instruments are non-diagnostic tools, and further development of clinical diagnostic criteria is needed to assess disordered internet use behaviors.

4.7. Limitations

There are a number of study limitations. First, the instruments were only tested among university students making it difficult to generalize the results to other age groups such as children and adults. Therefore, there is a need to explore the psychometric properties of these instruments among individuals across the lifespan and of other age groups. Second, the participants were recruited using a convenience sampling method, meaning the results are not generalizable to the whole population of university students in Malaysia. Third, the self-reported nature of the instruments could lead to biased findings due to participants providing socially desirable answers and experiencing recall bias. Fourth, as mentioned in Section 4.6, these instruments are non-diagnostic tools and cannot be used for diagnosis. Moreover, some of these behaviors are likely to be a consequence of other comorbid disorders. Therefore, future studies are needed to examine the association between problematic internet use and comorbid psychiatric disorders, which would likely expand the usefulness of these instruments. Fifth, the present sample had a gender imbalance (i.e., 72% female participants). Two reasons may explain the high ratio of female participants in the present study: (i) the proportion of female students in Malaysian public higher education institutions has been approximately 60% since 2012 [60]; (ii) females are more likely than males to participate in scientific studies [61]. Therefore, future psychometric evaluations of these instruments need to have more gender-balanced samples to corroborate the present study's findings. Sixth, the present sample was homogenous (given that all the participants were university students and predominantly female). Therefore, it is not known if the instruments evaluated in the present study can be applied to non-university students with diverse characteristics. Consequently, it is recommended that future studies should recruit more diverse samples (i.e., participants with a much broader demographic of education, age, gender, socioeconomic status, ethnicity, marital status, and type of job (e.g., knowledge worker versus manual laborer)) to further evaluate the psychometric properties of these instruments and their relationships with mental health. Given this limitation, the present study was unable to determine if Malaysian university students had higher levels of problematic internet use than non-university students.

5. Conclusions

The findings of the present study demonstrated that BSMAS, SABAS, IGDS9-SF, and NMPQ are valid and reliable instruments and are easy to use in assessing internet problem-related behaviors among university students in Malaysia. In addition, the psychometric properties of these instruments were found to be robust and valid. However, special attention is needed for Item 8 of IGDS9-SF (i.e., using gaming to escape) due to the misfit in the Rasch model. Furthermore, a significant association was found between internet use and psychological distress, which further confirms the concerns about internet use and addiction among university students.

Author Contributions: Conceptualization, S.E.H.T., W.Y.G., Y.-L.C., C.-Y.L. and A.H.P.; methodology, J.-S.C., R.K., I.P., S.R.N., C.-C.L. and M.D.G.; software, C.-Y.L.; validation, J.-S.C., R.K., I.P., S.R.N., Y.-L.C., A.H.P. and M.D.G.; formal analysis, C.-Y.L.; investigation, S.E.H.T., W.Y.G. and C.-Y.L.; resources, S.E.H.T., W.Y.G. and C.-Y.L.; data curation, S.E.H.T. and W.Y.G.; writing—original draft preparation, S.E.H.T., W.Y.G., Y.-L.C. and C.-Y.L.; writing—review and editing, J.-S.C., R.K., I.P., S.R.N., C.-C.L., A.H.P. and M.D.G.; visualization, C.-Y.L.; supervision, C.-Y.L.; project administration,

C.-Y.L.; funding acquisition, S.E.H.T., W.Y.G. and C.-Y.L. All authors have read and agreed to the published version of the manuscript.

Funding: Financial support from the Ministry of Science and Technology, Taiwan (MOST 110-2410-H-006-115; MOST 111-2410-H-006-100), the Higher Education Sprout Project, Ministry of Education to the Headquarters of University Advancement at National Cheng Kung University (NCKU), and the 2021 Southeast and South Asia and Taiwan Universities Joint Research Scheme (NCKU 31).

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki, and study approval was obtained from the Ethics Committee for Research Involving Human Subjects in Universiti Putra Malaysia (JKEUPM; Reference Number: JKEUPM-2021-455).

Informed Consent Statement: Informed consent using e-form was obtained from all subjects involved in the study.

Data Availability Statement: The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Acknowledgments: We thank all the participants to take part in the study.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Yam, C.W.; Pakpour, A.H.; Griffiths, M.D.; Yau, W.Y.; Lo, C.M.; Ng, J.; Lin, C.Y.; Leung, H. Psychometric testing of three Chinese online-related addictive behavior instruments among Hong Kong university students. *Psychiatr. Q.* **2019**, *90*, 117–128. [CrossRef] [PubMed]
2. Chen, C.Y.; Chen, I.H.; Hou, W.L.; Potenza, M.N.; O'Brien, K.S.; Lin, C.Y.; Latner, J.D. The relationship between children's problematic internet-related behaviors and psychological distress during the onset of the COVID-19 pandemic: A longitudinal study. *J. Addict. Med.* **2022**, *16*, e73–e80. [CrossRef] [PubMed]
3. Fernandes, B.; Uzun, B.; Aydin, C.; Tan-Mansukhani, R.; Vallejo, A.; Saldaña-Gutierrez, A.; Biswas, U.N.; Essau, C.A. Internet use during COVID-19 lockdown among young people in low- and middle-income countries: Role of psychological wellbeing. *Addict. Behav. Rep.* **2021**, *14*, 100379. [CrossRef] [PubMed]
4. Johnson, J. Internet Usage Worldwide—Statistics & Facts. Available online: https://www.statista.com/topics/1145/internet-usage-worldwide/#topicHeader__wrapper (accessed on 30 March 2022).
5. Department of Statistics Malaysia. ICT Use and Access by Individuals and Household Survey Report, Malaysia. Available online: https://www.dosm.gov.my/v1/index.php?r=column/cthemByCat&cat=395&bul_id=OWUvVnV5SHI2WFU2VFhnQ2ZjTm1Bdz09&menu_id=amVoWU54UTI0a21NWmdhMjFMMWcyZz09 (accessed on 30 March 2022).
6. Van Velthoven, M.H.; Powell, J.; Powell, G. Problematic smartphone use: Digital approaches to an emerging public health problem. *Digit. Health* **2018**, *4*, 2055207618759167. [CrossRef] [PubMed]
7. Van Deursen, A.J.; Helsper, E.J. Collateral benefits of Internet use: Explaining the diverse outcomes of engaging with the Internet. *New Media Soc.* **2018**, *20*, 2333–2351. [CrossRef]
8. Yang, G.; Cao, J.; Li, Y.; Cheng, P.; Liu, B.; Hao, Z.; Yao, H.; Shi, D.; Peng, L.; Guo, L.; et al. Association between internet addiction and the risk of musculoskeletal pain in Chinese college freshmen—A cross-sectional study. *Front. Psychol.* **2019**, *10*, 1959. [CrossRef]
9. Boer, M.; van den Eijnden, R.; Boniel-Nissim, M.; Wong, S.L.; Inchley, J.C.; Badura, P.; Craig, W.M.; Gobina, I.; Kleszczewska, D.; Klanšček, H.J.; et al. Adolescents' intense and problematic social media use and their well-being in 29 countries. *J. Adolesc. Health* **2020**, *66*, S89–S99. [CrossRef]
10. Huang, C. A meta-analysis of the problematic social media use and mental health. *Int. J. Soc. Psychiatry* **2022**, *68*, 12–33. [CrossRef] [PubMed]
11. Männikkö, N.; Ruotsalainen, H.; Miettunen, J.; Pontes, H.M.; Käriäinen, M. Problematic gaming behaviour and health-related outcomes: A systematic review and meta-analysis. *J. Health Psychol.* **2020**, *25*, 67–81. [CrossRef] [PubMed]
12. Yang, J.; Fu, X.; Liao, X.; Li, Y. Association of problematic smartphone use with poor sleep quality, depression, and anxiety: A systematic review and meta-analysis. *Psychiatry Res.* **2020**, *284*, 112686. [CrossRef] [PubMed]
13. Rodríguez-García, A.M.; Moreno-Guerrero, A.J.; López Belmonte, J. Nomophobia: An individual's growing fear of being without a smartphone—A systematic literature review. *Int. J. Environ. Res. Public Health* **2020**, *17*, 580. [CrossRef]
14. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*, 5th ed.; American Psychiatric Association: Washington, DC, USA, 2013.
15. Lim, P.K.; Amer Nordin, A.S.; Yee, A.; Tan, S.B. Prevalence of smartphone addiction in patients with depression and its association with depression severity: A cross-sectional study. *Int. J. Ment. Health Addict.* **2021**, *19*, 919–933. [CrossRef]
16. Nik Jaafar, N.R.; Bahar, N.; Ibrahim, N.; Baharudin, A.; Wan Ismail, W.S.; Sim, S.T.; Abdul Aziz, M.; Tan, K.A. Are Malaysian youths overdependent on the Internet? A narrative review. *Front. Psychiatry* **2021**, *12*, 710790. [CrossRef]
17. Baghaei, P. The Rasch model as a construct validation tool. *Rasch Meas. Trans.* **2008**, *22*, 1145–1146.

18. Beaton, D.E.; Bombardier, C.; Guillemin, F.; Ferraz, M.B. Guidelines for the process of cross-cultural adaptation of self-report measures. *Spine* **2000**, *25*, 3186–3191. [[CrossRef](#)] [[PubMed](#)]
19. Wild, D.; Grove, A.; Martin, M.; Eremenco, S.; McElroy, S.; Verjee-Lorenz, A.; Erikson, P.; ISPOR Task Force for Translation and Cultural Adaptation. Principles of good practice for the translation and cultural adaptation process for Patient-Reported Outcomes (PRO) measures: Report of the ISPOR Task Force for Translation and Cultural Adaptation. *Value Health* **2005**, *8*, 94–104. [[CrossRef](#)]
20. Schou Andreassen, C.; Billieux, J.; Griffiths, M.D.; Kuss, D.J.; Demetrovics, Z.; Mazzoni, E.; Pallesen, S. The relationship between addictive use of social media and video games and symptoms of psychiatric disorders: A large-scale cross-sectional study. *Psychol. Addict. Behav.* **2016**, *30*, 252–262. [[CrossRef](#)]
21. Csibi, S.; Griffiths, M.D.; Cook, B.; Demetrovics, Z.; Szabo, A. The psychometric properties of the Smartphone Application-Based Addiction Scale (SABAS). *Int. J. Ment. Health Addict.* **2018**, *16*, 393–403. [[CrossRef](#)]
22. Pontes, H.M.; Griffiths, M.D. Measuring DSM-5 internet gaming disorder: Development and validation of a short psychometric scale. *Comput. Hum. Behav.* **2015**, *45*, 137–143. [[CrossRef](#)]
23. Wu, T.Y.; Lin, C.Y.; Årestedt, K.; Griffiths, M.D.; Broström, A.; Pakpour, A.H. Psychometric validation of the Persian nine-item Internet Gaming Disorder Scale—Short Form: Does gender and hours spent online gaming affect the interpretations of item descriptions? *J. Behav. Addict.* **2017**, *6*, 256–263. [[CrossRef](#)] [[PubMed](#)]
24. Yildirim, C.; Correia, A.P. Exploring the dimensions of nomophobia: Development and validation of a self-reported questionnaire. *Comput. Hum. Behav.* **2015**, *49*, 130–137. [[CrossRef](#)]
25. Lin, C.Y.; Griffiths, M.D.; Pakpour, A.H. Psychometric evaluation of Persian Nomophobia Questionnaire: Differential item functioning and measurement invariance across gender. *J. Behav. Addict.* **2018**, *7*, 100–108. [[CrossRef](#)] [[PubMed](#)]
26. Lovibond, S.H.; Lovibond, P.F. *Manual for the Depression Anxiety & Stress Scales*, 2nd ed.; Psychology Foundation: Sydney, Australia, 1995.
27. Ahmad, N.; Roslan, S.; Othman, S.; Shukor, S.F.A.; Bakar, A.Y.A. The validity and reliability of psychometric profile for Depression, Anxiety and Stress scale (DASS21) instrument among Malaysian undergraduate students. *Int. J. Acad. Res. Bus. Soc. Sci.* **2018**, *8*, 812–827. [[CrossRef](#)]
28. Nunnally, J.C. *Psychometric Theory*, 2nd ed.; McGraw-Hill: New York, NY, USA, 1978.
29. Hayes, A.F.; Coutts, J.J. Use omega rather than Cronbach's alpha for estimating reliability. But *Commun. Methods Meas.* **2020**, *14*, 1–24. [[CrossRef](#)]
30. Nejati, B.; Fan, C.W.; Boone, W.J.; Griffiths, M.D.; Lin, C.Y.; Pakpour, A.H. Validating the Persian Intuitive Eating Scale-2 among breast cancer survivors who are overweight/obese. *Eval. Health Prof.* **2021**, *44*, 385–394. [[CrossRef](#)] [[PubMed](#)]
31. Lin, C.Y.; Broström, A.; Griffiths, M.D.; Pakpour, A.H. Psychometric evaluation of the Persian eHealth Literacy Scale (eHEALS) among elder Iranians with heart failure. *Eval. Health Prof.* **2020**, *43*, 222–229. [[CrossRef](#)] [[PubMed](#)]
32. Hair, J.F.; Babin, B.J.; Anderson, R.E.; Black, W.C. *Multivariate Data Analysis*, 8th ed.; Cengage: Boston, MA, USA, 2018.
33. Fan, C.W.; Chen, J.S.; Addo, F.M.; Adjaottor, E.S.; Amankwaah, G.B.; Yen, C.F.; Ahorsu, D.K.; Lin, C.Y. Examining the validity of the drivers of COVID-19 Vaccination Acceptance Scale using Rasch analysis. *Expert Rev. Vaccines* **2022**, *21*, 253–260. [[CrossRef](#)] [[PubMed](#)]
34. Poorebrahim, A.; Lin, C.Y.; Imani, V.; Kolvani, S.S.; Alaviyoun, S.A.; Ehsani, N.; Pakpour, A.H. Using Mindful Attention Awareness Scale on male prisoners: Confirmatory factor analysis and Rasch models. *PLoS ONE* **2021**, *16*, e0254333. [[CrossRef](#)] [[PubMed](#)]
35. Revelle, W. Psych: Procedures for Psychological, Psychometric, and Personality Research. R Package Version 2.2.3. Available online: <https://CRAN.R-project.org/package=psych> (accessed on 30 March 2022).
36. Rosseel, Y. Lavaan: An R package for structural equation modeling. *J. Stat. Softw.* **2012**, *48*, 1–36. [[CrossRef](#)]
37. Mair, P.; Hatzinger, R.; Maier, M.J. eRm: Extended Rasch Modeling. 1.0-2. Available online: <https://cran.r-project.org/package=eRm> (accessed on 30 March 2022).
38. Shin, N.Y. Psychometric properties of the Bergen Social Media Addiction Scale in Korean young adults. *Psychiatry Investig.* **2022**, *19*, 356–361. [[CrossRef](#)]
39. Ali, A.M.; Hendawy, A.O.; Abd Elhay, E.S.; Ali, E.M.; Alkhamees, A.A.; Kunugi, H.; Hassan, N.I. The Bergen Facebook Addiction Scale: Its psychometric properties and invariance among women with eating disorders. *BMC Womens Health* **2022**, *22*, 99. [[CrossRef](#)]
40. Vally, Z.; Alowais, A. Assessing risk for smartphone addiction: Validation of an Arabic Version of the Smartphone Application-Based Addiction Scale. *Int. J. Ment. Health Addict.* **2022**, *20*, 691–703. [[CrossRef](#)]
41. Ali, A.M.; Al-Amer, R.; Atout, M.; Ali, T.S.; Mansour, A.M.H.; Khatatbeh, H.; Alkhamees, A.A.; Hendawy, A.O. The nine-item Internet Gaming Disorder Scale (IGDS9-SF): Its psychometric properties among Sri Lankan students and measurement invariance across Sri Lanka, Turkey, Australia, and the USA. *Healthcare* **2022**, *10*, 490. [[CrossRef](#)]
42. Chen, I.-H.; Strong, C.; Lin, Y.-C.; Tsai, M.-C.; Leung, H.; Lin, C.-Y.; Pakpour, A.H.; Griffiths, M.D. Time invariance of three ultra-brief internet-related instruments: Smartphone Application-Based Addiction Scale (SABAS), Bergen Social Media Addiction Scale (BSMAS), and the nine-item Internet Gaming Disorder Scale- Short Form (IGDS-SF9) (Study Part B). *Addict. Behav.* **2020**, *101*, 105960. [[CrossRef](#)]

43. Chen, I.H.; Ahorsu, D.K.; Pakpour, A.H.; Griffiths, M.D.; Lin, C.Y.; Chen, C.Y. Psychometric properties of three simplified Chinese online-related addictive behavior instruments among mainland Chinese primary school students. *Front. Psychiatry* **2020**, *11*, 875. [[CrossRef](#)] [[PubMed](#)]
44. Whiting, A.; Williams, D. Why people use social media: A uses and gratifications approach. *Qual. Mark. Res.* **2013**, *16*, 362–369. [[CrossRef](#)]
45. Lin, C.Y.; Broström, A.; Nilsen, P.; Griffiths, M.D.; Pakpour, A.H. Psychometric validation of the Persian Bergen Social Media Addiction Scale using classic test theory and Rasch models. *J. Behav. Addict.* **2017**, *6*, 620–629. [[CrossRef](#)]
46. Lin, C.Y.; Imani, V.; Broström, A.; Nilsen, P.; Fung, X.C.; Griffiths, M.D.; Pakpour, A.H. Smartphone application-based addiction among Iranian adolescents: A psychometric study. *Int. J. Ment. Health Addict.* **2019**, *17*, 765–780. [[CrossRef](#)]
47. Ataş, A.H.; Çelik, B. Smartphone use of university students: Patterns, purposes, and situations. *Malays. Online J. Educ. Sci.* **2019**, *7*, 54–70. [[CrossRef](#)]
48. Ching, S.M.; Yee, A.; Ramachandran, V.; Lim, S.M.S.; Sulaiman, W.A.W.; Foo, Y.L.; Hoo, F.K. Validation of a Malay Version of the Smartphone Addiction Scale among medical students in Malaysia. *PLoS ONE* **2015**, *10*, e0139337. [[CrossRef](#)] [[PubMed](#)]
49. Poon, L.Y.J.; Tsang, H.W.H.; Chan, T.Y.J.; Man, S.W.T.; Ng, L.Y.; Wong, Y.L.E.; Lin, C.Y.; Chien, C.W.; Griffiths, M.D.; Pontes, H.M.; et al. Psychometric properties of the Internet Gaming Disorder Scale-Short-Form (IGDS9-SF): Systematic review. *J. Med. Internet Res.* **2021**, *23*, e26821. [[CrossRef](#)] [[PubMed](#)]
50. Ling, S.L.; Nik Jaafar, N.R.; Tan, K.A.; Bahar, N.; Baharudin, A.; Ahmad Tajjudin, A.I. Psychometric properties of the Malay version of the Internet Gaming Disorder Scale-Short Form (IGDS9-SF-M): Evidence from a sample of Malaysian undergraduates. *Int. J. Environ. Res. Public Health* **2021**, *18*, 2592. [[CrossRef](#)]
51. Kardefelt-Winther, D. A critical account of DSM-5 criteria for internet gaming disorder. *Addict. Res. Theory* **2015**, *23*, 93–98. [[CrossRef](#)]
52. Kardefelt-Winther, D. A conceptual and methodological critique of internet addiction research: Towards a model of compensatory internet use. *Comput. Hum. Behav.* **2014**, *31*, 351–354. [[CrossRef](#)]
53. Ko, C.H.; Yen, J.Y.; Chen, S.H.; Wang, P.W.; Chen, C.S.; Yen, C.F. Evaluation of the diagnostic criteria of internet gaming disorder in the DSM-5 among young adults in Taiwan. *J. Psychiatr. Res.* **2014**, *53*, 103–110. [[CrossRef](#)] [[PubMed](#)]
54. Charalambos, G.; Venetia, N.; Elissavet, V.; Vasilis, G.; Areti, L. Validity of the Greek NMP-Q and sociodemographic determinants of nomophobia among university students. *Int. J. Hum.-Comput. Interact.* **2022**, *1*–9. [[CrossRef](#)]
55. Galhardo, A.; Loureiro, D.; Raimundo, E.; Massano-Cardoso, I.; Cunha, M. Assessing nomophobia: Validation study of the European Portuguese version of the Nomophobia Questionnaire. *Community Ment. Health J.* **2020**, *56*, 1521–1530. [[CrossRef](#)] [[PubMed](#)]
56. Rangka, I.B.; Prasetyaningtyas, W.E.; Ifdil, I.; Ardi, Z.; Suranata, K.; Winingsih, E.; Sofyan, A.; Irawan, M.; Arjanto, P.; Muslifar, R.; et al. Measuring psychometric properties of the Indonesian version of the NoMoPhobia Questionnaire (NMPQ): Insight from Rasch measurement tool. *J. Phys. Conf. Ser.* **2018**, *1114*, 012127. [[CrossRef](#)]
57. Hussain, Z.; Griffiths, M.D. Problematic social networking site use and comorbid psychiatric disorders: A systematic review of recent large-scale studies. *Front. Psychiatry* **2018**, *9*, 686. [[CrossRef](#)] [[PubMed](#)]
58. Stavropoulos, V.; Vassallo, J.; Burleigh, T.L.; Gomez, R.; Colder, C.M. The role of internet gaming in the association between anxiety and depression: A preliminary cross-sectional study. *Asia-Pac. Psychiatry* **2022**, *14*, e12474. [[CrossRef](#)]
59. Lee, Z.H.; Chen, I.H. The association between problematic internet use, psychological distress, and sleep problems during COVID-19. *Sleep Epidemiol.* **2021**, *1*, 100005. [[CrossRef](#)] [[PubMed](#)]
60. Hirschmann, R. Number of Students Enrolled in Public Higher Education Institutions in Malaysia from 2012 to 2020, by Gender. 2022. Available online: <https://www.statista.com/statistics/794845/students-in-public-higher-education-institutions-by-gender-malaysia/#:~:text=Premium%20statistics-,Students%20in%20public%20higher%20education,Malaysia%202012%2D2020%2C%20by%20gender&text=In%202020%2C%20around%20234.08%20thousand,enrolled%20in%20public%20higher%20institutions> (accessed on 28 July 2022).
61. Galea, S.; Tracy, M. Participation rates in epidemiologic studies. *Ann. Epidemiol.* **2007**, *17*, 643–653. [[CrossRef](#)] [[PubMed](#)]