



## **Vol. 7 No. 1 (2023): AMERTA NUTRITION (Bilingual Edition)**

**Current Issue**



Volume 7, Issue 1, March 2023

p-ISSN: 2580-1163, e-ISSN: 2580-9776

# Amerta Nutrition



Published in collaboration with:

**IAGIKMI**

**&**

**UNIVERSITAS AIRLANGGA**

Accredited by the Ministry of Research and Higher Education Republic Indonesia  
Number SK: 10/E/KPT/2019



Amerta  
Nutr.

Volume 7

Issue 1

Page  
1-174

Surabaya,  
March 2023

p-ISSN : 2580-1163  
e-ISSN : 2580-9776

**Vol. 7 No. 1 (2023): AMERTA NUTRITION (Bilingual Edition)**

Published: 2023-03-03

**Front Matter****Front Matter Vol. 7 No. 1 March 2023**

📄 Abstract : 109

**Back Matter****Back Matter Vol. 7 No. 1 March 2023**

📄 Abstract : 69

**Original Articles****Association between Energy and Macronutrient Intake and Sleep Duration with Nutritional Status of New Students of the Faculty of Public Health, Universitas Airlangga during Online Courses**

Hubungan Asupan Energi dan Zat Gizi Makro serta Durasi Tidur dengan Status Gizi Mahasiswa Baru Fakultas Kesehatan Masyarakat Universitas Airlangga selama Kuliah Online

DOI : 10.20473/amnt.v7i1.2023.1-6

👤 Chrysoprase Thasya Abihail<sup>(1)</sup>, Ketut Herlin Simanoah<sup>(2)</sup>, Lailatul Muniroh<sup>(3)</sup>

(1) Departemen of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia ,

(2) Departemen of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia ,

(3) Departemen of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia

📄 1-6

📄 Abstract : 406

📄 PDF (English Version) : 46PDF (Versi Bahasa) : 84


[📄 PDF \(English Version\)](#)[📄 PDF \(Versi Bahasa\)](#)



## Correlation between Diabetes Mellitus Type 2, Cholesterol with Calcium Score in Patient with Hypertension and Obesity

Hubungan Diabetes Melitus, Kolesterol dengan Skor Kalsium pada Pasien Hipertensi dengan Status Gizi Obesitas

 DOI : 10.20473/amnt.v7i1.2023.7-13


 Saskia Dyah Handari <sup>(1)</sup>, Mirna Rahmasari <sup>(2)</sup>, Yurike Dhika Adhela <sup>(3)</sup>


(1) Departemen Kardiologi, Fakultas Kedokteran, Universitas Brawijaya, Malang, Indonesia ,

(2) Department of Community Nutrition, Faculty of Human Ecology, IPB University, Bogor, Indonesia ,

(3) Departemen Gizi, Fakultas Kesehatan Masyarakat, Universitas Airlangga, Surabaya, Indonesia

 7-13

 Abstract : 561

 PDF (English Version) : 38PDF (Versi Bahasa) : 100

 [PDF \(English Version\)](#)

 [PDF \(Versi Bahasa\)](#)



## Snacking Habits, Strict Diet, BMI, and Body Image of Adolescents in Three Sub-Districts in Depok and Bogor

Kebiasaan Jajan, Diet Ketat, Indeks Masa Tubuh dan Persepsi Body Image pada Remaja di Tiga Kecamatan di Depok dan Bogor

 DOI : 10.20473/amnt.v7i1.2023.14-19


 Lina Agestika <sup>(1)</sup>, Ratnayani Ratnayani <sup>(2)</sup>

(1) Nutrition Department, Faculty of Health Sciences and Technology, Binawan University, Jakarta, Indonesia ,

(2) Nutrition Department, Faculty of Health Sciences and Technology, Binawan University, Jakarta, Indonesia

 14-19

 Abstract : 269

 PDF (English Version) : 30PDF (Versi Bahasa) : 101



[PDF \(English Version\)](#)[PDF \(Versi Bahasa\)](#)

### Zinc Intake Affects Toddler Stunting: A Cross-Sectional Study on Toddlers Aged 3 Years

Asupan Zinc Berpengaruh pada Stunting Balita : Studi Belah Lintang pada Balita Usia 3 Tahun

DOI : 10.20473/amnt.v7i1.2023.20-26

Sri Priyantini <sup>(1)</sup>, Anisa Nurmalitasari <sup>(2)</sup>, Masyhudi AM <sup>(3)</sup>

(1) Departemen Ilmu Kesehatan Anak, Fakultas Kedokteran, Universitas Islam Sultan Agung, Semarang, Indonesia ,

(2) Program Studi Kedokteran Umum, Fakultas Kedokteran, Universitas Islam Sultan Agung, Semarang, Indonesia ,

(3) Departemen Ilmu Kesehatan Masyarakat, Fakultas Kedokteran, Universitas Islam Sultan Agung, Semarang, Indonesia

20-26

Abstract : 234

PDF (English Version) : 43PDF (Versi Bahasa) : 23

[PDF \(English Version\)](#)[PDF \(Versi Bahasa\)](#)

### Consumption Traditional Food Description in Children Under Five in the Coastal Family of Bengkulu City, Indonesia

Gambaran Konsumsi Makanan Tradisional pada Balita di Keluarga Pesisir di Kota Bengkulu, Indonesia

DOI : 10.20473/amnt.v7i1.2023.27-36

Betty Yosephin Simanjuntak <sup>(1)</sup>, Desri Suryani <sup>(2)</sup>, Miratul Haya <sup>(3)</sup>, Ali Khomsan <sup>(4)</sup>

(1) Politeknik Kesehatan Kementerian Kesehatan Bengkulu, Bengkulu, Indonesia ,

(2) Politeknik Kesehatan Kementerian Kesehatan Bengkulu, Bengkulu, Indonesia ,

(3) Politeknik Kesehatan Kementerian Kesehatan Bengkulu, Bengkulu, Indonesia ,

(4) Departemen Gizi Masyarakat, Fakultas Ekologi Manusia, IPB University, Bogor, Indonesia

📄 27-36

📄 Abstract : 241

📄 PDF (English Version) : 30PDF (Versi Bahasa) : 24

📄 PDF (English Version)

📄 PDF (Versi Bahasa)



### Low Birth Weight Related Factors at Kertek 2 Public Health Centre Wonosobo Regency

Faktor-Faktor yang Berhubungan dengan Kejadian Berat Badan Lahir Rendah di Puskesmas Kertek 2 Kabupaten Wonosobo

DOI : 10.20473/amnt.v7i1.2023.37-44

👤 Anggit Rizkika <sup>(1)</sup>, Mohammad Zen Rahfiludin <sup>(2)</sup>, Alfi Fairuz Asna <sup>(3)</sup>

(1) Faculty of Public Health, Diponegoro University, Semarang, Indonesia ,

(2) Faculty of Public Health, Diponegoro University, Semarang, Indonesia ,

(3) Faculty of Public Health, Diponegoro University, Semarang, Indonesia

📄 37-44

📄 Abstract : 224

📄 PDF (English Version) : 42PDF (Versi Bahasa) : 35

📄 PDF (English Version)

📄 PDF (Versi Bahasa)



### Food Consumption Pattern Affects Vitamin D Levels and Quality of Life in Children during the Second Growth Spurt Period

## Pola Konsumsi Makanan Mempengaruhi Kadar Vitamin D dan Kualitas Hidup Anak pada Masa Growth Spurt Kedua

DOI : 10.20473/amnt.v7i1.2023.45-53

Atina Husaana <sup>(1)</sup>, Siti Thomas Zulaikhah <sup>(2)</sup>, Ratnawati Ratnawati <sup>(3)</sup>

(1) Department of Pharmacology, Medical Faculty, Universitas Islam Sultan Agung, Semarang, Indonesia ,

(2) Department of Public Health, Medical Faculty, Universitas Islam Sultan Agung, Semarang, Indonesia ,

(3) Department of Public Health, Medical Faculty, Universitas Islam Sultan Agung, Semarang, Indonesia

45-53

Abstract : 362

PDF (English Version) : 61PDF (Versi Bahasa) : 33

[PDF \(English Version\)](#)

[PDF \(Versi Bahasa\)](#)



## The Association of Clean and Healthy Living Behavior along with Hemoglobin and Serum Ferritin Levels among Adolescent Girls

Hubungan Perilaku Hidup Bersih dan Sehat dengan Kadar Hemoglobin dan Feritin Serum pada Remaja Putri

DOI : 10.20473/amnt.v7i1.2023.54-62

Suci Amalia <sup>(1)</sup>, Sri Anna Marliyati <sup>(2)</sup>, Mira Dewi <sup>(3)</sup>, Dwinita Wikan Utami <sup>(4)</sup>

(1) Sekolah Tinggi Ilmu Kesehatan Holistik, Purwakarta, Indonesia ,

(2) Departemen Gizi Masyarakat, Fakultas Ekologi Manusia, IPB University, Bogor, Indonesia ,

(3) Departemen Gizi Masyarakat, Fakultas Ekologi Manusia, IPB University, Bogor, Indonesia ,

(4) Badan Riset Inovasi Nasional, Bogor, Indonesia

54-62

Abstract : 331

PDF (English Version) : 49PDF (Versi Bahasa) : 36

[PDF \(English Version\)](#)


[PDF \(Versi Bahasa\)](#)



## Nutritional Knowledge in Improving Immunity through Healthy Eating Habits during the Covid-19 Pandemic

Pengetahuan Terkait Gizi dalam Upaya Meningkatkan Imunitas melalui Kebiasaan Makan selama Pandemi Covid-19

 DOI : 10.20473/amnt.v7i1.2023.63-69


 Sri Wahyuni <sup>(1)</sup>, Sri Sumarmi <sup>(2)</sup>, Fathrizqita Aghnia Raudhany <sup>(3)</sup>, Trias Mahmudiono <sup>(4)</sup>

(1) Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia ,


(2) Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia ,

(3) Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia ,

(4) Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia

 63-69

 Abstract : 205

 PDF (English Version) : 28PDF (Versi Bahasa) : 33

 [PDF \(English Version\)](#)


 [PDF \(Versi Bahasa\)](#)



## Implementation of Premarital Counseling Description in Grobogan District, Central Java, Indonesia

Gambaran Implementasi Kursus Pranikah di Kabupaten Grobogan, Jawa Tengah, Indonesia

 DOI : 10.20473/amnt.v7i1.2023.70-78

 Sri Achadi Nugraheni <sup>(1)</sup>, Atik Mawarni <sup>(2)</sup>, Cahya Tri Purnami <sup>(3)</sup>, Sri Winarni <sup>(4)</sup>, Alfi Fairuz Asna <sup>(5)</sup>, Apoina Kartini <sup>(6)</sup>, Septo Pawelas Arso <sup>(7)</sup>,  
Novia Handayani <sup>(8)</sup>, Naila Fauziatin <sup>(9)</sup>

(1) Faculty of Public Health, Universitas Diponegoro, Semarang, Indonesia ,

(2) Faculty of Public Health, Universitas Diponegoro, Semarang, Indonesia ,

(3) Faculty of Public Health, Universitas Diponegoro, Semarang, Indonesia ,

(4) Faculty of Public Health, Universitas Diponegoro, Semarang, Indonesia ,

(5) Faculty of Public Health, Universitas Diponegoro, Semarang, Indonesia ,



(6) Faculty of Public Health, Universitas Diponegoro, Semarang, Indonesia ,  
(7) Faculty of Public Health, Universitas Diponegoro, Semarang, Indonesia ,  
(8) Faculty of Public Health, Universitas Diponegoro, Semarang, Indonesia ,  
(9) Faculty of Public Health, Universitas Diponegoro, Semarang, Indonesia  
📄 70-78

📄 Abstract : 212

📄 PDF (English Version) : 37PDF (Versi Bahasa) : 19

📄 PDF (English Version)

📄 PDF (Versi Bahasa)



### Trends Analysis Scope of Fe-3 Tablet Administration and ANC K4 Activities towards Pregnancy Complications in Surabaya in 2019

Analisis Kecenderungan Cakupan Pemberian Tablet Fe-3 dan Cakupan ANC K4 terhadap Kejadian Komplikasi Kehamilan di Surabaya Tahun 2019

doi DOI : 10.20473/amnt.v7i1.2023.79-87

👤 Sukma Arum Sekar Taji <sup>(1)</sup>, Lucia Yovita Hendrati <sup>(2)</sup>

(1) Department of Epidemiology, Biostatistics, Population Studies and Health Promotion, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia ,

(2) Department of Epidemiology, Biostatistics, Population Studies and Health Promotion, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia

📄 79-87

📄 Abstract : 179

📄 PDF (English Version) : 27PDF (Versi Bahasa) : 21

📄 PDF (English Version)


📄 PDF (Versi Bahasa)



### Total Flavonoid and Antioxidant Activity of Food Bar Torbangun – Katuk on The Effectiveness of Breast Milk Production


Total Flavonoid dan Aktivitas Antioksidan Food Bar Torbangun – Katuk terhadap Efektivitas Produksi ASI

 DOI : 10.20473/amnt.v7i1.2023.88-97


 Laeli Lutfiani <sup>(1)</sup>, Nanang Nasrulloh <sup>(2)</sup>

(1) Program Studi Gizi Program Sarjana, Fakultas Ilmu Kesehatan, Universitas Pembangunan Nasional Veteran Jakarta, Depok, Indonesia ,

(2) Program Studi Gizi Program Sarjana, Fakultas Ilmu Kesehatan, Universitas Pembangunan Nasional Veteran Jakarta, Depok, Indonesia

 88-97

 Abstract : 167

 PDF (English Version) : 35PDF (Versi Bahasa) : 73

 [PDF \(English Version\)](#)


 [PDF \(Versi Bahasa\)](#)



### Correlation Between Body Image, Eating Disorders, and Nutrient Adequacy Level with Nutritional Status of Adolescent Swimmers in Bogor City, Indonesia


Hubungan Persepsi Tubuh, Gangguan Makan, dan Tingkat Kecukupan Gizi dengan Status Gizi Atlet Renang Remaja di Kota Bogor, Indonesia

 DOI : 10.20473/amnt.v7i1.2023.98-111


 Rani Assyifa <sup>(1)</sup>, Hadi Riyadi <sup>(2)</sup>

(1) Department of Community Nutrition, Faculty of Human Ecology, IPB University, Bogor, Indonesia ,

(2) Department of Community Nutrition, Faculty of Human Ecology, IPB University, Bogor, Indonesia

 98-111

 Abstract : 382

 PDF (English Version) : 47PDF (Versi Bahasa) : 111

[PDF \(English Version\)](#)[PDF \(Versi Bahasa\)](#)

### **Apo-B Levels and Abdominal Aortic Wall Thickness in Hypercholesterolemic Rats Treated with Red Guava Fruit**

Kadar Apo-B dan Ketebalan Dinding Aorta Abdominalis Tikus Hiperkolesterolemia dengan Perlakuan Buah Jambu Biji Merah

DOI : 10.20473/amnt.v7i1.2023.112-119

Sugeng Maryanto <sup>(1)</sup>, Dian Oktianti <sup>(2)</sup>

(1) Department of Nutrition, Faculty of Health, Ngudi Waluyo University, Semarang, Indonesia ,

(2) Department of Pharmacy, Faculty of Health, Ngudi Waluyo University, Semarang, Indonesia

112-119

Abstract : 153

PDF (English Version) : 34PDF (Versi Bahasa) : 31

[PDF \(English Version\)](#)[PDF \(Versi Bahasa\)](#)

### **Risk Factors of Stunting, Iron Deficiency Anemia, and Their Coexistence among Children Aged 6-9 Years in Indonesia: Results from the Indonesian Family Life Survey-5 (IFLS-5) in 2014-2015**

Faktor Risiko Stunting, Anemia Defisiensi Besi, dan Koeksistensinya pada Anak Usia 6-9 Tahun di Indonesia: Hasil dari Indonesian Family Life Survey (IFLS-5) tahun 2014-2015

DOI : 10.20473/amnt.v7i1.2023.120-130

Mia Mustika Hutria Utami <sup>(1)</sup>, Lilik Kustiyah <sup>(2)</sup>, Cesilia Meti Dwiriani <sup>(3)</sup>

(1) Department of Community Nutrition, Faculty of Human Ecology, IPB University, Bogor, Indonesia ,

(2) Department of Community Nutrition, Faculty of Human Ecology, IPB University, Bogor, Indonesia ,

(3) Department of Community Nutrition, Faculty of Human Ecology, IPB University, Bogor, Indonesia

📄 120-130

📄 Abstract : 192

📄 PDF (English Version) : 87PDF (Versi Bahasa) : 89

📄 PDF (English Version)

📄 PDF (Versi Bahasa)



### Food Access Elderly Women Head Household in Kumpulrejo Salatiga, Indonesia

Akses Pangan Lansia Perempuan Kepala Keluarga di Kelurahan Kumpulrejo Salatiga, Indonesia

doi DOI : 10.20473/amnt.v7i1.2023.131-138

👤 Eza Media Arlan <sup>(1)</sup>, Theresia Pratiwi Elingsetyo Sanubari <sup>(2)</sup>, Firdhan Aria Wijaya <sup>(3)</sup>

(1) Program Studi Gizi, Fakultas Kedokteran dan Ilmu Kesehatan, Universitas Kristen Satya Wacana, Salatiga, Indonesia ,

(2) Program Studi Gizi, Fakultas Kedokteran dan Ilmu Kesehatan, Universitas Kristen Satya Wacana, Salatiga, Indonesia ,

(3) Program Studi Sosiologi, Fakultas Ilmu Sosial dan Ilmu Politik, Universitas Pattimura, Ambon, Indonesia

📄 131-138

📄 Abstract : 158

📄 PDF (English Version) : 24PDF (Versi Bahasa) : 20

📄 PDF (English Version)

📄 PDF (Versi Bahasa)


### Literature Review




**Literature Review: The Effect of Mediterranean Diet on Lipid Profile and Fasting Blood Glucose in Overweight Obese**



## Studi Literatur: Pengaruh Diet Mediterania terhadap Profil Lipid dan Glukosa Darah Puasa pada Orang Overweight atau Obesitas

 DOI : 10.20473/amnt.v7i1.2023.139-146

 Gustina Berta Uli <sup>(1)</sup>, Sekar Ramadhanti Asyahir <sup>(2)</sup>, Leny Budhi Harti <sup>(3)</sup>


(1) Program Studi Pendidikan Profesi Dietisien, Fakultas Ilmu Kesehatan, Universitas Brawijaya, Indonesia ,

(2) Program Studi Pendidikan Profesi Dietisien, Fakultas Ilmu Kesehatan, Universitas Brawijaya, Indonesia ,

(3) Departemen Gizi, Fakultas Ilmu Kesehatan, Universitas Brawijaya, Malang, Indonesia

 139-146

 Abstract : 427

 PDF (English Version) : 53PDF (Versi Bahasa) : 94

 [PDF \(English Version\)](#)

 [PDF \(Versi Bahasa\)](#)



## Literature Review: The Relationship between Dietary Diversity with Stunting in Underfive Children

Tinjauan Literatur: Hubungan antara Keragaman Pangan dengan Stunting pada Balita


 DOI : 10.20473/amnt.v7i1.2023.147-153

 Ulfa Al Uluf <sup>(1)</sup>, Alfadhila Khairil Sinatrya <sup>(2)</sup>, Siti Rahayu Nadhroh <sup>(3)</sup>


(1) Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia ,

(2) Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia ,

(3) Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia

 147-153

 Abstract : 180

 PDF (English Version) : 43PDF (Versi Bahasa) : 102

 [PDF \(English Version\)](#)


 [PDF \(Versi Bahasa\)](#)



## Literature Review: Food Supplement Intervention to Increase Z-Score Height for Age in Stunting Children

Tinjauan Literatur: Intervensi Suplemen Makanan untuk Meningkatkan Z-Skor PB/U pada Balita Stunting


 DOI : 10.20473/amnt.v7i1.2023.154-160

 Maya Fernandya Siahaan <sup>(1)</sup>, Auni Rahmatika <sup>(2)</sup>, Siti Rahayu Nadhiroh <sup>(3)</sup>


(1) Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia ,

(2) Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia ,

(3) Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia

 154-160

 Abstract : 165

 PDF (English Version) : 118PDF (Versi Bahasa) : 42

 [PDF \(English Version\)](#)

 [PDF \(Versi Bahasa\)](#)



## Ultra-Processed Food can be a Mediator between Food Security Status and Overweight or Obesity among Adults: A Literature Review

Makanan Ultra-Proses Berperan sebagai Mediator Hubungan Ketahanan Pangan dengan Status Kelebihan Gizi atau Obesitas pada Dewasa: Literature Review

 DOI : 10.20473/amnt.v7i1.2023.161-174

 Farah Faza <sup>(1)</sup>, Unun Fitri Febria Bafani <sup>(2)</sup>, Idri Iqra Fikha <sup>(3)</sup>

(1) Department of Nutrition and Health, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia ,

(2) Department of Nutrition, Faculty of Medicine, Universitas Indonesia, Jakarta, Indonesia ,

(3) Nutrition Science Undergraduate Study Program, STIKes Pekanbaru Medical Center, Riau, Indonesia

 161-174

 Abstract : 379



## Editorial Team



**Trias Mahmudiono S.KM., M.PH (Nutr), GCAS., Ph.D**

### Editor In Chief

Department of Health Nutrition, Faculty of Public Health, Universitas Airlangga, Indonesia

0000-0002-3128-2173

SQuNnN0AAAAJ

57189899256

5988526



**Dr. Beben Benyamin**

### Editorial Board

Biostatistics School of Health Sciences, University of South Australia, Australia

0000-0001-5608-2293

wMoqGWQAAAAJ

16314927500

-



**Assoc. Prof. Hazreen Bin Abdul Majid, Bsc. Dietetic.,**

**M.Nut.Dietetic., Ph.D**

### Editorial Board

Department Social and Preventive Medicine, Faculty of Medicine, University of Malaya, Malaysia

0000-0002-2718-8424

OsnphloAAAAJ

54893067500

-



**Dr. Agung Dwi Laksono, S.KM, M.Kes**

### Editorial Board

Center for Humanities, Health Policy, and Community Empowerment Research and Development Agency, Ministry of Health of the Republic of Indonesia., Indonesia

0000-0002-9056-0399

Hb0ypnwAAAAJ

56429657100

-



### Prof. Roger Hughes MPH, Ph.D

#### Editorial Board

Public Health and Health Systems, University of Tasmania, Australia

 0000-0002-5811-7960

-

7404299920



-



### Prof. Jörg-Ingolf Stein

#### Editorial Board

Department of Pediatrics III, Medizinische Universität Innsbruck, Austria

 -

-

6504189694



-



### Professor. C.A. Kalpana, M.Sc., Bed., M.Phil., PhD

#### Editorial Board

Avinashilingam Institute for Home Science and Higher Education for Women, India

 -

oo5KloUAAAAJ

-



-



### Assoc. Prof. Wantanee Kriengsinyos, Ph.D., RD

#### Editorial Board

Human Nutrition Division, Institute of Nutrition, Mahidol University, Thailand

 0000-0001-8262-5095

-

6506938692



-



### Febi Dwi Rahmadi, S. KM., M. Sc., Ph. D

#### Editorial Board

Centre for Environment and Population Health, Griffith School of Environment, Nathan Campus, Brisbane, Queensland, Australia

 0000-0002-9367-3452

-

55900228400



-





## Susy Katikana Sebayang, SP., M.Sc, Ph.D

### Editorial Board

Department of Biostatistic and Populations, Faculty of Public Health, PDD Banyuwangi, Universitas Airlangga, Indonesia

 0000-0003-0470-8308

y\_tZ-9IAAAAJ

24068188900

 6057476



## Dr. Luh Ade Ari Wiradnyani, M.Sc

### Editorial Board

SEAMEO Regional Center for Food and Nutrition, Indonesia

 0000-0002-4349-8588

-

57079273000

 -



## Dr. dr. Sri Adiningsih, MS., MCN

### Editorial Board

The National Sports Committee of Indonesia, East Java Province, Indonesia

 0000-0003-0875-1880

lwh\_Nd4AAAAJ

57214881157

 6072899



## Dr. Farida Wahyu Ningtyas, S.KM, M.Kes

### Editorial Board

Department of Health Nutrition, Faculty of Public Health, University of Jember, Indonesia

 0000-0002-2324-5137

JLcSwq8AAAAJ

-

 -



## Dr. Kadek Tresna Adhi, S.KM, M.Kes

### Section Editor in Community Nutrition

Department of Public Health and Preventive Medicine, Faculty of Medicine, Udayana University, Indonesia

 0000-0002-1963-448X

-

57211602808


 5974664



### Mahmud Aditya Rifqi, S.Gz, M.Si

#### Section Editor in Public Health Nutrition and Food Service Management

Department of Health Nutrition, Faculty of Public health, Universitas Airlangga, Indonesia

 0000-0003-4953-6159

aYihx7cAAAAJ

57208282840

 6006040



### Farapti dr., M.Gizi

#### Section Editor in Clinical Nutrition and Dietetics

Indonesian Medical Association, East Java Province, Indonesia

 0000-0001-9753-3855

\_4He0t8AAAAJ

57196396416

 6026885



### Dina Rahayuning Pangestuti, STP., M.Gizi

#### Section Editor in Food and Nutrition

Faculty of Public Health, Universitas Diponegoro, Indonesia

 -

IpoAW-QAAAAJ

57192903853


 6008377



### Fatqiatul Wulandari, S.Gz.

#### Editorial Assistant

Department of Health Nutrition, Faculty of Public Health, Universitas Airlangga, Indonesia

 0000-0002-1229-1496

-

 -



### Utari Gita Setyawati, S.Gz.

#### Editorial Assistant

Department of Health Nutrition, Faculty of Public Health, Universitas Airlangga, Indonesia

 -

-

 -

lecturers, students, public health nutritionists, registered dietitians, public health practitioners and other practitioners that focus on nutrition in Indonesia and worldwide.

## **Publications Ethics**

### **PUBLICATION ETHICS**

**Amerta Nutrition** ([pISSN 2580-1163](#), [e-ISSN 2580-9776](#)) is a double-blind peer-reviewed electronic journal. This letter explains the ethical rules of conduct established by all parties involved in the publication of articles into this journal, including the authors, the editor in chief, the editorial board, peer-reviewer as well as publisher in this case is Universitas Airlangga. This statement is adopted under [COPE's Best Practice Guidelines for Journal Editors](#).

### **ETHICAL GUIDELINE FOR JOURNAL PUBLICATION**

The publication of an article using peer-reviewed process in **Amerta Nutrition** is very important to build nutritional science. This is a real reflection of the work or the research of the authors and institutions from which they came from. The peer-reviewed article ensures the correct scientific method and maintains **Amerta Nutrition** quality standards. Therefore, it is very important for all parties involved in the Amerta Nutrition publications to maintain ethical standards of good worth as writers, editors, peer reviewers, publishers and society in general.

Universitas Airlangga as the publisher of **Amerta Nutrition** carries out its responsibility to maintain the quality standard of publishing an article in every publishing stage seriously and we understand the responsibilities and ethics that must be maintained. We are committed to keeping the Amerta Nutrition sponsorship, printing, distribution and commercialization of having no impact or effect on editorial decisions on an article.

### **DUTIES OF EDITOR**

#### **Publication Decisions**

The editor-in-chief of Amerta Nutrition is responsible for determining the manuscripts that have been submitted to be peer reviewed and if the manuscript is meeting the editorial policies and journal's requirement it will be published. Validation of research or literature review and the importance of the manuscript for scientific development, subsequent researchers as well as readers of Amerta Nutrition has always been a major consideration in decision making. Editors of Amerta Nutrition may refer to the Board of Editors directives and always refer to the rules and regulations that exist in the Republic of Indonesia regarding copyright and plagiarism. Chairman of the board of editors may authorize other members of the editor for decision-making by observing the peer review results.

#### **Equality**

The editor will always uphold the sense of fairness and fairness at all times in evaluating the manuscript regardless of race, sex, sexual orientation, religion and belief, ethics, citizenship or political ideology of the author.

#### **Confidentiality**

Editors and all editorial staff are not allowed to disclose information about manuscripts that have been signed by Amerta Nutrition to anyone other than to the corresponding author, editorial board, and publisher if feasible.

### **Disclosure and Conflicts of Interest**

Unpublished material from a manuscript that enters Amerta Nutrition is not allowed to be used by the editor as a form of his own research without written consent or written consent from the author.

## **DUTIES OF REVIEWERS**

### **Contribution to Editorial Decisions**

Peer review assists the editor in making editorial decisions through the editorial communications with the author may also assist the author in improving the paper.

The journal review uses a double-blind review. Both reviewers will review the same manuscript and the decision will be announced after all of the reviewers give the evaluation.

### **Promptness**

Any selected referee who feels unqualified to review the research reported in a manuscript or knows that its prompt review will be impossible should notify the editor and excuse himself from the review process.

### **Confidentiality**

Any manuscripts received for review must be treated as confidential documents. They must not be shown to or discussed with others except as authorized by the editor.

### **Standards of Objectivity**

Reviews should be conducted objectively. Personal criticism of the author is inappropriate. Referees should express their views clearly with supporting arguments.

### **Acknowledgment of Sources**

Reviewers should identify relevant published work that has not been cited by the authors. Any statement that an observation, derivation, or argument had been previously reported should be accompanied by the relevant citation. A reviewer should also call to the editor's attention any substantial similarity or overlap between the manuscript under consideration and any other published paper of which they have personal knowledge.

### **Disclosure and Conflict of Interest**

Privileged information or ideas obtained through peer review must be kept confidential and not used for personal advantage. Reviewers should not consider manuscripts in which they have conflicts of interest resulting from competitive, collaborative, or other relationships or connections with any of the authors, companies, or institutions connected to the papers.



## Literature Review: The Relationship between Dietary Diversity with Stunting in Underfive Children

### Tinjauan Literatur: Hubungan antara Keragaman Pangan dengan Stunting pada Balita

Ulfa Al Uluf<sup>1</sup>, Alfadhila Khairil Sinatrya<sup>1</sup>, Siti Rahayu Nadhiroh<sup>1\*</sup>

<sup>1</sup>Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia

#### ARTICLE INFO

Received: 14-02-2022

Accepted: 23-08-2022

Published online: 03-03-2023

\*Correspondent:

Siti Rahayu Nadhiroh

[stinadhiroh@fkm.unair.ac.id](mailto:stinadhiroh@fkm.unair.ac.id)

DOI:

10.20473/amnt.v7i1.2023.147-153

Available online at:

<https://e-journal.unair.ac.id/AMNT>

Keywords:

Dietary diversity, Underfive children, Stunting

#### ABSTRACT

**Background:** Stunting is a global health problem in children under five. Failure to grow, develop, and metabolize due to stunting can threaten a child's future. The diversity of food consumed daily can be a factor affecting the incidence of stunting.

**Objectives:** This literature aimed to determine the relationship between dietary diversity and the incidence of stunting in children under five.

**Discussion:** Ten articles met the criteria, with six articles stating that there was a relationship between dietary diversity and stunting, stating that the more diverse the food groups consumed, the less probability of children under five getting stunted.

**Conclusions:** This study concludes that dietary diversity is related to the incidence of stunting in children under five. This research can be used as a reference to maximize the availability of dietary diversity, especially in the locus area of stunting.

#### INTRODUCTION

Stunting is a world health problem experienced by more than a quarter (26%) of approximately 165 million children under the age of 5 worldwide. Indonesia ranks fifth with the most significant number of stunting<sup>1</sup>. In Southeast Asia, the prevalence of stunting has reached 14.9 million<sup>2</sup>. Stunting is when a toddler has a shorter length or height than his age. Stunting can be measured by length or height above -2 SD (standard deviance) of the WHO growth median in children under five<sup>3</sup>. Data from the Indonesian Nutrition Status Monitoring (PSG) in 2017, the stunting rate in children under five years is higher in the infant group (29.6%) compared to those under five years old (20.1%)<sup>4</sup>. Stunting not only has an impact in the short term but will also affect the subsequent life of toddlers. Among the impacts caused by stunting are a decrease in growth and development decline, cognitive abilities, intelligence, and endurance, and increased non-communicable diseases<sup>5</sup>.

Stunting can be interpreted as a condition of failure to thrive (body and brain) in children due to malnutrition for an extended period, from the fetus in the womb to the beginning of a child's life (the first 1000

days of birth). Inadequate access to nutritious food, inadequate consumption of vitamins and minerals, and poor diversity of food sources and animal protein are the causes of stunting<sup>6</sup>.

Public health related to nutrition depends on the quality of food consumed. Food quality describes all the nutrients the body needs in regulating the amount of food consumed and the ratio of types of food balanced in one plate. The more diverse and balanced the type and content of food consumed, the better the nutritional quality. No food has complete nutritional content or the right amount and type<sup>7</sup>. Several studies on food diversity and stunting say there is a relationship between diversity in food consumption and stunting in toddlers aged 6-24 months<sup>7</sup>. Research in the Cibungbulang Health Center area with a sample of 90 respondents said that 24.4% of children were stunted, and there was a relationship between food diversity and stunting<sup>8</sup>. Consumption of less diverse foods will impact the quality of nutrients and can result in a lack of fulfillment of daily nutrients. Lack of intake of these nutrients will hinder growth and trigger malnutrition, increasing the chance of stunting. Based on the description above, the authors were interested in

compiling a literature review on the relationship between dietary diversity and stunting in children under five.

#### DISCUSSIONS

Table 1 shows the results of the relationship between food diversity and stunting events in toddlers. The search results for literature articles found ten relevant articles. A total of 6 articles stated a relationship between dietary diversity and stunting, while four others said there was no relationship between dietary diversity and stunting.

Stunting is a condition of failure to thrive in children under five years of age (babies under five years of age). Multidimensional causes, including malnutrition at gestational age and poor parenting practices, limited ANC services, inadequate clean water and sanitation, and lack of household access to nutritious food cause stunting<sup>9</sup>. One of the nutritional problems several developing countries face is the lack of food diversity<sup>10</sup>. Food diversity is a type of food group that includes staple foods, side dishes, vegetables, fruit, water, and various types of food in each food group. The more diverse types of food consumed, the easier it is to meet nutritional needs<sup>11</sup>.

In this literature, six articles reported that there was a relationship between dietary diversity and stunting. In line with the results of this study, research conducted in Cimayang Village, Banten, stated that diversity in food consumption was associated with stunting in toddlers aged 6-24 months<sup>7</sup>. Another study shows that dietary diversity was associated with stunting. Poor dietary diversity is a risk factor for stunting<sup>12</sup>. A cross-sectional study in Aligarh also stated that dietary diversity was associated with stunting<sup>13</sup>. Research in the country parts of Myanmar<sup>14</sup> and Northwest Province, South Africa<sup>15</sup> found that stunting was related to food diversity. In Tanzania, 31% of children aged 6-23 months were found to be stunted.

A decrease in the amount of food consumed causes the occurrence of stunting. In addition, the consumption of animal protein was found in this study to reduce stunting<sup>16</sup>. Consuming a variety of foods can reduce the prevalence of stunting. Higher dietary diversity and variety of foods such as corn, fish, legume, and poultry appear to be beneficial for children's developmental growth of children <5 years of age in the Nouna area of Burkina Faso<sup>17</sup>.

Increasing the variety of food for children reduces the risk of stunting and promotes growth<sup>18</sup>. Higher dietary diversity was found to reduce the incidence of stunting and underweight in preschool children aged 4-5 years<sup>15</sup>. A study differentiates the relationship between food diversity in stunting toddlers based on where they live. The study results showed that

the diversity of foods for stunting toddlers in urban and rural areas was not much different<sup>19</sup>. Mothers of toddlers and their families needed to provide a more diverse diet for toddlers, especially types of fruit and vegetables, eggs, nuts, and seeds, and introduce various animal proteins such as meat and chicken liver.

Contrary to research that explains that dietary diversity was related to stunting, some studies stated there was no association between dietary diversity and toddler stunting. In this literature review, four articles stated that dietary diversity was not related to the incidence of stunting in toddlers. Research by Wirawan and Rahmawati (2016)<sup>20</sup> said there were differences in the types of food diversity used. In the research conducted, food diversity was collected based on food diversity at the household level, which was aimed at capturing the ability of households to access a variety of foods. In contrast, in several studies on the nutritional status of children under five, food diversity was used at the individual level (Individual Dietary Diversity Score). Individual dietary diversity score (IDDS). Research by Melaku et al. (2018) also stated that there was no relationship between dietary diversity and stunting caused by differences in sample size, sampling, and method analysis. They were mainly related to the method of analysis. Research by Nurmayasanti and Mahmudiono (2019) also stated that food diversity was not related to stunting or non-stunting probably because of several reasons: stunted mothers or caregivers have been exposed to information at the Integrated Service Post (Posyandu) about stunting and its prevention or efforts to grow to catch up with its growth, including providing food with balanced nutrition and mothers of toddlers, most of whom are homemakers, have much time to prepare food for toddlers and can regularly come to the Posyandu to monitor the growth of toddlers. This result was in line with research conducted by analyzing data from the 2012 Comprehensive Nutrition Survey in the State of Maharashtra, India, which surveyed 2,630 households where no relationship was found between the diversity of children's diets and stunting<sup>21</sup>.

On the contrary, in a literature review study by Purwoko, Triana, and Cahyaningrum (2020), as many as 17 articles stated that food diversity in toddlers was mostly stunting in the non-diverse category<sup>22</sup>. Data disaggregated by age showed that dietary diversity positively correlated with anthropometric status (HAZ) in children aged 24 months to 59 months<sup>23</sup>. Food diversity significantly affects stunting. Accordingly, there was an association between household-level food diversity based on energy consumption. The highest risk factors influencing stunting are exclusive breastfeeding and food diversity<sup>24</sup>.

**Table 1.** Relationship between food diversity and stunting incidence in toddlers

Article title	Sample	Method	Assessment method	Results
Mother's dietary diversity and association with stunting among children <2 years old in a low socioeconomic environment: A case-control study in an urban care setting in Dhaka, Bangladesh <sup>25</sup>	Mother of 296 children aged < 2 (148 in case group and 148 in control group) in Sait House Dhaka, Bangladesh	<ul style="list-style-type: none"> <li>- Design: Case-control</li> <li>- Exposure: a variety of foods within the 10 food groups 24 hours prior to measurement</li> <li>- Outcome: Stunting</li> <li>- Analysis: Chi-square test/Fisher exact test and Logistic regression</li> </ul>	<ul style="list-style-type: none"> <li>- Dietary diversity was measured using a questionnaire about 10 food groups consumed in the previous 24 hours. Food diversity was grouped into <math>\geq 5</math> food groups, and consumption &lt; 5 food groups</li> <li>- The case group was toddlers who are not wasting/underweight and have a length-for-age (LAZ) &lt; -2 SD. The control group was a wasting/underweight toddler who had a LAZ <math>\geq -1.00</math> z score</li> </ul>	Children whose mothers consumed <5 food groups were 1.7 times more likely to be stunted than children whose mothers consumed $\geq 5$ food groups (p-value= 0.04)
Child dietary diversity and food insecurity as a potential correlate of child anthropometric indicators in the context of the urban food system in the cases of north-central Ethiopia <sup>26</sup>	512 mothers with children aged 6 - 59 months in north-central Ethiopia	<ul style="list-style-type: none"> <li>- Design: Cross-sectional</li> <li>- Exposure: Diversity of food with 7 food groups diversity of food</li> <li>- Outcome: Stunting</li> <li>- Analysis: Generalized Linear Model (GLM)</li> </ul>	<ul style="list-style-type: none"> <li>- The diversity of children's diets was measured using the food groups recommended by IYCF, the Child Dietary Diversity Score (CDDS) consumed in the previous 24 hours and categorized as inadequate (0-3 food groups), moderate (4-5 food groups), good (6-7 food group).</li> <li>- Body length in children &lt;24 months (not yet able to stand / &lt;85 cm) was measured using a long wooden sliding board with the help of 2 measuring devices. Children <math>\geq 24</math> months were shaved using a height measuring board with the Frankfurt position and precision of 0.1 cm.</li> </ul>	Z-score height-for-age has a negative relationship with children's dietary diversity ( $\beta = -0.36$ )
Dietary diversity and nutritional status among children in rural Burkina Faso <sup>27</sup>	251 children aged 6 - 59 months in rural Burkina Faso	<ul style="list-style-type: none"> <li>- Design: Cross-sectional</li> <li>- Exposure: Diversity of food with 11 food groups consumed over the last 7 days</li> <li>- Outcome: Stunting</li> <li>- Analysis: Bivariate linear regression or logistic regression</li> </ul>	<ul style="list-style-type: none"> <li>- Dietary diversity was measured using 11 food groups consumed during the last 7 days.</li> <li>- Body length was measured lying on the back for children aged &lt;24 months and standing for children <math>\geq 24</math> months (Shirboard, weight and measure, Olney, MD, USA)</li> </ul>	<ul style="list-style-type: none"> <li>- Children with high dietary diversity have high HAZ</li> <li>- High food diversity can increase the z-score from 0.14 SD to 0.25 SD (p-value= 0.009) per increase in food diversity</li> <li>- Children with high food religiousness could reduce the risk of stunting 0.82 times (P = 0.07) per increase in food diversity.</li> </ul>
Household dietary diversity and child stunting in East Java, Indonesia <sup>19</sup>	768 households with children under 5 years in 8 urban and rural areas, both coastal and mountainous, in East Java, Indonesia	<ul style="list-style-type: none"> <li>- Design: Cross-sectional</li> <li>- Exposure: Dietary diversity with 12 food groups called the Household Food Diversity Score (HDDS)</li> <li>- Outcome: Stunting</li> </ul>	<ul style="list-style-type: none"> <li>- Dietary diversity was measured by the 12 food groups consumed in the previous 24 hours. The child's height is measured using a Vktech Stature Meter or microtoise with a precision of 0.1 cm.</li> </ul>	High dietary diversity was associated with a lower likelihood of stunting (p-value= 0.03)

Copyright ©2023 Faculty of Public Health Universitas Airlangga  
Open access under a CC BY – SA license | Jointly Published by IAGIKMI & Universitas Airlangga

How to cite: Al Uluf, U., Sinatrya, A. K., & Nadhroh, S. R. Literature Review: The Relationship between Dietary Diversity with Stunting In Underfive Children: Tinjauan Literatur: Hubungan antara Keragaman Pangan dengan Stunting pada Balita. Amerta Nutrition, 7(1), 147–153.

Article title	Sample	Method	Assessment method	Results
Feeding practices and growth among young children during two seasons in rural Ethiopia <sup>28</sup>	320 children aged 6-12 months in the harvest season and 312 children aged 6-12 months in the pre-harvest season in a rural area of southwest Ethiopia	<ul style="list-style-type: none"> <li>- Analysis: Logistic regression</li> <li>- Design: Cross-sectional</li> <li>- Exposure: Food diversity across the seven food groups was measured in the previous 24 hours</li> <li>- Outcome: Stunting</li> <li>- Analysis: Linear regression and logistic regression</li> </ul>	<ul style="list-style-type: none"> <li>- Food diversity is obtained by summarizing the food groups consumed in the previous 24 hours by children with 7 food groups.</li> <li>The child's body length was measured using a board in a recumbent position and recorded with a precision of 0.1 cm (SECA 210, Hamburg, Germany)</li> </ul>	Dietary diversity had a positive relationship with the long-for-age (LAZ) z score in the harvest season (p-value= 0.03)
Dietary diversity, parenting and stunting in toddlers aged 24-59 months <sup>29</sup>	100 children aged 24-59 months in Bayat District, Klaten Regency	<ul style="list-style-type: none"> <li>- Design: Cross-sectional</li> <li>- Exposure: Variety of foods with 9 food groups</li> <li>- Outcome: Stunting</li> <li>- Analysis: Chi-square and logistic regression</li> </ul>	<ul style="list-style-type: none"> <li>- Dietary diversity was measured using the previous 24-hour withdrawal method, and the data obtained were included in the IDDS (Individual Dietary Diversity Score) questionnaire, which consisted of 9 food groups. These results were categorized into foods that do not vary when the score is 0-5 and safe when the score is more than five food groups.</li> <li>- Height was measured using a microtoise with a precision of 0.1 cm.</li> </ul>	There was an association between dietary diversity and stunting (p-value= 0.029, OR = 3.213)
Food Availability and Diversity and Economic Level as Predictors of Toddler Nutritional Status <sup>20</sup>	115 families with toddlers in 57 districts in Malang	<ul style="list-style-type: none"> <li>- Design: Cross-sectional</li> <li>- Exposure: A variety of foods with 12 food groups</li> <li>- Outcome: Stunting</li> <li>- Analysis: Pearson/Spearmann and Fisher exact test</li> </ul>	<ul style="list-style-type: none"> <li>- Dietary diversity was measured using the 24-hour withdrawal method. The results were collected through the HDDS questionnaire and categorized into HDDS scores &lt;9 food groups and HDDS scores ≥9 food groups.</li> <li>- Height was measured using the German SECA microtoise brand model 206, and body length was measured with the German SECA model 2010 longboard</li> </ul>	There was no relationship between TB/U z-score and HDDS
Associations of childhood, maternal and household dietary patterns with childhood stunting in Ethiopia: Proposing an alternative and plausible dietary analysis method to dietary diversity scores <sup>30</sup>	3,788 mothers with children under 5 years of age in Ethiopia	<ul style="list-style-type: none"> <li>- Design: Cross-sectional</li> <li>- Exposure: A variety of foods with 12 food groups</li> <li>- Outcome: Stunting</li> <li>- Analysis: Chi-square and Poisson regression model</li> </ul>	<ul style="list-style-type: none"> <li>- Dietary diversity was measured using 12 food groups taken using the 24-hour withdrawal method and categorized into HDD ≤5 and &gt; 5 HDD scores.</li> <li>- The height or length of the child was measured using a wooden plank with a precision of 0.1 cm recommended by UNICEF. Children aged ≥24 months were measured in a standing position, and children aged &lt;24 were measured in a supine</li> </ul>	HDD scores had no relation with stunting (p-value= 0.624)

Copyright ©2023 Faculty of Public Health Universitas Airlangga  
Open access under a CC BY – SA license | Jointly Published by IAGIKMI & Universitas Airlangga

How to cite: Al Uluf, U., Sinatrya, A. K., & Nadiyah, S. R. Literature Review: The Relationship between Dietary Diversity with Stunting In Underfive Children: Tinjauan Literatur: Hubungan antara Keragaman Pangan dengan Stunting pada Balita. Amerta Nutrition, 7(1), 147-153.



Article title	Sample	Method	Assessment method	Results
Socioeconomic Status and Food Diversity in Stunting and Non-Stunting Toddlers Aged 24-59 Months in the Work Area of the Wilangan Health Center, Nganjuk Regency <sup>31</sup>	28 toddlers aged 24-59 months in each case group and control group (N = 56) in Nganjuk District	- Design: Case-control - Exposure: Food Diversity (IDDS) - Outcome: Stunting - Analysis: Chi-square	Dietary diversity was measured using the Individual Dietary Diversity Score (IDDS) form in which the data was obtained from 3x24 hour withdrawals (2 weekdays and 1 weekend) and was categorized into various if the IDDS score $\geq 4$ and did not vary when the IDDS score $< 4$ .	There was no association between dietary diversity and the incidence of stunting (p-value= 1,000), and it was not a risk factor for stunting under five (OR = 1,000)
Association between household dietary diversity and nutritional status of children (6–36 months) in Wenchi Municipality, Brong Ahafo Region, Ghana <sup>32</sup>	590 mothers of children aged 6-36 months in Wenchi Town, Brong Ahafo Region, Ghana	- Design: Cross-sectional - Exposure: A variety of foods with 12 food groups - Outcome: Stunting - Analysis: Chi-square and logistic regression	- Dietary diversity was measured using the 24-hour withdrawal method. The results were calculated with the Household Dietary Diversity Score (HDDS) and were categorized as low if the HDD score was 1-5 food groups and high if the HDD score was 6-12 food groups. - The child's height/length was measured lying on the back for children less than 24 months using an infantometer and measured standing for children aged 24 months and over with a precision of 0.1 cm in both measurements	There was no relationship between HDD and stunting categories (p-value= 0.409)

The advantage of this literature review was that it could formulate more deeply regarding the relationship between diversity and the incidence of stunting, which has been carried out in several studies both in Indonesia and outside Indonesia. The weakness of this literature review was that the studies studied came from various countries, so they have different characteristics, such as the type of food that was the basis for assessing food diversity. Then the research literature used two different approaches, the cross-sectional and case-control approaches to produce a further research analysis of each approach method. This research was limited to food diversity and did not explicitly discuss the frequency and quantity of the type of food consumed.

#### CONCLUSIONS

Based on the several studies described, out of 10 articles, six proved the relationship between dietary diversity and stunting, so this literature review study concludes that dietary diversity was related to stunting. The results of this study can be used as a reference for efforts to prevent and reduce stunting in toddlers by optimizing the availability of food diversity, especially in stunting locus areas.

#### ACKNOWLEDGEMENTS

Thanks to Dr. Siti Rahayu Nadhiroh, SKM., M.Kes as the supervisor of this writing and all Lecturers of the Public Health Masters Program specializing in Public Health Nutrition, Faculty of Public Health, Airlangga University.

#### Conflict of Interest and Funding Disclosure

All authors have no conflict of interest in this article. Article were independently funded by the author.

#### REFERENCES

1. UNICEF. Improving Child Nutrition. [Internet]. United Nations Children's Fund (UNICEF). 2013. Available from: [https://data.unicef.org/resources/improving-child-nutrition-the-achievable-imperative-for-global-progress/nutritionreport\\_april2013\\_final\\_29/](https://data.unicef.org/resources/improving-child-nutrition-the-achievable-imperative-for-global-progress/nutritionreport_april2013_final_29/)
2. UNICEF, WHO, World Bank Group. UNICEF/WHO/World Bank Group Joint Child Malnutrition Estimates. In 2018. Available from: <https://data.unicef.org/wp-content/uploads/2018/05/JME-2018-brochure-web.pdf>
3. The Ministry of Health of Indonesia. Buletin Stunting [Internet]. Kemenkes RI. 2018. Available from: <https://pusdatin.kemkes.go.id/folder/view/01/structure-publikasi-pusdatin-buletin.html>
4. The Ministry of Health of Indonesia. Buku saku pemantauan status gizi. Buku saku pemantauan status gizi tahun 2017 [Internet]. 2018;7–11. Available from: [https://kesmas.kemkes.go.id/assets/uploads/content/others/Buku-Saku-Nasional-PSG-2017\\_975.pdf](https://kesmas.kemkes.go.id/assets/uploads/content/others/Buku-Saku-Nasional-PSG-2017_975.pdf)
5. Trihono D. Pendek (Stunting) di Indonesia dan Solusinya [Internet]. Sudomo M, editor. Lembaga Penerbit Balitbangkes; 2015. 182 p. Available from: <http://repository.litbang.kemkes.go.id/3512/>
6. The Ministry of Health of Indonesia. Ini Penyebab Stunting pada Anak. Indonesia: Biro Komunikasi dan Pelayanan Masyarakat, Kementerian Kesehatan RI; 2018.
7. Wantina M, Rahayu LS, Yuliana I. Keragaman Konsumsi Pangan sebagai Faktor Risiko Stunting pada Balita Usia 6-24 Bulan. *J UHAMKA*. 2(2), 89–96 (2017).
8. Noor Prastia T, Listyandini R. Keragaman Pangan Berhubungan dengan Stunting pada Anak Usia 6-24 Bulan. *Hearty*. 8(1), 33–41 (2020).
9. Tim Nasional Percepatan Penanggulangan Kemiskinan. 100 Kabupaten/Kota Prioritas untuk Intervensi Anak Kerdil (Stunting) [Internet]. Jakarta; 2017. 42 p. Available from: <http://www.tnp2k.go.id/images/uploads/downloads/Buku-Ringkasan-Stunting.pdf>
10. Mahmudiono T, Sumarmi S, Rosenkranz RR. Household Dietary Diversity and Child Stunting in East Java, Indonesia. *Asia Pac J Clin Nutr*. 26(2), 317–25 (2017).
11. The Ministry of Health of Indonesia. Menteri Kesehatan Republik Indonesia. Kemenkes [Internet]. 2014 p. 1–96. Available from: [http://hukor.kemkes.go.id/uploads/produk\\_hukum/PMK\\_No\\_41\\_ttg\\_Pedoman\\_Gizi\\_Seimbang.pdf](http://hukor.kemkes.go.id/uploads/produk_hukum/PMK_No_41_ttg_Pedoman_Gizi_Seimbang.pdf)
12. Nai HME, Remyoet BS. Poor Dietary Diversity is Associated with Stunting among Children 6–23 Months in Area of Mergangsan Public Health Center, Yogyakarta. *J Nutr Sci Vitaminol (Tokyo)*. 66, S398–405 (2020).
13. Arulmohi M, Vinayagamoorthy V, R. DA. Physical Violence Against Doctors: A Content Analysis from Online Indian Newspapers. *Indian J Community Med*. 42(1), 147–50 (2017).
14. Hein AK, Hong SA, Puckpinyo A, Tejativaddhana P. Dietary Diversity, Social Support and Stunting among Children Aged 6–59 Months in an Internally Displaced Persons Camp in Kayin State, Myanmar. *Clin Nutr Res*. 8(4), 307 (2019).
15. Modjadji P, Molokwane D, Ukegbu PO. Dietary Diversity and Nutritional Status of Preschool Children in North West Province, South Africa: A Cross Sectional Study. *Children*. 7(10), 174 (2020).
16. Khamis AG, Mwanri AW, Ntwenya JE, Kreppel K. The Influence of Dietary Diversity on the Nutritional Status of Children between 6 And 23 Months of Age in Tanzania. *BMC Pediatr*. 19(1), 1–9 (2019).
17. Mank I, Vandormael A, Traoré I, Ouédraogo WA, Sauerborn R, Danquah I. Dietary Habits Associated with Growth Development of Children Aged < 5 Years in the Nouna Health and Demographic Surveillance System, Burkina Faso. *Nutr J*. 19(1), 1–14 (2020).

18. Busert LK, Neuman M, Rehfuess EA, Dulal S, Harthan J, Chaube SS, et al. Dietary Diversity is Positively Associated with Deviation from Expected Height in Rural Nepal. *J Nutr.* **146**(7), 1387–93 (2016).
19. Astuti DK, Sumarmi S. Keragaman Konsumsi Pangan pada Balita Stunting di Wilayah Pedesaan dan Perkotaan Kabupaten Probolinggo. *Media Gizi Indones.* **15**(1), 14–21 (2020).
20. Wirawan NN, Rahmawati W. Ketersediaan dan Keragaman Pangan serta Tingkat Ekonomi sebagai Prediktor Status Gizi Balita. *Indones J Hum Nutr.* **3**(1), 80–90 (2016).
21. Chandrasekhar S, Aguayo VM, Krishna V, Nair R. Household Food Insecurity and Children's Dietary Diversity and Nutrition in India. Evidence from the Comprehensive Nutrition Survey in Maharashtra. *Matern Child Nutr.* **1**, 1–8 (2017). Available from: <https://onlinelibrary.wiley.com/doi/epdf/10.1111/mcn.12447>
22. Purwoko SA, Triana NY, Cahyaningrum ED. Gambaran Keragaman Pangan, Pola Asuh Makan dan Hygiene Pada Balita Stunting. *J Kesehat.* **14**, 94–109 (2020).
23. Perkins JM, Jayatissa R, Subramanian S V. Dietary Diversity and Anthropometric Status and Failure among Infants and Young Children in Sri Lanka. *Nutrition.* **55–56**:76–83 (2018). Available from: <https://doi.org/10.1016/j.nut.2018.03.049>
24. Ersa DM, Hasnita E, Nurdin. Meta-Analisis Determinan Stunting pada Anak Usia di Bawah 5 Tahun di Asia. *J Hum Care.* **5**(4), 993–9 (2020).
25. Hasan M, Islam MM, Mubarak E, Haque MA, Choudhury N, Ahmed T. Mother's Dietary Diversity and Association with Stunting among Children <2 Years Old in a Low Socio-Economic Environment: A Case–Control Study in an Urban Care Setting in Dhaka, Bangladesh. *Matern Child Nutr.* **15**(2), 1–8 (2019).
26. Dinku AM, Mekonnen TC, Adilu GS. Child Dietary Diversity and Food (in)Security as a Potential Correlate of Child Anthropometric Indices in The Context of Urban Food System in the Cases of North-Central Ethiopia. *J Heal Popul Nutr.* **39**(1), 1–11 (2020).
27. Sié A, Tapsoba C, Dah C, Ouermi L, Zabre P, Bärnighausen T, et al. Dietary Diversity and Nutritional Status among Children in Rural Burkina Faso. *Int Health.* **10**(3), 157–62 (2018).
28. Wondafrash M, Huybregts L, Lachat C, Bouckaert KP, Kolsteren P. Feeding Practices and Growth among Young Children during Two Seasons in Rural Ethiopia. *BMC Nutr.* **3**(1), 1–10 (2017).
29. Widyaningsih NN, Kusnandar, Sappa A. Keragaman Pangan , Pola Asuh Makan dan Kejadian Stunting pada Balita Usia 24-59 Bulan. *J Gizi Indones.* **7** (1) (2018).
30. Melaku YA, Gill TK, Taylor AW, Adams R, Shi Z, Worku A. Associations of Childhood, Maternal and Household Dietary Patterns with Childhood Stunting in Ethiopia: Proposing an Alternative and Plausible Dietary Analysis Method to Dietary Diversity Scores. *Nutr J.* **17**(1), 1–15 (2018).
31. Nurmayasanti A, Mahmudiono T. Status Sosial Ekonomi dan Keragaman Pangan Pada Balita Stunting dan Non-Stunting Usia 24-59 Bulan di Wilayah Kerja Puskesmas Wilangan Kabupaten Nganjuk. *Amerta Nutr.* **3**(2), 114–21 (2019).
32. Wemakor A, Laari J. Association between Household Dietary Diversity and Nutritional Status of Children (6–36 Months) in Wenchi Municipality, Brong Ahafo Region, Ghana. *Nutrire.* 2018;43(1):1–10.