



Source details

Journal of Pakistan Association of Dermatologists

CiteScore 2021

0.2



Scopus coverage years: from 1999 to 2022

Publisher: Pakistan Association of Dermatologists

SJR 2021

0.127



ISSN: 1560-9014

Subject area: Medicine: Dermatology

Source type: Journal

SNIP 2021

0.127



[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology ✕

CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021 ▾

$$0.2 = \frac{90 \text{ Citations 2018 - 2021}}{414 \text{ Documents 2018 - 2021}}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022 ⓘ

$$0.3 = \frac{112 \text{ Citations to date}}{448 \text{ Documents to date}}$$

Last updated on 05 January, 2023 • Updated monthly

CiteScore rank 2021 ⓘ

| Category | Rank | Percentile |
|-------------|----------|------------|
| Medicine | | |
| Dermatology | #107/126 | 15th |

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)

About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

ELSEVIER

[Terms and conditions](#) ↗ [Privacy policy](#) ↗

Copyright © Elsevier B.V. ↗ . All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗.






Journal of Pakistan Association of Dermatologists

COUNTRY

Pakistan



SUBJECT AREA AND CATEGORY

Medicine
Dermatology

PUBLISHER

Pakistan Association of Dermatologists

H-INDEX

14

PUBLICATION TYPE

Journals

ISSN

15609014

COVERAGE


1999-2021

INFORMATION

[Homepage](#)
[How to publish in this journal](#)
zahidaraffad@yahoo.com

SCOPE

Information not localized

 Join the conversation about this journal










FIND SIMILAR JOURNALS ⓘ

options ⋮

1 2 3 4 5

< >



Journal of Pakistan Association of...

Q4 Dermatology

SJR 2021 0.13

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

`https://www.scimagojr.com`

SCImago Graphica



Explore, visually communicate and make sense of data with our [new data visualization tool](#).

**Journal of
Pakistan Association of Dermatologists**

JPAD
www.jpada.com.pk

SPECIAL FOCUS;

Tinea corporis caused by *microsporum canis* in children with HIV



A Publication of

PAKISTAN ASSOCIATION OF DERMATOLOGISTS

Vol. 32 No. 4 (2022): October - December



Published: 2022-12-08

Editorial

The importance of dermatological lesions in the agile diagnosis of Monkeypox

Gonzalo Dominguez Alvarado, Angie Marcela Arboleda Roca , Aymer Daniel Bolívar Cabrera , Gabrielle Yolmari Curicó Martínez , Laura Camila Millán Guerrero

662-664



PDF

Original Articles

The Spectrum of Dermatoses among Prisoners; A Retrospective Analysis

Hira Tariq

665-668



PDF

Basil Leaf Extract and Eugenol against Isolates of Candida sp. Causing Oral Candidiasis in HIV/AIDS

Dwi Murtiastutik, Emma Hidayati Sasmito, Diah Mira Indramaya, Sawitri Sawitri, Rahmadewi Rahmadewi, Afif Nurul Hidayati, Pepy Dwi Endraswari, Sudjarwo Sudjarwo, Budi Utomo

669-675

 PDF

Can Ki-67 immunohistochemistry marker differentiate mycosis fungoides from cutaneous lichen planus?

Mazaher Ramezani, Kambiz Kamyab Hesari, Negar Ramezanizoorab, Masoud Sadeghi, Mansour Rezaei, Sedigheh Khazaei

676-682

 PDF

Correlation between quality of life and clinical severity of melasma in Pakistani women

Maryam Qayyum, Saadiya Siddiqui, Mohsina, Mahmoona Ilyas, Atif Shahzad, Nadia Ali Zafar

Saadiya Siddiqui, Maryam Qayyum

683-689

 PDF

Association of Pruritus Visual Analogue Scale and Risk Factors in Adolescence Pediculosis Capitis in Two Public Boarding Schools, West Java

Arlene Rainamira, Firman Parrol, Yari Castiliani, Githa Rahmayunita, Mochamad Helmi Aziz, Kusmarinah Bramono, Sandra Widaty

690-695

 PDF

Scalp defects and their reconstruction in patients at Allama Iqbal Memorial Teaching Hospital, Sialkot

Sarfaraz Ahmad, Raffad, Zahida Rani, Atika Malik

696-700

 PDF

Novel trends in cutaneous manifestations of geriatric dermatoses in a tertiary care hospital, South India

Latest trends in geriatric dermatosis

Hasini budeda, Indira B, Darsan S, V. V. V. Satyanarayana, Suruthi Purushothaman

701-706

 PDF

Effect of Lactobacillus plantarum IS-10506 supplementation with oral metronidazole for the treatment of bacterial vaginosis : a randomized placebo-controlled clinical trial

Afif Nurul Hidayati, Ridha Ramadina Widiatma, Dwi Murtiastutik, Ingrid Suryanti Surono, Cita Rosita Sigit Prakoeswa, Lunardi Bintanjoyo

707-714

 PDF

The Platelet Rich Plasma (PRP) Preparation and Standardization of time interval for scalp hair loss

Munir Alam

715-718

 PDF

Review Articles

Zinc as an essential element for normal immune reactions and as a therapeutic agent for autoimmune diseases

Inas K. Sharquie, Khalifa E. Sharquie , Wasan W. Al-bassam

719-725

 PDF

Review of treatment modalities in the management of Genital Warts

Review of treatment modalities in the management of Genital Warts

Jai Singh Solanki, Swagat Waghmare, Manjyot Gautam, Praneet P. Awake

726-733

 PDF

Case Reports

Kaposi's sarcoma in a male with human immunodeficiency virus and condyloma acuminata: A case report

Minna Hasniah, Budi Eko Prasetyorini, Endra Yustin Ellistasari

734-740

 PDF

Tinea Corporis Caused by Microsporum Canis in Children with HIV

Sri Mariyani

741-744

 PDF

Recurrent herpes simplex in elderly with vitiligo vulgaris: A dilemmatic case

Novita Suprpto, dr. Lili Legiawati, dr. Rinadewi Astriningrum, dr. Shannaz Nadia Yusharyahya

745-747



[Make a Submission](#)



Recognized by HEC



Indexed

Information

[For Readers](#)

[For Authors](#)

[For Librarians](#)

[Open Journal Systems](#)

JPAD Team

Editorial Board

Pakistan

Ahmed Gardezi
Amer Ejaz
Arfan ul Bari
Asher Ahmad
Mashhood
Azim Jahangir Khan
Ghulam Mujtaba
Ijaz Ahmad
Jamil Ahmad Shaheen
Naseema Kapadia
Mansoor Dilnawaz
Muhammad Khalid
Muhammad Shahid
Nadia Ali Azfar
Naeem Raza

Nasser Rashid Dar
Rehanuddin
Safoora Aamir
Shahbaz Aman
Shahbaz Janjua
Shahid Jawaid Akhtar
Syed Dilawar Abbas
Rizvi
Syed Shamsuddin
Tahir Jamil Ahmad
Talat Akbar
Tariq Rashid
Yasmeena Khan
Zafar Iqbal Sheikh

Overseas

Aisha Sethi (USA)
Ayyaz Shah (USA)
Ian McColl (Australia)
Imtiaz Ahmad (UK)
Jyakar Thomas (India)
Khalid Al Aboud (KSA)
MR Namazi (Iran)

Peer Review Board

Prof. Shahbaz Aman
Dr. Atif Shahzad
Dr. Syed Saelah Batool

Prof. Muhammad Nadeem
Prof. Lamees Mahmood Malik
Prof. Mahmood Noor

Prof. Aneela Asghar Butt
Dr. Irfan ul Bari

Advisory Board

Prof. Tariq Rashid
Prof. Khawar Khurshid

Prof. Tahir Jamil Ahmad
Prof. Tariq Zaman

Prof. Azam Bukhari
Dr. Yasmeena Khan

[Make a Submission](#)



Recognized by HEC



Indexed

Information

[For Readers](#)

[For Authors](#)

[For Librarians](#)

[Open Journal Systems](#)

Platform &
workflow by
OJS / PKP

Effect of *Lactobacillus plantarum* IS-10506 supplementation with oral metronidazole for the treatment of bacterial vaginosis: A randomized placebo-controlled clinical trial

AN Hidayati^{*,†}, RR Widiatma^{*}, D Murtiastutik^{*}, IS Surono[§], CRS Prakoeswa^{*}, L Bintanjoyo^{*}

^{*} Department of Dermatology and Venereology, Faculty of Medicine, Universitas Airlangga, Dr. Soetomo General Academic Hospital, Surabaya, Indonesia.

[†] Universitas Airlangga Teaching Hospital, Surabaya, Indonesia.

[§] Department of Food Technology, Faculty of Engineering, Bina Nusantara University, Jakarta 11480, Indonesia.

Abstract

Background Replacement of *Lactobacillus* spp. by anaerobic and facultative bacteria is central in pathogenesis of bacterial vaginosis (BV). Oral metronidazole is currently recommended treatment. However, cure rate is variable. Probiotic supplementation was explored as alternative therapy. This study aimed to find out the effect of oral microencapsulated *Lactobacillus plantarum* IS-10506 supplementation on oral metronidazole 500 mg twice daily for the treatment of BV.

Methods Twenty nine females with BV were treated with oral metronidazole and randomized into probiotic (Pro) (n=14) and placebo (Pla) (n=15) groups. Pro group received microencapsulated *L. plantarum* IS-10506 supplementation at 0.9×10^9 CFU, twice daily. Pla group received placebo twice daily. Cure is defined from Amsel criteria and Nugent score 0-3. Metronidazole was stopped if subjects were cured. Probiotic and placebo were continued for 4 weeks. Cure rate and mean Nugent score were assessed at baseline, end of week 1, 2 and 4.

Results Mean Nugent score for Pro against Pla group were 8.07 vs. 8.07, 5.36 vs. 6.20, 4.07 vs. 4.93, and 3.57 vs. 4.33 respectively at baseline, end of week 1, 2 and 4. Cure rate for Pro against Pla group were 28.6% vs 20.0%, 50.0% vs 33.3%, 64.3% vs 40.0% respectively at end of week 1, 2 and 4. Mean Nugent score significantly decreased in both groups at end of week 1, 2 and 4 ($p \leq 0.05$). Although not statistically significant, mean Nugent score was lower and cure rate was higher in Pro than Pla group at end of week 1, 2 and 4 ($p > 0.05$). No adverse effect was recorded.

Conclusion Oral probiotic *L. plantarum* IS-10506 supplementation in addition to oral metronidazole was safe and potentially better than oral metronidazole alone for the treatment of BV. Studies with longer probiotic supplementation and more subjects may be required to demonstrate significant effect of this combination treatment on BV.

Key words

Bacterial vaginosis, probiotic, *Lactobacillus plantarum*, Nugent score, human health.

Introduction

Bacterial vaginosis (BV) is a vaginal disorder presenting with abnormal malodorous vaginal discharge associated with increased vaginal pH.^{1,2} BV most often affects females of reproductive age with prevalence ranging from

11.1-60.8% worldwide.¹ Although some patients may be asymptomatic, symptoms of BV may cause fear and embarrassment which lower patients' quality of life. BV may also lead to other serious complications including increased risk for acquisition of sexually transmitted infections such as *Neisseria gonorrhoea*,

Chlamydia and human immunodeficiency virus (HIV), also prematurity and low birth weight in infants born to pregnant women suffering from BV.^{2,3}

Douching of vagina, other vaginal infections, smoking, promiscuity and use of intrauterine devices were among the known risk factors for BV. However, the exact aetiology of BV is still uncertain. Dysbiosis of vaginal micro biome is thought to be central in the pathogenesis of BV. It is characterized by replacement of *Lactobacillus* spp. by anaerobic and facultative bacteria such as *Gardnerella vaginalis*, *Mobiluncus* spp., *Mycoplasma hominis*, *Bacteroides* spp., *Prevotella* spp. and *Atopobium vaginae*.^{1,2} The earliest event in the pathogenesis of BV is adhesion of *G. vaginalis* to the host cell. Subsequently, *G. vaginalis* produces sialidase and prolidase which attack vaginal mucosa resulting in increased release of proinflammatory cytokines and vaginolysin which initiates cellular death mechanisms. It also forms biofilms which act as defence against hydrogen peroxide and lactic acid produced by *Lactobacillus*, and act as environment for growth of other pathogens and exchange of genes encoding antimicrobial resistance. Other bacteria within the biofilms also act synergistically with *G. vaginalis*. *Prevotella* spp. producing fibrinolysin and collagenase to aid in the adhesion to vaginal mucosa and counter the mucosal defence system. Together with *Peptostreptococcus anaerobius*, *Prevotella* spp. also provides amino acids and nutrition for other

pathogens.^{3,4} Overgrowth of pathogens reduces nutritional support for lactobacilli. The enzymes released by these pathogens also impair vaginal barrier. The end results are reduced *Lactobacillus* spp., depleted lactic acid and hydrogen peroxide, increased vaginal pH and persistent infection.^{2,3}

Centre for Disease Control and Prevention currently recommends oral metronidazole 500mg twice a day for one week, intravaginal metronidazole gel once to twice a day for five days or intravaginal clindamycin cream once at night for one week for treatment of BV in non-pregnant females.³ However, these treatments showed variable efficacy with cure rate of 58-92%.⁵ Anukam *et al.* even reported a cure rate of 40% among BV patients treated with oral metronidazole alone.⁶ More than 50% of BV patients also experienced recurrence and reinfection within 6-12 months after treatment, probably due to antibiotic resistance. Oral antibiotics may also negatively influence normal gut microbiome, while intravaginal antibiotics may predispose patients to vulvovaginal candidiasis.^{5,7} Therefore, alternative therapies to restore normal predominantly-*Lactobacillus* spp. vaginal micro biome were being explored, including probiotics.³

Probiotics are oral or topical supplements containing live bacteria and/or yeasts which are beneficial for health.³ Probiotics containing *Lactobacillus* spp. have been studied for treatment of vaginal dysbiosis since they are the predominant commensal bacteria of vagina. They act to prevent pathogen adhesion by production of anti-biofilm biosurfactant and competition for binding site, and produce hydrogen peroxide against BV pathogens.³ Anukam *et al.* reported that probiotics containing *L. rhamnosus* GR-1 and *L. reuteri* RC-14 together with oral metronidazole increased cure rate of BV by 48% compared to

Address for correspondence

Dr. Afif Nurul Hidayati, SpKK(K), FINS DV, FAADV
Department of Dermatology and Venereology,
Faculty of Medicine, Universitas Airlangga,
Dr. Soetomo General Academic Hospital,
Jalan Mayjen Prof. Dr. Moestopo No. 6-8 Surabaya
60285, East Java, Indonesia.
Universitas Airlangga Teaching Hospital, Surabaya,
East Java, Indonesia.
Email: afif_nurulhidayati@fk.unair.ac.id

oral metronidazole alone.⁶ Cianci *et al.* reported that probiotic containing *L. plantarum*, which is another member of commensal vaginal bacteria, reduced risk of recurrent vaginal infections, although it wasn't statistically significant.⁸ Lin *et al.* also reported that probiotics containing *L. rhamnosus* and *L. plantarum* significantly decreased malodorous discharge, pruritus and Nugent scores in 19 BV patients.⁹

Lactobacillus plantarum IS-10506 is an indigenous Indonesian probiotic isolated from dadih, fermented buffalo milk.¹⁰⁻¹² As this species has interacted with other pathogenic and contaminant microbes in Indonesia, *L. plantarum* IS-10506 are expected to be beneficial and appropriate for use as probiotic in local settings.¹³⁻¹⁶ Microencapsulated formulation also enhanced the survival of *L. plantarum* during storage and transit in the upper gastrointestinal tract.¹⁴ This study aimed to evaluate the effect of probiotic containing microencapsulated *L. plantarum* IS-10506 as supplementation on oral metronidazole in the treatment of BV in Indonesian female patients.

Methods

Ethics Study protocol was approved by Ethics Committee of Dr. Soetomo General Academic Hospital, Surabaya (No. 1626/KEPK/XI/2019).

Study design and setting This was a randomized, double-blind, placebo-controlled trial to evaluate the effect of oral probiotic microencapsulated *L. plantarum* IS-10506 supplementation on oral metronidazole for BV treatment at the Dermatology and Venereology outpatient clinic of a Dr. Soetomo General Academic Hospital, Surabaya, Indonesia from November 2019 to January 2020.

Participants Eligible participants were females, aged 18-55 years, diagnosed with BV with apparently good health and willingness to

participate in the study and sign informed consent. Diagnosis of BV was established according to Amsel criteria and Nugent scoring system. Amsel criteria requires 3 out of 4 findings to diagnose BV, namely thin, whitish-grey, homogeneous vaginal discharge, fishy odor upon addition of 10% KOH solution to the discharge (Whiff test), vaginal pH>4.5, and presence of >20% clue cells in wet mount microscopy with NaCl 0.9% (100x magnification).^{2,3} Nugent scoring system evaluated three types of bacteria in Gram-stained vaginal discharge (1000x magnification, oil-immersion field): large Gram-positive rods or *Lactobacillus* morphotypes (score 0-4), small Gram-variable rods or *Gardnerella vaginalis* morphotypes (score 0-4), and curved Gram-variable rods or *Mobiluncus spp.* morphotypes (score 0-2). Total Nugent score was graded as: normal microflora (0-3), indeterminate microflora (4-6) and bacterial vaginosis (7-10).¹⁷ The exclusion criteria were pregnancy, history of antibiotics and/ or probiotics intake within previous 4 weeks and history of allergies to antibiotics and/or probiotics used in the study. The cure rate for subjects with BV treated with metronidazole alone was assumed to be 40% according to Anukam *et al.* and the expected difference in cure rates was 50%.⁶ Using 95% level of confidence and 80% power, the calculated sample size was 26.

Interventions and randomizations Participants were recruited by consecutive sampling and were randomly assigned into one of the two treatment arms, probiotic (Pro) and placebo (Pla) groups. Both groups were treated with oral metronidazole 500mg twice daily. Pro group was also given oral probiotic, consisting of 950 mg (0.9×10^9 CFU) microencapsulated *L. plantarum* IS-10506, twice daily. Pla group was also given oral placebo, consisting of cellulose (Avicel) and skim milk mixture which has identical color, taste, odor, shape and packaging

to the oral probiotic, twice daily. The probiotic or placebo was given 1 hour after metronidazole intake. The probiotic was packaged by the Pharmacy Department of our institution. Randomizations were carried out by the Pharmacy Department of our institution using a computer-generated scheme and participants were given numbers. Metronidazole and identical-looking probiotic or placebo capsules were supplied in numbered containers. Randomization data were kept in the pharmacy and were not disclosed until the study ended. Both researchers and participants were blinded to the treatment given. Amsel criteria, Nugent score and side effects were assessed at the end of week 1, 2 and 4. Oral metronidazole was stopped if participant was cured of BV, but oral probiotic or oral placebo was continued for 4 weeks. Cure was determined from Amsel criteria (presence of less than 3 out of 4 findings) and Nugent score of 0-3. The primary outcome of this study was the cure rate at the end of week 1,

2 and 4. The secondary outcome of this study was Nugent score and reported side effects at the end of week 1, 2 and 4. The research flow is illustrated in **Figure 1**.

Results

Twenty nine subjects met the inclusion criteria and were randomized into 14 subjects in the Pro group and 15 subjects in the Pla group. Fisher's exact test was used to compare baseline demographic and clinical characteristics between groups showing no statistically significant difference (**Table 1**). All subjects completed the study for 4 weeks. There was significant reduction of mean Nugent score from baseline at the end of week 1, 2 and 4 in both groups ($p \leq 0.05$) (**Table 2**). The mean Nugent score of Pro group was similar to Pla group at baseline, but consistently lower than Pla group at the end of week 1, 2 and 4.




Figure 1 Research flow.

Table 1 Baseline demographic and clinical characteristics.

| Characteristics | Frequency (%) | | p-value |
|----------------------------|------------------|------------------|---------|
| | Pro group (n=14) | Pla group (n=15) | |
| Age (years) | | | |
| 18-25 | 2 (14.2) | 3 (20.0) | 0.785 |
| 26-35 | 4 (28.6) | 6 (40.0) | |
| 36-45 | 4 (28.6) | 2 (13.3) | |
| 46-55 | 4 (28.6) | 4 (26.7) | |
| Education | | | |
| Elementary school | 0 | 2 (13.3) | 0.104 |
| Junior high school | 2 (14.2) | 0 | |
| Senior high school | 6 (42.9) | 10 (66.7) | |
| College | 6 (42.9) | 3 (20.0) | |
| Marital status | | | |
| Married | 12 (85.8) | 14 (93.3) | 0.598 |
| Unmarried | 2 (14.2) | 1 (6.6) | |
| Duration of illness (days) | | | |
| 1-14 | 8 (57.1) | 12 (80.0) | 0.249 |
| 15-30 | 6 (42.9) | 2 (13.3) | |
| > 30 | 0 (0.0) | 1 (6.6) | |
| Episode of BV | | | |
| Primary | 13 (92.9) | 14 (93.3) | 1.000 |
| Recurrent | 1 (7.1) | 1 (6.6) | |
| Risk factors | | | |
| Sexually active | 13 (92.9) | 12 (80.0) | 0.598 |
| Multiple sexual partner | 0 (0.0) | 1 (6.7) | |
| Vaginal douching | 9 (64.3) | 5 (33.3) | 0.143 |
| Contraceptive use* | 7 (50.0) | 4 (26.7) | 0.264 |
| Concomitant STI** | 4 (28.6) | 2 (13.3) | 0.390 |

BV=Bacterial vaginosis, Pla=placebo, Pro=probiotic, STI=sexually transmitted infections, * intrauterine device and hormonal, ** condyloma acuminata.

However, there was no statistically significant difference ($p>0.05$) (**Table 3 and Figure 2**). The cure rate of Pro group was also consistently greater than Pla group at the end of week 1 (28.6% vs. 20.0%), week 2 (50.0% vs. 33.3%) and week 4 (64.3% vs. 40.0%). However, there

was no statistically significant difference ($p>0.05$) (**Table 4**). In both the groups all subjects who were cured of BV at the end of week 1 and 2 showed no recurrence at the end of week 4. No participant reported any side effect throughout the study.

Table 2 Difference of Nugent score in both groups at baseline and on follow ups.

| Difference of Nugent score | Pro group (n=14) | | Pla group (n=15) | |
|----------------------------|------------------|----------|------------------|----------|
| | Mean±SD | p-value* | Mean±SD | p-value* |
| Baseline and end of week 1 | 2.72±2.867 | 0.005 | 1.87±2.475 | 0.011 |
| Baseline and end of week 2 | 4.00±2.680 | 0.003 | 3.13±2.973 | 0.003 |
| Baseline and end of week 4 | 4.50±2.133 | 0.002 | 3.73±2.374 | 0.001 |

Pla=placebo, Pro=probiotic, *Wilcoxon signed rank test.

Table 3 Mean Nugent score in both groups at baseline and on follow ups.

| Nugent score | Mean±SD | | p-value* |
|---------------|------------------|------------------|----------|
| | Pro group (n=14) | Pla group (n=15) | |
| Baseline | 8.07±0.475 | 8.07±0.883 | 0.865 |
| End of week 1 | 5.36±3.003 | 6.20±2.484 | 0.582 |
| End of week 2 | 4.07±2.759 | 4.93±2.658 | 0.407 |
| End of week 4 | 3.57±2.344 | 4.33±1.951 | 0.317 |

Pla=placebo, Pro=probiotic, *Mann-Whitney U test.




Figure 2 Mean Nugent score at baseline, week 1, week 2 and week 4 in both groups

Discussion

In this study, the mean Nugent score decreased significantly at each follow up compared to baseline in both Pro and Pla group. This study also showed increasing cure rate of BV at each follow up in both groups. The cure rate of BV patients treated with antibiotic only in this study (40.0% in Pla group) is similar to report by Anukam *et al.* (40%), but lower than reports by Li *et al.* (66.31%) and Munoz-Barreno *et al.* (74.6%). The cure rate of BV patients treated with antibiotic and probiotic combination in this study (64.3% in Pro group) is lower than reports by Anukam *et al.* (88%), Li *et al.* (78.38%) and Munoz-Barreno *et al.* (74.1%). These differences may be attributed to different route of administration (oral or intravaginal), types of antibiotics (such as metronidazole, tinidazole, clindamycin) and probiotics (such as *L. crispatus*, *L. rhamnosus*, *L. reuteri*, *L. acidophilus*, *L. brevis*, *L. salivarius*, and *L. plantarum*), and also to dose and duration of treatment in these studies.^{6,18,19} The low cure rate may also be caused by persistence of risk factors

for BV among the participants with treatment failure. Formation of biofilm and development of antibiotic resistance by BV pathogens may also contribute to treatment failure in this study.^{4,5}

In this study, Pro group showed lower mean Nugent score (3.57 vs. 4.33) and higher cure rate (64.3% vs. 40%) compared to Pla group at the end of week 1, 2 and 4. However, there were no statistically significant differences. All subjects in this study also reported no side effects. Munoz-Barreno *et al.* also reported no significant difference between pooled clinical cure rate of BV patients treated with antibiotic and probiotic combination or antibiotic alone.¹⁸ However, Anukam *et al.* reported significantly higher cure rate at day 30 in BV patients treated with oral metronidazole supplemented with probiotics containing *L. rhamnosus* GR1 and *L. reuteri* RC-14 compared to oral metronidazole alone.⁶ Li *et al.* also reported significantly higher cure rate in BV patients treated with antibiotics and probiotics compared to antibiotics alone.¹⁹ Wu *et al.* proposed that conflicting results on effects of probiotics in BV may be due to variation in probiotic species and route of administration.⁷ The importance of probiotic species is related to normal vaginal microbiome which varies according to ethnicity. *Lactobacillus crispatus* is reported to be higher in Asian and Caucasian females compared to African females.^{5,7} This species is also reported to confer protection against vaginal dysbiosis and improve stability of vaginal microbiome.²

Table 4 The cure rate of bacterial vaginosis in both groups at baseline and on follow ups.

| Time | Group | Cured* | Not cured | p-value** |
|---------------|-------|-----------|-------------|-----------|
| Baseline | Pro | 0 (0.0%) | 14 (100.0%) | 1.000 |
| | Pla | 0 (0.0%) | 15 (100.0%) | |
| End of week 1 | Pro | 4 (28.6%) | 10 (71.4%) | 0.682 |
| | Pla | 3 (20.0%) | 12 (80.0%) | |
| End of week 2 | Pro | 7 (50.0%) | 7 (50.0%) | 0.462 |
| | Pla | 5 (33.3%) | 10 (66.7%) | |
| End of week 4 | Pro | 9 (64.3%) | 5 (35.7%) | 0.272 |
| | Pla | 6 (40.0%) | 9 (60.0%) | |

Pla=placebo, Pro=probiotic, *Cured if negative Amsel’s criteria and Nugent score of 0-3, **Fisher’s exact test

Despite of 100% short term cure rate, 70-79.5% long term cure rate, and higher remission rate than placebo, studies on *L. crispatus* as probiotics in BV are relatively lacking compared to other species. Unsuitable probiotic species may explain the wide variation of BV cure rates between studies.⁷ Route of probiotic administration may also affect the cure rate of BV. Probiotics containing *L. reuteri* RC-14 and *L. rhamnosus* GR-1 showed better BV cure rate by intravaginal compared to oral route. This may be due to immediate replacement of BV pathogens by the probiotic species.⁷ On the other hand, oral probiotics require migration of lactobacilli through gut, perineum and vulva to vagina.²⁰ However, oral probiotics are developing more rapidly than intravaginal probiotics, because they are not regulated as drug but as food supplements.⁷

This study has some limitations including lack of assessment of biofilm formation and antimicrobial resistance and unavailability of intravaginal formulation of the probiotic used. Another limitation of this study is the short follow up period for subjects who were cured of BV. The follow up period completed at the end of week 4, which ranged from 14 to 21 days for subjects who were cured of BV at the end of week 1 or 2. Although all cured subjects showed no recurrence until the end of week 4, the follow up period is much shorter than average time to BV recurrence. Heczko *et al.* reported average time to BV recurrence of 47.3 days in subjects receiving placebo and 71.4 days in subjects receiving oral supplementation of *L. gasseri* 57C, *L. fermentum* 57A and *L. plantarum* 57B after stopping oral metronidazole.²¹ This short follow up period precludes accurate assessment of recurrence rate in this study.

Conclusion

Oral probiotic microencapsulated *L. plantarum*

IS-10506 supplementation in addition to oral metronidazole were safe and potentially better than oral metronidazole alone for treatment of BV. Studies with more subjects, also longer probiotic supplementation and follow up period may be required to demonstrate significant effect of this combination treatment on BV.

References

1. Manuputty AG, Matodiharjo S. The Profile of Bacterial Vaginosis in Academic Hospital Surabaya: A Retrospective Study. *Berk Ilmu Kesehatan Kulit dan Kelamin-Period Dermatology Venereol.* 2020;**32(2)**:141.
2. Saraf VS, Sheikh SA, Ahmad A, *et al.* Vaginal microbiome: normalcy vs. dysbiosis. *Arch Microbiol.* 2021;**203(7)**:3793–802.
3. Joseph RJ, Ser HL, Kuai YH, *et al.* Finding a balance in the vaginal microbiome: How do we treat and prevent the occurrence of bacterial vaginosis? *Antibiotics.* 2021;**10(6)**:1–39.
4. Hidayati AN, Liuwan CC. Peran Biofilm terhadap Infeksi Saluran Genital yang disebabkan oleh Vaginosis Bakterial (The Role of Biofilm in Genital Tract Infection Caused by Bacterial Vaginosis). *Berk Ilmu Kesehatan Kulit dan Kelamin-Period Dermatology Venereol.* 2019;**31(2)**:150–8.
5. Abou Chacra L, Fenollar F, Diop K. Bacterial Vaginosis: What Do We Currently Know? *Front Cell Infect Microbiol.* 2022;**11**:1–13.
6. Anukam K, Osazuwa E, Ahonkhai I, *et al.* Augmentation of antimicrobial metronidazole therapy of bacterial vaginosis with oral probiotic *Lactobacillus rhamnosus* GR-1 and *Lactobacillus reuteri* RC-14: randomized, double-blind, placebo controlled trial. *Microbes Infect.* 2006;**8(6)**:1450–4.
7. Wu S, Hugerth LW, Schuppe-Koistinen I, *et al.* The right bug in the right place: opportunities for bacterial vaginosis treatment. *Npj Biofilms Microbiomes.* 2022;**8(1)**:34.
8. Cianci A, Cicinelli E, De Leo V, *et al.* Observational prospective study on *Lactobacillus plantarum* P 17630 in the prevention of vaginal infections, during and after systemic antibiotic therapy or in

- women with recurrent vaginal or genitourinary infections. *J Obstet Gynaecol (Lahore)*. 2018;38(5):693–6.
9. Lin TC, Hsu IL, Tsai WH, *et al.* Improvement of bacterial vaginosis by oral lactobacillus supplement: A randomized, double-blinded trial. *Appl Sci*. 2021;11(3):1–21.
 10. Prakoeswa CRS, Bonita L, Karim A, *et al.* Beneficial effect of Lactobacillus plantarum IS-10506 supplementation in adults with atopic dermatitis: a randomized controlled trial. *J Dermatolog Treat*. 2022;33:1491-8.
 11. Prakoeswa CRS, Herwanto N, Prameswari R, *et al.* Lactobacillus plantarum IS-10506 supplementation reduced SCORAD in children with atopic dermatitis. *Benef Microbes*. 2017;8(5):833–40.
 12. Surono IS. Ethnic fermented foods and alcoholic beverages of Indonesia. In: Tamang JP, editor. *Ethnic Fermented Foods and Alcoholic Beverages of Asia*. Delhi: Springer India; 2016. p. 341–82.
 13. Adib A, H. Wahid M, Sudarmono P, *et al.* Lactobacillus plantarum PADA FESES INDIVIDU DEWASA SEHAT YANG MENGONSUMSI Lactobacillus plantarum IS-10506 DARI DADIH. *J Teknol dan Ind Pangan*. 2013;24(2):154–61.
 14. Surono I, Verhoeven J, Verbruggen S, *et al.* Microencapsulation increases survival of the probiotic Lactobacillus plantarum IS-10506, but not Enterococcus faecium IS-27526 in a dynamic, computer-controlled in vitro model of the upper gastrointestinal tract. *J Appl Microbiol*. 2018;124(6):1604–9.
 15. Collado MC, Surono IS, Meriluoto J, *et al.* Potential probiotic characters of Lactobacillus and Enterococcus strains isolated from traditional dadih fermented milk against pathogen intestinal colonization. *J Food Prot*. 2007;70(3):700–5.
 16. Collado MC, Surono I, Meriluoto J, *et al.* Indigenous dadih lactic acid bacteria: Cell-surface properties and interactions with pathogens. *J Food Sci*. 2007;72(3):89–93.
 17. Lim MLR, Lasala L. Effectiveness and safety of metronidazole plus probiotics versus metronidazole alone in the treatment of bacterial vaginosis: randomized controlled trial. *SPMC J Heal Care Serv*. 2015;1(1):31-7.
 18. Muñoz-Barreno A, Cabezas-Mera F, Tejera E, *et al.* Comparative effectiveness of treatments for bacterial vaginosis: A network meta-analysis. *Antibiotics*. 2021;10(8):1–16.
 19. Li C, Wang T, Li Y, *et al.* Probiotics for the treatment of women with bacterial vaginosis: A systematic review and meta-analysis of randomized clinical trials. *Eur J Pharmacol*. 2019;864:172660.
 20. Wirantari N, Hidayati AN. Peran Lactobacillus Pada Managemen Infeksi Endogen Genitalia Wanita. *Media Dermato-Venereologica Indones*. 2019;45(2):100–4.
 21. Heczko PB, Tomusiak A, Adamski P, *et al.* Supplementation of standard antibiotic therapy with oral probiotics for bacterial vaginosis and aerobic vaginitis: A randomised, double-blind, placebo-controlled trial. *BMC Womens Health*. 2015;15(1):1–12.