

DIGITAL HEALTH

Impact Factor: 4.687

5-Year Impact Factor: 4.249

[← Previous issue](#)

[Next issue →](#)

Volume 8, January-December 2022

[View issue contents](#) ▾

Select all Export selected citations

Original Research

 Open Access | Research article | First published January 4, 2022

[Participatory health research under COVID-19 restrictions in Bauchi State, Nigeria: Feasibility of cellular teleconferencing for virtual discussions with community groups in a low-resource setting](#)

Khalid Omer, Umaira Ansari, Amar Aziz, Khalid Hassan, Lami Aminati Bgeidam, [...] [View all](#) ▾

[Preview abstract](#) ▾

PDF / EPUB 

Original Research

 Open Access | Research article | First published January 5, 2022

[Promise and peril-defining ethical telehealth practice from the clinician and patient perspective: A qualitative study](#)

Amanda Jane Keenan , George Tsourtos, Jennifer Tieman

[Preview abstract](#) ✓

PDF / EPUB



Original Research

 Open Access | Research article | First published January 7, 2022

[Digitally enabled flash glucose monitoring for inpatients with COVID-19: Feasibility and pilot implementation in a teaching NHS Hospital in the UK](#)

Tim Robbins , Adam Hopper, Jack Brophy , Elle Pearson, [...]

[View all](#) ✓

[Preview abstract](#) ✓

PDF / EPUB



Original Research

 Open Access | Research article | First published January 20, 2022

[Online Public Attention of COVID-19 Vaccination in Mainland China](#)

Lisha Jiang , Qingxin Ma, Shanzun Wei, Guowei Che

[Preview abstract](#) ✓

PDF / EPUB



Original Research

 Open Access | Research article | First published January 20, 2022

[Risk-standardized sepsis mortality map of the United States](#)

Jiun-Ruey Hu, Chia-Hung Yo, Hsin-Ying Lee, Chin-Hua Su, Ming-Yang Su, [...]

Privacy

[Preview abstract](#) 

PDF / EPUB



Original Research

 Open Access | Research article | First published January 20, 2022

[Smart health via mHealth? Potentials of mobile health apps for improving prevention and adherence of breast cancer patients](#)

Stefanie Scholz , Laura Teetz

[Preview abstract](#) 

PDF / EPUB



Opinion Piece

 Open Access | Research article | First published January 21, 2022

[A unified health algorithm that teaches itself to improve health outcomes for every individual: How far into the future is it?](#)

Gaurav Laroia , Benjamin D Horne , Sean Esplin , Vasant K Ramaswamy

[Preview abstract](#) 

PDF / EPUB



Qualitative Study

 Open Access | Research article | First published January 24, 2022

[The feasibility and acceptability of digital technology for health and wellbeing in social housing residents in Cornwall: A qualitative scoping study](#)

Sarah Ann Buckingham , Tim Walker , Karyn Morrissey ,

[Preview abstract](#) 

Privacy

PDF / EPUB



Review Article

 Open Access | Review article | First published January 27, 2022

[The application of artificial intelligence and custom algorithms with inertial wearable devices for gait analysis and detection of gait-altering pathologies in adults: A scoping review of literature](#)

Ashley Cha Yin Lim , Pragadesh Natarajan , R Dineth Fonseka , Monish Maharaj , Ralph J Mobbs

[Preview abstract](#) 

PDF / EPUB



Original research

 Open Access | Research article | First published January 27, 2022

[CNN-based severity prediction of neurodegenerative diseases using gait data](#)

Çağatay Berke Erdaş , Emre Sümer , Seda Kibaroğlu

[Preview abstract](#) 

PDF / EPUB



Original Research



Open Access

Research article

First published January 28, 2022

[Comparison of the effectiveness of an e-health program versus a home rehabilitation program in patients with chronic low back pain: A double blind randomized controlled trial](#)

Inmaculada Carmen Lara-Palomo, Eduardo Antequera-Soler, [...]

[View all](#) [Preview abstract](#)

PDF / EPUB



Original research



Open Access

Research article

First published January 28, 2022

[The role of age and digital competence on the use of online health and social care services: A cross-sectional population-based survey](#)

T Heponiemi, A-M Kaihlanen, A Kouvonen, L Leemann, S Taipale, K Gluschkoff

[Preview abstract](#)

PDF / EPUB



Qualitative Study



Open Access

Research article

First published February 1, 2022

[Virtual Care Access and Health Equity during the COVID-19 Pandemic, a qualitative study of patients with chronic diseases from Canada](#)

Sophy Chan-Nguyen PhD, Benjamin Ritsma MD, Lisa Nguyen MD, [...]

[View all](#) [Preview abstract](#)

PDF / EPUB



Brief Communication



Open Access

Research article

First published February 2, 2022

[Privacy](#)

[Integration of new digital antenatal care tools using the WHO SMART guideline approach: Experiences from Rwanda and Zambia](#)

Rosemary Muliokela, Gilbert Uwayezu, Candide Tran Ngoc, María Barreix, [...]

[View all](#) ▾

[Preview abstract](#) ▾

PDF / EPUB



Original research

 Open Access | Research article | First published February 3, 2022

[A mobile app for home-based exercise in spinal cord injured persons: Proposal and pilot study](#)

Emiliana Bizzarini, Luca Chittaro, Mauro Frezza, Marika Polo, Cristina Malisan, [...]

[View all](#) ▾

[Preview abstract](#) ▾

PDF / EPUB



Original Research

 Open Access | Research article | First published February 4, 2022

[Social media-based intervention to promote HBV screening and liver cancer prevention among Korean Americans: Results of a pilot study](#)

Y. Alicia Hong, Soo Yee, Pramita Bagchi, Hee-soon Juon, Sojung Claire Kim, Daisy Le

[Preview abstract](#) ▾

PDF / EPUB



Quantitative Study

 Open Access | Research article | First published February 7, 2022

[Developing and validating a new scale to measure the acceptability of health apps among adolescents](#)

Privacy

Elizabeth Chen , Kathryn E. Moracco, Kirsten Kainz, Kathryn E. Muessig, Deborah F. Tate

[Preview abstract](#) 

PDF / EPUB



Review Article

 Open Access | Review article | First published February 7, 2022

[Patients' acceptance of video consultations in the mental health services: A systematic review and synthesis of qualitative research](#)

Anne Marie Moeller , Lone F. Christensen, Jens Peter Hansen, Pernille T. Andersen

[Preview abstract](#) 

PDF / EPUB



Special Collection: - Special Collection on Covid-19

 Open Access | Research article | First published February 7, 2022

[Rise of tele dermatology in the COVID-19 era: A pan-world perspective](#)

Piyu Parth Naik

[Preview abstract](#) 

PDF / EPUB



Original research

 Open Access | Research article | First published February 7, 2022

[Exploring healthcare professionals' perceptions of artificial intelligence: Piloting the Shinners Artificial Intelligence Perception tool](#)

Lucy Shinners , Sandra Grace, Stuart Smith, Alexandre Stephens, Christina Aggar

[Preview abstract](#) 

Privacy

PDF / EPUB



Original Research

 Open Access | Research article | First published February 8, 2022

[Mobile phone apps for family caregivers: A scoping review and qualitative content analysis](#)

Jamie Yea Eun Park , Christopher Shawn Tracy , Carolyn Steele Gray

[Preview abstract](#) 

PDF / EPUB



Original Research

 Open Access | Research article | First published February 9, 2022

[A pilot study using financial transactions' spatial information to define high-risk neighborhoods and distribution pattern of COVID-19](#)

Esmail Mohammadi , Mehrdad Azmin , Nima Fattahi , Erfan Ghasemi , [...]

[View all](#) 

[Preview abstract](#) 

PDF / EPUB



Review article

 Open Access | Review article | First published February 10, 2022

[COVID-19 infodemic and digital health literacy in vulnerable populations: A scoping review](#)

Mohamed-Amine Choukou , Diana C Sanchez-Ramirez , Margriet Pol , [...]

[View all](#) 

[Preview abstract](#) 

PDF / EPUB



Privacy

Review article

 Open Access | Review article | First published February 11, 2022

[Re-focusing explainability in medicine](#)

Laura Arbelaez Ossa , Georg Starke , Giorgia Lorenzini, Julia E Vogt, [...]

[View all](#) 

[Preview abstract](#) 

PDF / EPUB



Qualitative Study

 Open Access | Research article | First published February 14, 2022

[Digital Health Testbeds in Sweden: An exploratory study](#)

Md Shafiqur Rahman Jabin , Evalill Nilsson, Anna-Lena Nilsson, Patrick Bergman, Päivi Jokela

[Preview abstract](#) 

PDF / EPUB



Brief Communication

 Open Access | Research article | First published February 21, 2022

[Acceptance and use of telemedicine technology by health professionals: Development of a conceptual model](#)

Mohammed Roudi , Abdelmajid Elouadi, Amine Hamdoune

[Preview abstract](#) 

PDF / EPUB



Commentary

Privacy



Open Access

| Research article

| First published February 21, 2022

[Physician centricity in the deployment of digital solutions for neurological conditions](#)

Graham B Jones , Jason Freeman, Allison Mann, Amit Khanna

[Preview abstract](#)

PDF / EPUB



Review article



Open Access

| Review article

| First published February 23, 2022

[A scoping review of feasibility, cost-effectiveness, access to quality rehabilitation services and impact of telerehabilitation: A review protocol](#)

Eugene Nizeyimana , Conran Joseph, Quinette A Louw

[Preview abstract](#)

PDF / EPUB



Original research



Open Access

| Research article

| First published February 23, 2022

[‘Using humanity to change systems’ – understanding the work of online feedback moderation: A case study of Care Opinion Scotland](#)

Emma Berry , Zoë C Skea , Marion K Campbell, Louise Locock

[Preview abstract](#)

PDF / EPUB



Original Research



Open Access

| Research article

| First published February 25, 2022

DIGITAL HEALTH

Impact Factor: 4.687

5-Year Impact Factor: 4.249

Editorial board

 [Hide all](#)

Joint Editors-in-Chief

**Professor Theodoros N. Arvanitis, RT, DPhil, CEng,
FRSM**

University of Warwick, UK

Professor Thierry Moulin, MD, PhD

University of Franche-Comte,
France

Consulting Editor

Dr Samir Damani, MD, PharmD

MD Revolution, USA

Co-founding Editor

**Professor John Powell, MA MB BChir MSc
PGCMedEd PhD MRCPsych FFPH**

University of Oxford & University of
Manchester, UK

Associate Editors



Dr Helen Atherton, BSc, MSc, MPH, PhD

University of Warwick, UK

Professor Natale Daniele Brunetti, MD, PhD

University of Foggia, Italy

Dr Jennifer Graves, MD, PhD

University of California, San
Francisco, USA

Dr Shariful Islam, MBBS, MPH, PhD, FESC

Deakin University, Melbourne,
Australia

Dr Vincenzo Della Mea, MSc, PhD

University of Udine, Italy

**Dr Edward Meinert, MA, MSc, MBA, MPA, PhD,
CEng FBCS**

University of Plymouth, UK

Dr Joanna Milward

King's College London, UK

Dr Chris Paton, BMBS, BMedSci, MBA, FACHI

University of Oxford, UK

Dr Tim Robbins

University of Warwick, UK

Dr Aenor J Sawyer, MS

University of California, San
Francisco, USA

Dr Eline Smit

Editorial Board Members



Dr Urs-Vito Albrecht, MD, MPH

Hannover Medical School, Germany

**Professor Panagiotis D. Bamidis,
PhD**

Aristotle University of Thessaloniki, Greece

Privacy

Professor Alex Broom, MA, PhD

University of New South Wales, Australia

Dr Vasa Curcin

King's College London, UK

**Professor Kathleen Griffiths, BSc,
PhD**

Australian National University, Australia

**Professor Flis Henwood, BA, MSc,
PhD**

University of Brighton, UK

**Professor Deborah Lupton, BA,
LittB, MPH, PhD**

University of New South Wales (UNSW) Sydney,
Australia

**Dr Fadhila Mazanderani, BIT, MSc,
DPhil**

University of Edinburgh, UK

**Professor Anne Moen, RN, PhD,
FACMI**

University of Oslo, Norway

**Professor Jyotishman Pathak,
PhD**

Cornell University, USA

**Professor Catherine Pope, BA,
PhD**

Nuffield Department of Primary Care Health
Services, University of Oxford, UK

Professor Chunhua Weng

Columbia University, USA

Professor Sue Ziebland, BA, MSc

University of Oxford, UK

[Browse journal](#)

[All articles](#)

[Browse by year](#)

[Special collections index](#)



DIGITAL HEALTH

Impact Factor: 4.687

5-Year Impact Factor: 4.249

Journal description

DIGITAL HEALTH is a peer-reviewed open access journal which provides universally accessible and digestible content to all stakeholders involved in the digital healthcare revolution. It provides a unique forum for dissemination of high quality content applicable to researchers, clinicians and allied health practitioners, patients, social scientists, industry and government. Please see the Aims and Scope tab for further information.

This journal is a member of the [Committee on Publication Ethics \(COPE\)](#).

Why publish in Digital Health?

- Indexed in PubMed Central and Web of Science (ESCI)
- A leading Editorial Board of top researchers
- High-quality and efficient peer review
- An open access format ensuring maximum visibility
- Tracking the impact of your article with integrated Altmetric tools

Open access article processing charge (APC) information

As of 1st January 2022 the APC for this journal is 1600 USD.

The article processing charge (APC) is payable when a manuscript is accepted after peer review, before it is published. The APC is subject to taxes where applicable. Please see further details [here](#).


Submission information

Submit your manuscript today at <https://mc.manuscriptcentral.com/dhj>

Please see the Submission Guidelines tab for more information on how to submit your

[Privacy](#)

Use of mobile app to monitoring growth outcome of children: A systematic literature review

Digital Health
Volume 8: 1–10
© The Author(s) 2022
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/20552076221138641
journals.sagepub.com/home/dhj


Theresia Chrisanthy Kustiawan¹, Siti Rahayu Nadhiroh¹ , Roziana Ramli²
and Chaniphun Butryee³

Abstract

Advances in knowledge and technology have created opportunities to help monitor child growth. Thus, we conducted a systematic review to determine if the use of mobile apps resulted in improved growth outcomes for children. We include articles published related to children's growth with poor nutritional status. The relevant articles were searched from PubMed, ScienceDirect, Scopus, ProQuest, and Google Scholar. Twelve studies were identified, which is the use of the mobile app to monitor growth in undernutrition and obesity in children. Six studies found that the use of mobile apps improved undernutrition child growth and improved parents' and/or front health workers' knowledge to prevent, treat, and monitor children with undernutrition. Six studies stated that the use of mobile app helps overweight/obese children lose weight and motivate them to achieve ideal body weight. Mobile apps for monitoring the growth of children with various standards are likely a promising means for early detection of growth failure and guiding overweight/obese children in gaining normal weight. Studies with large sample sizes and long-term interventions and follow-ups are needed to help assess the effectiveness of mobile app intervention programs and their impact on multiple growth outcomes more comprehensively and accurately.

Keywords

Mobile app, children, growth outcome, undernutrition, obesity

Submission date: 11 March 2022; Acceptance date: 4 October 2022

Introduction

The early years of a child's life are the most important period for their health and development. Early child development sets the foundation for lifelong learning, behaviour, and health. The healthy growth and development of infants and young children are of paramount importance for children to develop their full physical and mental potentials.¹ Although child growth is a crucial period, more than 100 million children worldwide are failing to grow. In 2020, globally, WHO stated about 149.2 million children under the age of 5 years were stunted, 45.4 million wasted, and 38.9 million overweight.²

Child growth failure (CGF) is expressed as stunting, undernutrition, and obesity, and children under 5 years of age (0–59 months) are a specific subset of undernutrition excluding micronutrient deficiencies, that is characterized by the relationship between insufficient height and weight at a given age, and this subset is most universally described in terms of univariate 'growth standards',

for which age-specific heights and weights are compared to healthy reference populations.^{3,4} In a child with suspected impaired growth, a detailed evaluation should be conducted to identify the cause.⁵ Based on walker's research, risk factors for child growth are nutrition (childhood undernutrition, overnutrition, iodine deficiency, iron deficiency, and other nutritional factors), infectious disease, environmental exposures, and parenting factors.⁶ Therefore, a child with growth failure need to be treated as soon as possible.

¹Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia

²Department of Computer System and Technology, Faculty of Computer Science and Information Technology, Universiti Malaya, Kuala Lumpur, Malaysia

³Institute of Nutrition, Mahidol University, Nakhon Pathom, Thailand

Corresponding author:

Siti Rahayu Nadhiroh, Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Kampus C Unair Mulyorejo, Surabaya 60115, Indonesia.

Email: sitinadhiroh@fkm.unair.ac.id



Child with growth failure under treatment will have their growth to be assessed and recorded over time. This will allow the child to be monitored by comparing their growth with a normal range for other children of the same age and gender, relative to a reference population.⁷ Advances in knowledge and technology have created opportunities to help monitor the treatment progress of a child with poor nutritional status. Since child care is part of health service in that provided by the government and recorded, the use of technology to monitor child growth and development is needed. Emergence of smartphones and mobile devices has the potential to facilitate the use of technology in monitoring such as mobile applications. In 2021, the percentage of mobile device users reached 89.76% of the world's population and the percentage of internet users worldwide reached 59.5% of the global population; in total of 92.6% mobile device user who accessed the internet^{8,9} indicating mobile applications are accessible to anyone. Thus, mobile applications to monitor child growth provide many great benefits because it is easy to use, saves time, can be recorded automatically, and is available to millions of users. The objective of this systematic literature review was therefore to see the result of the use of mobile apps to monitor growth outcomes.

Methods

Search strategy

We identified the eligible literature through a systematic search in five electronic databases: PubMed, ScienceDirect, Scopus, ProQuest, and Google Scholar without time-window restriction. The search used a combination of keywords of mobile app, monitor, growth (anthropometric, growth, weight, length/height), and poor nutritional status in the child (underweight, stunting, wasting, malnutrition). After the abstracts were retrieved and screened, we evaluated the full text of the articles that related to mobile app use for monitoring and children's growth.

To read the article, first, pay attention to the title, then read the article and make notes. The article was retrieved and read by the first author. The first author and second author discussed the retrieved article, then the second author and the rest authors discussed and came to an agreement to use the article.

Inclusion and exclusion criteria

To be eligible for inclusion, the article must present the data from an observational study that includes (a) use of mobile app to monitor a child with poor nutritional status; (b) quantitative and descriptive result; (c) original research articles. The papers that are included are English papers.

Retrieved articles were excluded if (a) the outcome only had a descriptive result and/or was not defined/determined by number; (b) retrieved articles are non-English papers;

(c) monitored subjects above 19 years old; (d) non-original research articles and grey literature. As the use of mobile apps to monitoring malnutrition has been reviewed thoroughly elsewhere, the aim of the present review was to see the result of the use of mobile apps to monitoring growth outcomes which include nutritional status in a child such as stunting and malnutrition.

Data extraction

For all articles, the following data were independently extracted: year of publication, study design, participant sampling, country, number of participants, and growth outcome. All articles use mobile apps with technology features to monitor the treated child, which includes recorded assessment anthropometry of the monitored child, comparing the assessed growth data with a standard growth chart, and recommendations or tips to improve child conditions.

Result

The literature search identified 5346 records, of which 3723 were excluded after the removal of duplicates. Using the inclusion and exclusion criteria by screening the title and abstract for relevance to the research question, we selected 40 full texts for further assessment. A total of 22 studies were excluded after screening for relevant and sufficient information. In total, 12 studies were included in our final systematic review (Figure 1).

Mobile app for monitoring undernutrition in child

Six studies evaluated the result of using mobile applications to monitor undernutrition in a child at least with one indicator such as parents/caretaker knowledge or anthropometric scales (weight for age (WAZ), height for age (HAZ), weight for height (WHZ), and BMI for age (BMIZ)). The result of these studies shown in Table 1. indicated that mobile app usage for monitoring improved undernutrition child growth and parents' and front health workers' knowledge to prevent, treat, and monitor a child with malnutrition.¹⁰⁻¹⁵

Mobile app for monitoring obesity in child

Six studies evaluated the result of using mobile applications to monitor obesity in a child at least with one indicator such as knowledge or anthropometric scales (weight for height (WHZ), BMI for age (BIMZ)). The result of these studies shown in Table 2. stated that mobile app usage for monitoring helps overweight/obese children lose weight and motivates the child to achieve an ideal body weight.¹⁶⁻²¹

Discussion

The mobile phone application for monitoring growth in a child is mainly designed for two end users: healthcare




Figure 1. Flowchart for article selection.

workers or parents. Overall, the use of this technology by these two end users has a positive impact on child growth. However, the contexts of the impact differ between these end users.

The moderate underweight children were often misclassified as normal weight, which resulted in lost opportunities for early identification of malnutrition. The misclassification cases were mainly associated with the healthcare workers from community volunteers who have minimal training and supervision on the classification of children's growth status. The automated calculation to determine the children's growth status in mobile phone applications has helped untrained and less experienced healthcare workers to accurately classify the children's growth status. The visual display of the growth chart in the mobile phone application increases the understanding of inexperienced healthcare workers which results in a better response by scheduling follow-up counselling sessions/referrals. Other

than that, the mobile phone application also expedites data collection from community health posts and submission to higher-level healthcare providers compared to the conventional paper-based method allowing a quicker and effective intervention to help the children.

The mobile phone application for monitoring growth in child benefits parents in a variety of ways such as helping the parents to better understand and interpret the growth charts. Individual or dedicated visual feedback on a child's development makes it easier for parents to monitor their child's progress. During the visit, the mobile phone application also serves as a logbook or manual registration book, replacing the hand-held book. The hand-held book is easily lost or forgotten during the appointment, resulting in a lack of data measured.

The result of the current review is in line with other systematic reviews on using mobile apps to monitor nutritional outcomes. DiFillipo's review about the use of mobile apps

Table 1. Use of mobile app for monitoring undernutrition in children.

Authors (year)	Methodology	Sample and location	Number of participants	Population sample characteristic	Duration of mobile app usage	Features of the mobile application	Indicator	Outcomes
Patel et al. (2019) ¹⁰	Randomized controlled trial	Maharashtra, India	2501	Children and rural women	6 months	NR	HAZ	Reduce the prevalence of stunting at 18 months of age in the intervention clusters by 8% as compared with the control clusters.
Mistry et al. (2019) ¹¹	Cross-sectional survey	Bangladesh	3009	mother-children	3 months	NR	HAZ	The study revealed that the prevalence of stunting was significantly lower in areas where the intervention was delivered compared to the comparison areas (29% vs. 37%, $p < 0.001$).
Seyyedi et al. (2020) ¹²	Randomized controlled trial	Iran	110	Children under 3 years old	6 months	Nutrition status profile, food and child growth education	WAZ, WHZ	Children in the smartphone group intervention showed greater wasting status improvement (WHZ + 0.65 (95% Confidence Interval (CI) ± 0.16)) than children in the TAU group (WHZ + 0.31 (95% CI ± 0.21); $p = 0.011$) and greater reduction (89.6% vs. 51.5%; $p = 0.016$) of wasting caseness (i.e. WHZ < -2; yes/no).
Rokhaidah (2021) ¹³	Purposive sampling, Indonesia	Indonesia	60	Children	5 months	Food and child growth education	HAZ	The results of the statistical test with Mann-Whitney analysis showed a p value of 0.007, which means there was a significant difference between the independent practice of the control group and the intervention group at the time of the post test, so the use of the growth monitoring application has a good effect on the mother's independent practice in monitoring stunting.
Chanani et al. (2016) ¹⁴	Experimental	Mumbai	14	Frontline Worker	2 months	Nutrition status profile and record	WAZ, HAZ, WHZ	Comparing FWs ($N = 14$) who completed at least 40 screenings without and 40 with the calculator, the error rates

(continued)

Table 1. Continued.

Authors (year)	Methodology	Sample and location	Number of participants	Population sample characteristic	Duration of mobile app usage	Features of the mobile application	Indicator	Outcomes
Huda et al. (2018) ¹⁵	Mixed-methods pilot study	Bangladesh	340	Pregnant or recently delivered, lactating women	24 months	Food and child growth education	WAZ, WHZ	were 5.5% and 0.7%, respectively ($p < 0.0001$). The mobile calculator significantly reduces an important component of human error in using the WHO tables to calculate data and assess acute malnutrition at the community level. Less than 10% (22/275) of women reported difficulties understanding the voice messages or direct counselling through mobile phones. Approximately 87% (236/275) of women reported spending the cash to purchase food for themselves and their children. Mothers gave positive feedback on the program.

Table 2. Use of mobile app for monitoring obesity in children.

Authors (year)	Methodology	Sample and location	Number of participants	Population sample characteristic	Duration of mobile app usage	Features of the mobile application	Indicator	Outcomes
Nyström et al. (2017) ¹⁶	Randomization	Sweden	300	4–4.5 years old children	6 months	NR	WHZ	The intervention group increased their mean composite score from baseline to follow-up, whereas the control group did not ($+0.36 \pm 1.47$ compared with -0.06 ± 1.33 units; $p = 0.021$). This improvement was more pronounced among the children with an FMI above the median (4.11 kg/m^2) ($p = 0.019$).
Pretlow et al. (2015) ¹⁷	Pilot study	Seattle, Washington	43	Children and adolescent	20 weeks	Nutrition status profile, health consultation	WHZ, BMIZ	There was a significant decrease in % BMI over time of 7.1. There were significant improvements in participant ratings of self-esteem, control over food, and a reduction in turning to food when stressed. Males, younger participants, and participants with higher levels of program compliance achieved better weight loss.
Browne et al. (2020) ¹⁸	Randomized controlled trial	Ireland	20	Children and adolescent	8 weeks	NR	WHZ, BMIZ	The Child Behaviour Checklist (CBCL) mean total score was 71.7 (SD 3.1) in the intervention group vs 57.6 (SD 6.6) in the control group, t -test $p < 0.001$, the use of mobile health apps had a positive result.

(continued)

Table 2. Continued.

Authors (year)	Methodology	Sample and location	Number of participants	Population sample characteristic	Duration of mobile app usage	Features of the mobile application	Indicator	Outcomes
Johansson et al. (2020) ¹⁹	Two-arm feasibility trial	Sweden	28	Children	6 months	Nutrition status profile, food and child growth education	WHZ, BMIZ	At 6 months, the intervention group had a greater reduction of 0.24 units in BMI SDS than standard care (-0.23 vs. 0.01 , $p = 0.002$). The mHealth support system was a feasible and innovative treatment approach which, in addition to standard care, generated better treatment results than standard care alone.
Bovi et al. (2018) ²⁰	Randomized controlled trial	Italy	48	Children	12 months	Nutrition status profile, food and health education	Neck circumference	(1) 74% of children answered to >50% of WhatsApp messages. (2) 43.5% and 68.3% dropped out in IG and CG groups, respectively ($p = 0.14$). (3) IG showed a meaningful improvement in neck circumference excess (-69.4% in IG and $+9.45\%$ in CG; $p = 0.012$) with only minor trends for the other anthropometric parameters, and (4) for healthy behaviour (screen-time ($p = 0.16$); sweetened drinks ($p = 0.17$) and vegetables ($p = 0.09$) consumption).

(continued)

Table 2. Continued.

Authors (year)	Methodology	Sample and location	Number of participants	Population sample characteristic	Duration of mobile app usage	Features of the mobile application	Indicator	Outcomes
Cueto et al. (2019) ²¹	Retrospective cohort study	United States	1120	Children and adolescent	12–16 weeks	Nutrition status profile, health consultation	WHZ, BMIZ	Adjusted for commitment period, sex, and age group, the overall mean change in %BMIp95 was -0.21 (95% CI -0.25 to -0.17) per additional coaching session ($p < 0.001$). Among overweight and obese children using a mobile app-based health coaching and behaviour change program, increased engagement was associated with longer voluntary commitment periods, and increased number of coaching sessions was associated with decreased weight status.

Acronym: WAZ: weight for age; HAZ: height for age; WHZ: weight for height; BMIZ: BMI for age; NR: not reported.

to improve nutrition outcomes among adults resulting in behavioural changes in reviewed studies included increased adherence to diet monitoring with an app but decreased effort to continue diet without an app. Moreover, Langarizadeh's review about mobile apps for weight management in children and adolescents resulted in lower body mass, decrease in body weight, and improvement in physical activity.^{22,23}

Despite the fact that the mobile phone application for monitoring growth in a child provides various benefits to healthcare workers, the mobile phone application does not ease the workload of the healthcare workers when they are already overburdened with various tasks. Budget restrictions also make it difficult to provide healthcare workers with mobile phones. Thus, deploying the initiative on a large scale might be difficult without financial support and commitment from the government or a private organization. Issues related to the use of the digital tools that emerged during the studies included data loss as a result of technical difficulties with the server, a lack of mobile phone network coverage in remote areas, and the necessity to charge the phone on a regular basis. Furthermore, parents voiced concerns about the data acquired on their children, including concerns about its security and possible privacy risks.

Several aspects of the mobile phone application for monitoring growth in a child can be improved as outlined by the studies in this systematic review. The culture or local contexts need to be incorporated into the information provided in the mobile phone application to make it more appropriate for the local environment. Appropriate training is required, and complete deployment needs a pilot study to determine the appropriate local features. The ability to exchange information with a family doctor or other healthcare professional is important for some of the parents. The lack of mobile phone network coverage in remote areas requires the ability to directly transfer the data from the mobile phone to the computer via a cable.

Furthermore, validation of the data entered by parents, which is a significant first step, is needed as reported to be missing in many apps.²⁴ Not much research has been conducted which leads to limitations to this study. In addition, some of the studies in the current review presented in small sample size and short follow-up period. In future, more robust research works are necessary to address these limitations.

Conclusion

Mobile apps for monitoring the growth of children with various standards are likely a promising means to early detection of growth failure, guiding overweight/obese and undernutrition children in gaining normal weight. In the future, more research to validate the data self-monitored by using mobile apps should be conducted. Furthermore,

studies with large sample sizes and long-term interventions and follow-ups are needed to help assess the effectiveness of mobile app intervention programs and their impact on multiple growth outcomes more comprehensively and accurately.

Acknowledgement: Not Applicable.

Contributorship: SRN was responsible for conceptualizing the study and approved the final revision of the protocol. TCK was collecting the literature, writing the first draft, and the final revision of the review protocol. RR was writing the discussion section. CB assisted in conceptualization.

Declaration of conflicting interests: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Ethical approval: This is a scoping review of published literature and does not require ethical approval

Funding: The author(s) received no financial support for the research, authorship, and/or publication of this article.

Guarantor: Siti Rahayu Nadhiroh.

ORCID ID: Siti Rahayu Nadhiroh  <https://orcid.org/0000-0002-2870-6094>

References

1. de Onis M. Child growth and development. *Nutr Health Dev World* 2017; 119–141.
2. WHO. Levels and trends in child malnutrition. Key findings of the 2021. ISBN: 9789240025257. 2021.
3. Damris KK, Aaron EO-Z, Morenike OF, et al. Mapping child growth failure across low- and middle-income countries. *Nature* 2020; 577: 231–234.
4. WHO. WHO Child Growth Standards and the Identification of Severe Acute Malnutrition in Infants and Children: A Joint Statement by the World Health Organization and the United Nations Children's Fund. World Health Organization. 2009.
5. Rogol AD and Hayden GF. Etiologies and early diagnosis of short stature and growth failure in children and adolescents. *J Pediatr* 2014; 164: S1–S14. e6.
6. Walker SP, Wachs TD, Meeks GJ, et al. Child development: risk factors for adverse outcomes in developing countries. *The Lancet* 2007; 369: 145–157.
7. Humphrey G, Dobson R, Parag V, et al. See how they grow: testing the feasibility of a mobile app to support parents' understanding of child growth charts. Gopichandran V, editor. *PLOS ONE* 2021; 16: e0246045.
8. Turner A. How Many Smartphones Are In The World?. BankMyCell. 2021, <https://www.bankmycell.com/blog/how-many-phones-are-in-the-world>.

9. Johnson J. Digital users worldwide 2020. Statista, <https://www.statista.com/statistics/617136/digital-population-worldwide>.
10. Patel AB, Kuhite PN, Alam A, et al. M-SAKHI—Mobile health solutions to help community providers promote maternal and infant nutrition and health using a community-based cluster randomized controlled trial in rural India: A study protocol. *Matern Child Nutr* 2019; 15: e12850.
11. Mistry SK, Hossain M and Arora A. Maternal nutrition counselling is associated with reduced stunting prevalence and improved feeding practices in early childhood: a post-program comparison study. *Nutr J* 2019; 18: 47.
12. Seyyedi N, Rahimi B, Eslamlou HRF, et al. Smartphone-based maternal education for the complementary feeding of undernourished children under 3 years of age in food-secure communities: randomised controlled trial in Urmia, Iran. *Nutrients* 2020; 12: 587.
13. Rokhaidah R and Herlina H. Pengaruh Pelatihan Penggunaan Aplikasi Pemantau Pertumbuhan Anak Terhadap Praktik Mandiri Ibu Dalam Pemantauan Stunting. *Dunia Keperawatan: Jurnal Keperawatan dan Kesehatan* 2021; 9: 55.
14. Chanani S, Wacksman J, Deshmukh D, et al. M-health for improving screening accuracy of acute malnutrition in a community-based management of acute malnutrition program in Mumbai informal settlements. *Food Nutr Bull* 2016; 37: 504–516.
15. Huda TM, Alam A, Tahsina T, et al. Mobile-based nutrition counseling and unconditional cash transfers for improving maternal and child nutrition in Bangladesh: pilot study. *JMIR Mhealth Uhealth* 2018; 6: e156.
16. Nyström CD, Sandin S, Henriksson P, et al. Mobile-based intervention intended to stop obesity in preschool-aged children: the MINISTOP randomized controlled trial. *Am J Clin Nutr* 2017; 105: 1327–1335.
17. Pretlow RA, Stock CM, Allison S, et al. Treatment of child/adolescent obesity using the addiction model: a smartphone app pilot study. *Child Obes* 2015; 11: 248–259.
18. Browne S, Kechadi M-T, O'Donnell S, et al. Mobile health apps in pediatric obesity treatment: process outcomes from a feasibility study of a multicomponent intervention. *JMIR Mhealth Uhealth* 2020; 8: e16925.
19. Johansson L, Hagman E and Danielsson P. A novel interactive mobile health support system for pediatric obesity treatment: a randomized controlled feasibility trial. *BMC Pediatr* 2020; 20: 447.
20. Bovi APD, Di Michele L, Massa G, et al. P164 A personalized and non-automated mobile-based intervention in the management of paediatric obesity: preliminary results of a pilot study (PediaFit). *Dig Liver Dis* 2018; 50: e415.
21. Cueto V, Wang CJ and Sanders LM. Impact of a mobile app-based health coaching and behavior change program on participant engagement and weight status of overweight and obese children: retrospective cohort study. *JMIR Mhealth Uhealth* 2019; 7: e14458.
22. DiFilippo KN, Huang W-H, Andrade JE, et al. The use of mobile apps to improve nutrition outcomes: a systematic literature review. *J Telemed Telecare* 2015; 21: 243–253.
23. Langarizadeh M, Sadeghi M, As'habi A, et al. Mobile apps for weight management in children and adolescents; an updated systematic review. *Patient Educ Couns* 2021; 104: 2181–2188.
24. Akbar S, Coiera E and Magrabi F. Safety concerns with consumer-facing mobile health applications and their consequences: a scoping review. *J Am Med Inform Assoc* 2019; 27: 330–340.