The relationship between Big Five Personality Traits, eating habits, physical activity, and obesity in Indonesia based on analysis of the 5th wave Indonesia Family Life Survey (2014)

by Trias Mahmudiono

Submission date: 09-Mar-2023 04:51PM (UTC+0800)

Submission ID: 2032867924 **File name:** A8.pdf (590.29K)

Word count: 14759 Character count: 80925







OPEN ACCESS

Fanchang Kong. Fanchang Kong, 71
Central China Normal University, China

Petri Böckerman,

University of Jyväskylä, Finland

Beata Łubianka,

Jan Kochanowski University, Poland

111 ESPONDENCE Trias Mahmudiono

rias-m@fkm.unair.ac.id

This article was submitted to Eating Behavior. a section of the journal Frontiers in Psychology

RECEIVED 22 February 2022 ACCEPTED 30 June 2022 PUBLISHED 04 August 2022

<mark>2</mark> ATION Pristyna G, Mahmudiono T, Rifqi MA and Indriani D (2022) The relationship between Big Five Personality Traits, eating habits, physical activity, and esity in Indonesia based on analysis of the 5th wave Indonesia Family Life

13 ey (2014). Front. Psychol. 13:881436. doi: 10.3389/fpsyg.2022.881436

© 2022 Prist 6, Mahmudiono, Rifqi and Indriani. This is an open-access article distributed under the terms of nse (CC BY). The use, distribution reproduction in other forums is nitted, provided the original or(s) and the copyright owner(s) re credited and that the original publication in this journal is cited, in cordance with accepted academic ractice. No use, distribution or reproduction is permitted which does not comply with these terms

The relationship between Big Five Personality Traits, eating habits, physical activity, and obesity in Indonesia based on analysis of the 5th wave Indonesia Family Life Survey (2014)

Greena Pristyna¹, Trias Mahmudiono^{1*}, Mahmud A. Rifqi¹ and Diah Indriani²

6
Department of Nutrition, Faculty of Public Health, Airlangga University, Surabaya, Indonesia, ²Department of Biostatistics and Medical Demography, Faculty of Public Health, Airlangga University, Surabaya, Indonesia

study investigated the association between Big Five Personality Traits (Openness to new experience, Conscientiousness, Extraversion, Agreeableness, Neuroticism) and nutrition-related variables (eating habits, physical astivity, and obesity using Body Mass Index). We used secondary data from the Indonesia Family Life Survey (IFLS) wave 5 involving a total of 14,473 men and 16,467 women aged 15-101 years (mean = 37.34; SD = 14.916) in Indonesia that was selected by stratified random sampling conducted in the period 2014 to 2015. Data were collected through interviews with the Big Five Index 15 and a questionnaire similar to the Global Physical Activity Questionnaire which was translated into the Indonesian language, and based on measurements of height and weight. Analyses used binary logistic regression test controlled by socio-demographic factors (age, gender, glucation, occupation, and marital status) to determine the association between Big Five Personality Traits and eating habits (recommended and non-recongrended foods), physical activity, and obesity. Results showed that openness and extraversion were positively associated with recommended and non-recommended foods, conscientiousness vas positively associated with only recommended foods, agreeableness was positively associated with recommended foods, and negatively apociated with only one nonrecommended food. Whereas, neuroticism was positively associated with non-recommended foods and negatively associated with recommended foods. Openness (p = 0.010; OR = 1.015; 95% CI = 1.004–1.027) and conscientiousness (p < 0.001; OR = 1.045; 95% CI = $\frac{1.03}{10}$ 0-1.059) were associated with physical activity. Conscientiousness (p = 0.002); OR = 1.025; 95% CI = 1.009–1.041), extraversion (p < 0.001; OR = 1.079; 95% CI = 1.065-1.092), and neuroticism (p < 0.001; OR = 0.966; 95% CI = 0.953-0.978) were associated with obesity. Although some results were

different from previous studies, these associations can be used as strategies of behavioral change due to the adaptation of personality characteristics, which can be modified even though the personality tends to be persistent. Further studies are needed to understand other mechanisms that might underlie this association.

WORDS

Big Five Personality Traits, physical activity, obesity, Body Mass Index, Indonesia, eating habits

Introduction

Individuals need to maintain their health by showing healthy behavior in order to avoid various health problems. Poor eating habits can have a negative effect on an individual's nutritional status (Indonesian Ministry of Health [IMH], 2017); furthermore, a study published by 72 Lancet journal found that the food consumed daily was the leading cause of death compared to smoking and now has become one of the five causes of death worldwight (Indonesian Ministry of Health [IMH], 2019). Moreover, lack of physical activity is also a major risk factor for chronic diseases and premature death due to noncommunicable diseases (World Health Organization [WHO], 2018), although regular physical activity is also an only decrease the risk of various diseases but also can improve mental health and quality of life (World Health Organization [WHO], 2018).

Personality is a dynamic and complex physical and psychological system within the individual that underlies every avior (Ahdiyana, 2011; Riadi, 2012). Personality can be associated with several health conditions that can be explained by personality-related behavioral factors (Tiainen et al. 1913). As one of the personality theories that is often used, the Big Five Personality Traits or Five-Factor Model (FPM) developed by Costa and McCrae (1985) divided human 33 sonality into five major dimensions: Agreeableness (A), Extraversion (E), Neuroticism (N), Conscientiousness (C), and Openness to experience (O) (Feist et al., 2013), which were associated with several health-related behaviors.

Previous studies found an association between personality, particularly Big Five Personality Traits with nutrition-related behaviors (such as eating held) and physical activity) and nutritional status, but most of these studies were nonducted in Western countries, such as the study by Sutin et al. (2011) conducted in America, Terracciano et al. (2009) in Italy, Intiful et al. (2019) in Ghana, the study by Tiainen et 22 (2013) were conducted on people born in Helsinki, Finland, the meta-analysis by Sutin et al. (2016) consisting of the United States, England, and Japan, and many other studies. Similar to the significant cross-cultural differences between Western and Eastern countries, research conducted in Indonesia, one of

the Eastern countries, was scarcely found in a proposition of the Eastern countries, was scarcely found in a proposition of the Eastern countries, was scarcely found in a proposition of the Eastern countries, was scarcely found in a proposition of the Eastern countries, was scarcely found in a proposition of the Eastern countries of the Eastern article. The studies given in literature suggested a relationship between 127 ive Personality Traits and eating behavior or food choices, for example in a study of 288 students (121 males and 167 females) of Institut Pertanian Bogor aged 16-18 years old, the multiple linear regression tests results showed that extraversion who enjoys socializing ($\beta = 0.163$; p = 0.035), openness to experience who has creative and imaginative nature ($\beta = 0.208$; p = 0.013), and agreeableness who has friendly and environmentally sensitive nature ($\beta = 0.265$; p = 0.009) tend to choose to eat vegetables for various reasons (Purnamawati and Yuliati, 2016). In another study of 380 students of Universitas Islam Negeri Suska Riau aged 508-21 years old, the multiple regression test results showed conscientiousness (rxy = -1.230; p = 0.008) and neuroticism (rxy = 0.103; p = 0.022) was related to eating behavior (Nelvi, 2016). Additionally, extraversion was related to restrained eating, agreeableness was related to external and emotional eating, and openness to experience was related to emotional, external, and restrained eating (Nelvi, 2016). A study of 195 students (128 males arguer females) at several universities in South Tangerang linked the Big Five Personality Traits to healthy behavior, one of which was eating behavior, the multiple regression test results showed that personality as a whole did not have a significant influence on healthy behavior, but only openness (p = 0.018; B = 0.228) and openness among females (p = 0.002; B = 0.401) had a significant positive effect on healthy behavior based on the regression coefficient (Rahmadian, 2011). Unfortunately, in this study, health behavior (including eating behavior, exercise, smoking, and alcohol consumption) was measured as one, so there was no visible influence from the other four personalities that might only affect certain healthy behaviors (Rahmadian, 2011). As far as we could find, there were no studies that linked physical activity and nutritional status to Big Five Personality Traits in the Indonesian setting and these studies were conducted with a sample size of less than 400 respondents.

For this reason, 38 conducted this study, especially since evidence related to the association between personality traits, eating habits, physical activity, and nutritional status was scarcely found in a peer-reviewed article in the Indonesian

setting and this research was also conducted using secondary data analysis from the Indonesia Family Life Survey (IFLS) with a sample size of tens of thousands, so the results obtained in this study should better describe the existing situation. Therefore, this study aimed to analyze the relationship between Big Five Personality Traits, eating habits, physical activity, and obesity in Indonesia.

Materials and methods

Indonesia family life survey

Data collection

The study used a quantitative analytic cross sectional design from secondary data analysis of the 2014 Indonesia Family Life Survey (IFLS) or the 5th wave IFLS. IFLS represented about 83% of the Indonesian population (13 out of 26 provinces) in 1993 which was the result of collaboration bety 551 the non-profit RAND corporation and Survey Meter (Strauss et al., 2016). 1553 was a continuous longitudinal socio-economic and health survey based on a household sar 107 and had been carried out 5 times, IFLS1 (1993–1994), IFLS2 (1997–1998), 1099 (2000), IFLS4 (2007–2008), and IFLS5 (2014–2015) (Strauss et al., 2016).

Since IFLS was a continuous longitudinal survey, the sampling scheme of the first wave will determine the next wave. IFLS used a stratified random sampling technique where the sampling scheme was conducted randoml 68 the province to maximize population representation and capture the cultural and socio-economic diversity of Indonesia based on the Susenas (National rioeconomic Survey) in 1993 whose sampling frame was designed by the BPS (Central Bureau of Statistics) based on the 1990 sus (Frankenberg and Karoly, 1995). There were 321 EAs (Enumeration Areas) which were randomly selected from each of the 13 provinces with details of the sampling rate applied to each proving and resulted in a total EA distribution, and separated by sampling urban EAs and EAs in small provinces to facilitate urban-rural and Javaneseanese comparisons (Frankenberg and Karoly, 1995). In the selected EA, households were randomly selected based on the 1993 Susenas list, giving a total of 7,730 target sample households, and complete/partial interviews were obtained for 7,039 households, including 22,327 successfully interviewed individuals in IFLS1 (Frankenberg and Karoly, 1995).

In IFLS5, re-contact was established with respondents sampled since IFLS1 in 1993 with a re-contact rate of 92% of original IFLS1 dynasty households (Strauss et al., 2016). IFLS5 was conducted on the same set of health holds in the previous wave and the same IFLS distribution of 16,204 households and 50,148 individuals were interviewed with 2,662 people who had died since IFLS4 being interviewed with representatives who knew them well (Strauss et al., 2016).

Participants

Participants in this study were selected from all people aged ≥ 15 years taken from the IFLS5 dataset. They were chosen, as the information on certain study variables (eating habits and physical activity) was available only to them. From a total of 50,148 participants successfully interviewed on IFLS5, 19,208 participants (38%) were excluded from the study because they were under 15 years old and provided incomplete information regarding the required variables. If a value was missing or answered "don't know" in one of the studied variables, the respondent would be excluded from the study. Therefore, 30,940 120 cipants (53.2% female) aged 15–101 years in Indonesia were included in this study. The mean age was 37.34 years (SD=14.916).

Variables and measures

Data was collected using Computer Assisted Personal Interviewing (CAPI) system. This system that uses computers to manage survey questionnaires (all in the Indonesian language) so that interviewers can directly enter data during face-to-face interviews (Strauss et al., 241). The independent variable studied in this study was personality based on the Big Five Personality Traits. Variables related to sociodemographic characteristics were used as control variables. Whereas, the dependent variables studied were eating habits (recommended and non-recommended foods), physical activity, and obesity based on BMI.

Socio-demographic factors

The socio-demographic factors used in this study were age (continuous variable), gender, education (not attending school or attending school), occupation (unemployed, self-employed, or worker), and marital status (never married, married/living together, or divorced/separated). The occupation was defined as the respondent's primary employment status (Strauss et al., 2016). All these variables were obtained through interviews (Strauss et al., 2016).

Big five personality traits

Personality data were obtained to 35 µgh interviews (Strauss et al., 2016). The question used was the Big Five Index 15 (BFI 15), a subset of BFI 44 from John a 66 rivastava (1999), which consisted of 15 traits representing all two personality dimensions (openness to new experience, conscientiousness, extraversion, agreeableness, neuroticism) with the 115 µits for each personality dimension (Strauss et al., 2016). A short version of BFI, a 15-item instrument, used to measure personality dimensions, was first made available to SOEP (Socio-Economic Panel) 59 2005 (Dehne and Schupp, 2007). This instrument was used in many large-scale surveys, such as the German Socio-Economic Panel (GSOEP) (Strauss et al., 2016). The BF1 15 instrument

was translated into Indonesian and back into English several times until the English re-translation converged with the BFI 15 English (Strauss et al., 2016). Four lists of the same 15 words were made and the list used was randomly determined in CAPI (Strauss et al., 2016). A five-point ordinal scale was used to represent hours of the respondent believes that the trait represents them, 1 = disagree strongly, 2 = disagree a little, 3 = neither agree nor disagree, 4 = agree a little, and 5 = agree strongly (Strauss et al., 2016). In question (R) (reverse), the value was given in reverse from 5 to 1 (John et al., 1991). Scores on each personality were summed based on the five personality dimensions.

We tested the validity and reliability of the BFI 15 scale using SPSS version 20 with 30,940 respondents based on the number of samples that would be used in this study. Item validity was seen through the value of each item with Pearson product-moment. Items were considered valid if the significance value was p < 0.05. Based on these criteria, all item validity test results were considered valid. Meanwhile, the reliability test was performed using an internal consistency approach which was analyzed using the Cronbach Alpha formula. The determination of the high and low-reliability coefficient was based on Guilford's reliability coefficient classification (Sugiyono, 2011). The result of the reliability test showed that the alpha coefficient value was 0.402. Based on Guilford's reliability coefficient classification, a coefficient value of more than 0.40 to 0.70 means the variable has sufficient reliability category. Therefore, the scale was reliable enough to be used.

Eating habits

Eating habits data were obtained through interviews (Strauss et al., 2016). The foods were representative of foods rich in iron and vitamin A, two micronutrients considered deficient among the Indonesian population, and also include fast foods, fried foods, and sweet snacks to get at some unhealthy eating habits (Strauss et al., 2016). Eating habits were defined as the consumption frequency (number of days) by food type (17 types) in the past week which could be answered by never or between 1 to 7 days which were then classified as not consuming (answered never) and consuming (answered between 1 to 7 days) in this study. The 17 types of food we 17 weet potatoes, eggs, fish, meat (beef, chicken, pork, etc.), dairy, green leafy vegetables, banana, papaya, carrot, mango, sambal, rice, including five non-recommended foods gre instant noodle, fast food, soft drink (coca cola, sprite, etc.), fried snacks (tempe, tahu, bakwan, etc.), and sweet snacks (wajik, geplak, donuts, wafers, chocolate, etc.).

Physical activity

Physical activity data was obtained through interviews with a set of questions that were take from an international survey on physical activities about the types and times of

physical activities carried out in all parts of life (at work, home, and exercise) for at least 10 min continuously in the last 7 days (Strauss et al., 2016). The questionnaire consisted of similar questions as GPAQ but was simpler and translated into Indonesian, therefore the calculation and classification of physical activity were based on GPAQ (Global Physical Activity Questionnaire). However, because in the questionnaire activity duration was only asked for in a time frame, we chose the closest number from that time frame so that METs-minutes/week could be calculated. Respondents who answered < 30 min were counted as 15 min, >30 min counted as 60 min, <4 h counted as 3 h (180 min), and \geq 4 h counted as 5 h (300 min). In this study, the classification of high and moderate levels of physical activity based on reference values (GPAQ Analysis Guide Version 2 by World Health Organization [WHO], 2005) were grouped into one in high and moderate. Therefore, the level of physical activity was then classified into high and moderate, and low.

Obesity

To obtain BMI, tlas equired height data were obtained through measurement using a Seca plastic height board, model 213, measured to t 32 nearest millimeter, meanwhile, body weight was measured using a Camry model EB1003 digital scale, measured to the nearest tenth (one decimal point) of a kilogram (Strauss et al., 2016). Obesity of respondents aged ≥ 19 years was determined by BMI and < 19 years was determined by BMI-for-age (BMI/A) using 1030 Anthro plus software. The cut-off point used refers to the regulation of the Indonesian Ministry of Health which was based on the WHO reference 2007 for children aged 5 to 18 years (Regulation of the Indonesian Ministry of Health number 2 of 2020 Indonesian Ministry of Health [IMH], 2020) and based on the FAO/WHO provisions where for the benefit of Indonesia, the cut-off point was modified based on clinical experience and research results in several developing course ies to determine the BMI of individuals aged > 19 years (Regulation of the Indonesian Ministry of Health number 41 of 2014 Indonesian Ministry of Health [IMH], 2014). In this study, the classification of overweight and obesity based on reference values were grouped into one as obese. Therefore, obesity was categorized into: not obese for BMI: ≤25.0 and BMI/A: ≤+1 SD and obese for BMI: > 25.0 and BMI/A: > +1 SD.

Data analysis

The data that had been collected were tabulated and then tested for statistical correlation with binary logistic regression. First, the bivariate test was conditive between personality (each personality dimension based on the Big Five Personality Traits) and nutrition-related variables (eating habits, physical activity, and of 30 y), then continued with a multivariate test controlled by socio-demographic factors (sex, age, marital)

status, education, and occupation) conducted on significant variables in the bivariate test to examine the independent effect of personality. The covariates were predetermined based on previous studies that showed a confounding effect on relationship between the studied variables (Terracciano et al., 2009; Kye and Park, 2012; Mõttus et al., 2013; Tiainen et al., 2013; Sutin and Terracciano, 2015a,b; Sutin et al., 20 Gustavsen and Rickertsen, 2019; Pfeiler and Egloff, 2020). To examine whether the association between personality traits and nutrition-related variables varied by sex or age group (adolescent, adult, elderly), we also tested 47 is interaction, controlling for socio-demographic factors. Data analysis was carried out using Statistical Package for the Social Sciences (SPSS version 20). A significance level of 95% and an error rate of $\alpha = 5\%$ (0.05). Significant relationships between the tested variables were determined if the p < 0.05.

Ethical considerations

This study 6 ng secondary data analysis was granted ethical clearance by the Airlangga University Faculty of Dental Medicine Health Research Ethical Clearance Commission.

Results

Descriptive statistics

Ti02 ptal sample in this study included 30,940 respondents. All characteristics of study participants are presented in Tables 1.2

Of the 30,940 respondents aged 15-101 years, most were women (53.2%), adults (85.8%), married/living together (72.6%), attending school (96.2%), and working (45.2%), and the mean age was 37.34 years (SD = 14.916). Most of them consumed animal protein in the form of eggs, fish, and meat, rice as carbohydrate sources, green leafy vegetables, banana, sambal, and three non-recommended foods: fried snacks, instant noodles, and sweet snacks. Meanwhile, most of them did not consume dairy as animal protein sources, sweet potatoes as carbohydrate sources, papaya, mango, carrot, and two nonrecommended foods: fast food and soft drink. Respondents mainly consumed rice (99.8%) and did not consume fast food (88.9%). Most of the respondents had moderate and high levels of physical activity (55.2%) which was consistent with their nutritional status, most of them were classified as not obese (68.5%).

Table 2 shows that among the five dimensions of the Big Five Personality Traits, the dimension with the highest average score is agreeableness (11.70 \pm 1.533). Meanwhile, the dimension with the lowest average score is neuroticism (8.04 \pm 1.996).

TABLE 1 Characteristic of study participants.

Characteristics Engagement (v) Barrent and (0)				
Characteristics	Frequency (n)	Percentages (%)		
Gender				
Male	14,473	46.8		
Female	16,467	53.2		
Age				
Adoles cent (15-18 years)	2,868	9.3		
Adult (19-65 years)	26,553	85.8		
Elderly (>65 years)	1,519	4.9		
Marital status				
Married/living together	22,459	72.6		
Divorced/separated	2,378	7.7		
Never married	6,103	19.7		
Education				
Not attending school	1,191	3.8		
Attending school	29,749	96.2		
Occupation				
Unemployed	8,407	27.2		
Self-employed	8,548	27.6		
Worker	13,985	45.2		
Eating habits Sweet potatoes				
Not consuming	20,061	64.8		
Consuming	10,879	35.2		
Eggs				
Not consuming	5,342	17.3		
Consuming	25,598	82.7		
Fish				
Not consuming	6,097	19.7		
Consuming	24,843	80.3		
Meat (beef, chicken, pork, etc.)				
Not consuming	11,559	37.4		
Consuming	19,381	62.6		
Dairy				
Not consuming	21,201	68.5		
Consuming	9,739	31.5		
Green leafy vegetables				
Not consuming	3,691	11.9		
Consuming	27,249	88.1		
Banana				
Not consuming	15,007	48.5		
Consuming	15,933	51.5		
Papaya				
Not consuming	22,933	74.1		
Consuming	8,007	25.9		
Carrot				
Not consuming	19,996	64.6		
Consuming	10,944	35.4		
Mango				
Not consuming	20,401	65.9		
Consuming	10,539	34.1		
Instant noodle				
Not consuming	10,555	34.1		
Consuming	20,385	65.9		
0	.,			

(Continued)

TABLE 1 (Continued)

Characteristics	Frequency (n)	Percentages (%)
Fast food		
Not consuming	27,506	88.9
Consuming	3,434	11.1
Soft drink (Coca cola, sprite, etc.)		
Not consuming	24,922	80.5
Consuming	6,018	19.5
Sambal		
Not consuming	5,877	19.0
Consuming	25,063	81.0
Fried snacks (tempe, tahu, bakwan, etc.)		
Not consuming	10,777	34.8
Consuming	20,163	65.2
Rice		
Not consuming	53	0.2
Consuming	30,887	99.8
Sweet snacks (wajik, geplak, donuts, wafers, chocolate, etc.)		
Not consuming	14,642	47.3
Consuming	16,298	52.7
Physical Activity		
Low	13,872	44.8
Moderate and high	17,068	55.2
Obesity		
Not obesity	21,207	68.5
Obesity	9,733	31.5

TABLE 2 Characteristic of study participants based on Big Five Personality Traits.

Mean ± SD (min-max)
$11.11 \pm 2.006(3-15)$
$11.45 \pm 1.656(3-15)$
$10.33 \pm 1.998 (3\text{-}15)$
$11.70 \pm 1.533 (3\text{-}15)$
$8.04 \pm 1.996 (3\text{-}15)$

Regression results

To examine the relation hip between personality (each personality dimension based on the Big Five Personality Traits) and nutrition-related variables (eating habits, physical activity, and obesity), statistical tests were performed using binary logistic regression due to the large sample size, the data were not normally distributed, and assumption of proportional odds was not met. Supplementary Tables 1, 2 present the regression results on the bivariate test for each variable tested in this study.

From the eating habits of 17 types of food tested, including five non-recommended foods: instant noodles, fast food, soft drink (coca cola, sprite, etc.), fried snacks (tempe, tahu, bakwan, etc.), and sweet snacks (wajik, geplak, donuts, wafers, chocolate, etc.), Supplementary Table 1 shows that openness and extraversion had a significant association with eating habits of all food types except rice. Conscientiousness was associated with eating all food types except mango, rice, and two non-recommended foods (instant noodle and fritz) snacks). Agreeableness was associated with eating fish, green leafy vegetables, fruits (banana, papaya, mango), carrot, and four of the five non-recommended foods (fast food, soft drink, fried snacks, and sweet snacks). Meanwhile, neuroticism was unrelated to eating eggs, fish, sambal, and rice. Therefore, only rice consumption id not have a significant association with all dimensions of Big Five Personality Traits. In addition, the eating habit of green leafy vegetables, bananas, papaya, carr fast food, soft drink, and sweets snacks were associated with all dim 7 sions of the Big Five Personality Traits.

There was also a statistically significant associa between the level of physical activity with openness (p = 0.007), conscientiousness (p < 0.001), and agreeableness (p = 0.008). Conscientiousness was 1.070 times more likely to have more high physical activities than agreeableness (OR = 1.020; 95% CI = 1.005-1.035) and openness (OR = 1.016; 95% CI = 1.004-1.027). There also a statistically significant association between obesity and conscientiousness (p < 0.001), extraversion (p < 0.001), and neuroticism (p < 0.001). Extraversion was 1.093 times more likely to be obese than conscientiousness (OR = 1.052; 95% CI = 1.037-1.068). Meanwhile, neuroticism has a protective effect on obesity (OR = 0.977; 95% (300 0.965-0.989). Next step, a multivariate test controlled with socio-demographic factors (sex, age, marital status, education, and occupation) was conducted on the significant variables in the bivariate test to examine the independent effect of personality.

Supplementary Table 2 shows that there was still a statistically significant association between all food types tested with openness even after controlling for socio-demographic factors, which meant that openness was associated with both recommended and non-recommended foods. All food types showed a positive association, which meant respondents with a higher openness were more likely to consume the tested food type than those with a lower openness. The highest possibility was found in green leafy vegetables (OR = 1.107; CI = 1.088–1.126) and the lowest was in fried snacks (OR = 1.019; CI = 1.007–1.031). In addition, only sweet snacks where all control variables had a significant correlation with the association between openness and eating habits according to the food type.

Openness was also positively associated in adults for all food types tested, except eggs, meat, dairy, green leafy vegetables, fast food, and sweet snacks did not vary by age group. The association was stronger in the elderly between openness with sweet potatoes (OR = 1.067; CI = 1.020–1.117),

fish (OR = 1.112; CI = 1.058–1.168), and instant noodle (OR = 1.070; CI = 1.022–1.121) than in adults ([OR = 1.030; CI = 1.017–1.043]; [OR = 1.053; CI = 1.037–1.070], [OR = 1.056; CI = 1.042–1.070]), but no association was found in adolescents and the rest food types were not associated in adolescents and the elderly. In addition, the relationship between openness and all food types tested did not vary by sex, except fried snacks in males (OR = 1.026; CI = 1.008–1.044), and no association was found in females.

Supplementary Table 3 shows that after controlling for socio-demographic factors, only three non-recommended foods (fast food, soft drink, and sweet snacks) had no statistically significant association, while the others were positively associated with conscientiousness. The highest possibility was found in the green leafy vegetables (OR = 1.101; CI = 1.079-1.124) and the lowest was in sweet potatoes (OR = 1.018; CI = 1.003-1.032). Only meat where all control variables had a significant correlation with the association between conscientiousness and eating habits according to the food type.

Conscientiousness was also found to be positively related to all food types tested in adults, except green leafy vegetables, fast food, to start food, and sure the results also showed that conscientiousness was related to fish (OR = 1.056; CI = 1.008–1.106), meat (OR = 0.949; CI = 0.909–0.991), banana (OR = 1.049; CI = 1.007–1.093), and sambal (OR = 0.939; CI = 0.892–0.988) in adolescents and eggs (OR = 1.073; CI = 1.008–1.042), dairy (OR = 1.143; CI = 1.065–1.226), and papaya (OR = 1.075; CI = 1.006–1.148) in the elderly. The association between conscientiousness and all food types tested did not vary by sex, except dairy (OR = 1.051; CI = 1.028–1.074) and fast food (OR = 1.046; CI = 1.011–1.082) in males and soft drink in females (OR = 0.968; CI = 0.942–0.994).

Supplementary Table 4 shows that after controlling for socio-demographic factors, only fish had no statistically significant association, while the others were positively associated with extraversion, which meant that extraversion was associated with both recommended and non-recommended foods. The highest probability was found in fast food (OR = 1.081; CI = 1.062–1.101) and the lowest was in sweet potatoes, mango, and instant noodle (OR = 1.023; CI = 1.011–1.036). Only sweet snacks where all control variables had a significant correlation with the association between extraversion and eating habits according to the food type.

Extraversion was also positively associated in adults for all food types tested, except meat, dairy, and sweet snacks, which did not vary by age group. The relationship between extraversion and all food types tested did not vary by sex, except sweet potatoes (OR = 1.034; CI = 1.018–1.051), carrot (OR = 1.043; CI = 1.027–1.060), and mango (OR = 1.029; CI = 1.013–1.046) in females, also, no association was found in males. However, the results showed that this

relationship was stronger in females for eggs (OR = 1.065; CI = 1.043–1.087), meat (OR = 1.081; CI = 1.063–1.098), dairy (OR = 1.058; CI = 1.040–1.076), green leafy vegetables (OR = 1.041; CI = 1.016–1.066), instant noodle (OR = 1.026; CI = 1.009–1.043), fast food (OR = 1.092; CI = 1.066–1.118), soft drink (OR = 1.067; CI = 1.043–1.091), soft drink (OR = 1.067; CI = 1.043–1.091), soft drink (OR = 1.068; CI = 1.045–1.068), fried snacks (OR = 1.026; CI = 1.009–1.042), and sweet snacks (OR = 1.044; CI = 1.028–1.061) than in males. This showed that higher extraversion in females was more likely to consume all kinds of healthy and unhealthy foods than males.

Supplementary Table 5 shows that after controlling for socio-demographic factors, only carrots, fast food, fried snacks, and sweet snacks had no statistically significant association with agreeableness. All food types that had a significant association were positively associated with agreeableness and only soft drink (non-recommended food) was negatively associated. This meant respondents with higher agreeableness had a greater probability of not consuming soft drinks compared to those with lower agreeableness. The highest possibility was found in green leafy vegetables (OR = 1.041; CI = 1.018-1.064).

A negative association between agreeableness and soft consumption was found especially in adolescents (OR = 0.942; CI = 0.896-0.990) and in adults (OR = 0.977; CI = 0.957-0.997), but this association did not vary by sex. This negative association was also found in the consumption of fried snacks in males (OR = 0.976; CI = 0.954-0.999) and in adults (OR = 0.981; CI = 0.964-0.997), and sweet snacks in adults (OR = 0.981; CI = 0.966-0.997). In addition, agreeableness was associated in adults for green leafy vegetables (OR = 1.028; CI = 1.003-1.054), banana (OR = 1.027; CI = 1.011-1.044), papaya (OR = 1.024; CI = 1.006-1.043), and mango (OR = 1.018; CI = 1.001-1.035).

Supplementary Table 6 shows that after controlling for socio-demographic factors, only sweet potatoes, banana, papaya, mango, and fast food had no statistically significant association with neuroticism. Four non-recommended foods: instant noodle, soft drink, fried snacks, and sweet snacks were positively associated with neuroticism, while four recommended foods: meat, dairy, green leafy vegetables, and carrot were negatively associated. Fried snacks have the highest OR (OR = 1.021; CI = 1.009-1.033) and the lowest was green leafy vegetables (OR = 0.942; CI = 0.925-0.958). This meant that respondents with higher neuroticism increased their consumption of fried snacks by 1.021 times and decreased their consumption of green leafy vegetables by 0.942 times compared to those with lower neuroticism. Only meat and sweet snacks where all control variables had a significant correlation with the association between neuroticism and eating habits according to the food type.

Neuroticism was also associated in adults with all those four recommended foods (negative association) and four non-recommended foods (positive association)

with the same association. The association between neuroticism and sweet snacks was stronger in the elderly (OR = 1.081; CI = 1.025–1.140) than in adults (OR = 1.016; CI = 1.004–1.029), but no association was found in adolescents. Additionally, 16 roticism was also associated with soft drinks (OR = 1.024; CI = 1.001–1.047) and sweet snacks (OR = 1.018; CI = 1.001–1.034) in females, and fried snacks in males (OR = 1.029; CI = 1.011–1.048). The relationship between neuroticism with dairy, green leafy vegetables, and instant noodles did not vary by sex.

Table 3 shows that after controlling for socio-demographic factors, only agreeableness had a statistically significant association with physical activity. Openness (p = 0.010) and conscientiousness (p < 0.001) were positively associated with physical activity. Respondents with higher conscientiousness had a 1.045 times higher probability of engaging in more intense physical activity than those with lower conscientiousness, while openness had a lower probability of 1.015 times. In addition, all gender, marital status, education, and occupation were controlled with this association.

The relationship between physical activity at 11 greeableness did not vary by sex or age group. Meanwhile, physical activity 20 found to be associated with conscientiousness in adolescents (OR = 1.059; CI = 1.016-1.103) and adults (OR = 1.042; CI = 1.026-1.058), but not in the elderly. This association did not vary by sex. In addition, openness was associated with physical activity only in females (OR = 1.018; CI = 1.002-1.034) and the elderly (OR = 1.0123 CI = 1.017-1.115).

Table 4 shows that there was still association between conscientiousness (p = 0.002), extraversion (p < 0.001), and neuroticism (p < 0.001) with after controlling for socio-demographic factors. Extraversion and conscientiousness were positively associated with obesity, which meant respondents with higher extraversion were likely and conscientiousness was 1.025 times more likely and conscientiousness was 1.025 times more likely to be obese compared to respondents with lower scores. Meanwhile, neuroticism had a negative association or protective effect on obesity (OR = 0.966; CI = 0.953-0.978). In addition, all gender, marital status, age, education, and occupation (except self-employed) were correlated with this association.

Conscientiousness was associated with obesity only i23 males (OR = 1.031; CI = 1.010-1.051) and in adults (OR = 1.027; CI = 1.010-1.044). Neuroticism was associated with obesity only in males (OR = 0.935; CI = 0.916-0.955) and in adults (OR = 0.957; CI = 0.944-0.970). Meanwhile, version was associated with obesity in adolescents (OR = 1.069; CI = 1.012-1.128) and in adults (OR = 1.078; CI = 1.064-1.092), but not in the elderly. The association between obesity and extraversion also appeared to be stronger in females (OR = 1.083; CI = 1.066-1.101) than in males (OR = 1.074; CI = 1.053-1.015). This showed that women who have higher extraversion were more likely to be overweight or obese.

TABLE 3 Multivariate analysis between Big Five Personality Traits with physical activity and control variables.

Variables	Physical Activity			
	P-Value	OR	95% CI	
			Lower	Upper
Openness	0.010	1.015*	1.004	1.027
Age	0.075	1.002	1.000	1.004
Gender				
Male	< 0.001	1.218*	1.159	1.280
Female (referent)				
Maritalstatus				
Married/living together	< 0.001	1.325*	1.236	1.420
Divorced/separated	0.023	1.148*	1.019	1.292
Never married (referent)				
Education				
Not attending school	< 0.001	1.273*	1.122	1.445
Attending school (referent)				
Occupation				
Unemployed	< 0.001	0.702*	0.662	0.743
Self-employed	< 0.001	1.182*	1.116	1.251
Worker (referent)				
Conscientiousness	< 0.001	1.045*	1.030	1.059
Age	0.186	1.001	0.999	1.003
Gender				
Male	< 0.001	1.221*	1.162	1.283
Female (referent)				
Marital status				
Married/living together	< 0.001	1.298*	1.210	1.391
Divorced/separated	0.045	1.129*	1.003	1.271
Never married (referent)				
Education				
Not attending school	< 0.001	1.278*	1.127	1.450
Attending school (referent)				
Occupation				
Unemployed	< 0.001	0.710*	0.670	0.752
Self-employed	< 0.001	1.186*	1.121	1.256
Worker (referent)				
Agreeableness	0.286	1.008	0.993	1.023
Age	0.123	1.002	1.000	1.004
Gender				
Male	< 0.001	1.223*	1.164	1.285
Female (referent)				
Marital status				
Married/living together	< 0.001	1.321*	1.232	1.415
Divorced/separated	0.029	1.141*	1.013	1.284
Never married (referent)				
Education				
Not attending school	< 0.001	1.261*	1.112	1.431
Attending school (referent)				
Occupation				
Unemployed	< 0.001	0.700*	0.661	0.742
Self-employed	< 0.001	1.183*	1.117	1.252
Worker (referent)				

^{*}Significant at p < 0.05 using binary logistic regression test

TABLE 4 Multivariate analysis between Big Five Personality Traits with obesity and control variables.

obesity and control vari	obesity and control variables.				
Variables	Obesity				
	128 P-Value	OR	95% CI		
			Lower	Upper	
Conscientiousness	0.002	1.025*	1.009	1.041	
Age	< 0.001	1.011*	1.009	1.013	
Gender					
Male	< 0.001	0.429*	0.406	0.453	
Female (referent)					
Marital status					
Married/living together	< 0.001	2.500*	2.291	2.729	
Divorced/separated	< 0.001	1.593*	1.392	1.824	
Never married (referent)					
Education					
Not attending school	< 0.001	0.461*	0.400	0.532	
Attending school (referent)	-0.001	0.101	0.100	0.002	
Occupation					
Unemployed	0.004	0.911*	0.854	0.971	
Self-employed	0.272	1.035	0.974	1.099	
Worker (referent)	0.2/2	1.055	0.974	1.099	
		1.079*		1.092	
Extraversion	< 0.001		1.065		
Age	< 0.001	1.012*	1.010	1.014	
Gender					
Male	< 0.001	0.444*	0.420	0.469	
Female (referent)					
Marital status					
Married/living together	< 0.001	2.546*	2.333	2.779	
Divorced/separated	< 0.001	1.637*	1.429	1.875	
Never married (referent)					
Education					
Not attending school	< 0.001	0.480*	0.416	0.553	
Attending school (referent)					
Occupation					
Unemployed	0.008	0.916*	0.859	0.977	
Self-employed	0.468	1.023	0.962	1.087	
Worker (referent)					
Neuroticism	< 0.001	0.966*	0.953	0.978	
Age	< 0.001	1.011*	1.009	1.013	
Gender					
Male	< 0.001	0.421*	0.398	0.445	
Female (referent)					
Marital status					
Married/living together	< 0.001	2.519*	2.308	2.748	
Divorced/separated	< 0.001	1.601*	1.398	1.833	
Never married (referent)					
Education					
Not attending school	< 0.001	0.459*	0.399	0.529	
Attending school (referent)					
Occupation					
Unemployed	0.004	0.910*	0.853	0.970	
Self-employed	0.302	1.033	0.972	1.097	
		******	2200		

Significant at p < 0.05 using binary logistic regression test.

Worker (referent)

Discussion

Big five personality traits and eating habits

Eating habits were found related to personality as so personality was assessed as contributing to certain eating patterns (Möttus et al., 2013; Sutin and Terracciano, 2015b). Based on data on the consumption 347 types of food (including five types of non-recommended toods: instant numbers, fast food, soft drink, fried snacks, and sweet snacks) and personality based 121 e Big Five Personality Traits, also after controlling for socio-demographic factors (age, gender, education, occupation, and marital status), the following results were obtained. This study showed that rice was not associated with all dimensions of personality, which could be because the staple food of the Indonesians is rice and very few people do not eat it.

This study found openness was positively associated (especially in adults), both with recommended and non-recommended foods, which was positively associated with eating all food types tested, and the most likely positive association was with green leafy vegetables and the lowest was with fried snacks only a slight difference. This may indicate the nature of openness to accept 36 various types of food, both healthy and unhealthy. This study also found a positive association between openness and fried snacks in men. This might be because men tended to have higher ope 106 scores than women (Magee and Heaven, 2011; Armon et al., 2013; Intiful et al., 2019).

In previous studies, openness was associated with a Mediterranean-style diet and avoidance of sweet-based forms of diet (Mõttus et al., 2013), moreover, openness was found to be the most consistent and stroigs t result when associated with eating healthy foods (Mõttus et al., 2013; Tiainen et al., 2013). Openness was positively associated with eating plant-bassal food and fish (Pfeiler and Egloff, 2020), fruits and vegetables (Tiainen et al., 2013; Keller and Siegrist, 201729 onner et al., 2017), and was negatively associated with eating meat (Keller and Siegrist, 2015; Pfeiler and Egloff, 2020), sugary drinks (Keller and Siegrist, 2015), and potato chips (Conner et al., 2017), however, other studies showed that openness was associated with increased consumption of wine (Gustavsen and Rickertsen, 2019) and non-recommended products (Gacek et al., 2021). The different results can be explained by the characteristics of openness that are more open to new experiences, as well as greater curiosity and exploration, making them more open to change so that they are more open to habits that are considered sustainable, for example, it is easier to adopt new adaptive eating patterns and overcome taste aversions (Mõttus et al., 2013; Conner et al., 2017), including consuming healthier foods such as fruits and vegetables, t if healthy eating habit has been habituated or perceived to be more familiar, it may no longer match the characteristics of higher openness

(Mōttus et al., 2013). The different results were also possible due to the variation in the characteristics of the studied sample which may lead to different behavioral tendencies considering that openness may adopt healthy or unhealthy eating behaviors.

This study found that conscientiousness was positively associated with eating recommended foods (especially in adults and except mango, which was not associated), with the highest probability of consuming green leafy vegetables (did not vary by sex or age group). Conscientiousness was not associated with all non-recommended foods: fast food, soft did, instant noodle, fried snacks, and sweet snacks. This study also found a negative association between conscientiousness and meat in adolescents and soft drinks in women and a positive association between dairy and fast food in men.

Consistent with the previous studies, conscientiousness tends to choose healthy foods and foods recornsended in dietary guidelines (Keller and Siegrist, 2015), as in the study by Mõttus et al. (2013) higher conscientiousness was associated with a tendency to have a healthier diet. In the study by Gacek et al. (2021), there was an increase in eating most recommended products and a decrease in eating most nonrecommended products, and 1332e study by Ibigbami (2012), conscientiousness was found to be negatively associated with alcohol consumption, which was similar to previous studies, where it seemed that conscientiousness had a holistic effect pnly on alcohol use but on all aspects of their wellbeing. Conscientiousness was also found to be positively associated with eating plant-based food and fish (Pfeiler and Egloff, 2020), fruits (Keller and Siegrist, 2015), and the combined daily consumption of vegetables and fruits compared to lower conscientiousness (Conner et al., 2017). Conscientiousness was negatively associated with eating meat (Keller and Siegrist, 2015; Pfeiler and Egloff, 2020), sugary drinks, sweet and savory foods (Keller and Siegrist, 2015), and foods containing preservatives (among young women) (Jaworski and Rozenek, 316). Additionally, increasing conscientiousness also reduces he expected frequency of beer consumption (Gustavsen and Rickertsen, 2019). Meanwhile, lower conscientiousness tends to perceive their eating habits as more unhealthy and have poor eating habits (Lawler, 2018). This is due to conscientiousness tends to be more careful in choosing the food consumed (Keller and Siegrist, 2015; Nelvi, 2016). Furthermore, it is also due to the level of discipline and orderliness that affects adherence to healthy diets (such as Mediterranean diet) and avoidance of unhealthy foods (such as sweet foods) (Mõttus et al., 2013).

In this study, extraversion was positively associated with eating all types of food tested (except fish, which was not associated), which meant a positive association with eating both recommended and non-recommended foods (especially higher in women than men), with the highest probability of consuming fast food. This can be supported by previous studies, women scored higher on extraversion than men (Magee and Heaven, 2011; Armon et al., 2013; Purnamawati and Yuliati, 2016).

Previous studies showed that extraversion was negatively associated with eating carbohydrate-based food (Pfeiler and Egloff, 2020) and was positively associated with eating meat (Keller and Siegrist, 2015; Pfeiler and Egloff, 2020), plant-based food, and fish (Pfeiler and Egloff, 2020), fruits and vegetables (Purnamawati and Yuliati, 2016; Conner et al., 2017), wholegrain bread (Jaworski and Rozenek, 2016), sugary drinks, sweet and savory foods (Keller and Siegrist, 2015), alcohol (Ibigbami, 2012; Kye and Park, 2012), wine (Gustavsen and Rickertsen, 2019), and Mediterranean diet (Mõttus et al., 2013). Other studies on young women also found that individuals with higher extraversion tend to choose foods that are considered to ave a good impact on health (Jaworski and Rozenek, 2016). Based on those results, it can be seen that extraversion tends to increase consumption of both recommended and non-recommended foods. This was also seen in the study by Gacek et al. (2021) who suggested that more extroverted individuals tend to be more assertive, optimistic, open, sensation-seeking, and happy to socialize may increase consumption of most recommended products, but consumption of non-recommended products decreased with red meat and lard, while increased with sweets and alcohol. This may be due to extraversion who prefer to socialize and is more attached to social networks (Friedman et al., 2010) so they need energy from gathering with other people. Therefore, they regularly attend social situations and eat more often with other people, but the food selection is limited to the menu provided (Tiainen et al., 2013) so it is possible to frequently consume non-recommended foods which can have a negative impact on health (Keller and Siegrist, 2015). In this study, perhaps the biggest tendency was meeting other people in restaurants selling fast food, as this study found that the highest probability was fast food consumption.

In this study, agreeableness was positively associated with eating several recommended foods: fish, fruits (banana, papaya, mango, especially in adults), and most likely green leafy vegetables (especially in adults), and negatively associated with eating non-recommended foods: soft drink (especially in adolescents and adults), fried snacks in men and adults, and sweet gricks in adults. The rest of the food types were unrelated. The results of this study were in line wi 43 revious studies, higher agreeableness was associated with a healthier diet (Mõttus et al., 2013). In previous studies, agreeableness was found to be positively associated with eating plantbased food and fish (Pfeiler and Egloff, 2020) and fruit (in women) (Tiainen et al., 2013). Agreeableness was negatively associated with eating carbohydrate-based food (P31 r and Egloff, 2020), meat (Keller and Siegrist, 2015), and frequency of wine and beer consumption (Gustavsen and Rickertsen, 2019). However, other studies showed that the association between agreeableness and eating habits appears to be rather weak and inconsistent across studies (Pfeiler and Egloff, 2020).

Such agreeableness was found to be associated with increased consumption of non-recommended products (Gacek et al., 2021), and infrequent avoidance of salty foods (among young women) (Jaworski and Rozenek, 2016), and poor or unhealthy diet patterns (Kye and Park, 2028). It may be that agreeableness consumes less meat because ethical concerns based on high altruism and sympathy also apply to animals (Keller and Siegrist, 2015), or agreeableness was significantly negatively associated with alcohol consumption because individuals with high agreeableness are quite sociable and naturally have good and respectful natures, so they are easy to adopt other forms of coping strategies that could reduce the tendency to use or abuse alcohol (Ibigbami, 2012), or other traits such as obedience make them more likely to adhere to rules such as dietary guidelines, but other traits that tend to be easy to trust and have a high tolerance of others, happy to help, kind and gentle, make them easily influenced by their social environment which may force them to follow unhealthy dietary practices (Rahmadian, 2011; Kye and Park, 2012). These might contradict and confuse them, but as adults, they might begin to realize the benefits of dietary guidelines, therefore in this study the healthy eating habit was found particularly in higher agreeableness adults.

This study found that neuroticism was positively associated with eating almost all non-recommended foods (especially in adults): instant noodles, soft drinks, sweet snacks, and the highest probability was eating fried snacks (fast food was unrelated). Neuroticism was also positively associated with soft drinks and sweet snacks in women and fried snacks in men. In addition, neuroticism was negatively associated with eating several recommended foods (especially in adults): meat, dairy, carrot, and the highest probability of green leafy vegetable consumption, the rest of the food types were unrelated.

Consistent with this finding, neuroticism was found to be associated with poorer diet quality (Tiainen et al., 2013). In previous studies, neuroticism was positively associated with eating carbohydrate-based food and meat (Pfeiler and Egloff, 2020), sweet and savory foods (Keller and Siegrist, 2015), soft drinks (for women) (Tiainen et al., 2013), French fries (for men) (Conner et al., 2017), and non-recommended products (including sweets) (Gacek et al., 2021). Neuroticism was negatively associated with plant-based food and fish (Pfeiler and Egloff, 2020), vegetables (in women) (Tiainen et al., 2013), recommended whole grain cereals (Gacek et al., 2021), and the Mediterranean diet (Mõttus et al., 2013) which dietary recommendations encourage eating healthy foods. This can happen because individuals with high neuroticism have low selfesteem, high sensitivity, and are emotionally unstable (Gacek et al., 2021), which causes their mood to fluctuate often, also easily feel upset and disturbed, which is then overcome by eating because they react to eating behavior as a form of reassurance over negative emotions (Elfhag and Morey, 2008; Nelvi, 2016), thereby ignoring healthy food choices. It can be worse, especially in productive age or adulthood, where stress levels are higher and they are more prone to depression. Meanwhile, individuals with emotional stability (low levels of neuroticism) tend to choose foods considered to have a good impact on health (Jaworski and Rozenek, 2016).

Big five personality traits and physical activity

In this study, only openness, conscientiousness, and agreeableness were associated with physical activity. However, after controlling for age, gen 224 education, occupation, and marital status, only openness and conscientiousness remained associated with physical activity, but 42 e did not seem to moderate this association. Meanwhile, the association between agreeableness and physical activity was weakened by these sociodemographic factors so that it no longer had a statistically significant association.

This study found textraversion was not associated and conscientiousness was positively associated with physical activity. Street differences were found in previous findings that high extravers and conscientiousness (as well as low neuroticism) were found to be associated with higher or more active levels 83 physical activity (Rhodes, 2006; Rhodes and Smith, 2006; Allen and Laborde, 2014; Stephan 411, 2014b; Mõttus et al., 2016). Individuals with higher extraversion and conscientiousness tend to have more energy (Terracciano et al., 2013) and an active nature is theorized as a key aspect of extraversion and conscientiousness that represents their tendency to be busy and energetic (Rhodes, 2006). Thus, physical activity can be used as an activity in accordance with the energy pacity they need (Stephan et al., 2014b). Individuals with low extraversion may be less likely to engage in physical activity because they find it less enjoyable and have a lower level of control over it than people with higher extraversion (Rhodes, 2006). Meanwhile, individual 18 who lack order and self-discipline (low conscientiousness) are likely to have more difficulty maintaining their physical activity plans (Rhodes, 2006). Extraversion and conscientiousness also largely have a positive association with activity variability within individuals (Mõttus et al., 2016). The difference in results on extraversion found in this study may be due to the fact that Indonesians have very low levels of physical activity compared to other countries (Althoff et al., 2017), so although individuals with high extraversion have more energy and like to gather with other people, this is directed toward activities other than physical activity and their social environment also do not support physical activity due to the 199 lentary habits of the community. Therefore, extraversion was not associated with physical activity in this study.

This study also found that individuals with higher conscientiousness in adolescents and adults tended to be more

active, but there was no association in the elderly. This might be because age was positively correlated with a decreased physical activity where children were more active than adults (Miles, 2007) and physical activity steadily decreased in the later stages of life (Wilson and Dishman, 2015). However, age was also related to personality such as conscientiousness which tended to increase with age (Brummett et al., 2006). These the paps became very contradictory in old age, thus negating the relationship between conscientiousness and physic tivity in the elderly.

This study found that openness was positively associated with physical activity (especially in women and the elderly), which was consistent with previous findings. Stephan et al. (2014a) showed that not only extraversion, but openness was also associated with the possibility of being involved in variation physical, cognitive, and social activities, and also found a consistent pattern in two different Western societies (United States and France), that there were intrinsic tendencies toward extraversion and openness to having a more active lifestyle. These characteristics are perhaps more dominant in women and older people, as they tend to prefer to gather, talk, and do activities together, including physical activities. Openness characteristics which have a tendency to new ideas and experiences and also prefer variety are characteristics of individuals who are open and can lead to active involvement in various types of activities (Stephan et al., 2014a) thus enabling individuals to have a higher level of physical activity because physical activity can encourage a tendency to explore and seek new and varied experiences (Stephan et al., 2014b).

In this study, agreeableness was found to be unrelated to physical activity. The result was consistent with other studies that also found no association between agreeableness and physical activity, perhaps suggesting that the tendency to trust and help others is unrelated to motivation and involve that in physical activity (Courneya and Hellsten, 1998; Sutin et al., 2016; Intiful et al., 2019). However, in the study by Stephan et al. (2014a), agreeableness showed a positive association with variations in the type of activity in one of the samples tested. Physical activity also encourages a tendency to be exposed to various bodily, 11 otional, and social experiences which can also produce social interactions that may contribute to the maintenance of a prosaic orientation that characterizes agreeableness (Stephan et al., 2014b).

In this study, neuroticism was unrelated to physical activity, but in previous studies, neuroticism was negatively associated with physical activity in one of the samples tested in the studie 130 Stephan et al. (2014a,b). Overall, neuroticism seems to be negatively associated with physical activity but has little impact (Rhodes and Smith, 2006). Theoretically, neuroticism with a tendency to feel insecure and jealous may look to others for motivation to behave (Rhodes, 2006) for example in physical activity. Feelings of worry about appearance, feelings of obligation, and feelings of guilt if they do not exercise become central to their motivation for physical activity.

(Ingledew and Markland, 2008), but due to their low self-control which is reflected by impulsivity (Elfhag and Morey, 2008), it is difficult for them to maintain the intention and schedule in exercising. This may have a contradictory impact on neuroticism and therefore neuroticism was unrelated to physical activity.

The results of this study were slightly different from previous studies. This may be due to variations and differences in the characteristics of the respondents studied since a study related to the association between Big Five Personality Traits and physical activity has never been studied before in Indonesians.

Big five personality traits and obesity

In this study, both before and after being controlled by socio-demographic factors, conscientiousness, extraversion, and neuroticism were associated with obesity, it seemed that all control variables moderated this associated working as self-employed. Meanwhile, openness and agreeableness were not associated with obesity.

The positive association between conscientiousness and obesity (especially in women and adults) found in this st was not unexpected. Compared to other dimensions, low conscientiousness and high neuroticism had the strongest associations v92 BMI and obesity (Sutin and Terracciano, 2015b). But based on a meta-analysis by Jokela et al. (2013), only conscientiousness was consistently associated with obesity. Inconsistent results were found for other dimensions in previous studies. Previous studies showed that conscientiousness was negatively associated with nutritional status, which meant that low conscientiousness was associated with the gher nutritional status (obesity) in adult samples (Magee and Heaven, 2011; Sutin et al., 2011; Sutin and Terracciano, 2015b; Pfeiler and Egloff, 2020) and pler people (Mõttus et al., 2013). Conscientiousness was also found to be negatively associated wi43 several measures of body weight (Armon et al., 2013). Conscientiousness was associated with healthier body weight, possibly because high conscientiousness was characterized as conscientious, efficient, orderly, obedient, organized, and disciplined individuals, so they were more likely to have behaviors that support a healthy weight or maintain the ed body shape, such as eating habits, following a 56 al plan, eating at the same time every day, and exercising (Brummett et al., 2006; Terracciano et al., 2009; Sutin et al., 2011; Sutin and Terracciano, 2015b). Meanwhile, low conscientiousness had difficulties in self-control, so they tend to overeat, lack physical activity, and had poor eating habits such as chaotic eating schedules, which can then lead to obesity (Magee and Heaven, 2011; Sutin and Terracciano, 2015 34 Although conscientiousness was associated with healthier eating habits and more physically active found in this study, the different results in association with nutritional status may be due

to other mechanisms linking this association. For example, an imbalance in total energy intake and energy expenditure that was not investigated in this study or other possibilities due to physiological factors such as health conditions or hist 108 of disease. Moreover, several previous studies found an association between personality dimensions and several physiological and genetic mechanisms that can then affect body composition (Kakizaki et al., 2008), such a circulating levels of 110 p (Sutin et al., 2013) and FTO genetic variant had little effect on the relationship between personality traits with underweight and overweight (Terracciano et al., 2009). Therefore, future studies may further investigate other mechanisms that might under this association.

This study found that neuroticism was negatively associated with obesity (especially in men and adults). However, previous ge lies found inconsistent findings such as a negative association between neuroticism and nutritional status (Kakizaki et al., 2008) as well as a positive association (Sutin et al., 2011; Pfeiler and Egloff, 2020). Individuals with low conscientiousness or high neuroticism were prone to obesity (Sutin et al., 2011). High neuroticism was characterized by individuals who tend to have 101 tional vulnerability, fear, anxiety, and depression related to eating behavior for comfort due to negative emotions, poor eating habits, avoidance of physical activity, and impulsiveness associated with emotional eating can lead to obesity (Elfhag and Morey, 2008; Magee and Heaven, 2011). This can be seen in low self-discipline (conscientiousness) and impulsiveness (neuroticism) which reflect poor individual self-control (Elfhag and Morey, 2008). Moreover, this impulsivity trait was the strongest predictor of tritional status which was found in many studies (Terracciano et al., 20125 However, Terracciano et al. (2009) found that lean people scored higher on neuroticism, but scored lower on impulsivity. The inconsistent results on neuroticism and obesity might be explained by the nature of their susceptibility to depression, especially in productive age or adulthood. Depression can trigger individuals to eat more or less, so it was strongly associated with emotion and nutritional status (Yilmaz and Köse, 2020). This can lead individuals to have 17 more or less nutritional status. As in this study, although neuroticism was found to be positively associated with eating non-recommended foods and negatively associated with eating recommended foods, but neuroticism was negatively associated with obesity, this might be related to their susceptibility to depression so that it affected the nutritional status tended to be not obese.

In this study, extraversion was positively associated with sesity (especially in adolescents, adults, and higher in women). This finding was consistent with other studies that found a positive associated with other studies that found a positive associated between extraversion and nutritional status (Magee and Heaven, 2011; Sutin et al., 2011), but another study found to extraversion was negatively associated with BMI among women but not among men (Sutin and Terracciano,

2015b). Brummett et al. (2006) showed that men with higher extraversion had a higher BMI explained by extraversion traits that are adventurous, fun-seeking, sociable, and gregarious, which may lead them to consume more high-calorie foods and/or consume more alcohol. Armon et al. (2013) offered an alternative explanation that the tendency of individuals with high positive moods to be more careless, feel less vulnerable to unwanted health conditions, and adopt unhealthy behaviors that lead to overweight. Although in this study extraversion was associated with eating recommended and non-recommended foods (especially higher in women), which reflected their lack of attention to the food they consumed, with then may have an impact on obesity, in another study, extraversion was associated with greater physical activity (Hoyt et al., 2009), which means they actually have the potential to be able to maintain a healthy weight.

Our findings could be supported by findings from previous studies that obesity affects British and Indonesian women more than men (Mihardja and Soetrisno, 2012), and women tended to consume more sweet foods such as candy and chocolate (Conner et al., 2017). Another study also found that men had higher quality food intake and eating habits than women (Puspadewi, 2014), but other studies found the opposite (Möttus et al., 2013). In addition, the increase in BMI occurs mainly in young and middle adults, then after passing the age of 60 years, body weight decreases rapidly and returns to a value comparable to that of young adults (Sutin et al., 2011; Suiraoka, 2012; Chaudhuri and Bandopadhyay, 2014).

This study found no association between obesity with openness and agreeableness. Previous studies showed inconsistent findings, which suggested that nutritional status was unrelated to agreeableness (Pfeiler and Egloff, 2020) and openness (Sutin 21 Terracciano, 2015a; Pfeiler and Egloff, 2020), positively associated with openness (Sutin et al., 2011), also negatively (Sutin et al., 2011; Sutin and Terracciano, 2015a) and positively associated with agreeableness (Magee and Heaven, 2011). Openness was characterized by an open and curios nature may be easier to adopt a new adaptive eating pattern, but pshaps if the eating habit has been habituated or perceived to be more familiar, it may no longer match the characteristics of higher openness (Mõttus et al., 2013). Meanwhile, agreeableness was characterized by individuals who are warm, empathetic and need the consideration of others, which may lead to contradictory eating behaviors that can then affect body weight (Armon et al., 2013). This may be what causes the different findings in various studies or even the lack of association between nutritional status with openness

Differences in results with previous studies may be due to variations in the sample tested and methodological differences such as the composition of the Big Five Personality Trait questions used differently from previous studies. Due to inconsistent findings across many studies, it is difficult to

draw conclusions. Thus, further study is needed to describe a more specific mechanism regarding the association between personality and nutritional status.

What can we learn?

In this study, openness was found to be positively associated with both recommended 57 non-recommended foods (especially in adults), positively associated with physical activity (especially in women and the elderly), but not associated with obesity. This can be explained by the fact that even though they were more active, the nature of individuals with high openness tend to like variety and be open to new experiences and ideas, which allows them to more easily follow new healthy or unhealthy dietary guidelines, which leads to openness did not determine nutritional status. Therefore, individuals with high openness should be directed to adaptive behavior patterns that come with a lot of variety so that they can meet their needs for variety and their interest in new experiences, such as food menus or healthy types of diets that regularly changed various physical activities. Meanwhile, low openness maybe enjoys a more fixed schedule, not too varied, and a selection of menus and activities that are more familiar to them due to their more conservative and conventional tendencies.

Conscientiousness was positively associated with eating almost all recommended foods (especially in adults) and not associated with all non-recording ended foods in this study. Moreover, conscientiousness was positively associated with physical activity (especially in adolescents and adults), but also positively associated with obesity (especially in women and adults), which was an unexpected result. This may be because in this study conscientiousness was not associated with eating non-recommended foods, so they could have high or low levels of consumption, although conscientiousness increased the possibility of consuming recommended foods and physical activity, which then impacted obesity, or there may be another mechanism underlying this relationship (such as their knowledge of healthy foods which was not examined in this study). Therefore, individuals with high conscientiousness can be directed to adaptive behavior with clear and organized guidelines and schedules so that it will be easier to adopt because it is in line with their conscientious, disciplined, organized, and obedient nature. Meanwhile, low conscientiousness may need behavioral strategies, implementation intentions, and behavior change goals such as increasing physical activity (Rhodes, 2006).

This study found that extraversion was positively associated with eating recommended and notification (especially higher in women), was not associated with physical activity and was positively associated with obesity (especially in adolescents, adults, and higher in women). This was possible because food choices are disregarded by individuals with high

extraversion due to their tendency to like to gather and even eat with other people more often, which may limit their food choices (Tiainen et al., 2013), in addition to being more careless and inattentive their attention to unwanted health conditions (Armon et al., 2013) which then has an impact on obesity. Therefore, individuals with high extraversion may be more suited to group activities and activities that require a lot of energy as it suits their sociable and energetic nature to make it easier for them to adopt adaptive behavior. Meanwhile, low extraversion may be able to take advantage of behavior change strategies that do not require much energy and do not involve many people, such as exercising at home.

Agreeableness was positively associated with eating several recommended foods and negatively associated with eating non-recommended foods (especially in adults). However, agreeableness was unrelated to physical activity and obesity. This may be because the nature of individuals with high agreeableness who like to help and easily trust others has no effect on their motivation and involvement physical activities (Courneya and Hellsten, 1998; Sutin et al., 2016; Intiful et al., 2019). In addition, their tendency to obey and trust others leads them to be easily influenced by their social environment which can lead them to adopt healthy or unhealthy dietary guidelines depending on the individual who directs them or individuals they trust, causing contradictory eating behaviors and agreeableness did not determine nutritional status. Therefore, individuals with high agreeableness can be directed to adaptive behavior by involving the people around them, especially those they trust, so that they can adopt the behavior more easily. Meanwhile, low agreeableness may benefit from behavior change strategies related to independence and avoid the involvement of others, as they tend to be suspicious and critical of others, such as preparing their wn healthy meals.

This study found that neuroticism was positively associated with eating four of the five non-recommended foods and negatively associated with four recommended foods (especially in adults), and the rest was unrelated. Moreover, neuroticism was found to be unrelated to physical activity and negatively associated with respectively in men and adults). This may be because individuals with high neuroticism tend to be unstable, emotional, and impulsive, which indicates their low self-control, so they find it harder to maintain the intention to do physical activity, but in addition, they may engage in physical activity because they feel insecure or see others as motivation to exercise (Rhodes, 2006), so neuroticism was unrelated to physical activity in this study. Additionally, individuals with high neuroticism susceptibility to stress may cause them to tend to have low intakes in this study which might protect them from obesity even though they had poor quality food choices. However, if poor ring habits are taken frequently and continuously, it will have a negative impact on their nutritional and health status. Therefore, individuals with higher neuroticism may be able to involve friends or family with lower

neuroticism in making behavioral changes, such as exercising regularly (Rhodes, 2006), and can benefit from menu planning and regular meal times.

91 Strengths and limitations

The strengths of this study included a large study with over 30,000 respondents so that the results can be more representative of the study population. Our statistical analysis also included control variables to obtain the independent effect of personality on nution, related variables. In addition, since evidence related to the association between personality traits, eating habits, physical activity, and nutritional status was scarcely found in peer-reviewed articles in the Indonesian setting, this study is, therefore, a relatively new and pioneering study linking personality and nutrition in an Indonesian setting. Limitations included limited data quality control and the availability of personality and nutrition-related variables, as we used secondary data. Not all samples were used in this study due to data completeness, but among the excluded respondents, there was no indication that they had a BMI that tended to be higher or lower, so there was no systematic bias. The physical activity duration data processing also used an approach method because IFLS5 had only time frame data, which might lead to overestimation or underestimation. Another limitation, measuri 80 obesity using BMI has a weakness because it is unable to distinguish between body fat and fat-free mass (Johansson et al., 2009). The absence of body composition in assessing obesity leads to a risk of bias, therefore more precise measurements of obesity are recommended, such as fat mass and waist of 53 m ference (Johansson et al., 2009). Lastly, this study was a cross-sectional study, so it was not possible to draw conclusions about causality.

Conclusion

After controlling for socio-demographic variables, openness was positively associated with recommended and nonrecommended foods, and physical activity, and not associated with obesity. Conscientiousness was positively associated with almost all recommended food types, physical activity, and obesity, and not associated with all non-recommended foods. Extraversion was positively associated with recommended and non-rece79 mended foods and obesity, and not associated physical activity. Agreeableness was positively associated with several recommended foods, atively associated with one non-recommended food, and not associated with physical activity and obesity. Neuroticism was positively associated with almost all non-recommended foods, negatively associated with four recommended foods and obesity, was also not associated with physical activity. Behavior change can take into account individual characteristics such as personalized strategies with personality because the adaptation of personality characteristics can be modified even though the personality tends to be persistent. This strategy can be used to improve nutritional status or public health because personality, particularly conscientiousness, was found to be consistent with previous studies across western and eastern countries that conscientiousness had a relationship with eating habits and physical activity. Additionally, higher extraversion (especially in women) was found to be more likely to be careless in food choices and more prone to obesity. Therefore, the understanding of personality needs to be improved so that health and nutrition improvement interventions are more effective and allow for sustainability as the interventions or programs are most suited to the needs of individual preferences. Programs to prevent obesity, improve physical activity, and healthy eating habits should be tailored to accommodate individuals with low conscientiousness personalities especially women with higher extraversion. This information can be useful not only to health professionals, but also to teachers, psychologists, and the community who target the promotion of healthy living or the prevention of nutrition and health problems.

62 Data availability statement

Publicly available datasets were analyzed in this study. This data can be found here: https://www.rand.org/well-being/social-and-behavioral-policy/data/FLS/iFLS/ifls5.html

Author contributions

GP and TM conceived the project and research question.
GP conducted the data analyses and wrote the original draft of the manuscript. TM and MR reviewed and edited to manuscript. TM was responsible for funding the publication of the manuscript. All authors contributed to data processing and read and agreed to the published version of the manuscript.

Funding

This article publication was funded by the Riset Kolaborasi Indonesia–WCU (World Class University), 2021, number: 154/UN3.15/LT/2021.

Acknowledgments

This article was derived from GP's undergraduate the grateful to all supervisors for guiding this article. We are grateful to RAND which have provided the Indonesia Family Life Survey (IFLS) dataset.

1 Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All daims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: https://www.frontiersin.org/articles/10.3389/fpsyg,2022.881436/full#supplementary-material

References

Ahdiyana, M. (2011). Perilaku Organisasi. Yogyakarta: Yogyakarta State University.

Allen, M. S., and Laborde, S. (2014). The role of personality in sport and physical activity. Curr. Direct. Psychol. Sci. 23, 460–465. doi: 10.1177/0963721414550705

Althoff, T., Sosiè, R., Hicks, J. L., King, A. C., Delp, S. L., and Leskovec, J. (2017). Large-scale physical activity data reveal worldwide activity inequality. *Nature* 547, 336–339. doi: 10.1038/nature23018

Armon, G., Melamed, S., Shirom, A., Shapira, I., and Berliner, S. (2013). Personality traits and body weight measures: concurrent and across-time associations. *Eur. J. Pers.* 27, 398–408. doi:10.1002/per.1902

Brummett, B. H., Babyak, M. A., Williams, R. B., Barefoot, J. C., Costa, P. T., and Siegler, I. C. (2006). NEO personality domains and gender predict levels and trends in body mass index over 14 years during midlife. *J. Res. Pers.* 40, 222–236. doi: 10.1016/j.jrp.2004.12.002

Chaudhuri, A., and Bandopadhyay, A. K. (2014). Obesity, a burning problem. IOSR J. Pharm. 3, 1–5. doi: 10.9790/3013-04101-05

Conner, T. S., Thompson, L. M., Knight, R. L., Flett, J. A. M., Richardson, A. C., and Brookie, K. L. (2017). The role of personality traits in young adult fruit and vegetable consumption. Front. Psychol. 8:119. doi: 10.3389/fpsyg.2017.00119

Costa, P. T. and McCrae, R. R. (1985). The NEO Personality Inventory Manual. Odessa, FL: Psychological Assessment Resources.

Courneya, K. S., and Hellsten, L. A. (1998). Personality correlates of exercise behavior, motives, barriers and preferences: an application of the five-factor model. Pers. Individ. Differ. 24, 625–633. doi: 10.1016/S0191-8869(97)00231-6

Dehne, M., and Schupp, J. (2007). Persönlichkeitsmerkmale im Soziooekonomischen Panel (SOEP) – Konzept, Umsetzung und empirische Eigenschaften. DIW Research Note 26. Berlin: DIW Berlin.

Elfhag, K., and Morey, L. C. (2008). Personality traits and eating behavior in the obese: poor self-control in emotional and external eating but personality assets in restrained eating. *Eat. Behav.* 9, 285–293. doi: 10.1016/j.eatbeh.2007.10.003

Feist, J., Feist, G. J., and Roberts, T.-A. (2013). Theories of Personality, 8th Edn. New York, NY: McGraw-Hill.

Frankenberg, E., and Karoly, L. (1995). The 1993 Indonesian Family Life Survey: Overview and Field Report. Santa Monica: RAND. Available online at: https://www.rand.org/well-being/social-and-behavioralpolicy/data/FLS/IFLS/download.html (accessed January 14, 2021).

Friedman, H. S., Kern, M. L., and Reynolds, C. A. (2010). Personality and health, subjective well-being, and longevity. *J. Pers.* 78, 179–216. doi: 10.1111/j.1467-6494. 2009.00613.x

Gacek, M., Kosiba, G., and Wojtowicz, A. (2021). Personality determinants of diet quality among polish and spanish physical education students. *Int. J. Environ. Res. Public Health* 18, 466. doi: 10.3390/ijerph18020466

Gustavsen, G. W., and Rickertsen, K. (2019). Personality traits and consumption of wine and beer. J. Wine Econ. 14, 392–399. doi: 10.1017/jwe.2019.34

Hoyt, A. L., Rhodes, R. E., Hausenblas, H. A., and Giacobbi, P. R. (2009). Integrating five-factor model facet-level traits with the theory of planned behavior and exercise. *Psychol. Sport Exerc.* 10, 565–572. doi:10.1016/j.psychsport.2009.02.009

Ibigbami, O. I. (2012). The Relationship between Personality Traits, Alcohol and Risky Sexual Behaviour in An Undergraduate Student Population in South-Western Nigeria. Thesis. If: Obsemi Awolowo University.

Indonesian Ministry of Health [IMH] (2014). Regulation of the Indonesian Ministry of Health number 41 of 2014 regarding The Balanced Nutrition Guidelines. Jakarta: Indonesian Ministry of Health.

Indonesian Ministry of Health [IMH] (2017). Penilaian Status Gizi. Available online at: http://bppsdmk.kem.kes.go.id/pusdiksdmk/wp-content/uploads/2017/11/PENILAIAN-STATUS-GIZI-FINAL-SC.pdf (accessed September 17, 2020).

Indonesian Ministry of Health [IMH] (2019). Peneliti: Pola Makan Buruk/Menu Tidak Sehat Penyebab Satu dari Lima Kematian. Available online at: http://p2ptm.kemkes.go.id/artikel-sehat/peneliti-pola-makan-burukmenutidak-sehat-penyebab-satu-dari-lima-kematian (accessed January 7, 2020).

Indonesian Ministry of Health [IMH] (2020). Regulation of the Indonesian Ministry of Health number 2 of 2020 Regarding Child Anthropometry Standards. Jakarta: Indonesian Ministry of Health.

Ingledew, D. K., and Markland, D. (2008). The role of motives in exercise participation. *Psychol. Health* 23, 807–828. doi: 10.1080/08870440701405704

Intiful, F. D., Oddam, E. G., Kretchy, I., and Quampah, J. (2019). Exploring the relationship between the big five personality characteristics and dietary habits among students in a Ghanaian University. *BMC Psychol.* 7:10. doi: 10.1186/s40359-019-0286-z

Jaworski, M., and Rozenek, H. (2016). Selected personality traits and frequency of taking some healthy eating behaviours in young women. J. Public Health Nurs. Med. Rescue 191, 42–48.

Johansson, E., Böckerman, P., Kiiskinen, U., and Heliövaara, M. (2009). Obesity and labour market success in finland: the difference between having a high BMI and being fat. *Econ. Hum. Biol.* 7, 36–45. doi: 10.1016/j.ehb.2009.0

John, O. P., Donahue, E. M., and Kentle, R. L. (1991). The Big Five Inventory-Versions 4a and 54. Berkeley, CA: University of California.

John, O. P., Naumann, L. P., and Soto, C. J. (2008). "Paradigm shift to the integrative big five trait taxonomy: history, measurement, and conceptual issues," in *Handbook of Personality: Theory and Research*, eds O. P. John, R. W. Robins, and L. A. Pervin (New York, NY: Guilford Press), 114–158.

John, O. P., and Srivastava, S. (1999). "The big-five trait taxonomy: history, measurement, and theoretical perspectives," in *Handbook of Personality: Theory and Research*, Vol. 2, eds L. A. Pervin and O. P. John (New York: Guilford Press), 102–138.

Jokela, M., Hintsanen, M., Hakulinen, C., Batty, G. D., Nabi, H., Singh-Manoux, A., et al. (2013). Association of personality with the development and persistence of obesity: a meta-analysis based on individual-participant data. *Obesity Rev.* 14, 315–323. doi: 10.1111/obr.12007

Kakizaki, M., Kuriyama, S., Sato, Y., Shimazu, T., Matsuda-Ohmori, K., Nakaya, N., et al. (2008). Personality and body mass index: a cross-sectional analysis from the Miyagi Cohort Study. *J. Psychosomatic Res.* 64, 71–80. doi: 10.1016/j. jpsychores.2007.07.008

Keller, C., and Siegrist, M. (2015). Does personality influence eating styles and food choices?? Direct and indirect effects. *Appetite* 84, 128–138. doi: 10.1016/j. appet.2014.10.003

Kye, S. Y., and Park, K. (2012). Psychosocial factors and health behavior among Korean adults: a cross-sectional study. Asian Pacific J. Cancer Prevent. 13, 49–56. doi: 10.7314/APJCP.2012.13.1.049

Lawler, T. R. (2018). Personality Types and Self-Reported Eating Habits. Honors theses. Ohio: University of Dayton.

Magee, C. A., and Heaven, P. C. L. (2011). Big-five personality factors, obesity and 2-Year weight gain in Australian adults. *J. Res. Pers.* 45, 332–335. doi: 10.1016/j.jrp.2011.02.009

Mihardja, L., and Soetrisno, U. (2012). Prevalence and Determinant Factors for Overweight and Obesity and Degenerative Diseases Among Young Adults in Indonesia. JAFES, Vol. 27. Available online at https://www.asean-endocrinejournal.org/index.php/JAFES/article/view/17/30 (accessed November 20, 2021).

Miles, L. (2007). Physical activity and health. Br. Nutr Foundation Nutr Bull. 32, 314–363. doi: 10.1111/j.1467-3010.2007.00668.x

Mõttus, R., Epskamp, S., and Francis, A. (2016). Within- and between individual variability of personality characteristics and physical exercise. *J. Res. Pers.* 69, 139–148. doi: 10.1016/j.jrp.2016.06.017

Möttus, R., McNeill, G., Jia, X., Craig, L. C. A., Starr, J. M., and Dear, I. J. (2013). The associations between personality, diet and body mass index in older people. Health Psychol. 32, 353–360. doi: 10.1037/a0025537

Nelvi, R. (2016). Hubungan antara Dimensi Kepribadian Big Five dengan Perilaku Makan pada Mahasiswa UIN Suska Riau di Pekanbaru Riau. J. Psikologi 12:47. doi: 10.24014/jp.y1211.3008

Pfeiler, T. M., and Egloff, B. (2020). Personality and eating habits revisited: associations between the big five, food choices, and body mass index in a representative australian sample. *Appetite* 149:104607. doi: 10.1016/j.appet.2020. 104607

Purnamawati, M., and Yuliati, L. N. (2016). Pengaruh Kepribadian dan Situasi terhadap Pemilihan Makanan pada Mahasiswa Tingkat Pertama. *Jur. Ilm. Kel. Kons* 9, 231–242. doi: 10.24156/jikk.2016.9.3.231

Puspadewi, R. H. (2014). Persepsi tentang Pangan Sehat, Pemilihan Pangan dan Kebiasaan Makan Sehat pada Mahasiswa. Undergraduate thesis. Bogor: Institut Pertanian Bogor.

Rahmadian, S. (2011). Faktor-Faktor Psikologis yang Mempengaruhi Perilaku Sehat Mahasiswa Beberapa Penguruan Tinggi di Tangerang Selatan. Undergraduate thesis. Iakarta: UIN Svarif Hidavatullah Iakarta.

Rhodes, R. E. (2006). The built-in environment: the role of personality and physical activity. Exerc Sport Sci. Rev. 34, 83–88. doi: 10.1249/00003677-200604000-00008

Rhodes, R. E., and Smith, N. E. I. (2006). Personality correlates of physical activity: a review and meta-analysis. *Br. J. Sports Med.* 40, 958–965. doi: 10.1136/bism.2006.028860

Riadi, E. (2012). Pengaruh kepribadian, kecerdasan emosional, dan perilaku kewargaan organisasi terhadap kinerja kepala sekolah menengah atas dan kejuruan di kota tanggrang. J. Manajemen Pendidikan 3, 432–441.

Stephan, Y., Sutin, A. R., and Terracciano, A. (2014b). Physical activity and personality development across adulthood and old age: evidence from two longitudinal studies. *J. Res. Pers.* 49, 1–7. doi: 10.1016/j.jrp.2013.

Stephan, Y., Boiché, J., Canada, B., and Terracciano, A. (2014a). Association of personality with physical, social, and mental activities across the lifespan: findings

from US and French samples. Br. J. Psychol. 105, 564-580. doi: 10.1111/bjop. 12056

Strauss, J., Witoclar, F., and Sikoki, B. (2016). The Fifth Wave of the Indonesia Family Life Survey (IFLS5): Overview and Field Report. WR-1143/1-NIA/NICHD. Available online at: https://www.rand.org/well-being/social-and-behavioral-policy/data/FLS/IFLS/download.html (accessed January 14, 2021).

Sugiyono (2011). Metode Penelitian Kuantitatif Kualitatif dan Ré-D. Bandung:

Suiraoka, I. P. (2012). 9 Penyakit Degeneratif dari Perspektif Preventif (Mengenal, Mencegah dan Mengurangi Faktor Resiko 9 Penyakit Degeneratif). Available online at: http://repository.poltekkes-denpasar.ac.id/3290/1/9%20-%20buku% 20penyakit%20degeneratif.pdf (accessed September 30, 2020).

Sutin, A. R., Ferrucci, L., Zonderman, A. B., and Terracciano, A. (2011). Personality and obesity across the adult life span. J. Pers. Soc. Psychol. 101, 579–592. doi: 10.1037/a0024286

Sutin, A. R., Stephan, Y., Luchetti, M., Artese, A., Oshio, A., and Terracciano, A. (2016). The five-factor model of personality and physical inactivity: a meta-analysis of 16 samples. J. Res. Pers. 63, 22–28. doi: 10.1016/j.jrp.2016.0 5001

Sutin, A. R., and Terracciano, A. (2015a). Five-factor model personality traits and the objective and subjective experience of body weight. *J. Pers.* 84, 102–112. doi: 10.1111/jopy.12143

Sutin, A. R., and Terracciano, A. (2015b). Personality traits and body mass index: modifiers and mechanisms. *Psychol. Health* 31, 259–275. doi: 10.1080/08/70446.2015.1082561

Sutin, A. R., Zonderman, A. B., Uda, M., Deiana, B., Taub, D. D., Longo, D. L., et al. (2013). Personality traits and leptin. Psychosomatic Med. 75, 505–509. doi: 10.1097/psy.0b013e3182919ff4

Terracciano, A., Schrack, J. A., Sutin, A. R., Chan, W., Simonsick, E. M., and Ferrucci, L. (2013). Personality, metabolic rate and aerobic capacity. *PLoS One* 8:e54746. doi: 10.1371/journal.pone.0054746

Terracciano, A., Sutin, A. R., McCrae, R. R., Deiana, B., Ferrucci, L., Schlessinger, D., et al. (2009). Facets of personality linked to underweight and overweight. *Psychosomatic Med.* 71, 682–689. doi: 10.1097/PSY.0b013e3181a 2925b

Tiainen, A.-M. K., Männistö, S., Lahti, M., Blomstedt, P. A., Lahti, J., Perälä, M.-M., et al. (2013). Personality and dietary intake – findings in the Helsinki birth cohort study. *PLoS One* 8:e68284. doi: 10.1371/journal.pone.006

Wilson, K. E., and Dishman, R. K. (2015). Personality and physical activity: a systematic review and meta-analysis. *Pers. Individ. Differ.* 72, 230–242. doi: 10.1016/j.paid.2014.08.023

World Health Organization [WHO] (2005). Global Physical Activity Questionnaire (GPAQ) Instrument and Analysis Guide v2. Available online at: https://lh-hsrc.pnu.edu.sa/wp-content/uploads/2019/04/Global-Physical-Activity-Questionnaire-GPAQ.pdf (Accessed February 9, 2021).

World Health Organization [WHO] (2018). Physical Activity for Health. WHO Press. Available online at: https://apps.who.int/gb/ebwha/pdf_files/WHA71/A71_18-en.pdf (Accessed January 2, 2021).

Yilmaz, H. O., and Köse, G. (2020). How does emotional appetite and depression affect BMI and food consumption? *Progr. Nutr.* 22:e2020088. doi: 10.23751/pn. v2214.10750

The relationship between Big Five Personality Traits, eating habits, physical activity, and obesity in Indonesia based on analysis of the 5th wave Indonesia Family Life Survey (2014)

ORIGINALITY REPORT SIMILARITY INDEX **INTERNET SOURCES PUBLICATIONS** STUDENT PAPERS **PRIMARY SOURCES** constellation.uqac.ca Internet Source Maria Gacek, Agnieszka Wojtowicz, Adam Popek. "Personality Determinants of Diet Health Quality among an Elite Group of Polish Team Athletes", International Journal of Environmental Research and Public Health, 2022 **Publication** Submitted to Regis College 1 % Student Paper microdata.worldbank.org 4 Internet Source Submitted to The University of Buckingham <1% 5 Student Paper Submitted to Universitas Airlangga Student Paper

7	Internet Source	<1%
8	Galit Armon, Samuel Melamed, Arie Shirom, Itzhak Shapira, Shlomo Berliner. "Personality Traits and Body Weight Measures: Concurrent and Across–Time Associations", European Journal of Personality, 2020 Publication	<1%
9	Submitted to Mahidol University Student Paper	<1%
10	Submitted to Universiti Putra Malaysia Student Paper	<1%
11	Yannick Stephan, Angelina R. Sutin, Antonio Terracciano. "Physical activity and personality development across adulthood and old age: Evidence from two longitudinal studies", Journal of Research in Personality, 2014 Publication	<1%
12	oamjms.eu Internet Source	<1%
13	epub.uni-regensburg.de Internet Source	<1%
14	academic.oup.com Internet Source	<1%
15	journals.plos.org Internet Source	<1%

16	www.tandfonline.com Internet Source	<1%
17	Sri Intan Zakaria, Sofa Dewi Alfian, Neily Zakiyah. "Determinants of Cardiovascular Diseases in the Elderly Population in Indonesia: Evidence from Population-Based Indonesian Family Life Survey (IFLS)", Vascular Health and Risk Management, 2022	<1%
18	bmcgeriatr.biomedcentral.com Internet Source	<1%
19	www.unica.it Internet Source	<1%
20	Anik R. Patel, Andrew B. Breck, Michael R. Law. "The impact of pharmacy-based immunization services on the likelihood of immunization in the United States", Journal of the American Pharmacists Association, 2018 Publication	<1%
21	"Chapter 301826 PAI", Springer Science and Business Media LLC, 2020 Publication	<1%
22	"Sport and Exercise Psychology", Springer Science and Business Media LLC, 2023 Publication	<1%
23	Sigrid Bjornelv, Hans M. Nordahl, Turid Lingaas Holmen. "Psychological factors and	<1%

weight problems in adolescents. The role of eating problems, emotional problems, and personality traits: the Young-HUNT study", Social Psychiatry and Psychiatric Epidemiology, 2010

Publication

2015
Publication

24	Yannick Stephan, Antonio Terracciano, Martina Luchetti, Damaris Aschwanden et al. "Physical Activity and Sedentary Behavior During COVID-19: Trajectory and Moderation by Personality", Social Psychological and Personality Science, 2020 Publication	<1%
25	eujournal.org Internet Source	<1%
26	Ryan E. Rhodes. "The Built-In Environment: The Role of Personality and Physical Activity", Exercise and Sport Sciences Reviews, 04/2006	<1%
27	www.scielo.br Internet Source	<1%
28	Carmen Keller, Michael Siegrist. "Does personality influence eating styles and food choices? Direct and indirect effects", Appetite,	<1%

29	Sapto Adi, Dian Mawarni, Siti Istiqomah. "The Calculation of The Need for Medical Record Personel Based on The Full-Time Equivalent Method in The Outpatient Registration Department of Public Health Center", Jurnal Kesehatan Prima, 2021	<1%
30	kjfm.or.kr Internet Source	<1%
31	Magalie Dubois, Lara Agnoli, Jean-Marie Cardebat, Raúl Compés et al. "Did Wine Consumption Change During the COVID-19 Lockdown in France, Italy, Spain, and Portugal?", Journal of Wine Economics, 2021 Publication	<1%
32	Toshiaki Aizawa. "Regional Disparity in the BMI distribution of Indonesian People: New Evidence 'Beyond the Mean'", Bulletin of Indonesian Economic Studies, 2018 Publication	<1%
33	Submitted to University College London Student Paper	<1%
34	Yohanes Sondang Kunto, Hilde Bras. "Sibling Inequalities in Overweight and the Role of Mother's Education: Evidence From the Indonesian Family Life Survey", Food and Nutrition Bulletin, 2021 Publication	<1%

35	files.osf.io Internet Source	<1%
36	ro.uow.edu.au Internet Source	<1%
37	Jokela, M., M. Hintsanen, C. Hakulinen, G. D. Batty, H. Nabi, A. Singh-Manoux, and M. Kivimäki. "Association of personality with the development and persistence of obesity: a meta-analysis based on individual-participant data: Personality and obesity", Obesity Reviews, 2013. Publication	<1%
38	Tiia Kekäläinen, Tuija H. Tammelin, Martin S. Hagger, Taru Lintunen et al. "Personality, motivational, and social cognition predictors of leisure-time physical activity", Psychology of Sport and Exercise, 2022 Publication	<1%
39	discovery.ucl.ac.uk Internet Source	<1%
40	journal.frontiersin.org Internet Source	<1%
41	www.depts.ttu.edu Internet Source	<1%
42	www.midus.wisc.edu Internet Source	<1%

43	Angelina R. Sutin, Antonio Terracciano. "Five- Factor Model Personality Traits and the Objective and Subjective Experience of Body Weight", Journal of Personality, 2016 Publication	<1%
44	Submitted to Aston University Student Paper	<1%
45	bmcpublichealth.biomedcentral.com Internet Source	<1%
46	doi.org Internet Source	<1%
47	www.sciencegate.app Internet Source	<1%
48	Submitted to Taipei Medical University Student Paper	<1%
49	Submitted to University of Strathclyde Student Paper	<1%
50	repository.uin-suska.ac.id Internet Source	<1%
51	c-changeprogram.org Internet Source	<1%
52	refubium.fu-berlin.de Internet Source	<1%
53	Joko Mulyanto, Dionne S Kringos, Anton E Kunst. "Socioeconomic inequalities in	<1%

healthcare utilisation in Indonesia: a comprehensive survey-based overview", BMJ Open, 2019

<1%

<1%

<1%

<1%

Publication

- Kumiko Ohara, Tomoki Mase, Katsuyasu Kouda, Chiemi Miyawaki, Katsumasa Momoi, Tomoko Fujitani, Yuki Fujita, Harunobu Nakamura. "Association of anthropometric status, perceived stress, and personality traits with eating behavior in university students", Eating and Weight Disorders Studies on Anorexia, Bulimia and Obesity, 2019
- Nurul Widyaningrum, Jiang Yu. "Tobacco Use Among the Adult Muslim Population in Indonesia: A Preliminary Study on Religion, Cultural, and Socioeconomic Factors", Journal of Drug Issues, 2018

René Mõttus, Geraldine McNeill, Xueli Jia, Leone C. A. Craig, John M. Starr, Ian J. Deary. "The associations between personality, diet and body mass index in older people.", Health Psychology, 2013

Publication

Tiia Kekäläinen, Antonio Terracciano, Sarianna Sipilä, Katja Kokko. "Personality traits and physical functioning: a cross-sectional

multimethod facet-level analysis", European Review of Aging and Physical Activity, 2020

Publication

58	Submitted to University of Limerick Student Paper	<1%
59	Submitted to University of Stirling Student Paper	<1%
60	e-space.mmu.ac.uk Internet Source	<1%
61	ruj.uj.edu.pl Internet Source	<1%
62	www2.mdpi.com Internet Source	<1%
63	Paans, Nadine P.G., Mariska Bot, Deborah Gibson-Smith, Willem Van der Does, Philip Spinhoven, Ingeborg Brouwer, Marjolein Visser, and Brenda W.J.H. Penninx. "The association between personality traits, cognitive reactivity and body mass index is dependent on depressive and/or anxiety status", Journal of Psychosomatic Research, 2016. Publication	<1%
64	kops.uni-konstanz.de Internet Source	<1%

link.springer.com

65

66 www.fas.harvard.edu

Internet Source

www.journalijar.com

Internet Source

67

68 ARIEF ANSHORY YUSUF, PHOEBE

KOUNDOURI. "Willingness to pay for water and location bias in hedonic price analysis: evidence from the Indonesian housing market", Environment and Development Economics, 2005

Publication

Gabriela Rossiter Stux Veiga, Bruna Merten Padilha, Nassib Bezerra Bueno, Jessiane Rejane Lima Santos et al. "Salutogenesis, nutritional status and eating behaviour: a systematic review", Public Health Nutrition, 2021

Publication

Imelda. "Cooking that kills: Cleaner energy access, indoor air pollution, and health",

Journal of Development Economics, 2020

Publication

Li He, Kun Wang, Tianyang Li, Jiangyin Wang, Yuting Wang, Zixian Zhang, Yuanyang Wu, Shuo Zhang, Siqing Zhang, Hualei Yang. "The <1%

<1%

<1%

71

Positive Impact of Having Served as a Danwei Leader on Post-retirement Life Satisfaction: Experiences in China", Frontiers in Psychology, 2021

Publication

Selly Ruth Defianna, Ailiana Santosa, Ari <1% 72 Probandari, Fatwa Sari Tetra Dewi. "Gender Differences in Prevalence and Risk Factors for Hypertension among Adult Populations: A Cross-Sectional Study in Indonesia", International Journal of Environmental Research and Public Health, 2021 Publication Supa Pengpid, Karl Peltzer. "The prevalence of <1% edentulism and their related factors in Indonesia, 2014/15", BMC Oral Health, 2018 Publication Submitted to University of West London <1% Student Paper hrep-website.s3.ap-southeast-1.amazonaws.com Internet Source neardb.near-aging.se 76 Internet Source pdfs.semanticscholar.org Internet Source

- Bridget A Hearon, Thomas J Harrison. "Not the exercise type? Personality traits and anxiety sensitivity as predictors of objectively measured physical activity and sedentary time", Journal of Health Psychology, 2020 Publication
- < | %

Johansson, E.. "Obesity and labour market success in Finland: The difference between having a high BMI and being fat", Economics and Human Biology, 200903

Publication

<1%

- Kitae Sohn. "The fatter are happier in Indonesia", Quality of Life Research, 2016
- <1%
- Modou Diop, Manuel Ruiz Adame, Alessio Gaggero. "Influence of personality traits, gender, and socio economic factors in obesity in the English Longitudinal Study of Ageing", Personality and Mental Health, 2022
- <1%

Sandra Düzel, Johanna Drewelies, Sarah E.
Polk, Carola Misgeld et al. "No Evidence for a
Boost in Psychosocial Functioning in Older
Age After a 6-Months Physical Exercise

<1%

Intervention", Frontiers in Human Neuroscience, 2022

Publication

84	Tiainen, Anna-Maija K., Satu Männistö, Marius Lahti, Paul A. Blomstedt, Jari Lahti, Mia-Maria Perälä, Katri Räikkönen, Eero Kajantie, and Johan G. Eriksson. "Personality and Dietary Intake – Findings in the Helsinki Birth Cohort Study", PLoS ONE, 2013. Publication	<1%
85	Yannick Stephan, Angelina R. Sutin, Martina Luchetti, Damaris Aschwanden, Vincent Cabibel, Antonio Terracciano. "Measures of physical performance as mediators between personality and cognition in two prospective studies", Archives of Gerontology and Geriatrics, 2023 Publication	<1%
86	digitalcommons.wayne.edu Internet Source	<1%
87	eduvest.greenvest.co.id Internet Source	<1%
88	gradworks.umi.com Internet Source	<1%
89	journals.rcni.com Internet Source	<1%

90	Internet Source	<1%
91	pure.uvt.nl Internet Source	<1%
92	research.ou.nl Internet Source	<1%
93	watermark.silverchair.com Internet Source	<1%
94	www.plosone.org Internet Source	<1%
95	"The Wiley-Blackwell Handbook of Individual Differences", Wiley, 2011	<1%
96	Angelina R. Sutin, Yannick Stephan, Lei Wang, Shoumin Gao, Ping Wang, Antonio Terracciano. "Personality traits and body mass index in Asian populations", Journal of Research in Personality, 2015 Publication	<1%
97	Cillian P. McDowell, Kathryn E. Wilson, Derek C. Monroe, Cathal McCrory, Rose Anne Kenny, Matthew P. Herring. "Physical activity partially mediates associations between "Big" personality traits and incident generalized anxiety disorder: Findings from the irish	<1%

longitudinal study on ageing", Journal of Affective Disorders, 2020

Publication

98	Dominik Siemon, Jörn Wessels. "Performance prediction of basketball players using automated personality mining with twitter data", Sport, Business and Management: An International Journal, 2022 Publication	<1%
99	Ewa Piechurska-Kuciel. "The Big Five in SLA", Springer Science and Business Media LLC, 2020 Publication	<1%
100	Kakizaki, M "Personality and body mass index: A cross-sectional analysis from the Miyagi Cohort Study", Journal of Psychosomatic Research, 200801	<1%
101	Kristina Elfhag, Lesley C. Morey. "Personality traits and eating behavior in the obese: Poor self-control in emotional and external eating but personality assets in restrained eating", Eating Behaviors, 2008 Publication	<1%
102	Michał Seweryn Karbownik, Łukasz Mokros, Maria Dobielska, Mateusz Kowalczyk, Edward	<1%

Maria Dobielska, Mateusz Kowalczyk, Edward Kowalczyk. "Association Between Consumption of Fermented Food and FoodDerived Prebiotics With Cognitive Performance, Depressive, and Anxiety Symptoms in Psychiatrically Healthy Medical Students Under Psychological Stress: A Prospective Cohort Study", Frontiers in Nutrition, 2022

Publication

- Putu Wuri Handayani, Puspa Indahati Sandhyaduhita, Achmad Nizar Hidayanto, Ave Adriana Pinem et al. "chapter 14 Integrated Hospital Information System Architecture Design in Indonesia", IGI Global, 2020
- <1%

Publication

Stephan, Yannick, Julie Boiché, Brice Canada, and Antonio Terracciano. "Association of personality with physical, social, and mental activities across the lifespan: Findings from US and French samples", British Journal of Psychology, 2013.

<1%

Publication

Yannick Stephan, Angelina R. Sutin, Martina Luchetti, Louis Hognon, Brice Canada, Antonio Terracciano. "Personality and self-rated health across eight cohort studies", Social Science & Medicine, 2020

<1%

Publication

Yi Liao, Xiaoyu Cheng, Wei Chen, Xiaowei Peng. "The Influence of Physical Exercise on

<1%

Adolescent Personality Traits: The Mediating Role of Peer Relationship and the Moderating Role of Parent-Child Relationship", Frontiers in Psychology, 2022

Publication

107	Yu Bai, Yanjun Li. "More suffering, more involvement? The causal effects of seismic disasters on social capital", World Development, 2021 Publication	<1%
108	diginole.lib.fsu.edu Internet Source	<1%
109	diposit.ub.edu Internet Source	<1%
110	dspace.lib.cranfield.ac.uk Internet Source	<1%
111	e-journal.unair.ac.id Internet Source	<1%
112	etd.uum.edu.my Internet Source	<1%
113	gevhernesibedergisi.com Internet Source	<1%
114	hub.hku.hk Internet Source	<1%
115	lumsa.it Internet Source	<1%

116	psycnet.apa.org Internet Source	<1%
117	psyct.psychopen.eu Internet Source	<1%
118	uir.unisa.ac.za Internet Source	<1%
119	www.apa.org Internet Source	<1%
120	www.nature.com Internet Source	<1%
121	www.nepjol.info Internet Source	<1%
122	www.parinc.com Internet Source	<1%
123	"Poster session I", Acute Cardiac Care, 2009	<1%
124	Angelina R. Sutin, Antonio Terracciano. "Personality traits and body mass index: Modifiers and mechanisms", Psychology & Health, 2015 Publication	<1%
125	Marius Lahti, Katri Räikkönen, Sakari Lemola, Jari Lahti et al. "Trajectories of physical growth and personality dimensions of the Five-Factor	<1%

Model.", Journal of Personality and Social Psychology, 2013

Publication

Zidan Yang, Guanghua Lei, Xiaoxiao Li, Yilun Wang, Zikun Xie, Xiurui Zhang, Yuchen He, Yilin Xiong, Tubao Yang. "Does symptomatic knee osteoarthritis increase the risk of all-cause mortality? Data from four international population-based longitudinal surveys of aging", Clinical Rheumatology, 2019

<1%

"Handbook of Personality and Self - Regulation", Wiley, 2010

<1%

Publication

Publication

Eurika Zebadia, Trias Mahmudiono,
Dominikus Raditya Atmaka, Mira Dewi, Siti
Helmyati, Cindra Tri Yuniar. "Factors
Associated with Minimum Acceptable Diet in
6–11-Month-Old Indonesian Children Using
the 2017 IDHS", Open Access Macedonian
Journal of Medical Sciences, 2021

<1%

129

Publication

Nicholas P. Tan, Tamlin S. Conner, Haisu Sun, Steven Loughnan, Luke D. Smillie. "Who Gives a Veg? Relations Between Personality and Veg*nism", Appetite, 2021

<1%

R E Rhodes. "Personality correlates of physical activity: a review and meta-analysis", British Journal of Sports Medicine, 12/1/2006

<1%

Publication

Sattler, Sebastian, and Reinhard Schunck.

"Associations Between the Big Five
Personality Traits and the Non-Medical Use of
Prescription Drugs for Cognitive
Enhancement", Frontiers in Psychology, 2016.

Publication

<1%

Taiwo A. Alatishe, Olukayode Abayomi,
Babatunde Suleiman, Olaitan Oladele, Adeoye
Oyewole. "Personality traits, alcohol use and
problem drinking among underraduates in
southwest Nigeria", Journal of Ethnicity in

<1%

Publication

Tamara M. Pfeiler, Boris Egloff. "Personality and attitudinal correlates of meat consumption: Results of two representative German samples", Appetite, 2018

<1%

Publication

Exclude quotes

On

Substance Abuse, 2022

Exclude matches

Off

The relationship between Big Five Personality Traits, eating habits, physical activity, and obesity in Indonesia based on analysis of the 5th wave Indonesia Family Life Survey (2014)

	<u> </u>	
GRADEMARK REPORT		
FINAL GRADE	GENERAL COMMENTS	
/0	Instructor	
7 0		
PAGE 1		
PAGE 2		
PAGE 3		
PAGE 4		
PAGE 5		
PAGE 6		
PAGE 7		
PAGE 8		
PAGE 9		
PAGE 10		
PAGE 11		
PAGE 12		
PAGE 13		
PAGE 14		
PAGE 15		
PAGE 16		
PAGE 17		