

DE GRUYTER

2018 · VOLUME 29 · ISSUE 6  
ISSN 0792-6855 · e-ISSN 2191-0296

# JOURNAL OF BASIC AND CLINICAL PHYSIOLOGY AND PHARMACOLOGY

EDITOR-IN-CHIEF  
*Michal Horowitz*

DE  
|  
G

[www.degruyter.com/jbcpp](http://www.degruyter.com/jbcpp)



 Requires Authentication | Published since December 1, 1986

# Journal of Basic and Clinical Physiology and Pharmacology

ISSN: 2191-0286

Editor-in-chief: Ugo Oliviero

Managing Editor: Alberto Marra

[Submit manuscript](#)

[OVERVIEW](#) [LATEST ISSUE](#) [ISSUES](#) [RANKING](#) [SUBMIT](#) **[EDITORIAL](#)**

## Editorial

### Editor-in-Chief:

Ugo Oliviero (Federico II University, Naples, Italy)

### Deputy Editor:

Alberto M. Marra (Federico II University, Naples, Italy and University of Heidelberg, Germany)

### Associate/Section Editors:

*Emergency Medicine*: Giorgio Bosso (S. Maria delle Grazie Hospital, Pozzuoli, Naples)

*Oncology:* Evelyne Bischof (prev.Ewelina Biskup; University Hospital Basel, Switzerland, Shanghai University of Medicine & Health Sciences, Shanghai, China)

*Hematology and Coagulation disorders:* Pablo Demelo-Rodriguez (G. Marangon Hospital and Universidad Complutense de Madrid, Spain)

*Vascular Medicine:* Antonio Valvano (Legnano Hospital, Legnano, Italy)

*Gastroenterology:* Theodor Voiosu (University of Bucharest, Bucharest, Romania)

*Liver Disease:* Andrei Voiosu (University of Bucharest, Bucharest, Romania)

*Neurology and Cerebrovascular:* Lorenzo Falsetti (Azienda Ospedaliero-Universitaria "Ospedali Riuniti" di Ancona, Italy)

*Gender Medicine:* Valeria Raparelli (University of Ferrara, Ferrara, Italy)

*Endocrinology:* Ieva Ruza, (University of Riga, Riga, Latvia)

*Diabetology and Metabolism:* Mariarosaria De Luca (Federico II University, Naples)

*Cardiovascular Diseases:* Andrea Salzano (Glenfield General Hospital, University of Leicester, Leicester, UK)

*Heart Failure:* Antonio Cittadini (Federico II University of Naples, Naples, Italy)

*Respiratory Medicine:* Salvatore Torrisi (University of Catania, Catania, Italy)

*Geriatrics:* Leonardo Bencivenga (Federico II University, Naples, Italy)

*Immunology:* Gilda Varricchi (Federico II University, Naples, Italy)

*Rheumatology:* Domenico Sambataro (Artroreuma, Catania, Italy)

*Basic Science:* Francesca Vinchi (New York Blood Center, New York, USA), Roberta D'Assante (Federico II, Naples),

*Urology, Andrology and Nephrology:* Felice Crocetto (Federico II University, Naples, Italy)

**Editorial Office:**

E-mail: [jbcpp.editorial@degruyter.com](mailto:jbcpp.editorial@degruyter.com)

---



Published by De Gruyter

## Volume 32 Issue 6

November 2021

Issue of Journal of Basic and Clinical Physiology and Pharmacology

[Submit manuscript](#)

### CONTENTS

### JOURNAL OVERVIEW

[Publicly Available](#) December 8, 2021

#### Frontmatter

Page range: i-ii

[Cite this](#)[Download PDF](#)

#### Review

[Requires Authentication](#) February 8, 2021

##### Historical account of endocrinal disorders in Unani medicine

Md Anzar Alam, Mohd Aleemuddin Quamri, Ghulamuddin Sofi

Page range: 1013-1019

[More ▾](#)[Cite this](#)

#### Original Articles

[Requires Authentication](#) February 9, 2021

##### Impact of sleep disturbances and autonomic dysfunction on the quality of life of patients with fibromyalgia

Ruchi Singh, Nirendra Kumar Rai, Aakarsh Rastogi, Chiranjeevi Endukuru, Ankur Joshi, Sudhanshu Shekhar Mishra

Page range: 1021-1029

[More ▾](#)[Cite this](#)[Requires Authentication](#) March 12, 2021

##### Effects of date seeds administration on anxiety and depressive symptoms in streptozotocin-induced diabetic rats: biochemical and behavioral evidences

Samir Bikri, Youssef Aboussaleh, Assia Berrani, Ismail Louragli, Affaf Hafid, Soukaina Chakib, Ahmed Aharni

Page range: 1031-1040

[More ▾](#)[Cite this](#)[Requires Authentication](#) February 8, 2021

##### Piroxicam reduces acute and chronic pain response in type 1 diabetic rats

Abbas Alimoradian, Fatemeh Samimi, Hadise Asljalal, Seied Amirhossein Latifi, Mehdi Salehi, Maryam Khodaei, Jamal Amri

Page range: 1041-1046

[More ▾](#)[Cite this](#)[Requires Authentication](#) February 8, 2021

##### Protective effect of *Elaeagnus argentea* in maternal obesity-induced deficit cognitive function of the offspring

Azrina Zainal Abidin, Nurain Syahirah Rosli, Santhra Segaran, Azmiza Syawani Jasni, Hasnah Bahari

Page range: 1047-1055

[More ▾](#)[Cite this](#)[Requires Authentication](#) February 8, 2021

##### Pharmacological investigations on efficacy of Phlorizin a sodium-glucose co-transporter (SGLT) inhibitor in mouse model of intracerebroventricular streptozotocin induced dementia of AD type

Reena Rani, Amit Kumar, Amteshwar Singh Jaggi, Nirmal Singh

Page range: 1057-1064

[More ▾](#)[Cite this](#)[Requires Authentication](#) December 17, 2020

##### Concurrent administration of acetaminophen and ethanol: impact on mouse liver and testis

Jonah Sydney Aprioku, Precious Gospel

Page range: 1065-1074

[More ▾](#)[Cite this](#)



Requires Authentication November 19, 2020

### Electrical vestibular nerve stimulation as an adjunctive therapy in the management of type 2 diabetes

Sai Sathesh Kumar Goothy, Jason McKeown, Paul D. McGeoch, Srilatha B, Rajagopalan Vijayaraghavan, Ravikanth Manyam, Mahadik Vijay Khanderao

Page range: 1075-1082

More ▾

Cite this

Requires Authentication February 9, 2021

### Effect of roxithromycin on contractile activity of gastrointestinal smooth muscles in colitic rats

Vijay L. Kumar, Abhimanu Pandey, Hilal Ahmad

Page range: 1083-1086

More ▾

Cite this

Requires Authentication February 8, 2021

### Anti-inflammatory and analgesic effects of methanol extract of red cultivar *Allium cepa* bulbs in rats and mice

Adeoye Joshua Oyewusi, Olayinka Ayotunde Oridupa, Adebowale Bernard Saba, Ibiranke Kofoworola Oyewusi, Jonny Olufemi Olukunle

Page range: 1087-1092

More ▾

Cite this

Requires Authentication October 12, 2020

### Prophylactic effect of aquatic extract of stevia on acetic acid induced-ulcerative colitis in male rats: a possible role of Nrf2 and PPAR $\gamma$

Abeer F. Mostafa, Mahmoud M. Elalfy, Ahmed Shata, Mona G. Elhadidy

Page range: 1093-1104

More ▾

Cite this

Requires Authentication December 14, 2020

### Molecular docking study of flavonoid compounds for possible matrix metalloproteinase-13 inhibition

Amir Taherkhani, Shirin Moradkhani, Athena Orangi, Alireza Jalalvand, Zahra Khamverdi

Page range: 1105-1119

More ▾

Cite this

Requires Authentication December 14, 2020

### Inhibition of key enzymes linked to snake venom induced local tissue damage by kolaviron

Azubulike Ikechukwu Okafor, Elewechi Onyike

Page range: 1121-1130

More ▾

Cite this

Requires Authentication February 8, 2021

### Hepatoprotective effects of *Quassia amara* stem bark against cadmium-induced toxicity in male Wistar rats

Olawale O. Obembe, Taofeek O. Usman, Yinusa Raji

Page range: 1131-1136

More ▾

Cite this

Requires Authentication February 10, 2021

### Curative effects of the aqueous extract of *Tithonia diversifolia* (Hemsl.) A. gray (Asteraceae) against ethanol induced-hepatotoxicity in rats

Ivan Stève Tsopmeji Nguépi, Florence Tsofack Nguéguim, Raceline Kamkumo Gounoue, Adolphe Mbatchou, Theophile Dimo

Page range: 1137-1143

More ▾

Cite this

Requires Authentication February 15, 2021

### The effect of deer antler from East Kalimantan to increase trabecular bone density and calcium levels in serum on osteoporotic mice

Retno Widayati, Suciati Suciati, Dewi Melani Haryadi, Hsin-I Chang, IPG Ngurah Suryawan, Nurliana Tarigan

Page range: 1145-1150

More ▾

Cite this

## Case Report

Requires Authentication April 22, 2021

### A case of statin-induced liver injury with positive rechallenge with a second statin. Is there a class effect?

Giovanna Onfiani, Fabio Nascimbeni, Francesca Carubbi

Page range: 1151-1155

More ▾

Cite this

Retno Widyowati\*, Suciati Suciati, Dewi Melani Haryadi, Hsin-I Chang, IPG Ngurah Suryawan and Nurliana Tarigan

# The effect of deer antler from East Kalimantan to increase trabecular bone density and calcium levels in serum on osteoporotic mice

<https://doi.org/10.1515/jbcpp-2020-0140>

Received May 7, 2020; accepted September 23, 2020;

published online February 15, 2021

## Abstract

**Objectives:** Glucocorticoid-induced osteoporosis (dexamethasone) is a primary cause of secondary osteoporosis by the decreasing formation and increasing resorption activities. Previously, the *in vitro* study showed that 70% ethanol and aqueous extract of deer antler have increased alkaline phosphatase in osteoblast cell that known as marker of bone formation. The mind of this study is to analyze the effect of deer antlers in increasing the bone trabecular density of osteoporosis-induced male mice.

**Methods:** This study used a post-test control group design. A total of 54 healthy male mice were randomly divided to nine groups, i.e., healthy control, osteoporotic, positive control, 70% ethanol (4, 8, and 12 mg/kg BW), and aqueous extracts (4, 8, and 12 mg/kg BW) of deer antler groups. All of the interventions were given 1 mL of test sample for 4 weeks orally. The bone densities were determined using histomorphometry by Image J and Adobe Photoshop. The statistical data were performed using SPSS 23 and statistical significance was set at  $p < 0.05$ .

**Results:** The results showed that alendronate group, 70% ethanol, and aqueous extract groups increased bone density

and calcium levels in serum ( $p < 0.05$ ) compared to osteoporotic group in dose dependent manner. It indicated that 70% ethanol and aqueous extract of deer antler stimulating bone turnover and aqueous extract showed the highest.

**Conclusions:** Dexamethasone induction for 4 weeks caused osteoporotic mice and the administration of 70% ethanol and aqueous extracts of deer antler from East Kalimantan increased trabecular bone density and calcium levels in dose dependent manner.

**Keywords:** bone density; bone turnover; calcium level; deer antler.

## Introduction

Osteoporosis is considered to be a skeletal problem due to impaired bone strength that results in an increased risk of fracture [1]. Novel epidemiological studies have shown that osteoporosis becomes a primary public health problem not only for women population, but also for men population. It is estimated that the total amount of hip fractures in women and men in 2025 will be alike [2, 3]. In men, the distribution of osteoporosis prevalence is bimodal, showing that the initial peak (<50 year old) is mostly due secondary osteoporosis, while the later peak (>60 year old) is mostly categorized as primary osteoporosis [4]. Correspond to the World Health Organization (WHO), by applying the standard from The International Society for Clinical Densitometry, it is estimated that 1–2 million men in the United States have osteoporosis (T-score  $< -2.5$ ) and 8–13 million have osteopenia (T-score between  $-1.0$  and  $-2.5$ ), or their prevalence are 6% for osteoporosis and 47% for osteopenia [3, 5, 6]. In aging population, morbidity and mortality from hip fractures are higher in men than in women with fatality rates, among over 75 years is 20.7% in men vs. 7.5% in women [7]. The causes of osteoporosis in men are relative to hormonal, genetics, environmental, other specific disease factors, and along 50% of men with secondary osteoporosis [6, 8]. The three primary causes of secondary osteoporosis in men are glucocorticoid excess, hypogonadism, and alcohol abuse. The precaution and therapy of osteoporosis disease according to the Recommendation of American College of

\*Corresponding author: Retno Widyowati, Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Universitas Airlangga, Nanizar Zaman Joenoes Building, Campus C, Mulyorejo, Surabaya, Indonesia, Phone: +62 81615886978, E-mail: rr-retno-w@ff.unair.ac.id. <https://orcid.org/0000-0002-6166-1289>  
Suciati Suciati, Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Universitas Airlangga, Nanizar Zaman Joenoes Building, Campus C, Mulyorejo, Surabaya, Indonesia  
Dewi Melani Haryadi, Department of Pharmaceutics, Faculty of Pharmacy, Universitas Airlangga, Nanizar Zaman Joenoes Building, Campus C, Mulyorejo, Surabaya, Indonesia  
Hsin-I Chang, Department of Biochemical Science and Technology, National Chiayi University, Chiayi, Taiwan, P. R. China  
IPG Ngurah Suryawan and Nurliana Tarigan, UPTD Pembibitan dan Inseminasi Buatan, Dinas Peternakan dan Kesehatan Hewan Provinsi Kalimantan Timur, Penajam Paser Utara, Indonesia

Rheumatology Ad Hoc Committee is using supplementation with vitamin D and calcium, anti-resorptive agents, calcitonin, gonadal sex hormone replacement, and modifying lifestyle risk factors [9]. Antiresorptive agents that are currently widely used are nitrogenous bisphosphonates which contain alendronate and risedronate, and also calcitonin, estrogen, and raloxifene selective estrogen receptor modulators. These agents increase bone strength and degrade the risk of fractures to varying degrees [10, 11]. Some clinical evidences recommend a role for phytoestrogen in the therapy of osteoporosis [12–15]. Previous studies showed that 70% ethanol and aqueous extracts of deer antler from East Kalimantan increased the alkaline phosphatase (ALP) and mineralization activities of 7F2 cell that is an osteoblast cell lines [16, 17]. These extracts contain several major constituents such as protein, lipids, ash, calcium, collagen, chondroitin sulfate and glucosamine [18, 19]. Based on these data, it is necessary to prove whether the 70% ethanol and aqueous extracts of deer antler can increase the bone density of osteoporotic mice.

## Materials and methods

### Materials

Deer antler of *Rusa unicolor* was collected in the middle of March 2017 in UPTD (Technical Implementation Service Unit) of East Kalimantan, Indonesia and voucher specimens were deposited at the UPTD of East Kalimantan, Indonesia. The experimental animal used was male mice (*Mus musculus*) obtained from the Animal Laboratory of the Faculty of Pharmacy, Universitas Airlangga, Surabaya (No.2.KE.176.09.2019). The mice were 5 months old, healthy and weighed of  $19.208 \pm 10.265$  g. The materials were Dexamethasone tablets (Generic, Indonesia), Alendronate® (Novell Pharma, Indonesia), Ethanol pro analysis (Merck, Indonesia), HCl Ketamine (Kepro BV, Indonesia), and CMC-Na.

### Extraction of *R. unicolor* antlers

*R. unicolor* antler powder was received from UPTD of East Kalimantan, Indonesia. The 991 g of powder was extracted with 70% ethanol – water (2.0 L × 3) using maceration method. The 70% ethanol solution was concentrated using BUCHI rotary evaporator to get 70% ethanol extract (Et-TL, 35.0 g). In addition, the deer antler powder (430 g) was extracted with 100% water (1.0 L × 3) by applying continuous percolation method. The water solution was dried by freeze dried to get aqueous extract (A-TL, 6.1 g).

### Trabecular bone density and calcium levels in serum

This study applied a posttest control group design. The 54 healthy male mice were randomly divided into nine groups. There were healthy control group (without dexamethasone induction), osteoporotic group (induction with dexamethasone and without extracts),

positive control group (alendronate suspension), the 70% ethanol extract groups of deer antler (4, 8, and 12 mg/kg BW), and the aqueous extract groups of deer antler (4, 8, and 12 mg/kg BW). First of all, the mice were induced by 1 mL dexamethasone (0.0029 mg/20 g BW/day) orally for four weeks to obtain osteoporotic conditions [20]. Then the mice were carried out with or without 1 mL extracts orally for four weeks. After 4 weeks, they were sacrificed with anesthesia using HCl ketamine at 10 mg/g BW, i.p. [21] and their femur bones and blood were taken. Blood sampling was performed to measure the calcium levels in serum, whereas femoral bones were taken as the material for histological preparations. The femoral bones were immediately fixed in 10% neutral-buffered formalin and placed in decalcifying solution for 24 h at 37 °C, continuous with being dehydrated and embedded in paraffin. The proximal femur section was stained with a Hematoxylin-eosin (HE) staining. Then, histomorphometry observations of the percentage of trabecular bone density using the Optic Lab microscope and computer software of Image J and Adobe Photoshop were performed [22]. Calculation of trabecular bone density was obtained by dividing the area of observed trabecular bone ( $\mu\text{m}^2$ ) with the area of entire measurement area (trabecular bone and bone marrow space). The formula for calculating bone density was as follows [23]:

$$\begin{aligned} \text{Trabecular Bone Density \% BV/TV} \\ = \frac{\text{the area of observed trabecular bone ( T )}}{\text{the area of entire measurement area ( T + TS )}} \times 100\% \end{aligned}$$


The observations were made in the metaphyseal region that approached epiphyseal line and it was in cortisol bone region [23]. The bone density values were obtained in units of % BV/TV (% Bone volume/Tissue volume).

### Statistical analysis

The experiments were carried out for three more consecutive times using similar sample. It was then presented as means  $\pm$  standard deviations. The statistical data were performed using SPSS 23 and statistical significance was set at  $p < 0.05$ .


## Results

The result of dexamethasone induction for 4 weeks caused osteoporotic occurred in mice due to decreased trabecular bone density and characterized by a change in the vertebrae posture into kyphosis (Figure 1). Changes in kyphosis




**Figure 1:** Healthy mice (A) and osteoporotic mice (B), arrow point marks changes in vertebrae posture to kyphosis.





**Figure 2:** Histomorphometry of the trabecular area from control group (A) and osteoporotic group (B) by hematoxylin-eosin staining. T = Trabecular Bone, ST = Bone Marrow, TK = Cortical Bone, GE = Epiphyseal Line (100× Magnification).



**Figure 3:** Trabecular bone density levels of several groups; healthy (H), osteoporotic (O), positive control (A, alendronate), 70% ethanol extracts (Et at 4, 8, and 12 mg), and aqueous extracts (EA at 4, 8, and 12 mg) after intervention, (\* $p < 0.05$  compare to osteoporotic group).

posture occurred in all groups (O, A, Et, and EA) except the healthy group (H).

The occurrence of osteoporosis in mice due to dexamethasone induction was not only observed visually, but histomorphometry observation was also carried out on the trabecular area of mice (Figure 2). It could be seen that the trabecular femur area of osteoporotic group was narrower than healthy group. Then, these results were measured in percentage of trabecular bone density (Figure 3).

Dexamethasone induction for 4 weeks in mice reduced the trabecular bone density (O) to  $23.79 \pm 1.23\%$ , while the healthy group (H) was  $17.48 \pm 1.57\%$  (Figure 3). These results were statistically analyzed using SPSS 23 and the  $\alpha$  value was  $< 0.05$ , it showed that osteoporotic mice group suffered osteoporosis disease.

Observation of anti-osteoporosis activity test from 70% ethanol and aqueous extracts of deer antlers was done through histomorphometry calculation of the average trabecular bone density (%). The results showed that trabecular bone density of the osteoporotic, positive control, 70% ethanol extract at 4, 8, and 12 mg/kg BW and aqueous extract at 4, 8, and 12 mg/kg BW groups were  $17.48 \pm 1.57\%$ ;  $35.09 \pm 1.53\%$ ;  $22.39 \pm 2.36\%$ ;  $26.42 \pm 2.47\%$ ;  $31.83 \pm 3.84\%$ ;  $30.66 \pm 2.65\%$ ;  $33.84 \pm 1.53\%$ ; and  $35.64 \pm 2.42\%$  respectively (Figure 3). Furthermore, a statistical analysis test was performed using SPSS 23 ( $\alpha < 0.05$ ) and the results showed that 70% ethanol extract and aqueous from deer antlers at three concentrations had anti-osteoporosis activity by increasing the value of trabecular bone density, significantly, compared to the osteoporotic group.

**Table 1:** Calcium levels in serum of several groups after intervention (\* $p < 0.05$ ).

Groups	Calcium levels in serum, ppm
Healthy	$9.65 \pm 0.02^*$
Osteoporotic	$9.15 \pm 0.04$
Positive control	$9.89 \pm 0.05^*$
70% ethanol extract	
4 mg/g BW	$9.43 \pm 0.14^*$
8 mg/g BW	$9.51 \pm 0.42^*$
12 mg/g BW	$9.73 \pm 0.04^*$
Aqueous extract	
4 mg/g BW	$9.62 \pm 0.22^*$
8 mg/g BW	$9.74 \pm 0.24^*$
12 mg/g BW	$9.87 \pm 0.12^*$

Calcium levels in the obtained blood/serum were measured by using a spectrophotometer, as shown in Table 1. The results were analyzed using SPSS with a  $p$  value of  $< 0.05$  and showed that the value of calcium levels in serum increased in the 70% ethanol and aqueous extract of deer antlers at dose dependent manner compared to the osteoporotic group.

## Discussion

Deer antlers have been highly used in traditional oriental medicine and several studies have examined the benefits



of potential effects. An evaluation of the effect of deer antler on physical growth and bone development is necessary and relevant for the amount of bioactive substances in deer antler, which are assured to have bone-strengthening assets [24]. Therefore, we assessed bone development parameter (femoral bone densitometry) and bone-related enzyme (calcium). Calcium in serum directly affects bone calcification and dissolution and can be measured as parameters related to bone metabolism.

Before anti-osteoporosis activity was carried out, mice were induced by dexamethasone for 14 days so that the mice suffered osteoporosis disease. Dexamethasone is a glucocorticoid drug that can directly inhibit osteoblast activity and inhibit the production of sex hormones that affect bone formation. The use of dexamethasone for a long time (4 weeks at a dose of 0.0029 mg/20 g BW rat/day) causes a decrease in the average percentage of trabecular bone density [20].

The effect of dexamethasone induction in this study has been seen by changing the vertebrae posture into the kyphosis bone, as in Figure 1, and was supported by histomorphometry observations in the trabecular area of mice, as in Figure 2. The figure showed that the trabecular area of osteoporotic group was narrower than the area of healthy group. Then, histomorphometry results were measured in percentage of trabecular bone density, as in Figure 3, that showed that there was a decrease in the percentage of trabecular bone density in osteoporotic group compared with the healthy group. Low total bone density is the causes of osteoporosis and has several cytokines such as IL,  $1\beta$ -11, and TNF $\alpha$  that stimulate aromatase activity of osteoblast cells [25] and have effect to intestinal metabolism of phyto-testosterone [26]. Ma et al. (2011) reported that glucocorticoid had increased the expression and signaling activity of  $\beta$ 2-adrenergic receptors in osteoblast. These stimulations inhibited osteoblast proliferation, stimulated osteoclastogenesis and increased regulation of nuclear factor- $\kappa$ B ligand expression [27].

Dexamethasone-induced mice were given a test treatment for 4 weeks, then their calcium levels in serum were observed and histophotometry of their trabecular bone density were examined. Calcium is a mineral found in bones (99%). In the study of ovariectomy osteoporotic rat models showed that calcium levels in serum decreased to 7.26 mg/dL compared to normal rat, which was 8.35 mg/dL [28]. Glucocorticoid induction can reduce calcium levels in serum by decreasing the absorption of calcium from intestine and inhibiting calcium reabsorption in kidney tubules thereby increasing the excretion of calcium through urine [29].

Based on the statistical analysis, there were significant differences of trabecular bone density between positive

control, the 70% ethanol extract and the aqueous extract groups toward osteoporotic group. This study showed that the level of trabecular bone density of osteoporosis group (after intervention with dexamethasone for 4 weeks) was lower than other groups. There was an increase of trabecular bone density in positive control group. There is not too much information regarding the mechanism but it is possible that the alendronate mechanism of action may be indirect. After 12 months of therapy, alendronate was found to stimulate bone mass in femoral neck and lumbar spine in androgen replacing men with long-term hypogonadism. After 6 months of alendronate treatment, urinary deoxypyridinoline which marker of bone resorption was decreased significantly [30]. This condition will result in the balance of bone remodeling and increase osteocalcin as serum marker of bone formation [31].

Shimon et al. (2005) informed that alendronate at 10 mg daily in osteoporotic men with long-standing hypogonadism for 6–12 months increased lumbar-spine bone mineral density significantly ( $p < 0.005$ ) [30]. Alendronate is an anti-resorptive agent that hampers farnesyl diphosphate (FPP) synthase, thereby blocking the prenylation of small signaling proteins that are important for osteoclast function and viability [32, 33]. Revell (1986) reported that histomorphometry method could be used to show a great correlation between the actual bone volume (0.998) [34].

Several components in deer antlers have started a direct modulation effect on bone growth. Deer antlers have been declared to contain essential amino acids, hyaluronic acid, chondroitin sulfate, collagen, polysaccharides, glycosaminoglics, a number of fatty acids (C18: 3-omega-6 fatty acids), phosphorus, zinc, iron, and calcium [35]. Chondroitin sulfate in deer antlers is a major glycosaminoglycan [36, 37] which is water soluble and shows an increased growth effect on osteoblast cells [38] and believed to manage water storage, differentiation, and proliferation of chondrocytes in cartilage tissue.

Overall, the positive effects observed from deer antler extracts from East Kalimantan on bone development in this study are appropriate with some fundamental of traditional Chinese medicine. The proper mechanism of action and the biologically active substances responsible for these effects will involve further research to be explained.

## Conclusions

Dexamethasone induction for 4 weeks caused osteoporotic mice, markedly by the occurrence of kyphosis and the

narrowing of the trabecular area. The administration of 70% ethanol and aqueous extracts of deer antler from East Kalimantan increased trabecular bone density and calcium levels in dose dependent manner. Thus, the extracts stimulated bone turnover.

**Acknowledgments:** The authors are grateful for access to Animal Laboratory of the Faculty of Pharmacy, Universitas Airlangga and would like to express their sincere thanks to Research and Innovation Institute of Universitas Airlangga and the dean of Faculty of Pharmacy, Universitas Airlangga for supporting fund.

**Research funding:** This research was supported by research mandatory of Airlangga University (No. 886/UN3/2018) and research collaboration between Faculty of Pharmacy Airlangga University, UPTD of East Kalimantan and National Chiayi University, Taiwan.

**Author contributions:** All authors have accepted responsibility for the entire content of this manuscript and approved its submission. RW designed the research concepts, SC designed methods, DW edited the article, HC designed discussion, NS and NT provided the deer antlers.

**Competing interests:** The authors declared that no conflict of interest in this article.

**Informed consent:** Not applicable.

**Ethical approval:** This study was approved by ethics commission of Faculty of Veterinary Medicine, Universitas Airlangga with No. 2.KE.176.09.2019.

## References



- Agrawal VK, Gupta DK. Recent update on osteoporosis. *Int J Med Sci Publ Health* 2013;2:164–8.
- Eiben G, Dey DK, Rothenberg E, Steen B, Björkelund C, Bengtsson C, et al. Obesity in 70-year-old Swedes: secular changes over 30 years. *Int J Obes* 2005;29:810–7.
- Gennari L, Bilezikian JP. Osteoporosis in men. *Endocrinol Metab Clin N Am* 2007;36:399–419.
- Sözen T, Özişik L, Başaran NÇ. An overview and management of osteoporosis. *Eur J Rheumatol* 2017;4:46–56.
- Kling JM, Clarke BL, Sandhu NP. Osteoporosis prevention, screening and treatment: a review. *J Women Health (Larchmt)* 2014;23:563–72.
- Licata A. Osteoporosis in men: suspect secondary disease first. *Cleve Clin J Med* 2003;70:247–54.
- Siddapur PR, Patil AB, Borde VS. Comparison of bone mineral density, T-score and serum zink between diabetic and non diabetic postmenopausal women with osteoporosis. *J Lab Phys* 2015;7:43–8.
- Kotwal N, Upreti V, Nachankar A, Kumar KVSH. A prospective, observational study of osteoporosis in men. *Indian J Endocrinol Metab* 2018;22:62–6.
- American College of Rheumatology Ad Hoc Committee on Glucocorticoid-Induced Osteoporosis. Recommendations for the prevention and treatment of glucocorticoid-induced osteoporosis. *Arthritis Rheum* 2001;44:1496–503.
- Drake MT, Clarke BL, Khosla S. Bisphosphonates: mechanism of action and role in clinical practice. *Mayo Clin Proc* 2008;83:1032–45.
- Epstein S. The roles of bone mineral density, bone turnover and other properties in reducing fracture risk during antiresorptive therapy. *Mayo Clin Proc* 2005;80:378–88.
- Al-Anazi AF, Qureshi VF, Javaid K, Qureshi S. Prevention effects of phytoestrogens against postmenopausal osteoporosis as compared to the available therapeutic choices: an overview. *J Nat Sci Biol Med* 2011;2:154–63.
- Arjmandi BH. The role of phytoestrogens in the prevention and treatment of osteoporosis in ovarian hormone deficiency. *JACN* 2001;20:3985–4025.
- Uesugi T, Fukui Y, Yamori Y. Beneficial effects of soybean isoflavon supplementation on bone metabolism and serum lipids in postmenopausal Japanese women. A four weeks study. *JACN* 2002;21:97–102.
- Atkinson C, Compston JE, Day NE, Dowsett M, Bingham SA. The effects of phytoestrogen isoflavons on bone density in women: a double-blind, randomized, placebo-controlled trial. *Am L Clin Nutr* 2004;79:326–33.
- Widyowati R, Suciati, Haryadi DM, Chang H, Suryawan IPGN, Utama AW. The effect of *Rusa unicorn* antler extracts from East Kalimantan in bone turnover cell models. *Turk J Pharm Sci* 2020;17:440–5.
- Gong W, Li F. Cervi cornu pantotrichum aqueous extract promote osteoblasts differentiation and bone formation. *Biomed Res* 2014;25:249–52.
- Sui Z, Zhang L, Huo Y, Zhang Y. Bioactive components of velvet antlers and their pharmacological properties. *J Pharmaceut Biomed Anal* 2014;87:229–40.
- Kawtikwar PS, Bhagwat DA, Sakarkar DM. Deer antler-traditional use and future perspectives. *Indian J Trad Knowl* 2010;9:245–51.
- Hadiwidjojo MGH. Aktivitas antiosteoporosis ekstrak etanol 70% dan hasil fraksinasi *Spilanthes acmella* dalam meningkatkan kepadatan tulang trabecular femur mencit jantan [Thesis]. Fakultas Farmasi Universitas Airlangga; 2014.
- Li-Xia S, Zhang JC, Wu J, Hashimoto K. Antidepressant effects of ketamine on depression-like behavior in juvenile mice after neonatal dexamethasone exposure. *Clin Psychopharmacol Neurosci* 2014;12:124–7.
- Egan KP, Brennan TA, Pignolo RJ. Bone histomorphometry using free and commonly available software. *Histopathology* 2012;61:1168–73.
- Duque G, Watanabe K. Osteoporosis research animal models. London Dordrecht Heidelberg, New York: Springer; 2011.
- Sim JS, Sunwoo HH. Antler nutraceuticals for the newly emerging functional food market in North America: ASPT research update. In: Sim JS, Sunwoo HH, Hudson RJ, Jeon BT, editors *Antler science and product technology*. Edmonton Canada: ASPTRC; 2001.
- Shozu M, Simpson ER. Aromatase expression of human osteoblast-like cell. *Mol Cell Endocrinol* 1998;139:117–29.
- Chiechi LM, Micheli L. Utility of dietary phytoestrogens in preventing postmenopausal osteoporosis. *Curr Top Nutraceutical Res* 2005;3:15–28.

27. Ma Y, Nyman JS, Tao H, Moss HH, Yang X, Elefteriou F. B2-Adrenergic receptor signaling in osteoblasts contributes to the catabolic effect of glucocorticoids on bone. *Endocrinology* 2011;152:1412–22.
28. Elkomy MM, Elsaid FG. Anti-osteoporotic effect of medical herbs and calcium supplementation on ovariectomized rats. *JOBAZ* 2015;72:81–8.
29. Zaqqa D, Jackson RD. Diagnosis and treatment of glucocorticoid-induced osteoporosis. *CCJM* 1999;66:221–30.
30. Shimon I, Eshed V, Doolman R, Sela B-A, Karasik A, Vered I. Alendronate for osteoporosis in men with androgen-related hypogonadism. *Osteoporos Int* 2005;16:1591–6.
31. Karsenty G, Oury F. Regulation of male fertility by the bone derived hormone osteocalcin. *Mol Cell Endocrinol* 2014;382:1–13.
32. Orwoll E, Ettinger M, Weiss S, Miller P, Kendler D, Graham J, et al. Alendronate for the treatment of osteoporosis in men. *N Engl J Med* 2000;343:604–10.
33. Melsen F, Melsen B, Mosekilde L, Bergmann S. Histomorphometric analysis of normal bone from the iliac crest. *Acta Pathol Microbiol Scand* 1978;86:70–81.
34. Revell P. Histomorphometry of bone. *J Clin Pathol* 1983;36:1323–31.
35. Hemmings SJ, Song X. The effects of elk velvet antler consumption on the rat: development, behavior, toxicity and the activity of liver  $\gamma$ -glutamyltranspeptidase. *CBP Part C* 2004;138:105–12.
36. Sunwoo HH, Sim LYM, Nakano T, Hudson RJ, Sim JS. Glycosaminoglycans from growing antlers of wapiti (*Cervus elaphus*). *Can J Anim Sci* 1997;77:715–21.
37. Sunwoo HH, Nakano T, Hudson RJ, Sim JS. Isolation, characterization and localization of glycosaminoglycans in growing antlers of wapiti (*Cervus elaphus*). *CBP Part B* 1998;120:273–83.
38. Sunwoo HH, Nakano T, Sim JS. Effect of water-soluble extract from antler of wapiti (*Cervus elaphus*) on the growth of fibroblasts. *Can J Anim Sci* 1997;77:343–5.

## Journal of Basic and Clinical Physiology and Pharmacology

### COUNTRY

Germany



### SUBJECT AREA AND CATEGORY

 Biochemistry, Genetics and Molecular Biology  
 ↳ Physiology

 Medicine  
 ↳ Medicine (miscellaneous)

 Pharmacology, Toxicology and Pharmaceutics  
 ↳ Drug Discovery  
 ↳ Pharmacology

### PUBLISHER

Walter de Gruyter GmbH

### H-INDEX

36

### PUBLICATION TYPE

Journals

### ISSN

07926855, 21910286



### COVERAGE

1985-1988, 1990-2021

### INFORMATION

[Homepage](#)
[How to publish in this journal](#)
[m.horowitz@mail.huji.ac.il](mailto:m.horowitz@mail.huji.ac.il)

### Quartiles







# Source details

## Journal of Basic and Clinical Physiology and Pharmacology

Formerly known as: Reviews in Clinical and Basic Pharmacology

Scopus coverage years: from 1985 to 1988, from 1990 to Present

Publisher: Walter de Gruyter

ISSN: 0792-6855 E-ISSN: 2191-0286

Subject area: Pharmacology, Toxicology and Pharmaceutics: Pharmacology

Pharmacology, Toxicology and Pharmaceutics: Drug Discovery

Biochemistry, Genetics and Molecular Biology: Physiology

CiteScore 2021

2.5



SJR 2021

0.347



SNIP 2021

0.728



Source type: Journal

[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

### i Improved CiteScore methodology



CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021

$$2.5 = \frac{1,062 \text{ Citations 2018 - 2021}}{421 \text{ Documents 2018 - 2021}}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022

$$3.5 = \frac{1,521 \text{ Citations to date}}{429 \text{ Documents to date}}$$

Last updated on 05 March, 2023 • Updated monthly

### CiteScore rank 2021

Category	Rank	Percentile
Pharmacology, Toxicology and Pharmaceutics	#203/303	33rd
Pharmacology		
Pharmacology, Toxicology and Pharmaceutics	#109/154	29th
Drug Discovery		

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)