

Intended for healthcare professionals

Journal of Public Health Research

About this journal

The Journal of Public Health Research is an online Open Access, peer-reviewed scholarly journal in the field of public health science. The aim of the journal is to stimulate debate and dissemination of knowledge in the public health field in order to improve efficacy, effectiveness and efficiency of public health interventions to improve health outcomes of populations. This aim can only be achieved by adopting a global and multidisciplinary approach.

More



All Issues

Current Issue

Submit Paper

Latest Articles

Most Read

Most Cited

Trending on Altmetric

Articles most recently published online for this journal.

Privacy

Academic Stress in Adolescent Students of an Islamic-Based School: The Correlation with Parenti...

Intan Nurul Dwi Utari, Achir Yani S. Hamid

First published: 16 Jun 2022

[View More](#)

Tell us what you think



We're continually looking for ways to enhance the site to make sure you're having the best experience. We greatly value your feedback!

[Feedback](#)

Editorial Team

Editor

- Professor Luigi Barberini
University of Cagliari and University Hospital of Cagliari, Italy

[View All](#)

Email Alerts


Sign up and receive alerts:

- With the latest Table of Contents
- When new articles are published online

[Sign Up](#)

More from this Journal

Related Journals



[Asia-Pacific Journal of Public Health](#)



[Perspectives in Public Health](#)



[Public Health Reports](#)



[Scandinavian Journal of Public Health](#)

Journal of Public Health Research

Table of Contents

Volume 11 Issue 2, April - June 2022

Research Article



Correlates of attitudes toward COVID-19-related public health policies and prevention practices in six states

Karen Glanz, Jade Avelis, Pui L Kwong, John H Holmes

First Published June 30, 2022

[Abstract](#)
> [Preview](#)



Perspectives and Debates



What do we know about corporate social responsibility and stakeholders physical activity? A Public Health Perspective

Malin Johansson, Anne-Sofie Hiswåls, Lena Svennberg, Gloria Macassa

First Published June 30, 2022

[Abstract](#)
> [Preview](#)



Research Article





Digital environment: An evolutionary component in environmental health

Afiqah Syamimi Masrani, Nik Rosmawati Nik Husain

First Published June 30, 2022

[Abstract](#)
> [Preview](#)



Research Article



Adverse and other unintended consequences of setting-based interventions to prevent illicit drug use: A systematic review of reviews

Renke L Biallas, Eva Rehfuess, Jan M Stratil

First Published June 30, 2022

[Abstract](#)
> [Preview](#)



Review



Antibacterial, antifungal, antiviral, and photocatalytic activities of TiO₂ nanoparticles, nanocomposites, and bio-nanocomposites: Recent advances and challenges

Mohammad Reza Amiri, Mehran Alavi, Mojtaba Taran, Danial Kahrizi

First Published June 30, 2022

[Abstract](#)
> [Preview](#)



Original Article



Tensions between infection control regulations and the home care working environment: An interview study with health care providers early during the COVID-19 pandemic

Sofia Tavemark, Helle Wijk, Cecilia Pettersson

First Published June 30, 2022

[Abstract](#)
> [Preview](#)



Research Article



Husband's involvement in wife's antenatal care visits in Indonesia: What factors are related?

Agung Dwi Laksono, Ratna Dwi Wulandari, Noor Edi Widya Sukoco, Suharmiati Suharmiati

First Published June 30, 2022

[Abstract](#)
> [Preview](#)



Original Article



The prevalence of gaming and gambling in a child and adolescent psychiatry unit

Frida André, Anders Håkansson, Björn Axel Johansson, Emma Claesdotter-Knutsson

First Published June 30, 2022

[Abstract](#)
> [Preview](#)



Brief Report



Journal of Public Health Research



Editorial Board

 [Hide All](#)

Editor in Chief

Luigi Barberini Department of Medical Sciences and Public Health, University of Cagliari, Italy

Associate Editors

Laura Atzori Department of Medical Sciences and Public Health, University of Cagliari, Italy

Antonella Balestrieri Department of Medical Sciences and Public Health, University of Cagliari, Italy

Doris Barcellona Department of Medical Sciences and Public Health, University of Cagliari, Italy

Cristina Cabras Department of Pedagogy, Psychology, Philosophy, University of Cagliari, Italy

Francesco Campanella Department of Occupational and Environmental Medicine, Epidemiology and Hygiene, Worker Compensation Authority (INAIL), Italy

Emanuele Cannizzaro Department of Health Promotion, Maternal and Infant Care, University of Palermo, Italy

Alessandro Carriero Faculty of Medicine and Surgery, University of Eastern Piedmont, Italy

Mauro Carta	Department of Medical Sciences and Public Health, University of Cagliari, Italy
Flaviana Cau	Department of Medical Sciences and Public Health, University of Cagliari, Italy
Giulia Cossu	Department of Medical Sciences and Public Health, University of Cagliari, Italy
Daniela Fanni	Department of Medical Sciences and Public Health, University of Cagliari, Italy
Claudia Fattuoni	University of Cagliari Department of Chemical and Geological Sciences, University of Cagliari, Italy
Pierpaolo Ferrante	Department of Occupational and Environmental Medicine, Epidemiology and Hygiene, Worker Compensation Authority (INAIL), Italy
Clara Gerosa	Department of Medical Sciences and Public Health, University of Cagliari, Italy
Dmitri Grapov	University of California, Davis, CA, USA
Danial Kahrizi	Razi University, Iran
Jutta Lindert	University of Emden, Germany
Federico Meloni	Department of Medical Sciences and Public Health, University of Cagliari, Italy
Olga Mulas	Department of Medical Sciences and Public Health, University of Cagliari, Italy
Marcello Nonnis	Department of Pedagogy, Psychology and Philosophy, University of Cagliari, Italy
Germano Orrù	Department of Medical Sciences and Public Health, University of Cagliari, Italy
Monica Piras	Department of Medical Sciences and Public Health, University of Cagliari, Italy
	Department of Medical Sciences and Public Health University of Cagliari, Italy

Luca Saba Department of Medical Sciences and Public Health, University of Cagliari, Italy

Salvatore Sardo Department of Medical Sciences and Public Health, University of Cagliari, Italy

Alessandra Scano Department of Medical Sciences and Public Health, University of Cagliari, Italy

More about this journal



Description



Aims and Scope



Abstracting/Indexing



Submit Paper

Husband's involvement in wife's antenatal care visits in Indonesia: What factors are related?

Journal of Public Health Research
2022, Vol. 11(2), 1–8
© The Author(s) 2022
DOI: 10.1177/22799036221104156
journals.sagepub.com/home/phj
SAGE

Agung Dwi Laksono^{1,2}, Ratna Dwi Wulandari^{2,3}, Noor Edi Widya Sukoco¹ and Suharmiati Suharmiati¹

Abstract

Background: Involving husbands in maternal and child health programs can reduce maternal morbidity and mortality. The study aimed to analyze the factors related to the husband's involvement in antenatal care (ANC) visits in Indonesia.

Methods: The study was a cross-sectional study. The research employed secondary data from the 2017 Indonesian Demographic and Health Survey. The analysis units were wives aged 15–49 years old, married, and pregnant in the past 5 years, and the final samples obtained were 14,319 respondents. In addition to the husband's involvement were residence, age, education, occupation, wealth, and parity. The study used a binary logistic regression test in the final stage.

Results: Husbands who lived in urban areas have odds of being involved in ANC visits 1.2 times greater than those in rural areas. The better the husband's education was, the more involved they were in ANC. The results show husbands who work in any field were better at being involved in ANC. Poorer husbands had odds of being engaged in ANC visits 2.0 times more likely than the husbands' lowest group. The most prosperous husbands have odds of being involved in ANC visits 5.4 times than the poorest husbands. The husbands' wealth is better, the more the husbands were engaged in ANC. The more children were born, the less frequent the husbands' involvement in ANC is.

Conclusion: The study concluded five variables associated with husbands' participation in ANC in Indonesia, including residence, education, occupation, wealth, and parity.

Keywords

Husband's involvement, maternal care, maternal health, antenatal care, pregnancy care, family factors, public health

Date received: 13 October 2021; accepted: 14 February 2022

Introduction

The Maternal Mortality Rate (MMR) in Indonesia has significantly decreased. Based on the 2010 Population Census, the Indonesian MMR was 346 maternal deaths per 100,000 live births. The 2015 SUPAS (*Survei Penduduk Antar Sensus/Inter-Census Population Survey*) data showed the MMR in Indonesia dropped to 305 maternal deaths per 100,000 live births. Although it is pretty significant, this figure has not met the 2015 Indonesian MDGs target, 102 maternal deaths per 100,000 live births.¹ Moreover, in the Sustainable Development Goals (SDGs), the MMR target is 70 per 100,000 live births by 2030.²

Compared to other South East Asia region countries, the Indonesian MMR has the highest record. The Indonesian MMR was nine times higher than Malaysia,

five times higher than Vietnam, and almost two times higher than Cambodia. The World Health Organization (WHO) estimated the MMR in developed countries was at 12 maternal deaths per 100,000 live births. While in developing countries, there were 239 maternal deaths per

¹National Research and Innovation Agency, Republic of Indonesia, Jakarta, Indonesia

²The Airlangga Centre for Health Policy (ACeHAP) Research Group, Surabaya, Indonesia

³Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia

Corresponding author:

Ratna Dwi Wulandari, Faculty of Public Health, Universitas Airlangga, Universitas Airlangga Campus C Mulyorejo, Surabaya 60115, Indonesia.
Email: ratna-d-w@fkm.unair.ac.id



100,000 live births. On the other hand, the WHO estimates the MMR was 216 per 100,000 live birth globally.^{3,4}

The government's primary strategy is reducing maternal morbidity and mortality by conducting antenatal care (ANC). ANC is an early strategy to monitor and maintain mothers' and fetuses' health and safety, detect all pregnancy complications, take necessary actions, respond to complaints, prepare for birth, and promote a healthy lifestyle. Antenatal care is essential to detect and prevent all unexpected conditions during pregnancy.^{5,6} ANC should be conducted at least four times throughout pregnancy, according to the Indonesia Ministry of Health: once in the first trimester, once in the second trimester, and twice in the third trimester.⁷⁻⁹ The 2018 Indonesian Basic Health Survey reported 86.0% of the first ANC visit in Indonesia. Meanwhile, the percentage of the fourth ANC visit was 74.1%.¹⁰ More than a quarter of pregnant women in Indonesia have not optimally done an ANC visit.

In the early 2000 period, the government issued a "Suami Siaga" policy (literally known as "Alert Husband"). The policy release addresses the high MMR in Indonesia. "Suami Siaga" is a national campaign to promote husbands' participation in Indonesia's maternal and child health programs. The previous evaluation in 2012 showed that husbands' involvement was proven beneficial in increasing ANC's involvement in Indonesia.¹¹

In many countries, including Indonesia, males or husbands play as the primary decision-makers in the household.¹² They commonly decide to allow wives or women access to maternal care services.¹³ The majority of husbands supported their partners to allocate funds for maternal care services but were less involved in labor preparedness and postnatal care.¹⁴

Several studies conducted in various countries showed that one of the ANC barriers was poor husbands' support and involvement in the antenatal and labor processes. The poor of husbands' permission is why women delay maternal health care.¹⁵⁻¹⁸ Some research in Lao People's Democratic Republic (Lao PDR) and Bangladesh found that husband involvement is the determinant of a continuum for maternal care. Husbands' support in accompanying their wives to receive health services positively correlates with maternal and newborn health services.^{13,19}

Indonesia recently has thousands of cultures that believe in their local medical knowledge. Local medical knowledge is sometimes contradictory to modern medical knowledge.²⁰ Pregnancy and childbirth are always judged as domestic affairs, specifically women's experiences. In the Papua Muyu tribes, pregnant women must be exiled to a separate hut far from home because blood affects families' health after childbirth.²¹ It is different from the Gayo female tribes in Aceh, who felt ashamed and scared during pregnancy. Gayo women must hide their pregnancy because they were afraid of spiritual fetal kidnapping by evils as her stomach suddenly can be deflated by itself.²²

Husband's participation is essential in encouraging their wives to utilize maternal care services. Therefore, research on husbands' involvement in maternal care is required. This study aimed to analyze the aspects of the husband's participation in the wife's antenatal care visits in Indonesia. Based on the background narration, what factors can affect the husband's involvement in ANC in Indonesia?

Materials and methods

Study design

The study was a cross-sectional study that used secondary data from the 2017 Indonesian Demographic and Health Survey (IDHS) (Female file: IDIR71FL_Individual Recode). The 2017 IDHS utilized stratified two-stage sampling. Stage 1: Using a systematic proportionate to size probability with the number of households from the 2010 population census listing, choose various census blocks. The survey utilizes an implicit stratification technique based on urban and rural areas to sort census blocks depending on their wealth.²³

Setting

To address the high MMR, Indonesian public health Institution actively supported (since the beginning of the 2000s) the involvement of husbands in the ANC through a national campaign called "Suami Siaga."¹¹

Participants

The study population was women of childbearing age, aged 15–49 years old, and pregnant in the last 5 years before the interview. The study employed 14,319 respondents as the samples.

Outcome variables

The study used husbands' involvement (Yes or Not) in ANC as the binary outcome variable. The related survey question was "Does husband accompany his wife in ANC visit?"

Independent variables

We focused on husband characteristics to avoid collinearity problems and according to paper aims. The explanatory variables were: residence (urban or rural), husband age (as continuous), an education level (no education, primary, secondary, and higher education), occupation (unemployed, professional/technical, managers/administrators, clerical, sales, services, agricultural self-employed, industrial workers, other), wealth status (poorest, poorer, middle, richer, richest), and parity (primiparous = 1, multiparous = 2–4, grand multiparous > 4).




Figure 1. Distribution map of the percentage of husband's involvement in antenatal care by the province. Indonesia, 2017.

The urban or rural place was based on the Central Bureau of Statistics requirement. The husband's age is the respondents' acknowledgment of their husbands' last birthday. Primary education refers to junior high school, secondary to senior high school, and higher to tertiary education. The survey assessed wealth status through types of furniture and their prices. It counts a television, a bicycle or a car, and household goods, such as drinking water sources, bathroom amenities, and building materials for flooring—the study calculated assessment of this variable with the principal component analysis. National wealth quintiles were arranged based on household scores for each household and then divided into the same five categories, each contributing 20% of the population based on the distribution.²⁴

Data analysis

The first step used chi-square to select the categorical variables and the T-test to choose the continuous ones. Both tests identified the relationship between the independent variables and the husbands' involvement in ANC as the dependent variable. This study employed Binary Logistic Regression to determine predictors because of the dependent variable's nature. The research performed all statistical analyzes with IBM SPSS Statistics 21 software.

In addition, the study used ArcGIS 10.3 (ESRI Inc., Redlands, CA, USA) to produce a distribution map of the percentage of husbands' involvement in antenatal care by the province in Indonesia. The Indonesian Bureau of

Statistics provided a shapefile of administrative border polygons for the study.

Ethical approval

The study used secondary data from the 2017 IDHS for its materials analysis. The 2017 IDHS removed all respondents' identities from the dataset. Respondents signed written consent forms to participate in this study, and children's parents or guardians signed consent forms (under 16 years). For this study, the author obtained permission to use data from the website <https://dhsprogram.com>.

The 2017 IDHS follows the Standard DHS survey methodology established by ICF International's Institutional Review Board under The Demographic and Health Surveys (DHS) Program (DHS-7) and evaluated and approved by the ORC Macro IRB in 2002. The Institutional Review Board of ICF International follows the US Department of Health and Human Services' "Protection of Human Subjects" requirements (45 CFR 46). DHS surveys that meet the criteria are labeled DHS-7 Program authorized, and the paperwork for approval is included.

Results

This study presents the distribution map of the portion of husband's participation in antenatal care by the province in Indonesia in Figure 1. The figure shows a random distribution pattern. The analysis results show that nationally, the

Table 1. Descriptive statistic of husband's involvement in ANC ($n = 14,319$). Indonesia, 2017.

Characteristics	<i>n</i>	Husband's involved in ANC visit (%)		<i>p</i> -value
		No	Yes	
Type of place of residence				<0.001*
Urban	7154	16.5%	83.5%	
Rural	7165	31.6%	68.4%	
Age (mean)	14,319	(36.05)	(34.71)	<0.001*
Education level				<0.001*
No education	204	51.0%	49.0%	
Primary	3703	38.0%	62.0%	
Secondary	8239	20.9%	79.1%	
Higher	2173	9.8%	90.2%	
Occupation type				<0.001*
Not work	111	36.0%	64.0%	
Professional/technical	1186	15.6%	84.4%	
Managers and administration	262	13.7%	86.3%	
Clerical	981	12.9%	87.1%	
Sales	1931	17.4%	82.6%	
Services	2536	21.0%	79.0%	
Agricultural—self employed	3539	37.2%	62.8%	
Industrial worker	3594	23.3%	76.7%	
Other	179	20.7%	79.3%	
Wealth status				<0.001*
Poorest	3568	45.4%	54.6%	
Poorer	2846	25.8%	74.2%	
Middle	2740	18.1%	81.9%	
Richer	2647	14.8%	85.2%	
Richest	2518	8.1%	91.9%	
Parity of wife				<0.001*
Primiparous	4403	16.6%	83.4%	
Multiparous	8904	25.0%	75.0%	
Grandemultiparous	1012	48.2%	51.8%	

* $p < 0.001$.

average percentage of husbands' involvement in antenatal care is 75.927%. Meanwhile, as the center of government, the island of Java offers husband's participation in antenatal care is better than in other regions.

Table 1 shows the descriptive statistics of the husband's involvement in ANC and other related variables. The study discovered that husbands involved in ANC visits were dominant in both residence types: urban and rural areas. Meanwhile, the average age of husbands involved in ANC visits was slightly younger than those not engaged in ANC.

On the other hand, husbands of all education levels are dominantly involved in ANC visits. Except for the no education husband was not engaged in ANC visits. Moreover, husbands with any work type, wealth status, and parity are dominantly involved in ANC visits.

Table 2 displays co-linearity test results of the husband's involvement in ANC and other related variables. It shows that all variables' tolerance value, between the dependent and independent variables, is more significant

than 0.10, while the variance inflation factor (VIF) value is less than 10.00. It indicated there are no multicollinearity symptoms in the regression model.

The binary logistic regression test results were employed to determine variables influencing the husband's ANC involvement in Indonesia. Table 3 shows that husbands who lived in urban areas have odds of being involved in ANC visits 1.188 times (95% CI 1.078–1.308) more likely than those in rural areas. Meanwhile, the study found husband's age did not correlate with the wife's ANC in Indonesia.

Results found husbands who had primary education have odds of being involved in ANC visits 1.357 times (95% CI 1.004–1.834) more likely than husbands with no education. Husbands with secondary education have odds of being involved in ANC visits 1.967 times (95% CI 1.456–2.657), more likely than those with no education. Husbands with higher education have odds of being involved in ANC visits 3.264 times (95% CI 2.320–4.593) more likely than husbands with no education.

Table 2. Results for the co-linearity test of husband's involvement in ANC ($n = 14,319$). Indonesia, 2017.

Variables	Collinearity statistics	
	Tolerance	VIF
Type of place of residence	0.673	1.485
Age	0.697	1.435
Education level	0.624	1.603
Occupation type	0.945	1.058
Wealth status	0.601	1.664
Parity	0.728	1.374

*Note: Dependent variable: husband's involvement in ANC.

Table 3. Binary logistic regression of husband's involvement in ANC ($n = 14,319$). Indonesia, 2017.

Predictors	Husband's involved in ANC visit ($n = 14,319$)			p-value
	AOR	95% CI		
		Lower bound	Upper bound	
Type of place of residence				
Urban	1.19	1.08	1.31	<0.001***
Rural	—	—	—	—
Age	0.99	0.99	1.00	0.102
Education level				
No education	—	—	—	—
Primary	1.36	1.00	1.83	0.047*
Secondary	1.97	1.46	2.66	<0.001***
Higher	3.26	2.32	4.59	<0.001***
Occupation				
Did not work	—	—	—	—
Professional/technical	1.91	1.22	3.01	0.005**
Managerial and administration	2.02	1.15	3.55	0.014*
Clerical	2.14	1.35	3.41	0.001**
Sales	2.28	1.47	3.53	<0.001***
Services	2.20	1.42	3.39	<0.001***
Agricultural—self employed	1.89	1.23	2.90	0.004**
Industrial worker	2.14	1.40	3.29	0.001**
Other	1.62	0.92	2.86	0.097
Wealth status				
Poorest	—	—	—	—
Poorer	2.00	1.79	2.24	<0.001***
Middle	2.84	2.50	3.23	<0.001***
Richer	3.24	2.81	3.74	<0.001***
Richest	5.42	4.52	6.51	<0.001***
Parity				
Primiparous	—	—	—	—
Multiparous	0.64	0.58	0.71	<0.001***
Grand multiparous	0.34	0.28	0.41	<0.001***

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Based on occupation type, professional/technical workers have odds of being involved in ANC visits 1.914 times (95% CI 1.217–3.010) more likely than unemployed husbands. Husbands who worked as managers and administrators have odds of being involved in ANC visits 2.023 times (95% CI 1.151–3.554), a higher chance than unemployed husbands. Clericals have odds of being engaged in ANC visits 2.142 times (95% CI 1.346–3.408), more likely than husbands who did not work. Husbands who worked as a salesman have odds of being involved in ANC visits 2.227 times (95% CI 1.468–3.532) more likely than those who did not work. Service providers have odds of being engaged in ANC visits 2.196 times (95% CI 1.424–3.385) more chances than unemployed husbands. Husbands who worked in agriculture have odds of being involved in ANC visits 1.887 times (95% CI 1.229–2.899) more likely than husbands who did not work. Besides, husbands who were industrial workers have odds of being involved in ANC visits 2.139 times (95% CI 1.393–3.285) more likely than husbands who had no job. These findings showed that husbands with employment types listed in the table were more likely to be involved in ANC. However, husbands who had other jobs did not differ from husbands who did not work. Based on the analysis results, husbands who work as salesmen have the highest probability of being involved in ANC visits.

Based on wealth status, Table 3 further illustrates that husbands in lower wealth status have odds of being involved in ANC visits 2.003 times (95% CI 1.789–2.243) more likely than the weakest groups of husbands. Those with median wealth status have odds of being involved in ANC visits 2.840 times (95% CI 2.499–3.229) more chances than the lowest groups of husbands. Wealthier husbands have odds of being engaged in ANC visits 3.242 times (95% CI 2.811–3.740), more likely than the lowest groups of husbands. Husbands with the most prosperous wealth status have odds of being involved in ANC visits 5.424 times (95% CI 4.516–6.514) the chances than the lowest groups of husbands. It shows that the more prosperous the husbands' wealth status is, the more they get involved in ANC.

Table 3 informs that husbands who had multiparous wives have odds of being involved in ANC visits 0.641 times (95% CI 0.575–0.714) less likely than those who had primiparous wives. Husbands with grand multiparous wives have odds of being involved in ANC visits 0.341 times (95% CI 0.284–0.408) fewer chances than husbands who had primiparous wives. It suggests that the more children wives had ever been born, the more passive husbands' involvement in ANC was.

Discussion

Husbands' support for their partners is a good practice during married life. During pregnancy and childbirth,

husbands who support their partners are considered modern men at home and handle tasks apart from their usual duties. It allows their wives to have enough rest during pregnancy.²⁵ Husbands' active involvement was strongly correlated with alertness and increased wives' knowledge of newborn danger signs.²⁶ However, a literature review with a narrative approach to husbands' feelings found a different condition. Although husbands want to get involved, support their wives, and learn about fetal health, they often feel excluded from making perinatal screening decisions.²⁷ It signals that awareness of husbands' involvement and maternal care providers need to be improved.

Husbands' involvement in ANC in Indonesia is more concentrated in Java as the government center. This situation indicates that the implementation of the "*Suami Siaga*" (Alert Husband) policy which seeks to increase husbands' involvement in the ANC, has not been spread evenly across all regions in Indonesia. The government still has to work harder to disseminate the policy.

Furthermore, the study finds five variables associated with husbands' involvement in ANC in Indonesia: (1) the type of residence, (2) education level, (3) types of work, (4) wealth status, (5) parity.

- (1) Husbands who lived in urban areas were more likely to be involved in ANC visits than rural areas. The Indonesian community tends to have more open access in urban areas, including information and health services.²⁸⁻³⁰ Regions with dominant urban regions have better ANC utilization.^{8,31,32}
- (2) The better husbands' education level was, the more they got involved in ANC. This finding is similar to Lao PDR and Pakistan's research conclusion, informing that husbands with higher education levels correspond to their involvement in the maternal process and newborn and child health services.^{19,33} In line with these findings, another study found husbands who had no formal education were 0.02 times more likely to decide maternal services for wives than those who attended formal education.³⁴ A study in Nigeria discovered that husbands' involvement in maternity and delivery processes was minimal. In addition to the low educational level, this was also due to prevailing social norms. Nigerian husbands assumed that pregnancy and childbirth were under women's business.¹⁶ Several previous studies that took other health fields' themes found education a positive determinant,³⁵⁻³⁹ while poor education hinders achieving better output.⁴⁰⁻⁴² A slightly different finding was informed in a study in the Oromia Region-Ethiopia, telling that many husbands know obstetric danger signs. On the other

hand, the study found that the husband had low participation. The situation indicates their low willingness to donate blood during their wife's antenatal, delivery, and postpartum.¹⁷

- (3) Husbands who were employed in jobs listed were more likely to be involved in ANC, while those who had no job and more time are not a determinant of their involvement in the maternal process. Other studies in Northern Nigeria also found husbands' occupation proves one of the determinants of women's participation in maternal services. Husbands' work determines the affordability to pay healthcare costs.⁴³
- (4) The better husband's wealth status was, the more involvement in ANC was. Previous studies in Pakistan, Eritrea, and Ethiopia similarly revealed that the better husbands' wealth status was, the more they got involved in ANC.^{33,34,44} Encouraging husbands with good wealth status could reduce delayed ANC visits.⁴⁴ Several previous studies have informed that wealth status is a positive determinant of better health sector outcomes. Conversely, poor wealth status is often associated with lower effects.^{45,46}
- (5) Husbands who had wives with more births participated less in ANC. Husbands got involved more actively when their wives gave birth to their first child. In short, parity becomes one of the determinants in husbands' involvement in maternal care services. These findings were similar to the previous study in Yangon, Myanmar. Husbands with high parity wives tend to have low participation in the ANC. In addition to parity, some other variables as determinants were educational level, marriage status, decision-making level in the family, exposure to maternal health education, and perception of risk during pregnancy and childbirth.¹⁴

Policymakers in Indonesia should consider the findings of this research while planning their policies; they may help accelerate the husband's involvement in antenatal care visits.

Strength and limitation

The study's strength is the use of big data as material for analysis. However, this study also has limitations. The study was conducted with a quantitative approach not to cover the phenomena of local Indonesian values and culture that could potentially influence husbands' involvement in their wives' ANC, for example, concerning the patriarchal social system, the value of children, the value of family, and the value of pregnancy.^{22,47}

Acknowledgements

The author would like to thank the Inner City Fund for allowing the use of the 2017 IDHS.

Contributions

ADL designed study concepts, prepared data, conducted data analysis and wrote manuscript drafts. RDW analyzed results, wrote manuscript drafts, and finalized manuscripts. NEWS and SS conducted data analysis, wrote draft manuscripts, and finalized the manuscripts.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

Significance for public health

In the early 2000s, the Indonesian government launched a national campaign (Suami Siaga) to encourage husbands to participate in maternal and child health programs in order to reduce maternal morbidity and mortality. The findings of this study could be used to send specific messages to the 25% of Indonesian husbands who are not yet actively participating in pregnancy by highlighting the socio-demographic predictors of husbands' involvement in antenatal care.

References

1. Data and Information Center Ministry of Health. Mother's day: maternal health situation [Internet]. Jakarta, <http://www.kemkes.go.id/download.php?file=download/pusdatin/infodatin/infodatin-ibu.pdf> (2014, accessed 20 September 2021).
2. Susiana S. Maternal mortality rate: causes and efforts to address it. *Br INFO Br Study Actual Strateg Issues* 2015; XI(24): 13–18.
3. World Health Organization. Trends in maternal mortality: 1990 to 2015: estimates by WHO, UNICEF, UNFPA, World Bank Group and the United Nations Population Division [Internet]. Geneva, https://apps.who.int/iris/bitstream/handle/10665/194254/9789241565141_eng.pdf;jsessionid=AB201B62E0913D576E5CF9430F900F98?sequence=1 (2015, accessed 2 September 2021).
4. Achadi EL. Maternal and Neonatal Death in Indonesia (Kematian Maternal dan Neonatal di Indonesia) [Internet]. Rakerkernas. Jakarta, <https://docplayer.info/116404935-Kematian-maternal-dan-neonatal-di-indonesia-endang-l-achadi-fakultas-kesehatan-masyarakat-universitas-indonesia.html> (2019, accessed 2 September 2021).
5. Masruroh YUSUFA, Rohmah N, Pakki IB, et al. Neonatal death incidence in healthcare facility in Indonesia: does antenatal care matter? *Indian J Forensic Med Toxicol* 2021; 15(1): 1265–1271.
6. Hijazi HH, Alyahya MS, Sindiani AM, et al. Determinants of antenatal care attendance among women residing in highly disadvantaged communities in northern Jordan: a cross-sectional study. *Reprod Health* 2018; 15(1): 106.
7. Wulandari RD, Putri NK and Laksono AD. Socioeconomic disparities in antenatal care utilisation in urban Indonesia. *Int J Innov Creat Chang* 2020; 14(2): 498–514.
8. Laksono AD, Rukmini R and Wulandari RD. Regional disparities in antenatal care utilization in Indonesia. *PLoS One* 2020; 15(2): e0224006.
9. Denny HM, Laksono AD, Matahari R, et al. The determinants of four or more antenatal care visits among working women in Indonesia. *Asia Pac J Public Health* 2022; 34(1): 51–56.
10. National Institute of Health Research and Development of The Indonesia Ministry of Health. The 2018 Indonesia Basic Health Survey (Riskesdas): National Report [Internet]. Jakarta, https://kesmas.kemkes.go.id/assets/upload/dir_519d41d8cd98f00/files/Hasil-riskesdas-2018_1274.pdf (2019, accessed 2 September 2021).
11. Kurniati A, Chen C-M, Efendi F, et al. Suami SIAGA: male engagement in maternal health in Indonesia. *Health Policy Plan* 2017; 32(8): 1203–1211.
12. Megatsari H, Laksono AD, Herwanto YT, et al. Does husband/partner matter in reduce women's risk of worries? Study of psychosocial burden of covid-19 in Indonesia. *Indian J Forensic Med Toxicol* 2021; 15(1): 1101–1106.
13. Rahman AE, Perkins J, Islam S, et al. Knowledge and involvement of husbands in maternal and newborn health in rural Bangladesh. *BMC Pregnancy Childbirth* 2018; 18(1): 247.
14. Wai KM, Shibanuma A, Oo NN, et al. Are husbands involving in their spouses' utilization of maternal care services? A cross-sectional study in Yangon, Myanmar. *PLoS One* 2015; 10(12): e0144135.
15. Sumankuuro J, Mahama MY, Crockett J, et al. Narratives on why pregnant women delay seeking maternal health care during delivery and obstetric complications in rural Ghana. *BMC Pregnancy Childbirth* 2019; 19(1): 260.
16. Sharma V, Leight J, Giroux N, et al. "That's a woman's problem": A qualitative analysis to understand male involvement in maternal and newborn health in Jigawa state, northern Nigeria. *Reprod Health* 2019; 16(1): 143.
17. Gize A, Eyassu A, Nigatu B, et al. Men's knowledge and involvement on obstetric danger signs, birth preparedness and complication readiness in Burayu town, Oromia region, Ethiopia. *BMC Pregnancy Childbirth* 2019; 19(1): 515.
18. Uldbjerg CS, Schramm S, Kaducu FO, et al. Perceived barriers to utilization of antenatal care services in northern Uganda: a qualitative study. *Sex Reprod Healthc* 2020; 23: 100464.
19. Sakuma S, Yasuoka J, Phongluxa K, et al. Determinants of continuum of care for maternal, newborn, and child health services in rural Khammouane, Lao PDR. *PLoS One* 2019; 14(4): e0215635.
20. Laksono AD, Wulandari RD, Nantabah ZK, et al. The concept of illness among ethnic groups in Indonesia: a meta-ethnographic study. *Syst Rev Pharm* 2020; 11(9): 584–591.
21. Laksono AD, Soerachman R and Angkasawati TJ. Case Study of Muyu Ethnic's Maternal Health in Mindiptara District-Boven Digoel (Studi Kasus Kesehatan Maternal

- Suku Muyu di Distrik Mindiptana, Kabupaten Boven Digoel). *J Reprod Heal* 2016; 07(3): 145–155.
22. Pratiwi NL, Fitrianti Y, Nuraini S, et al. Concealed pregnant women or Kemel of Gayo ethnic in Blang Pegayon District, Gayo Lues District, Aceh. *Bull Heal Syst Res* 2019; 22(2): 81–90.
 23. National Population and Family Planning Board. Statistics Indonesia, Ministry of Health, The DHS Program. The 2017 Indonesia Demographic and Health Survey [Internet]. Jakarta, <https://www.dhsprogram.com/pubs/pdf/FR342/FR342.pdf> (2018, accessed 3 September 2021).
 24. Wulandari RD, Qomarrudin MB, Supriyanto S, et al. Socioeconomic disparities in hospital utilization among elderly people in Indonesia. *Indian J Public Health Res Dev* 2019; 10(11): 2192.
 25. Kashaia DK, Mselle LT and Mkoka DA. Husbands' experience and perception of supporting their wives during childbirth in Tanzania. *BMC Pregnancy Childbirth* 2020; 20(1): 85.
 26. Zaman SB, Gupta RD, Al Kibria GM, et al. Husband's involvement with mother's awareness and knowledge of newborn danger signs in facility-based childbirth settings: a cross-sectional study from rural Bangladesh. *BMC Res Notes* 2018; 11(1): 286.
 27. Cheng ER, McGough H and Tucker Edmonds B. Paternal preferences, perspectives, and involvement in perinatal decision making. *Obstet Gynecol Surv* 2019; 74(3): 170–177.
 28. Laksono AD, Wulandari RD and Soedirham O. Urban and rural disparities in hospital utilization among Indonesian adults. *Iran J Public Health* 2019; 48(2): 247–255.
 29. Wulandari RD and Laksono AD. Urban-rural disparity: the utilization of primary health care center among elderly in East Java, Indonesia. *J Adm Kesehat Indones* 2019; 7(2): 147–154.
 30. Seran AA, Laksono AD, Sujoso ADP, et al. Does contraception used better in urban areas? An analysis of the 2017 IDHS (Indonesia Demographic and Health Survey). *Syst Rev Pharm* 2020; 11(11): 1892–1897.
 31. Laksono AD, Wulandari RD and Soedirham O. Regional disparities of health center utilization in rural Indonesia. *Malaysian J Public Heal Med* 2019; 19(1): 158–166.
 32. Wulandari RD, Laksono AD and Rohmah N. Urban-rural disparities of antenatal care in South East Asia: a case study in the Philippines and Indonesia. *BMC Public Health* 2021; 21(1): 1221.
 33. Maken ZH, Nasir Idrees I, Zahid A, et al. Factors influencing father's antenatal and perinatal involvement in maternal health care. *J Matern Fetal Neonatal Med* 2018; 31(19): 2569–2575.
 34. Kifle MM, Kesete HF, Gaim HT, et al. Health facility or home delivery? Factors influencing the choice of delivery place among mothers living in rural communities of Eritrea. *J Health Popul Nutr* 2018; 37(1): 22.
 35. Megatsari H, Laksono AD, Ibad M, et al. The community psychosocial burden during the COVID-19 pandemic in Indonesia. *Heliyon* 2020; 6(10): e05136.
 36. Ipa M, Widawati M, Laksono AD, et al. Variation of preventive practices and its association with malaria infection in eastern Indonesia: findings from community-based survey. *PLoS One* 2020; 15(5): e0232909.
 37. Wulandari RD and Laksono AD. Education as predictor of the knowledge of pregnancy danger signs in Rural Indonesia. *Int J Innov Creat Chang* 2020; 13(1): 1037–1051.
 38. Kusriani I and Laksono AD. Regional disparities of stunted toddler in Indonesia. *Indian J Forensic Med Toxicol* 2020; 14(3): 1685–1691.
 39. Seran AA, Antaria MD, Haksama S, et al. Disparities of the use of hormonal and non-hormonal contraceptive drugs in urban and rural areas in Indonesia and the world. *Syst Rev Pharm* 2020; 11(9): 66–73.
 40. Laksono AD and Wulandari RD. The barrier to Maternity Care in rural Indonesia. *J Public Health* 2022; 30(1): 135–140.
 41. Rohmah N, Yusuf A, Hargono R, et al. Determinants of teenage pregnancy in Indonesia. *Indian J Forensic Med Toxicol* 2020; 14(3): 2080–2085.
 42. Andayani Q, Koesbardiati T, Prahastuti AD, et al. The barrier to access health insurance for maternity care: case study of female workers in Indonesia. *Med Leg Update* 2021; 21(2): 926–932.
 43. Adewemimo AW, Msuya SE, Olaniyan CT, et al. Utilisation of skilled birth attendance in Northern Nigeria: a cross-sectional survey. *Midwifery* 2014; 30(1): e7–e13.
 44. Tesfaye G, Loxton D, Chojenta C, et al. Delayed initiation of antenatal care and associated factors in Ethiopia: a systematic review and meta-analysis. *Reprod Health* 2017; 14(1): 150.
 45. Laksono AD, Wulandari RD, Ibad M, et al. The effects of mother's education on achieving exclusive breastfeeding in Indonesia. *BMC Public Health* 2021; 21(1): 14.
 46. Wulandari RD and Laksono AD. Determinants of knowledge of pregnancy danger signs in Indonesia. *PLoS One* 2020; 15(5): e0232550.
 47. Sakina AI and A DHS. Highlighting patriarchal culture in Indonesia (Menyoroti Budaya Patriarki di Indonesia). *Share Soc Work J* 2017; 7(1): 71–80.