

FABAD

**JOURNAL of
PHARMACEUTICAL
SCIENCES**

ISSN 1300-4182
e-ISSN: 2651-4648
www.fabad.org.tr

Volume: 46 • Issue: 1 • March 2021

An Official Journal of The Society of Pharmaceutical Sciences of Ankara (FABAD)

Author Checklist (<https://dergi.fabad.org.tr/en/yazar-kontrol-listesi/>)

Sample Issue (<https://dergi.fabad.org.tr/en/ornek-yayinlar/>)

Archive (<https://dergi.fabad.org.tr/en/archive/>)

☎ +90 (312) 305 30 17 (tel:+90 (312) 305 30 17)

✉ fabadankara@gmail.com (mailto:fabadankara@gmail.com)

Editorial Board

Publisher

Ayşegül KÖROĞLU (Ankara University, Department of Pharmaceutical Botany, Ankara, Turkey)

Editor in Chief

Nesrin GÖKHAN KELEKÇİ (Hacettepe University, Department of Pharmaceutical Chemistry, Ankara, Turkey)

Co-Editors

Selen ALP

(Ankara University, Department of Pharmaceutical Chemistry, Ankara, Turkey)

Fatma Sezer ŞENOL

(Gazi University, Department of Pharmacognosy, Ankara, Turkey)

DENİZ

(Gazi University, Department of Pharmaceutical Technology, Ankara, Turkey)

Sibel İLBASMIŞ TAMER

Technical Editors

Gökçen TELLİ

(Hacettepe University, Department of Pharmacology, Ankara, Turkey)

Vahap Murat KUTLUAY

(Hacettepe University, Department of Pharmacognosy, Ankara, Turkey)

Biostatistics Editor

Hatice Yağmur Zengin

Hacettepe University, Faculty of Medicine, Department of Biostatistics, Ankara, Turkey

Language Edition: The compliance of the language with the spelling rules is evaluated by the field editors and technical editors through the Grammarly program.

Editorial Board

- Almira RAMANAČIENĖ** Vilnius University, Nanotechnology and Material Sciences Center, Vilnius, Latvia
- Alper GÖKBULUT** Ankara University, Department of Pharmacognosy, Ankara, Turkey
- Ashok K. SHAKYA** Al Ahliyya Amman University, Department of Pharmaceutical Chemistry, Amman, Jordan
- Aygin EKİNCİOĞLU** Hacettepe University, Department of Clinical Pharmacy, Ankara, Turkey
- Ayşe KURUÜZÜM UZ** Hacettepe University, Department of Pharmacognosy, Ankara, Turkey
- Bharat JHNAWAR** Lovely Professional University, Pharmaceutical Sciences, Punjab, India
- Bülent KIRAN** Ege University, Department of Pharmacy Management, İzmir, Turkey
- Ceyda Tuba ŞENGEL** Ankara University, Department of Pharmaceutical Technology, Ankara, Turkey
- TÜRK** Chung Yuan Christian University, Biomedical Engineering, Taoyuan, Taiwan
- Chia-Yi TSENG** Gazi University, Department of Pharmacognosy, Ankara, Turkey
- Didem DELİORMAN** Ege University, Department of Pharmaceutical Technology, Izmir, Turkey
- ORHAN** Ankara University, Department of Biochemistry, Ankara, Turkey
- Emel Öykü ÇETİN** G. D'Annunzio" University, Department of Pharmaceutical Chemistry, Chieti-Pescara, Italy
- UYANIKGİL** University of Burgundy, French Institute for Medical and Health Research, Dijon, France
- Filiz BAKAR ATEŞ** Istanbul University, Department of Pharmaceutical Chemistry, Ankara, Turkey
- Francesco EPIFANO** Gazi University, Department of Pharmaceutical Chemistry, Ankara, Turkey
- Gerard LIZARD** Ege University, Department of Pharmaceutical Toxicology, Izmir, Turkey
- Gökçe CİHAN ÜSTÜNDAĞ** King Saud University, Department of Pharmacognosy, Riyadh, Saudi Arabia
- Gökçen EREN** Yeditepe University, Department of Pharmacognosy, Istanbul, Turkey
- Hande GÜRER ORHAN** University of Mississippi, National Center for Natural Product Research, USA.
- Hasan Abougazar** Ankara University, Department of Pharmacology, Ankara, Turkey
- YUSUFOĞLU** Karadeniz Technical University, Department of Pharmaceutical Chemistry, Trabzon, Turkey
- Hasan KIRMIZİBEKMEZ** Anadolu University, Department of Pharmaceutical Chemistry, Eskisehir, Turkey
- İkhlas KHAN** Hacettepe University, Department of Pharmacology, Ankara, Turkey
- Işıl ÖZAKCA GÜNDÜZ** Ankara University, Department of Pharmaceutical Chemistry, Ankara, Turkey
- İnci Selin DOĞAN** University of Health Sciences, Department of Pharmaceutical Toxicology, Ankara, Turkey
- Leyla YURTTAŞ** Afyonkarahisar University of Health Sciences, Department of Pharmaceutical Toxicology, Afyonkarahisar, Turkey
- Melike H. ÖZKAN** Marmara University, Department of Clinical Pharmacy, Istanbul, Turkey
- Meltem ÜNLÜSOY** Chung Yuan Christian University, Department of Bioscience Technology, Taoyuan, Taiwan
- Merve BACANLI** Afyonkarahisar University of Health Sciences, Department of Pharmaceutical Technology, Afyonkarahisar, Turkey
- Merve BECİT** University of Messina, Department of Chemistry and Biology, Messina, Italy
- Mesut SANCAR** Istanbul University, Department of Pharmacy Management, Ankara, Turkey
- Ming-Wei CHAO** Hacettepe University, Department of Pharmaceutical Technology, Ankara, Turkey
- Muharrem ÖLÇER** Hacettepe University, Department of Pharmaceutical Toxicology, Ankara, Turkey
- Natalizia MICELI** The Assam Kaziranga University, Department of Pharmaceutical Sciences, Assam, India
- Özlem Nazan ERDOĞAN** Anadolu University, Department of Pharmaceutical Microbiology, Ankara, Turkey
- Sevda ŞENEL** Gazi University, Department of Pharmaceutical Technology, Ankara, Turkey
- Sevtap AYDIN DİLSİZ** Hacettepe University, Department of Biochemistry, Ankara, Turkey
- Suryakanta SWAIN** Marmara University, Department of Pharmaceutical Toxicology, Istanbul, Turkey
- Şükrü BEYDEMİR** Gazi University, Department of Analytical Chemistry, Eskisehir, Turkey
- Tuba İNCEÇAYIR** Hanoi University of Pharmacy, Department of Analytical Chemistry and Toxicology, Hanoi, Vietnam
- Tuğba TÜYLÜ** University of Graz Institute of Pharmaceutical Sciences, Department of Pharmacognosy, Graz, Austria
- KÜÇÜKKILINÇ**
- Tuğçe YEŞİL**

FABAD | Journal of Pharmaceutical Sciences© All right reserved 2021

Read More by Acme Themes (<http://www.acmethemes.com/>)

Author Checklist (<https://dergi.fabad.org.tr/en/yazar-kontrol-listesi/>)

Sample Issue (<https://dergi.fabad.org.tr/en/ornek-yayinlar/>) Archive (<https://dergi.fabad.org.tr/en/archive/>)

+90 (312) 305 30 17 (tel:+90 (312) 305 30 17) ✉ fabadankara@gmail.com (<mailto:fabadankara@gmail.com>)

2020 Volume 45, Issue 3

This issue is published electronically on December 2020

CONTENTS

Research Articles

Anticancer Property of Orthosiphon stamineus Benth. Extracts in Different Solvent Systems against T47D Human Breast Cancer Cell Lines

Lusiana ARIFIANTI* , Sukardiman SUKARDIMAN** , Niken INDRIYANTI*** , Retno WIDYOWATI****°

* ORCID:0000-0003-1704-7908, Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Universitas Airlangga, Indonesia

** ORCID:0000-0001-9689-5088, Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Universitas Airlangga, Indonesia

*** ORCID:0000-0003-3733-5018, Departement of Pharmacology, Faculty of Pharmacy, Mulawarman University, Indonesia

**** ORCID:0000-0003-0572-7551, Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Universitas Airlangga, Indonesia

° Corresponding Author; Retno Widyowati

Tel. +6231 5933150, Fax. +6231 5935249, e-mail: rr-retno-w@ff.unair.ac.id

Pages 187-194, **Abstract | PDF (628 KB)**

<http://dergi.fabad.org.tr/pdf/volum45/Issue3/187-194.pdf>



(https://dergi.fabad.org.tr/wp-content/uploads/2021/06/FABAD_46-1_Kapak.png)

ISSN: 1300-4182
e-ISSN: 2651-4648



(<https://www.fabad.org.tr/en/>)

Click for the website of the FABAD Society (The Ordinary General Assembly of the Society of Pharmaceutical Sciences of Ankara)

Archive Search

For Authors

Evaluation and Refereeing

Process

(<https://dergi.fabad.org.tr/en/evaluation-and-refereeing-process/>)

Sample Issue and Templates

(<https://dergi.fabad.org.tr/en/ornek-yayinlar/>)

Checklist

(<https://dergi.fabad.org.tr/en/yazar-kontrol-listesi/>)

Copyright Release Form

(<https://dergi.fabad.org.tr/en/yayin-hakki-devir-formu/>)

Article Submission

(<https://dergipark.org.tr/tr/journal/3647/submission/step/manuscript/new>)

Archive

(<https://dergi.fabad.org.tr/en/archive/>)

Issue List

2023 Cilt 48, Sayı 1

(<https://dergi.fabad.org.tr/en/2023-cilt-48-sayi-1/>)

2022 Volume 47, Issue 3

(<https://dergi.fabad.org.tr/en/2022-volume-47-issue-3/>)

2022 Volume 47, Issue 2

(<https://dergi.fabad.org.tr/en/2022-cilt-47-sayi-2/>)

Cytotoxic Effects of Methanolic Extract of *Ziziphora tenuior* L. on the Growth of the Lung Cancer Cell Line

Christine BATENI* , Mohammad TAHERI** , Masomeh BORBOR*** , Sima NASRI****°

* ORCID: 0000-0003-1325-7836, Department of Biology, Payame Noor University, Tehran, Iran

** ORCID: 0000-0002-9510-5125, Department of Biotechnology, Research Laboratory, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran

*** ORCID: 0000-0003-0922-3510, Department of Biology, Payame Noor University, Tehran, Iran

**** ORCID: 0000-0001-7849-6315, Department of Biology, Payame Noor University, Tehran, Iran

° Corresponding Author: Sima Nasri

Phone number: 0098 2177312752, e-mail: s_nasri@pnu.ac.ir

Pages 195-204, [Abstract](#) | [PDF \(249 KB\)](#)

(<http://dergi.fabad.org.tr/pdf/volum45/issue3/195-204.pdf>)

Synthesis and Anti-Inflammatory Activity of Some Novel Quinazolinone Derivatives

Neha KRISHNARTH*° , Santosh Kumar VERMA* , Anurag Chaudhary***

* ORCID:0000-0002-2137-2028, Faculty of Pharmaceutical Sciences, Motherhood University, Vill. Karoundi, India.

** ORCID:0000-0002-7490-5289, Faculty of Pharmaceutical Sciences, Motherhood University, Vill. Karoundi, India.

*** ORCID: 0000-0002-1657-811X, Department of Pharmaceutical Technology, Meerut Institute of Engineering and Technology, India.

° Corresponding Author; Neha Krishnarth (Assistant Professor)

Phone: 0591-2450793,+919412391802, Fax : 0591-2452207e-mail:

nkrishnarth@gmail.com

Pages 205-210, [Abstract](#) | [PDF \(215 KB\)](#)

(<http://dergi.fabad.org.tr/pdf/volum45/issue3/205-210.pdf>)

2022 Volume 47, Issue 1
(<https://dergi.fabad.org.tr/en/2022-volume-47-issue-1/>)

2021 Volume 46, Issue 3
(<https://dergi.fabad.org.tr/en/2021-volume-46-issue-3/>)

2021 Volume 46, Issue 2
(<https://dergi.fabad.org.tr/en/2021-volume-46-issue-2/>)

2021 Volume 46, Issue 1
(<https://dergi.fabad.org.tr/en/2021-volume-46-issue-1/>)

2020 Volume 45, Issue 2
(<https://dergi.fabad.org.tr/en/2020-volume-45-issue-2/>)

2020 Volume 45, Issue 1
(<https://dergi.fabad.org.tr/en/2020-volume-45-issue-1/>)

2019 Volume 44, Issue 3
(<https://dergi.fabad.org.tr/en/2019-volume-44-issue-3/>)

2019 Volume 44, Issue 2
(<https://dergi.fabad.org.tr/en/2019-volume-44-issue-2/>)

2019 Volume 44, Issue 1
(<https://dergi.fabad.org.tr/en/2019-volume-44-issue-1/>)

Synthesis of New 2-(4-Oxothiazolidin-2-ylidene)-acetamides as Potential Antimicrobial Agents

Volodymyr HORISHNY* , Taras CHABAN** , Vasyl MATIYCHUK***^o

* ORCID: 0000-0002-5868-4105, Department of Pharmaceutical, Organic and Bioorganic Chemistry, Danylo Halytsky Lviv National Medical University, Ukraine

** ORCID: 0000-0003-0618-275X, Department of General, Bioinorganic, Physical and Colloidal Chemistry, Danylo Halytsky Lviv National Medical University, Ukraine

*** ORCID: 00000-0001-8077-2139, Department of Organic Chemistry, Ivan Franko National University of Lviv, Ukraine

^o Corresponding Author; Vasyl MATIYCHUK,

Tel.: +38 067 675-16-83 e-mail: v_matiychuk@ukr.net

Pages 211-218, [Abstract](#) | [PDF \(255 KB\)](#)

(<http://dergi.fabad.org.tr/pdf/volum45/Issue3/211-218.pdf>)

Improvement of the Antimicrobial Activity of Moxifloxacin Using W/O Microemulsion System for Skin Infections

Gülşah EREL-AKBABA* , ismail ÖZTÜRK** , Zeynep AY-ŞENYİĞİT***^o

* ORCID: 0000-0003-3287-527, Department of Pharmaceutical Biotechnology, Faculty of Pharmacy, Izmir Katip Celebi University, 35620, Izmir, Turkey

** ORCID: 0000-0002-2669-3090, Department of Pharmaceutical Microbiology, Faculty of Pharmacy, Izmir Katip Celebi University, 35620, Izmir, Turkey

*** ORCID: 0000-0002-4920-2469, Department of Pharmaceutical Technology, Faculty of Pharmacy, Izmir Katip Celebi University, 35620, Izmir, Turkey

^o Corresponding Author; Zeynep Ay-Şenyiğit, Ph.D.

Tel: +90 232 329 35 11 / 6106, GSM: +90 533 649 10 18, Fax: +90 232 325 40 42, e-mail:

zeynep.senyigit@ikcu.edu.tr

Pages 219-227, [Abstract](#) | [PDF \(701 KB\)](#)

(<http://dergi.fabad.org.tr/pdf/volum45/Issue3/219-227.pdf>)

Review Articles

Evaluation of Hepatotoxicity Risk of Green Tea

Methiye MANCAK KARAKUŞ* , Nurgün KÜÇÜKBOYACI***^o

* ORCID: 0000-0002-0085-7552, Gazi Üniversitesi, Eczacılık Fakültesi, Farmakognozi Anabilim Dalı, 06330 Ankara, Türkiye

** ORCID: 0000-0001-5489-3367, Gazi Üniversitesi, Sağlık Bilimleri Enstitüsü, Ankara, Türkiye

^o Corresponding Author: Prof. Dr. Nurgün KÜÇÜKBOYACI

Tel.: +90 312 2023177, e-mail: nurgunkucukboyaci@gmail.com

Pages 229-241, [Abstract](#) | [PDF \(241 KB\)](#)

(<http://dergi.fabad.org.tr/pdf/volum45/Issue3/229-241.pdf>)

2018 – Volume 43, Issue 3
(<https://dergi.fabad.org.tr/en/2018-volume-43-issue-3/>)

2018 – Volume 43, Issue 2
(<https://dergi.fabad.org.tr/en/2018-volume-43-issue-2/>)

2018 – Volume 43, Issue 1
(<https://dergi.fabad.org.tr/en/2018-volume-43-issue-1/>)

2017 – Volume 42, Issue 3
(<https://dergi.fabad.org.tr/en/2017-volume-42-issue-3/>)

2017 – Volume 42, Issue 2
(<https://dergi.fabad.org.tr/en/2017-volume-42-issue-2/>)

2017 – Volume 42, Issue 1
(<https://dergi.fabad.org.tr/en/2017-volume-42-issue-1/>)

2016 – Volume 41, Issue 1
(<https://dergi.fabad.org.tr/en/2016-volume-41-issue-1/>)

2015 – Volume 40, Issue 1
(<https://dergi.fabad.org.tr/en/2015-volume-40-issue-1/>)

[Click for other Issue](http://dergi.fabad.org.tr/tr/archiv)
(<http://dergi.fabad.org.tr/tr/archiv>
e).

The Role of Folate Metabolism in Neurodegenerative Diseases

Nihan İŞBİLEN* , Tuba TÜYLÜ KÜÇÜKKİLİNÇ**°

* ORCID No: 0000-0003-2627-2728, Hacettepe Üniversitesi Eczacılık Fakültesi 06100 Ankara

** ORCID No: 0000-0003-1566-0717, Hacettepe Üniversitesi Eczacılık Fakültesi Biyokimya Anabilim Dalı 06100 Ankara

° Corresponding Author: Tuba TÜYLÜ KÜÇÜKKİLİNÇ

Tel.: 3051499, e-mail: ttuylu@hacettepe.edu.tr

Pages 243-252, [Abstract](#) | [PDF \(621 KB\)](#)

(<http://dergi.fabad.org.tr/pdf/volum45/issue3/243-252.pdf>)

3D Printing Technology in Pharmaceutical Delivery System: Recent Advancement in Innovative Approach

Nupur ANAND* , Disha SINGH** , Gurjeet SINGH*** , Lakhvir KAUR****° , R.K DHAWAN***** and Narinder KAUR*****

* ORCID: 0000-0003-2612-3212, Department of Pharmaceutics, Khalsa College of Pharmacy, Amritsar, Punjab, India

** ORCID: 0000-0002-5120-4148, Department of Pharmaceutics, Khalsa College of Pharmacy, Amritsar, Punjab, India

*** ORCID: 0000-0003-4399-4693, Department of Pharmaceutics, Khalsa College of Pharmacy, Amritsar, Punjab, India

**** ORCID: 0000-0001-8091-2365, Department of Pharmaceutics, Khalsa College of Pharmacy, Amritsar, Punjab, India

***** ORCID:0000-0001-5130-327X, Department of Pharmacology, Khalsa College of Pharmacy, Amritsar, Punjab, India

***** ORCID:0000-0002-3465-4286, Department of Pharmaceutical Chemistry, Khalsa College of Pharmacy, Amritsar, Punjab, India

° Corresponding Author: Dr. Lakhvir Kaur

e-mail: lakhvir86@gmail.com

Pages 253-268, [Abstract](#) | [PDF \(345 KB\)](#)

(<http://dergi.fabad.org.tr/pdf/volum45/issue3/258-268.pdf>)

A Review on Christia vespertilionis: A New Emerging Medicinal Plant

Suganya MURUGESU* , Vikneswari PERUMAL**^o , Tavamani BALAN*** , Nur Amiera Syakira HAMZAN**** , Nur Syazwani Syasya SHAHRIM***** , Nur Syakirah Ramlah SHARRIZAL***** , Lee Wing HIN*****

* ORCID: 0000-0002-0867-1090, Faculty of Pharmacy and Health Sciences, Universiti Kuala Lumpur Royal College of Medicine Perak, Ipoh, Perak, Malaysia.

** ORCID: 0000-0002-4097-7727, Faculty of Pharmacy and Health Sciences, Universiti Kuala Lumpur Royal College of Medicine Perak, Ipoh, Perak, Malaysia.

*** ORCID: 0000-0002-2126-0798, Faculty of Pharmacy and Health Sciences, Universiti Kuala Lumpur Royal College of Medicine Perak, Ipoh, Perak, Malaysia.

**** ORCID: 0000-0002-0463-5906, Faculty of Pharmacy and Health Sciences, Universiti Kuala Lumpur Royal College of Medicine Perak, Ipoh, Perak, Malaysia.

***** ORCID: 0000-0003-2856-5401, Faculty of Pharmacy and Health Sciences, Universiti Kuala Lumpur Royal College of Medicine Perak, Ipoh, Perak, Malaysia.

***** ORCID: 0000-0001-9685-2344, Faculty of Pharmacy and Health Sciences, Universiti Kuala Lumpur Royal College of Medicine Perak, Ipoh, Perak, Malaysia.

***** ORCID: 0000-0003-0212-6523, Faculty of Pharmacy and Health Sciences, Universiti Kuala Lumpur Royal College of Medicine Perak, Ipoh, Perak, Malaysia.

^o Corresponding Author: Vikneswari PERUMAL

Tel: 0(605) 243 2635 / 2637, Fax: 0(605) 243 2636, email: vikneswari@unikl.edu.my

Pages 269-277, Abstract | PDF (167 KB)

(<http://dergi.fabad.org.tr/pdf/volum45/issue3/269-277.pdf>)

Novel Coronavirus Disease (COVID-19): Causes, Pathogenesis and Efforts of Treatment

Mohammed TAHA* ,** ^o , Loreen HABOUB**

* ORCID: 0000-0001-9350-2111, Department of Pharmacology and Medical Sciences, Faculty of Pharmacy, Al-Azhar University of Gaza, Palestine


** ORCID: 0000-0003-2415-7291, Inserm, U1068, Centre Recherche en Cancérologie de Marseille (CRCM), CNRS, Faculty of Medicine, Aix-Marseille University, Marseille, France

^o Corresponding Author: Mohammed TAHA

GSM: 0033 (0)7 58 67 70 25, mohammed.taha@inserm.fr ; m.taha@alazhar.edu.ps

Pages 279-296, Abstract | PDF (1,065 KB)

(<http://dergi.fabad.org.tr/pdf/volum45/issue3/279-296.pdf>)

 Journal of Pharmaceutical Sciences (<https://dergi.fabad.org.tr/en/author/journal-of-pharmaceutical-sciences/>)

 Saturday December 12th, 2020 (<https://dergi.fabad.org.tr/en/turkish-2020-volume-45-issue-3/>)

Previous: 2020 Volume 45, Issue 2
(<https://dergi.fabad.org.tr/en/2020-volume-45-issue-2/>)

Next: 2021 Volume 46, Issue 1
(<https://dergi.fabad.org.tr/en/2021-volume-46-issue-1/>)

FABAD | Journal of Pharmaceutical Sciences© All right reserved 2021

Read More by Acme Themes (<http://www.acmethemes.com/>)

Anticancer Property of *Orthosiphon stamineus* Benth. Extracts in Different Solvent Systems against T47D Human Breast Cancer Cell Lines

Lusiana ARIFIANI^{*}, Sukardiman SUKARDIMAN^{**}, Niken INDRIYANTI^{***}, Retno WIDYOWATI^{****}

Anticancer Property of *Orthosiphon stamineus* Benth. Extracts in Different Solvent Systems against T47D Human Breast Cancer Cell Lines

SUMMARY

Solvent system is an important factor in extraction process in order to obtain compounds that have pharmacological activity. The aim of this research is to develop a comprehensive extraction methods by modification of solvents used that might produce compounds possessing pharmacological activity for anticancer. In this study, *Orthosiphon stamineus* Benth. extract was used as sample in different solvent systems to observe their metabolite profiles. Extraction carried out using sonication techniques with ethanol solvents in three types of concentrations (96%, 70% and 50%). Then, the extracts introduced into anticancer activity profiles in order to find its active compounds. The anticancer activity had explored against breast cancer cells (T47D) using the MTT assay and doxorubicin as a positive control. The best IC_{50} value obtained from the 50% ethanol extract of *Orthosiphon stamineus* Benth. Based on the scanning chromatogram at 366 nm using Thin Layer Chromatography, each sample contains sinensetin and rosmarinic acid. The largest percentage of rosmarinic acid area was found on 70% ethanol extract of *Orthosiphon stamineus* Benth, while the highest percentage of sinensetin was found on 50% ethanol extract of *Orthosiphon stamineus* Benth. Thus, it can be concluded that sinensetin which has the most influence on anticancer activity.

Key Words: *Orthosiphon stamineus*, breast cancer, rosmarinic acid, sinensetin, T47D, anticancer

***Orthosiphon stamineus* Benth'in farklı Çözücü Ekstrelerinin T47D İnsan Meme Kanseri Hücre Hattına Karşı Antikanser Özelliği**

ÖZ

Solvent sistemi, farmakolojik aktiviteye sahip bileşiklerin elde edilmesinde ekstraksiyon sürecinde önemli bir faktördür. Bu araştırmanın amacı, antikanser farmakolojik aktiviteye sahip bileşikler için çözücülerini modifiye ederek kapsamlı bir ekstraksiyon yöntemi geliştirmektir. Bu çalışmada *Orthosiphon stamineus* Benth. ekstraktı metabolit profillerini gözlemlemek için farklı solvent sistemlerinde numune olarak kullanılmıştır. Ekstraksiyon, üç farklı konsantrasyonda (% 96, % 70 ve % 50) etanol çözücülerle sonikasyon teknikleri kullanılarak gerçekleştirildi. Daha sonra ekstraktlar, aktif bileşiklerini bulmak için antikanser aktivite profillerine sokuldu. Antikanser aktivitesi, meme kanseri hücreleri (T47D) kullanılarak MTT analizi ile araştırılmıştır, pozitif kontrol olarak doksorubisin kullanılmıştır. *Orthosiphon stamineus* Benth'in elde edilen en iyi IC_{50} değeri % 50 etanol ekstraktında bulunmuştur. İnce Tabaka Kromatografisi kullanılarak 366 nm'de elde edilen kromatogramına göre, her numune sinensetin ve rosmarinik asit içermektedir. Rosmarinik asit alanının en büyük yüzdesi *Orthosiphon stamineus* Benth'in % 70 etanol ekstresinde bulunurken, en yüksek sinensetin yüzdesi *Orthosiphon stamineus* Benth'in % 50 etanol ekstresinde bulunmuştur. Böylece antikanser aktivitesi üzerinde en fazla etkiye sahip olanın sinensetin olduğu sonucuna varılabilir.

Anahtar Kelimeler: *Orthosiphon stamineus*, meme kanseri, rosmarinik asit, sinensetin, T47D, antikanser

Received: 23.12.2019

Revised: 29.04.2020

Accepted: 02.05.2020

^{*} ORCID:0000-0003-1704-7908, Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Universitas Airlangga, Indonesia

^{**} ORCID:0000-0001-9689-5088, Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Universitas Airlangga, Indonesia

^{***} ORCID:0000-0003-3733-5018, Departement of Pharmacology, Faculty of Pharmacy, Mulawarman University, Indonesia

^{****} ORCID:0000-0003-0572-7551, Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Universitas Airlangga, Indonesia

^o Corresponding Author; Retno Widyowati

Tel. +6231 5933150, Fax. +6231 5935249, e-mail: rr-retno-w@ff.unair.ac.id

INTRODUCTION

Recently, the efforts to treat cancer patients have focused on inhibiting the growth or killing cancer cells. The investigations to find an ideal drug that targets cancer cells with minimal side effects are ongoing. Some Indonesia medicinal plants may have a potential bioactive compound to be developed into an ideal drug for cancer. One of them is *Orthosiphon stamineus* Benth.

Orthosiphon stamineus Benth. (Lamiaceae) is a Indonesia plant native, traditionally used for diuretics, rheumatism, diabetes and hypertension (Fei *et al.*, 2010). Previous study reported the activity of this plant extract inhibits human oral cancer cells (Younis *et al.*, 2013). The methanol extract of this plant enhanced tamoxifen on breast cancer cell (MCF 7) proliferation (Sahib *et al.*, 2009). In addition, 200 mg/kg ethanol extract of this plant showed no tumor cell growth compared to control group using xenograph method of tumor models that transplanted with breast cancer cells (MCF7) and colon cancer cells (HCT116) (Ahmad *et al.*, 2010).

The bioactive compound will become lead compound to find an effective drug for cancer. One of the bioactive compounds found in this plant is sinensetin (5,6,6,7,3',4'-pentamethoxy flavone). It also contains orthosiphonol D, orthosiphonol E (Takeda *et al.*, 1993), orthosiphonol A, orthosiphonol B, 3'-hydroxy-5,6,7,4'-tetramethoxyflavone, neoorthosiphonol A, neoorthosiphonol B, α -amyrin, β -amyrin, maslinic acid, urosolic acid, orthosiphonone A, orthosiphonone B, myo-inositol, β -caryophyllene, caffeic acid, sinensetin, tetra-methyl scutellarein, eupatorin, cirsimaritin, acetovanillochromene, orthochromene A, methylripariochromene, agermacrene-D, β -selinen, α -cadinol, choline, betaine, O-cyamene-terpineol, lyrol, valencene, naphthalin, camphor, α -elemene (Singh *et al.* 2015), 5,6,7,8-tetra hydroxy-6-methoxy-flavones (Hossain *et al.*, 2008), potassium, flavonol glycosides, caffeic acid (rosmarinic acid) (Sumaryono, *et al.*, 1991), essential oils, diterpenes, lipophilic flavones such as eupatorin, (6-hydroxy-5,7,4-trimethoxy flavone), and TMF (3'-hydroxy-5, 6, 7, 4' tetramethoxy flavone) (Awale *et al.*, 2001), triterpenes such as betulinic, ursolic, oleanolic acids, β sitosterol (Tezuka *et al.*, 2000) and flavonoids such as 5-hydroxy-6,7,30,40-tetrametoxiflavone, salvigenin, 6-hydroxy-5,7,40-trimethoxyflavone, 5,6,7,30-tetramethoxy-40-hydroxy-8-Cprenylflavone (Hossain and Rahman, 2015). The chloroform extract of this plant contained 1.48% of sinensetin, 2.26% of eupatorin, and 0.58% of 30-hydroxy- 5,6,7,40-tetrametoksiflavon (Mohamed *et al.*, 2013, Yam *et al.*, 2012).

Previous studies showed that the 50% methanol

extract of this plant using freeze-dried and spray-dried methods contained protein, polysaccharides and saponins (Siddiqui *et al.*, 2009). Research on metabolite profiles in this plants had been carried out using chromatographic and spectroscopic techniques combined with chemometrics (Akowuah *et al.*, 2004; Sumaryono, *et al.*, 1991; Saidan *et al.*, 2015a). Ethanol extract using maceration method contained high phenolics and flavonoids, (rosmarinic acid and eupatorin) as antioxidants, while 50% ethanol and methanol extracts using Soxhlet contained high protein and glycosaponin. Water extracts using reflux and maceration showed high polysaccharides (Saidan *et al.*, 2015b).

Solvent system is an important factor in extraction process in order to obtain compounds that have pharmacological activity. In this study, the effect of different solvents on the metabolites profile in each extract had been determined. Extractions with 96, 70 and 50% ethanol were carried out according to the previous study with different method (Arifianti *et al.*, 2014). Arifianti had extracted this plant using maceration method while this study used ultrasonic method to accelerate the extraction process and % yield with optimum results at same concentrations. It correlated to the anticancer activity (breast cancer) and their secondary metabolites are responsible for their activity.

MATERIAL AND METHODS

General Experimental Procedures and Materials

The plant was extracted on CAMAG ultrasonic and then evaporated by BUCHI rotary evaporator. The metabolite profiles of *Orthosiphon stamineus* Benth leaves were measured on a CAMAG Scanner 3 Densitometer and Linomat 5. The solvent extracts were combination between ethanol p.a (Merck) and aquadest. Sinensetin and rosmarinic acid from Sigma used as standard. The material used for breast cancer activity of T47D cells were RPMI 1640 medium (Gibco, Invitrogen), Fetal Bovine Serum (FBS, Sigma), Penicillin-Streptomycin (Sigma), Amphotericin B (Sigma), Dimethyl sulfoxide (DMSO, Sigma), Phosphate Buffer Saline (PBS, Sigma), 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT, ThermoFisher) and sodium dodecylsulfate (Sigma). The cancer cells inhibition was determined with Robonik Elisa Reader.

Plant Materials

Orthosiphon stamineus Benth leaves were obtained on late April 2018 from Balai Materia Medika, Malang and voucher specimens were deposited in Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Universitas Airlangga.

Extraction of plant materials

Extraction was carried out by ultrasonic method according to the previous published method by modification (Juliana *et al.*, 2019). The dried leaves of *Orthosiphon stamineus* Benth (50 g) were extracted with 250 mL of each 96%, 70% or 50% ethanol (3 x 10 minutes) using CAMAG ultrasonic. Then the extracts were separated by filtration. The residue was re-extracted by using same procedure (3 times repeated). The filtrates were evaporated by BUCHI rotary evaporator to dryness under *vacuo* to get 96%, 70% or 50% ethanol extracts. The extracts were used to examine bioassay activity and phytochemical analysis. It performed by thin layer chromatography (TLC) to observe the sinensetin and rosmarinic acid profile.

Phytochemical analysis of plant extracts using TLC

The 10 mL of each 96%, 70%, 50% ethanol extracts of *Orthosiphon stamineus* Benth, sinensetin and rosmarinic acid were applied on a pre-coated TLC plate, silica gel 60F 254 (10 cm x 20 cm) as 7 mm bands using a Camag automatic TLC sampler (Linomat 5) spray-on band applicator equipped with 100 mL syringe, and the space between two spots was 2 mm of the plate. The extracts were each applied duplicates on the plate. The TLC plates were developed with chloroform-ethyl acetate in a ratio of 6:4 as mobile phase. Then, they were identified using a UV lamp at 365 nm. The phytochemical were analyzed based on chromatogram pattern using a CAMAG TLC Scanner 3 Densitometer and winCATS software, using a deuterium light source, the slit dimension was 6.00 x 0.45 mm. Peak areas were recorded and the spot of sinensetin and rosmarinic acid in the sample were confirmed by comparing the RF and spectra of the spot with that of sinensetin and rosmarinic acid standard (Arifianti *et al.*, 2014, Hossain and Ismail, 2016).

Cell line

The T47D Human breast cancer cell lines were obtained from the CCRC (Cancer Chemoprevention Research Center), Gajah Mada University, Indonesia and a modification method described by Freshney Method (Freshney, 2005). The T47D Human breast cancer cells were maintained in RPMI 1640 that contained 10% of FBS, 2% of Penicillin-Streptomycin and 1% of Amphotericine B. It was stored at 37°C with humidified atmosphere of 5% CO₂ (Eppendorf). The cells were routinely observed to keep them from contamination.

Measurement of inhibition of cancer cell by MTT method

The MTT method used was a method that has been modified by Freshney (Freshney, 2005). The 5 x 10⁴ cells/wells of T47D cells with or without samples (96%, 70%, 50% ethanol extracts of *Orthosiphon stamineus* Benth) were cultured in RPMI 1640 medium that contained 10% of FBS and 1 % (v/v) of penicillin-streptomycin into 96 well plate then incubated for 24 hours at 37°C and 5% CO₂ (70-80% confluent). The samples were dissolved in DMSO and further diluted with medium to make series of concentrations (15 – 1,000 mg/mL). The final concentration of DMSO in the test solution should not more than 1%. Control cell was treated with 1% DMSO. Cells were then treated with a serial dilution of tested samples. The doxorubicin concentrations of 2.5-100 mg/ml were used as positive control. After 24 h incubation, 0.5 mg/ml of MTT was added to each well and incubated for 4 hours. Then, the stopper solution (sodium dodecylsulfate 10% in 0.1 N HCl) was added to dissolve the formazan crystal and incubated overnight at room temperature and dark. Finally, the cells viability was measured using ELISA reader at 1 570 nm. The absorbance of each well then converted into percentage of viable cells using calculation below and the IC₅₀ values were determined by Probit analysis using SPSS software. Experiments were done in triplicates.

$$\%cellviability = \frac{samp\ absorbance - medium\ control\ absorbance}{cell\ control\ absorbance - medium\ contro\ labsorbance} \times 100\%$$

RESULT AND DISCUSSION

Extraction

Several ways can be do to obtain phytochemicals from plants, one of which is extraction. Extraction efficiency is influenced by the chemical properties of the compound, the extraction method used, the particle size of the sample, the solvent used, and the presence of disturbing substances (Stalikas, 2007). Conventional extraction techniques are often associated with long heating times and a risk of bioactive

compounds degradation. This has led to sophisticated techniques such as ultrasonic extraction which are efficient in terms of extraction time and solvent consumption. In view of this method, ultrasonic cavitation produces shock waves that are able to disrupt the external structure of plant samples and release plant bioactives effectively (Budynas & Nisbett, 2008; Flores & Liang, 1994).

Ultrasonic extraction using ultrasonic frequencies at >20 kHz can accelerate the contact time between

samples and solvents at room temperature. It causes the mass transfer of bioactive compounds from plant cells to solvent to be faster. Sonication relies on sound energy that causes the cavitation. It forms small bubbles due to ultrasonic frequency transmission to help the diffusion of solvents into plant cell walls (Ashley *et al.*, 2001).

General extraction parameters such as concentration and ratio of solvents using ultrasonic method were first optimized. The solvent used was ethanol because it is non-toxic, good polarity for the sound energy and ultrasonic frequencies, so it is able to dissolve interesting bioactive compounds (Xiao *et al.*, 2008). The extraction yield depends on the solvent with various polarity, pH, temperature, extraction time, and sample composition. At the effect of the same extraction time and temperature, the solvent and sample composition are the most important parameters. In this study, *Orthosiphon stamineus* Benth. extracts were obtained using ethanol and water at various concentrations (50%, 70%, and 96%). Their extraction yields were ranged between 15.64%, 12.39% and 3.44%, respectively (Table 1). The results of extraction with various solvents decreased in the following order: 50% ethanol > 70% ethanol > 96% ethanol. It showed that the extraction yield increases with increasing polarity of the solvent used in the extraction. Increasing the concentration of water in the solvent can increase the extraction yield. Compounds other than sinensetin and rosmarinic acid in the extract, it may have been extracted and contributed to higher yields. This might be caused by higher solubility of protein and carbohydrate in water-ethanol than pure ethanol (Zieliński and Kozłowska, 2000). The use of a combination of water in organic solvents can facilitate the extraction of water-soluble chemicals and/or organic solvents. This might be the reason why the ethanol extract yield is 50% higher than other extracts. The results of this study are in accordance with the results of extraction in *Limnophila aromatica* (Do *et al.*, 2014) and several medicinal plants (Sultana *et al.*, 2009).

Table 1. % yield of *Orthosiphon stamineus* Benth extracts


Sample	% yield
96% ethanol extract	3.444
70% ethanol extract	12.390
50% ethanol extract	15.636

The addition of water in the extraction solvent showed that the extraction yield is improved, because the presence of water increased heating efficiency due to its high dielectric constant (Sato & Buchner, 2004), and increased the permeability of plant matrices to encourage mass transfer and diffusion of bioactive compounds (Boeing *et al.*, 2014). The effects of aqueous ethanol have different effects, the optimum solvent concentration was found to be 50% (v/v) ethanol.

The phytochemical analysis

A number of sinensetin and rosmarinic acid found in the leaves of *Orthosiphon stamineus* Benth. TLC-densitometry is the current method for the quantization of some flavonoids and caffeic acid derivatives in pharmaceutical formulations. It is quickly gaining widespread acceptance in pharmaceutical analysis. This is due to simplicity, accuracy, cost effectiveness and possibility of simultaneous determination of a number of samples on a single TLC plate. HPTLC allows the identification and quantification of more than 20 samples in the same chromatographic process and requires more than 2 hours. Whereas TLC takes only 15-30 minutes because it does not require conditioning steps, such as in HPLC, and is cheaper.

A fingerprint chromatography was performed to describe components found in sinensetin & rosmarinic acid-rich extract using TLC densitometry according to the modified method of Hossain and Ismail (2016) as well as Hossain and Ismail (2009). Samples (96, 70 and 50% ethanol extracts of *Orthosiphon stamineus* Benth) and standard (sinensetin and rosmarinic acid) were explored on a silica gel GF 254 and developed with chloroform-ethyl acetate in a ratio of 6:4. under UV observation at 366 nm by TLC visualizer (Fig. 1). The advantage of using the TLC-visualizer method is easy, fast, accurate, inexpensive and most suitable for natural material analysis. A sample chromatogram showed the presence of spots of the same color and at the same R_f value as the standard (Fig.1). A single peak at R_f 0.49 was observed in the chromatogram of sinensetin standard and R_f 0.06 as rosmarinic acid standard. At 70% ethanol extract had more intensity spot because it contained highest rosmarinic acid compare than other extracts.



Sinensetin Ros. Ac. 50% 50% 70% 70% 96% 96%

Figure 1. The chromatograms of samples and standard without any spray reagent, in UV light at 366 nm

TLC analysis results showed bright blue fluorescent spots and the ultraviolet spectrophotometry showed the spectra images that were identical between samples with standards spectra. Two spectra are said to be identical if they have a MF (Match Factor) price > 95. In this study, the price of a match factor from the sample against sinensetin and rosmarinic acid standard obtained 0.99559 and 0.99985, respectively, so that they can be said to be identical (Fig. 2). The presence of sinensetin and rosmarinic acid in samples were proven by comparison of standards spectra with components that are separate from the samples UV-VIS spectra. Figure 2a showed the standard UV-VIS spectra of sinensetin (black) with samples, while in figure 2b was rosmarinic acid (pink) with samples. It can be observed the presence of sinensetin and rosmarinic acid peak in a sample at the same Rf value. It showed the similarity of spotting between sinensetin and rosmarinic acid standard in each sample. This data is supported by a standard spectrum profile of rosmarinic acid that has similarities with the spectrum of 96, 70 and 50% ethanol extracts of *Orthosiphon stamineus* Benth. Furthermore, a similar spectrum image was also obtained between sinensetin standard and them, but there was a slight shift in the sinensetin spectrum of 96% ethanol extract of *Orthosiphon stamineus* Benth (Fig.2).




Figure 2. The UV-VIS Spectra overlay results of sinensetin (a, black) and rosmarinic acid (b, pink), together with samples of 50 % ethanol extract (blue), 70% ethanol extract (green) and 96% ethanol extract (yellow).

Quantitative determination was done by TLC-densitometry using the calibration curve method. The calibration curve was performed by the winCATS software program. In table 2, the percentage data of each chromatogram area referred to the sinensetin and rosmarinic acid standard. The largest percentage of ros-

marinic acid area was found on 70% ethanol extract (74.61 ± 0.03), while the highest percentage of sinensetin was found on 50% ethanol extract (32.97 ± 0.06) of *Orthosiphon stamineus* Benth. So 70% ethanol is the best solvent system for extracting rosmarinic acid while 50% ethanol for sinensetin.

Table 2. Peak identification by TLC-densitometry

Samples	Start position	Start Height	Max position	Max Height	Max %	End Position	End Height	Area	% Area	Mean % area
50% EtOH Ex_1	0.06 Rf	0.4 AU	0.11 Rf	102.3 AU	72.22%	0.16 Rf	0.0 AU	1330.3 AU	65.85%	67.04±0.02
	0.48 Rf	4.3 AU	0.55 Rf	39.4 AU	27.78%	0.60 Rf	2.9 AU	689.9 AU	34.15%	32.97±0.06
50% EtOH Ex_2	0.06 Rf	0.6 AU	0.11 Rf	103.4 AU	73.36%	0.14 Rf	1.2 AU	1338.7 AU	68.22%	
	0.49 Rf	3.3 AU	0.55 Rf	37.6 AU	26.64%	0.59 Rf	3.2 AU	623.7 AU	31.78%	
70% EtOH Ex_1	0.06 Rf	1.0 AU	0.12 Rf	192.9 AU	80.66%	0.18 Rf	0.0 AU	2774.9 AU	76.45%	74.61±0.03
	0.49Rf	7.0AU	0.56 Rf	46.3 AU	19.34%	0.60 Rf	2.8 AU	855.0 AU	23.55%	22.15±0.47
70% EtOH Ex_2	0.06 Rf	0.2 AU	0.12 Rf	191.2 AU	76.66%	0.15 Rf	1.2 AU	2821.9 AU	72.77%	
	0.50 Rf	5.9 AU	0.56 Rf	46.0 AU	18.45%	0.61 Rf	0.3 AU	804.1 AU	20.74%	
96% EtOH Ex_1	0.07Rf	0.0AU	0.11 Rf	73.6 AU	72.24%	0.15 Rf	0.1 AU	892.7AU	70.10%	71.37±0.02
	0.54Rf	6.3 AU	0.57 Rf	28.3 AU	27.76%	0.61 Rf	2.7 AU	380.8 AU	29.90%	28.63±0.02
96% EtOH Ex_2	0.07Rf	0.3AU	0.12 Rf	75.4 AU	74.20%	0.18 Rf	0.0 AU	942.1 AU	72.64%	
	0.56Rf	4.6AU	0.60Rf	26.2 AU	25.80%	0.64 Rf	0.4 AU	354.9 AU	27.36%	

Anticancer activity

The anticancer properties of the ethanol extracts of 96, 70 and 50% of *Orthosiphon stamineus* Benth. were determined by MTT test. This test was chosen because it is reliable, simple, applies to a variety of cells, and can be done in microtiter plates. The test was based on the reaction of colorimetry of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide with the enzyme dehydrogenase in living cells to form a colored formazan corresponding to a viable cell numbers (McCauley et al., 2013).

Table 3. IC₅₀ value of the samples on breast cancer activity

Sample	IC ₅₀ (ppm)
96% ethanol extract	259.016 ± 18.3
70% ethanol extract	390.521 ± 14.5
50% ethanol extract	159.049 ± 12.9
Doxorubicin	63.916 ± 5.5

In this study, the anticancer activities of the extracts were examined against human breast carcinoma cells (T47D). Based on the bioactivity results, the best IC₅₀ value was obtained from 50% ethanol extract of *Orthosiphon stamineus* Benth. (table 3) and TLC-densitometry results also showed that this extract contains the highest sinensetin. Sinensetin was able to inhibit proliferation of gastric cancer cell, arterial blood gas (ABG) cancer cells by apoptosis mechanism through P53 and P21 regulation cell using Western Blot Technique (Dong et al., 2011). While rosmarinic acid had known to prevent cell damage caused by free radicals, thereby reducing the risk of cancer and osteosclerosis (Fernando et al., 2016) and is a major compound of polyphenol that can be used as a nutraceutical prod-

uct that helps improve body immunity in cancer patients (Moore et al., 2016). Therefore, the 50% ethanol extract from this plant showed the highest cytotoxic activity against T47D breast cancer cells compared to other extracts and sinensetin has an important role for anticancer properties in the extracts.

CONCLUSION

The 50% ethanol extract from *Orthosiphon stamineus* Benth showed the highest cytotoxic activity against T47D breast cancer cells compared to other extracts. This extract contains highest of sinensetin (32.97±0.06) compared to other extracts. This compound may be responsible for anticancer properties in the extracts.

Acknowledgements

The authors would like to acknowledge Faculty of Pharmacy, Universitas Airlangga for providing the facilities to do the research and funding Penelitian Dosen Pemula (PDP) Project.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

REFERENCES

Ahmad, M. K., Ismail, Z., Aisheh, A. A., Majid, A. M. S. A., & Majid A. (2010). *Orthosiphon stamineus* leaf extract can prevent the development of colorectal and breast cancer in vivo. *Cancer Prevention Research*, 3(12), A54.

Akowuah, A. G., Zhari, I., Norhayati, I., Sadikun, A., & Khamsah, S. (2004). Sinensetin, eupatorin, 3'-hydroxy-5, 6, 7, 4'-tetramethoxyflavone and rosmarinic acid contents and antioxidative effect of *Orthosiphon stamineus* from Malaysia. *Food Chemistry*, 87(4), 559-566.

- Arifianti, L., Oktarina, R. D., & Kusumawati, I. (2014). Pengaruh jenis pelarut pengekstraksi terhadap kadar sinensetin dalam ekstrak daun *Orthosiphon stamineus* Benth. *Planta Husada*, 2(1), 1-4.
- Ashley, K., Andrews, R. N., Cavazosa, L., & Demange, M. (2001). Ultrasonic extraction as a sample preparation technique for elemental analysis by atomic spectrometry. *Journal of Analytical Atomic Spectrometry*, 16, 1147-1153.
- Awale, S., Tezuka, Y., Banskota, A. H., Kouda, K., Tun, K. M., & Kadota, S. (2001). Five novel highly oxygenated diterpenes of *Orthosiphon stamineus* from Myanmar. *Journal Natural Product*, 64(5), 592-596.
- Boeing, J. S., Barizão, É. O., Silva, B. C., Montanher, P. F., Almeida, V. D. C., & Visentainer, J. V. (2014). Evaluation of solvent effect on the extraction of phenolic compounds and antioxidant capacities from the berries: application of principal component analysis. *Chemistry Central Journal*, 8(1), 48.
- Budynas, R. G., & Nisbett, J. K. (2008). *Shigley's mechanical engineering design* (8th ed.). New York: McGraw-Hill.
- Do, Q. D., Angkawijaya, A. E., Tran-Nguyen, P. L., Huynh, L. H., Soetaredjo, F. E., Ismadji, S., & Ju, Y. (2014). Effect of extraction solvent on total phenol content, total flavonoid content, and antioxidant activity of *Limnophila aromatica*. *Journal of Food and Drug Analysis*, 22(3), 296-302.
- Dong, Y., Ji, G., Cao, A., Shi, J., Shi, H., Xie, J., & Wu, D. (2011). Effects of sinensetin on proliferation and apoptosis of human gastric cancer AGS cells. *Zhongguo Zhong Yao, Za Zhi*, 36(6), 790-794.
- Fei, Y. M., Vuanghao, L., Muhammad, S. I., Ziad, A. O., Fung, A. L., Noersal, R., Fawzy, A. M., Zuhair, A. G., Rusliza, B., Amirin, S., & Zaini, A. M. (2010). HPLC and anti-inflammatory studies of the flavonoid rich chloroform extract fraction of *Orthosiphon Stamineus* leaves. *Molecules*, 15, 4452-4466.
- Fernando, P. M. D. J., Piao, M. J., Kang, K. A., Ryu, Y. S., Hewage, S. R. K. M., Chae, S. W., Hyun, J. W. (2016). Rosmarinic acid attenuates cell damage against UVB radiation-induced oxidative stress via enhancing antioxidant effects in human HaCaT cells. *Biomolecules & Therapeutics*, 24(1), 75-84.
- Floros, J. D., & Liang, H. (1994). Acoustically, assisted diffusion through membranes and biomaterials. *Food Technology and Biotechnology*, 48(12), 79-84.
- Freshney, R.I. (2005). *Culture of animal cell: A manual of basic technique*. 5th Ed. John Wiley and Sons: New York. 120-135.
- Hossaina, M. A., Ismail, Z., Rahman, A., & Kang, S. C., (2008). Chemical composition and anti-fungal properties of the essential oils and crude extracts of *Orthosiphon stamineus* Benth. *Industrial Crops and Products*, 27(3), 328-334.
- Hossain, M. A., & Ismail, Z. (2009). High performance thin layer chromatographic determination of caffeic acid and rosmarinic acid from the leaves of *Orthosiphon stamineus*. *Indonesian Journal of Chemistry*, 9(1), 137-141.
- Hossain, M. A., & Ismail, Z. (2016). Quantification and enrichment of sinensetin in the leaves of *Orthosiphon stamineus*. *Arabian Journal of Chemistry*, 9(2), S1338-S1341.
- Hossain, M. A. & Rahman, S. M. M. (2015). Isolation and characterisation of flavonoids from the leaves of medicinal plant *Orthosiphon stamineus*. *Arabian Journal of Chemistry* 8, 218-221.
- Juliana, S., Suciati, S. & Indrayanto, G. (2019). Sterol and triterpene profiles of the callus culture of *Solanum mammosum*. *Makara J. Sci.*, 23(2), 72-78.
- McCauley, J., Zivanovic, A., Skropeta, D. (2013). Bioassays for anticancer activities. *Methods in Molecular Biology*, 1055, 191-205
- Mohamed, E. A. H., Yam, M. F., Ang, L. F., Mohamed, A. J. & Asmawi, M. A. (2013). Antidiabetic properties and mechanism of action of *Orthosiphon stamineus* Benth bioactive sub-fraction in streptozotocin-induced diabetic rats. *Journal of Acupuncture and Meridian Studies*, 6(1), 31-40.
- Moore, J., Yousef, M., & Tsiani, E. (2016). Anticancer effects of rosemary (*Rosmarinus officinalis* L.) extract and rosemary extract polyphenols. *Nutrients*, 8(11), 731, 1-32.
- Sahib, I., Ismail, Z., Othman, N. H., & Majib, A. (2009). *Orthosiphon stamineus* methanolic extract enhances the anti proliferative effect of tamoxifen on human hormone dependent breast cancer. *International Journal of Pharmacology*, 5(4), 273-276.
- Saidan, N. H., Aisha, A. F., Hamil, M. S., Malik, A. M., & Ismail, Z. (2015a). A novel reverse phase high performance liquid chromatography method for standardization of *Orthosiphon stamineus* leaf extracts. *Pharmacognosy Research*, 7(1), 23-31.
- Saidan, N. H., Hamil, M. S. R., Memon, A. H., Abdelbari, M. M., Hamdan, M. R., Mohd, K. S., Majid, A. M. S. A., & Ismail, Z. (2015b). Selected metabolites profiling of *Orthosiphon stamineus* Benth leaves extracts combined with chemometrics analysis and correlation with biological activities, *BMC Complementary and Alternative Medicine*, 15(350), 1-12

- Sato, T., & Buchner, R. (2004). Dielectric relaxation processes in ethanol/water mixtures. *The Journal of Physical Chemistry A*, 108(23), 5007-5015.
- Siddiqui, M. J., Hafizoh, S. N., Ismail, Z., Sahib, H., Helal, M., & Majid, A. M. S. A. (2009) Analysis of total proteins, polysaccharides and glycosaponins contents of *Orthosiphon stamineus* Benth. in spray and freeze dried methanol:water (1:1) extract and its contribution to cytotoxic and antiangiogenic activities. *Pharmacognosy Research*, 1(5), 320-326.
- Singh, M. K., Gidwani, B., Gupta, A., Dhongade, H., Kaur, C. D., Kashyap, P. P. & Tripathi, D. K. (2015). A review of the medicinal plants of Genus *Orthosiphon* (Lamiaceae). *International Journal of Biological Chemistry*, 9(6), 318-331.
- Stalikas, C. D. (2007). Extraction, separation, and detection methods for phenolic acids and flavonoids. *J. Sep Sci.*, 30, 3268-3295.
- Sultana, B., Anwar, F., & Ashraf, M. (2009). Effect of extraction solvent/technique on the antioxidant activity of selected medicinal plant extracts. *Molecules*, 14, 2167-2180.
- Sumaryono, W., Proksch, P., Wray, V., Witte, L., & Hartmann, T. (1991). Qualitative and quantitative analysis of the phenolic constituents from *Orthosiphon aristatus*. *Planta Medica*, 57, 176-80.
- Takeda, Y., Matsumoto, T., Terao, H., Shingu, T., Futatsuishi, Y., Nohara, T., & Kajimoto, T. (1993). Orthosiphon D and E, minor diterpenes from *Orthosiphon stamineus*. *Phytochemistry*, 33(2), 411-415.
- Tezuka, Y., Stampoulis, P., Banskota, A., Awale, S., Tran, K., & Saiki, I. (2000). Constituents of the Vietnamese medicinal plant *Orthosiphon stamineus*. *Chemical and Pharmaceutical Bulletin*, 48, 1711-1719.
- Yam, M. F., Mohamed, E. A. H., Ang, L. F., Pei, L., Darwis, Y., Mahmud, R., Asmawi, M. Z., Basir, R. & Ahmad, M. (2012). A simple isocratic HPLC method for the simultaneous determination of sinensetin, eupatorin, and 30-hydroxy-5,6,7,40-tetramethoxyflavone in *Orthosiphon stamineus* extracts. *Journal of Acupuncture and Meridian Studies*, 5(4), 176-181.
- Younis, L., Abu-Hassan, M. I., Zahar, F. F., & Husin, M. F. M. (2013). The use of *Orthosiphon stamineus* anticancer action against human oral squamous cell carcinoma. *The Open Conference Proceedings Journal*, 4, 217.
- Xiao, W., Han, L., & Shi, B. (2008). Microwave-assisted extraction of flavonoids from *Radix astragali*. *Separation and Purification Technology*, 62(3), 614-618.
- Zieliński, H., & Kozłowska, H. (2000). Antioxidant activity and total phenolics in selected cereal grains and their different morphological fractions. *Journal of Agricultural and Food Chemistry*, 48, 2008-2016.

Fabad Journal of Pharmaceutical Sciences

COUNTRY

Turkey



Universities and research institutions in Turkey



Media Ranking in Turkey

SUBJECT AREA AND CATEGORY

Pharmacology, Toxicology and
Pharmaceutics
↳ Pharmaceutical Science

PUBLISHER

Society of Pharmaceutical
Sciences of Ankara

H-INDEX

18

PUBLICATION TYPE

Journals

ISSN

13004182

COVERAGE

1999-2001, 2003-2021

INFORMATION


[Homepage](#)

[How to publish in this journal](#)

fabadnkar@gmail.com

Quartiles


Pharmaceutical Science



SJR

2013	0.111
2014	0.107
2015	0.111
2016	0.101
2017	0.112
2018	0.125
2019	0.152
2020	0.150
2021	0.150

Total Documents





Source details

Fabad Journal of Pharmaceutical Sciences

Scopus coverage years: from 1999 to 2001, from 2003 to 2022

Publisher: Society of Pharmaceutical Sciences of Ankara

ISSN: 1300-4182

Subject area: Pharmacology, Toxicology and Pharmaceutics: Pharmaceutical Science

Source type: Journal

[View all documents >](#)

[Set document alert](#)

[Save to source list](#)

CiteScore 2021

0.8



SJR 2021

0.150



SNIP 2021

0.276



[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology



CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021

$$0.8 = \frac{81 \text{ Citations 2018 - 2021}}{100 \text{ Documents 2018 - 2021}}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022

$$0.9 = \frac{74 \text{ Citations to date}}{85 \text{ Documents to date}}$$

Last updated on 05 March, 2023 • Updated monthly

CiteScore rank 2021

Category	Rank	Percentile
Pharmacology, Toxicology and Pharmaceutics	#115/171	33rd
Pharmaceutical Science		

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)