

# Antropometric study on the parameters of double-bundle anterior cruciate ligament reconstruction using magnetic resonance imaging

*by* Mohammad Zaim Chilmi

---

**Submission date:** 15-Sep-2022 12:44PM (UTC+0800)

**Submission ID:** 1900231135

**File name:** Prof\_DNU,\_dr\_MCH,\_dr\_JUP,\_dr\_TYO\_EurAsian\_JBS.pdf (230.33K)

**Word count:** 3105

**Character count:** 17083



## Antropometric study on the parameters of double-bundle anterior cruciate ligament reconstruction using magnetic resonance imaging

Dwikora Novembri Utomo<sup>1</sup>, I Gusti Ngurah Dodo M. Ranuh<sup>1</sup>, Mohammad Zaim Chilmi<sup>1</sup>, Jeffry Andrianus<sup>1</sup>, Andre Triadi Desnantyo<sup>1</sup>, Rosy Setiawati<sup>2\*</sup>

<sup>1</sup> Department of Orthopedics & Traumatology, Faculty of Medicine, Universitas Airlangga, Surabaya 60131, INDONESIA

<sup>2</sup> Department of Radiology, Faculty of Medicine, Universitas Airlangga, Surabaya 60131, INDONESIA

\*Corresponding author: [rosy-s@fk.unair.ac.id](mailto:rosy-s@fk.unair.ac.id)

### Abstract

**Background:** The reconstruction techniques of double-bundle anterior cruciate ligament (ACL) can restore normal kinematics on the knees and its stability. Data regarding the reconstruction parameters of double-bundle ACL and ACL footprint size on femur and tibia have never existed in Indonesia. **Purpose:** This study aims to obtain the reconstruction parameters' value of double-bundle ACL to determine the parameter value suitability with a minimum size, and the correlation of these parameters with age, sex, ethnicity, and body mass index (BMI). **Methods:** This study employed an analytic observational study with a cross-sectional design. In this study, eight parameters were measured, then analyzed based on age, gender, ethnicity, and BMI. The measurements were carried out using OsiriX program. **Result:** The average width of lateral condylar ridge amounted to 15.734 mm; the average diameter of ACL femoral footprint reached 15.85 mm; the average surface of ACL femoral footprint reached 72.72mm<sup>2</sup>; the average diameter of ACL tibial footprint 16.04 mm; the average surface of ACL tibial footprint amounted to 72.69mm<sup>2</sup>, and the average width of intercondylar notch reached 21.29 mm. In addition, no significant correlation was identified between age, sex, ethnicity, and BMI to the double-bundle ACL reconstruction parameters. **Conclusion:** Double-bundle ACL reconstruction can be performed because the reconstruction parameters' values have exceeded the minimum size. Subjects' characteristics based on age, sex, ethnicity, and BMI do not affect these parameters' values. The surface measurements of ACL femoral footprint and ACL tibial footprint significantly correlate with age.

**Keywords:** anthropometry, ACL footprint, double-bundle ACL, MRI

Utomo DN, M. Ranuh IND, Chilmi MZ, Andrianus J, Desnantyo AT, Setiawati R (2020)  
Antropometric study on the parameters of double-bundle anterior cruciate ligament reconstruction using magnetic resonance imaging. Eurasia J Biosci 14: 3165-3168.

© 2020 Utomo et al.

This is an open-access article distributed under the terms of the Creative Commons Attribution License.

### INTRODUCTION

The significant occurrence of anterior cruciate ligament (ACL) injuries needs to be a concern at this time. The quite large morbidity of ACL injuries in the community, especially athletes, becomes a concern at present. Data in the United States shows out of 200,000 cases on ACL injuries, half of which require reconstructive surgery (Singh, 2018). A meta-analysis study indicated that ACL injury incidence reached 68.6 cases per 100,000 people which resulted in ACL rupture as the most common injury in the orthopedics field (Sanders, et al. 2016). ACL injuries can also cause other tissue injuries and degenerative joint disease, with the result a major surgery is required (Mardina, et al. 2014).

ACL's anatomical position requires to identify and understand well to obtain maximum ACL reconstruction

results for the patients. ACL ligament anatomy consists of two bundles, i.e., anterolateral and posteromedial bundles, attached to the inferoposterior area of the lateral condylar ridge, and the cartilage surface of posterior side on the femur to between the tibial eminences. These matters are substantial to carry out ACL reconstruction as anatomical as possible.

There are two types of ACL reconstruction techniques; single-bundle and double-bundle techniques. The double-bundle reconstruction technique has the advantages to support anteroposterior and rotational stability compared to single-bundle reconstruction, which only supports anteroposterior stability. In this case, the double-bundle

Received: October 2019

Accepted: March 2020

Printed: September 2020

technique can be an option in ACL rupture reconstruction because it can provide greater results than a single-bundle technique (Siebold, 2014). Besides, a reconstruction that is identical to anatomical conditions is expected to restore normal ACL kinematics and maintain knee condition as excellent as possible for the long term (Hamer, et al. 1999; Shiberu, & Tamiru, 2016)..

Double-bundle reconstruction has a certain indication, i.e., the width of the lateral condylar ridge is more than 14 mm to obtain the bone distance between two tunnels at least two mm (Siebold, 2014). In addition, it is also required to identify intercondylar notch width more than 12 mm, thus the visualization of ACL insertion location can be observed clearly, and no impingement occurs during ACL graft insertion (Siebold, 2014). Graft impingement on intercondylar notch is one of the troubling complications in double-bundle reconstruction (Yasuda, et al. 2011). as a result, intercondylar notch width is one of the determinants of whether double-bundle ACL reconstruction can be performed or not.

The parameter data of ACL reconstruction regarding ACL insertion size in the lateral condylar ridge width on the distal femur, intercondylar notch width, and ACL footprint size on femur and tibia have never existed in Indonesia, especially in Dr. Soetomo General Hospital Surabaya. Based on the background, this study aims to obtain patients' ACL insertion measurement in the Dr. Soetomo General Hospital as a basis for double-bundle ACL reconstruction. In this study, the evaluation is carried out using MRI. The procedure for using the MRI does not require an invasion of the patient's body, so it is greater than the ligament biopsy procedure (Setiawati, et al. 2017. Sukmaningtyas, H., Pandelaki, J., Astuti, M. D. K., Fauziah, Riwanto, & Muttaqin, 2018). Based on the explanation above, this study was conducted to obtain the parameters' value of double-bundle ACL reconstruction to determine parameter value suitability with a minimum size, and identify the correlation of these parameters with age, sex, ethnicity, and BMI.

## METHODS

This study was an observational analytic study with a cross-sectional design. This study involved eight variables which were studied, and then analyzed based on age, sex, ethnicity, and BMI. The subjects were knee MRI examinations evaluated on sagittal, axial, and coronal sections with inclusion criteria: patients with 17-50 years old, patients with no ACL structure rupture, patients with no degeneration of distal femur and proximal tibia, and patients that had never experienced fracture on the distal femur and proximal tibia. The samples were the entire knee MRI results of patients at Dr. Soetomo General Hospital which fulfilled the inclusion criteria and excepted from exclusion criteria. The minimum sample size was calculated with the

**Table 1.** The Average of Double-Bundle ACL Reconstruction Parameters

Measurement	Size (Mean ± SD)
The width of the lateral condylar ridge	15.734 mm ± 0.4836
The width of ACL femoral footprint	7.0158 mm±0.47632
The length of ACL femoral footprint	15.8540 mm±1.76179
The surface of ACL femoral footprint	72.7202 mm <sup>2</sup> ±0.69432
The width of ACL tibial footprint	8.1563 mm±0.02559
The length of ACL tibial footprint	16.0429 mm±1.62422
The surface of ACL tibial footprint	72.6968 mm±0.66420
Intercondylar notch width	21.290 mm±2.8080

sample size formula for anthropometric study. The grouping was carried out with consecutive sampling by picking research samples that fit the inclusion criteria and excepted from exclusion criteria.

MRI electronic data obtained were then accessed using the OsiriX program to carry out measurements with the program's digital ruler on the width of lateral intercondylar ridge, ACL insertion on proximal tibia, and femoral intercondylar notch, as well as the surface area of ACL footprint on femur and tibia. In carrying out the measurement, the researchers were assisted by a radiology consultant in the musculoskeletal field to reduce bias.

## RESULTS

**Table 1** contains the measurement results in the average of lateral condylar ridge width, the width, length, and surface of ACL femoral and tibial footprints, as well as the intercondylar notch width of 65 samples used.

In this study, the average width of the lateral condylar ridge reached 15.734 mm. On ACL femoral footprint, the average width reached 7.0158 mm; the average length reached 15.8540 mm; and average surface reached 72.7202 mm<sup>2</sup>. Meanwhile, On ACL tibial footprint, the average width amounted to 8.1563 mm; the average length amounted to 16.0429 mm; and average surface amounted to 72.6968 mm<sup>2</sup>. In addition, the average of intercondylar notch width reached 21.290 mm.

Age variable correlation was obtained using the statistical test of the Pearson Correlation Test. It was obtained that age did not affect the diameter of the lateral condylar ridge with p-value = 0.085. There was a significant correlation between age and the surface of the ACL femoral footprint with p-value = 0.016. There was a significant correlation between age and the surface of the ACL tibial footprint with p-value = 0.014. There was no significant correlation between age and intercondylar notch diameter with p-value = 0.138.

The gender variable correlation was obtained using the Mann-Whitney analysis test. It was found that there was no correlation between sex and the width of the lateral condylar ridge (p = 0.379). There was no correlation between gender and the width of ACL femoral footprint (p = 0.806). There was no correlation between sex and the length of ACL femoral footprint (p = 0.517). There was no correlation between gender and the surface of ACL femoral footprint (p = 0.879). There

was no correlation between gender and the width of ACL tibial footprint ( $p = 0.194$ ). There was no correlation between sex and the length of ACL tibial footprint ( $p = 0.412$ ). There was no correlation between gender and the surface of ACL tibial footprint ( $p = 0.879$ ). There was no correlation between sex and intercondylar notch width ( $p = 0.682$ ).

The ethnicity variable correlation was obtained using the Mann-Whitney analysis test. There was no correlation between ethnicity and the width of the lateral condylar ridge ( $p = 0.734$ ). There was no correlation between ethnicity and the width of ACL femoral footprint ( $p = 0.621$ ). There was no correlation between ethnicity and the length of ACL femoral footprint ( $p = 0.822$ ). There was no correlation between ethnicity and the surface of ACL femoral footprint ( $p = 0.984$ ). There was no correlation between ethnicity and the width of ACL tibial footprint ( $p = 0.494$ ). There was no correlation between ethnicity and the surface of ACL tibial footprint ( $p = 0.392$ ). There was no correlation between ethnicity and intercondylar notch width ( $p = 0.479$ ).

BMI variable correlation was obtained using ANOVA analysis test, in which it was identified that there was no difference between intercondylar notch width and division based on BMI ( $p = 0.059$ ). No difference was identified between the width of ACL femoral footprint and the division based on BMI ( $p = 0.079$ ). There was no difference between the length of ACL femoral footprint and the division based on BMI ( $p = 0.649$ ). Surface differences were identified between the ACL femoral footprint and the division based on age ( $p = 0.026$ ). There was no difference between the length of ACL tibial footprint and the division based on BMI ( $p = 0.713$ ). There were significant differences between the width of ACL tibial footprint and the division based on height ( $p = 0.036$ ). There was no difference between the width of ACL tibial footprint and the division based on BMI ( $p = 0.723$ ).

## DISCUSSION

In this study, it was identified that there were no differences in the width of the lateral condylar ridge on the distal femur compared based on ethnicity, gender, age, and BMI. This study suggested that the average width of the lateral condylar ridge reached 15.72 mm. The size of the lateral condylar ridge was considered as crucial matter because the surface area of the ACL femoral footprint was affected by lateral condylar ridge and cartilage inferior border on joint surfaces. These matters would affect ACL reconstruction to generate as anatomical as possible (Bhattacharyya, 2018). Studies on the width of the lateral condylar ridge and its effects on ACL were still minimal, thus it was difficult to provide comparative views between the results of this study and previous studies.

Intercondylar notch width did not indicate any difference when compared based on ethnicity, gender, age, and BMI. This study exposed that the width range of intercondylar notch amounted to 16.7-28.4 mm, with an average of 21.15 mm. It indicated that intercondylar notch size had fulfilled the functional size, in which the minimum width of the intercondylar notch is 12 mm (Yasuda, et al. 2011. Marzo, et al. 1992).

In this study, the parameters of ACL footprint on distal femur measured did not suggest any differences in the size of ACL femoral footprint when compared based on ethnicity, gender, age, and BMI. In addition, the average surface of ACL footprint on the distal femur amounted to 72.66 mm, with an average length of 15.80 mm and an average width of 6.98 mm. It was identified that the ACL footprint on the distal femur was influenced by ACL insertion size (Siebold, 2014. Yasuda, et al. 2011 Mochizuki, et al. 2014 Ferretti, et al. 2012. Mochizuki, et al. 2014). Furthermore, identification found that age and sex affected ACL insertion size (Siebold, 2014. A study stated that men's ACL insertion size was longer and wider than women's (Amis, & Jakob, 1998. Colombet, et al. 2006). However, this study showed no differences in size, both in length and width, of ACL footprint on distal femur when compared with age and sex factors.

The parameters of ACL tibial footprint, which were then compared based on ethnicity, gender, age, and BMI, did not suggest any difference in the size of ACL tibial footprint. The mean surface of ACL footprint on proximal tibia reached 72.6 mm. This value was greatly smaller than the size of previous studies, in which the tibial insertion was wider in men than women, with a surface range from 114 mm<sup>2</sup>-229 mm<sup>2</sup> (Siebold, et al. 2008). ACL insertion size varied among individuals, and it could depend on the knee size of each individual (Odensten, & Gillquist, 1985).

This study needs to develop by involving a larger number of samples. Further research is required with a larger and wider range of samples to confirm the existence of covariation and to obtain the references of condylar ridge's geometric size, ACL femoral footprint, ACL tibial footprint, and intercondylar notch which can be generalized.

## CONCLUSION

Double-bundle ACL reconstruction can be carried out because the reconstruction parameters' values performed on the subject have exceeded the minimum size. Subjects' characteristics based on age, gender, ethnicity, and BMI do not affect these parameters' values except the surface measurements of ACL femoral footprint and ACL tibial footprint have a significant correlation with age.



**ACKNOWLEDGEMENT**

This study conducted with self-funding by authors.  
Dwikora Novembri Utomo gave critical comments when

the study conducted. All authors have agreed on the final form of the article manuscript.

**REFERENCES**

- Amis, A. A., & Jakob, R. P. (1998). Anterior cruciate ligament graft positioning, tensioning and twisting. *Knee Surgery, Sports Traumatology, Arthroscopy*, 6(1), S2-S12.
- Bhattacharyya, R., Ker, A., Fogg, Q., Spencer, S. J., & Joseph, J. (2018). Lateral Intercondylar Ridge: Is it a reliable landmark for femoral ACL insertion?: An anatomical study. *International Journal of Surgery*, 50, 55-59.
- Colombet, P., Robinson, J., Christel, P., Franceschi, J. P., Djian, P., Bellier, G., & Sbihi, A. (2006). Morphology of anterior cruciate ligament attachments for anatomic reconstruction: a cadaveric dissection and radiographic study. *Arthroscopy: The Journal of Arthroscopic & Related Surgery*, 22(9), 984-992.
- Ferretti, M., Doca, D., Ingham, S. M., Cohen, M., & Fu, F. H. (2012). Bony and soft tissue landmarks of the ACL tibial insertion site: an anatomical study. *Knee Surgery, Sports Traumatology, Arthroscopy*, 20(1), 62-68.
- Harner, C. D., Baek, G. H., Vogrin, T. M., Carlin, G. J., Kashiwaguchi, S., & Woo, S. L. (1999). Quantitative analysis of human cruciate ligament insertions. *Arthroscopy: The Journal of Arthroscopic & Related Surgery*, 15(7), 741-749.
- Mardina, Z., Fitriana, N., Siswanto, R., Oktavina, O., Zahra, N., Widiyanti, P.,... & Langenati, R. (2014). The Influence of Braiding Angle Variation in Braided-Twisted Fiber Scaffold Based Poly L-Lactic Acid for Anterior Cruciate Ligament Reconstruction Application. In *Advanced Materials Research* (Vol. 845, pp. 925-928). Trans Tech Publications Ltd.
- Marzo, J. M., Bowen, M. K., Warren, R. F., Wickiewicz, T. L., & Altchek, D. W. (1992). Intraarticular fibrous nodule as a cause of loss of extension following anterior cruciate ligament reconstruction. *Arthroscopy: The Journal of Arthroscopic & Related Surgery*, 8(1), 10-18.
- Mochizuki, T., Fujishiro, H., Nimura, A., Mahakkanukrauh, P., Yasuda, K., Muneta, T., & Akita, K. (2014). Anatomic and histologic analysis of the mid-substance and fan-like extension fibres of the anterior cruciate ligament during knee motion, with special reference to the femoral attachment. *Knee Surgery, Sports Traumatology, Arthroscopy*, 22(2), 336-344.
- Odensten, M., & Gillquist, J. (1985). Functional anatomy of the anterior cruciate ligament and a rationale for reconstruction. *The Journal of bone and joint surgery. American volume*, 67(2), 257-262.
- Sanders, T. L., Maradit Kremers, H., Bryan, A. J., Larson, D. R., Dahm, D. L., Levy, B. A., ... & Krych, A. J. (2016). Incidence of anterior cruciate ligament tears and reconstruction: a 21-year population-based study. *The American journal of sports medicine*, 44(6), 1502-1507.
- Setiawati, R., Utomo, D. N., Rantam, F. A., Ifran, N. N., & Budhiparama, N. C. (2017). Early graft tunnel healing after anterior cruciate ligament reconstruction with intratunnel injection of bone marrow mesenchymal stem cells and vascular endothelial growth factor. *Orthopaedic Journal of Sports Medicine*, 5(6), 2325967117708548.
- Shiberu, T., & Tamiru, S. (2016). Effect of Intra Spacing on Yield and Yield Components of Carrot (*Daucus Carota* L. Sub Sp. *Sativus*). *Current Research in Agricultural Sciences*, 3(1), 1-6.
- Siebold, R. (2014). *ZSDBARISR. Anterior Cruciate Ligament Reconstruction, A Practical Surgical Guide*. Springer. 283-90.
- Siebold, R., Ellert, T., Metz, S., & Metz, J. (2008). Tibial insertions of the anteromedial and posterolateral bundles of the anterior cruciate ligament: morphometry, arthroscopic landmarks, and orientation model for bone tunnel placement. *Arthroscopy: The Journal of Arthroscopic & Related Surgery*, 24(2), 154-161.
- Singh, N. (2018). International epidemiology of anterior cruciate ligament injuries. *Orthopedic Res Online J*, 1.
- Sukmaningtyas, H., Pandelaki, J., Astuti, M. D. K., Fauziah, D., Riwanto, I., & Muttaqin, Z. (2018). Diagnostic Value of Fractional Anisotropy in Detecting Hippocampal Sclerosis: A Study on Intractable Mesial Temporal Lobe Epilepsy with Normal MRI. *Hiroshima Journal of Medical Sciences*, (67), 191-198.
- Yasuda, K., van Eck, C. F., Hoshino, Y., Fu, F. H., & Tashman, S. (2011). Anatomic single-and double-bundle anterior cruciate ligament reconstruction, part 1: basic science. *The American journal of sports medicine*, 39(8), 1789-1800.

# Antropometric study on the parameters of double-bundle anterior cruciate ligament reconstruction using magnetic resonance imaging

## ORIGINALITY REPORT

9%

SIMILARITY INDEX

5%

INTERNET SOURCES

8%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 Yong-Beom Park, Chul-Won Ha, Hyung-Joo Kim, Yong-Geun Park. "Preoperative prediction of anterior cruciate ligament tibial footprint size by anthropometric variables", *Knee Surgery, Sports Traumatology, Arthroscopy*, 2016  
Publication 2%
- 2 Huizhi Wang, Zhuoyue Zhang, Yang Qu, Qinyi Shi, Songtao Ai, Cheng-Kung Cheng. "Correlation between ACL size and dimensions of bony structures in the knee joint", *Annals of Anatomy - Anatomischer Anzeiger*, 2022  
Publication 1%
- 3 [link.springer.com](https://link.springer.com)  
Internet Source 1%
- 4 Anna Starzyńska, Magdalena Kaczoruk-Wieremczuk, Michele Antonio Lopez, Pier Carmine Passarelli, Paulina Adamska. "The

Growth Factors in Advanced Platelet-Rich Fibrin (A-PRF) Reduce Postoperative Complications after Mandibular Third Molar Odontectomy", International Journal of Environmental Research and Public Health

Internet Source

5

Condro Utomo, Zulfikar Achmad Tanjung, Redi Aditama, Rika Fithri Nurani Buana et al. "Complete Mitochondrial Genome Sequence of the Phytopathogenic Basidiomycete Ganoderma boninense Strain G3", Microbiology Resource Announcements, 2019

Publication

1 %

6

[digitalcommons.wayne.edu](https://digitalcommons.wayne.edu)

Internet Source

1 %

7

[cmuj.cmu.ac.th](https://cmuj.cmu.ac.th)

Internet Source

1 %

8

Akira TANI, Tomoki MOCHIZUKI. "Review: Exchanges of volatile organic compounds between terrestrial ecosystems and the atmosphere", Journal of Agricultural Meteorology, 2021

Publication

<1 %

9

T. P. Branch, R. Siebold, H. I. Freedberg, C. A. Jacobs. "Double-bundle ACL reconstruction demonstrated superior clinical stability to single-bundle ACL reconstruction: a matched-pairs analysis of instrumented tests of tibial

<1 %

anterior translation and internal rotation laxity", Knee Surgery, Sports Traumatology, Arthroscopy, 2010

Publication

10

Yong-Beom Park, Young-Suk Song, Su-Cheol Kim, Yong-Geun Park, Chul-Won Ha. "The size of tibial footprint of anterior cruciate ligament and association with physical characteristics in Asian females", Archives of Orthopaedic and Trauma Surgery, 2015

Publication

<1 %

11

[www.ppjpi.unair.ac.id](http://www.ppjpi.unair.ac.id)

Internet Source

<1 %

12

Birte Technau, Ester Rut Unnsteinsdóttir, Fokje L. Schaafsma, Susanne Kühn. "Plastic and other anthropogenic debris in Arctic fox (*Vulpes lagopus*) faeces from Iceland", Polar Biology, 2022

Publication

<1 %

13

[journal.fkm.ui.ac.id](http://journal.fkm.ui.ac.id)

Internet Source

<1 %

Exclude quotes Off

Exclude matches < 10 words

Exclude bibliography On



# Antropometric study on the parameters of double-bundle anterior cruciate ligament reconstruction using magnetic resonance imaging

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---