

Policy to Decrease Low Birth Weight in Indonesia: Who Should Be the Target?

by Ratna Dwi Wulandari

Submission date: 07-Apr-2023 02:47PM (UTC+0800)

Submission ID: 2058231666

File name: rease_Low_Birth_Weight_in_Indonesia_Who_Should_Be_the_Target.pdf (2.51M)

Word count: 6703

Character count: 36763

Article

Policy to Decrease Low Birth Weight in Indonesia: Who Should Be the Target?

Ratna Dwi Wulandari ^{1,2,*}, Agung Dwi Laksono ^{2,3} and Ratu Matahari ⁴

- ¹ Faculty of Public Health, Universitas Airlangga, Surabaya 60115, Indonesia
² The Airlangga Centre for Health Policy (ACeHAP), Surabaya 60115, Indonesia
³ National Research and Innovation Agency, the Republic of Indonesia, Jakarta 10340, Indonesia
⁴ Faculty of Public Health, Ahmad Dahlan University, Yogyakarta 55164, Indonesia
* Correspondence: ratna-d-w@fkm.unair.ac.id

Abstract: The study aimed to analyze the target of the policy to decrease low birth weight (LBW) in Indonesia. This cross-sectional study used a sample of live births in last five years preceding the survey of birth weight. Data collection took place from July to September 2017. The weighted sample size was 17,848 participants. The variables analyzed included residence, age, marital status, education, employment, parity, and wealth. The study employed binary logistic regression in the final stage to determine the target of policy regarding LBW. The results showed that women in urban areas were 1.200 times more likely to deliver babies with LBW than women in rural areas. All age groups were less likely to deliver babies with LBW than those aged 45–49. The study also found all marital statuses had a lower likelihood of providing babies with LBW than those who had never been in a marriage. Women of all education levels had a greater risk of giving birth to babies with LBW than women with higher education levels. Unemployed women had 1.033 times more chances of delivering babies with LBW than employed women. Primiparous women were 1.132 times more likely to give birth to babies with LBW than multiparous women. Overall, the women in all wealth status categories had a higher probability of delivering babies with LBW than the wealthiest groups. The study concluded that policymakers should target women who live in urban areas, are old, have never been married, have low education, and are unemployed, primiparous, and poor to decrease LBW cases in Indonesia.

Keywords: low birth weight; maternal and child health; maternal health; public health; nutrition; health policy



Citation: Wulandari, R.D.; Laksono, A.D.; Matahari, R. Policy to Decrease Low Birth Weight in Indonesia: Who Should Be the Target? *Nutrients* **2023**, *15*, 465. <https://doi.org/10.3390/nu15020465>

Academic Editor: Haider Mannan

Received: 2 December 2022

Revised: 6 January 2023

Accepted: 12 January 2023

Published: 16 January 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

A newborn should have an average body weight of ≥ 2500 g. The baby is categorized as having a low birth weight if its weight is < 2500 g (1500–2499 g). Birth weight is crucial for infant development because it is associated with infant mortality [1,2]. A study states that birth weight is an essential indirect indicator of maternal nutrition and a predictive predictor of potential infant mortality and malnutrition. Due to these issues, researchers nowadays measure birth weight as an indicator of health. Babies born with low birth weight are at high risk of neonatal mortality. They are also at risk of experiencing stunting, delays in brain nerve development, and other diseases at this stage of development [3,4]. Another impact of LBW is increasing neonatal deaths globally by around 60–80% [5].

The incidence of LBW babies is more than 20 million births each year globally. Overall, the global prevalence shows that the LBW rate in developed regions was 27.2%, compared to 17.3% in Asia, 5.6% in Central Asia, and 27.2% in Southern Asia [3]. Low birth weight became the focus of the World Health Organization and United Nations Children's Fund (UNICEF) in 1992, and then the WHO targeted ending malnutrition by 2025 globally. Based on the WHO's data, 15% of babies worldwide experienced LBW, and more than half were

babies born in Asia [6]. Risk factors related to the incidence of LBW include socioeconomic and maternal characteristics. Socioeconomics deals with the type of residence, occupation type, parents' educational status, and wealth index [7]. At the same time, maternal factors include preterm delivery, a history of low birth weight, maternal age, height, hemoglobin level, lack of iron supplementation, and frequency of Antenatal Care (ANC) visits [8]. LBW is also related to exclusive breastfeeding behavior, which significantly supports a baby's growth and development in the first two years [4].

A previous study showed that other factors that influence the incidence of LBW, including cultural practices, are crucial to be addressed by health workers in providing maternal and child health care services. Indonesia is rich in artistic methods for treating diseases and caring for mothers and newborns, which can cause maternal and infant mortality [9]. The trend of LBW babies in Indonesia in 2018 was 6.2%. This figure had decreased by 4% from 2013, when the rate was 10.2% [10]. Although Indonesia has experienced a decrease in the number of LBW babies, this percentage has not met the target reduction of cases of 3% per year [10]. The situation is also related to the number of stunting incidents resulting from LBW. Indonesia is a developing country ranked fifth for the highest LBW babies out of 88 countries worldwide [11]. The incidence of stunting in Indonesia reached 37.8% in 2015 and 31% in 2018. This decrease in percentage has not yet attained the annual reduction target of 7.3% [12].

Based on the factors related to LBW, the Indonesian government strives and is committed to accelerating nutrition levels through the 'Scaling Up Nutrition (SUN)' program. The government includes long-term, medium-term, and short-term development plans regulated by the law. In the long-term development plan (2005–2025), the Indonesian government focuses on the first 1000 days of life through the fulfilment of nutrition for the womb until the baby is two. The government carries out cross-sectoral synergies in concrete steps to improve food production, processing, and consumption that meet nutritional needs. Such actions increase nutrition awareness, promotion, education regarding physical and health activities, and access to quality science- and technology-based nutrition services [13,14]. Furthermore, the Indonesian government has initiated the 'Desa Siaga' or Alert Village program since 1994. It is a community engagement program that seeks to help pregnant women by maximizing the resources available, such as access to transportation, costs, and social support in their living areas [15].

The most critical part of implementing the policy for LBW prevention is educating pregnant women and young women (adolescents) who will prepare for pregnancy. Health workers can attempt to minimize beliefs in cultural behavior that can endanger pregnant women's and babies' health and safety and improve the quality of baby feeding, especially exclusive breastfeeding [16,17]. Advocacy and dissemination of information to the community by healthcare workers will help empower knowledge of LBW prevention [9]. Based on the background, the study aimed to analyze the target of the policy to decrease LBW cases in Indonesia.

2. Materials and Methods

2.1. Data Source and Study Design

This cross-sectional study analyzed secondary data from the 2017 Indonesia Demographic and Health Survey (IDHS). It used a sample of live births in the last five years preceding the survey, comprising birth weight, written records, and maternal memories. The author used the 'IDIR71FL_Individual Recode' files.

The stratified two-stage sampling design used in the 2017 IDHS was as follows: Stage 1: selecting several census blocks systematically proportional to size probability with the number of households resulting from the 2010 population census listing. In this example, an implicit stratification procedure based on urban and rural regions was used for sorting census blocks based on the wealth index category of the 2010 population census data. Stage 2 picks 25 ordinary households in each census block based on updating the households in each census block [18]. The 2017 IDHS collected data from July to September

2017. The 2017 IDHS investigated the babies' mothers as respondents among live births within the last five years. Around 94% of mothers reported their babies' birth weight; the study selected as many as 17,848 participants as the sample in the final process.

2.2. Variables

The study used LBW as a dependent variable. It defined LBW as a birth weight of fewer than 2500 g regardless of gestational age [18]. Apart from LBW as the dependent variable, the study analyzed seven independent variables: the type of residence, age group, marital status, education level, employment status, parity, and wealth status.

Following the Indonesian Central Bureau of Statistics, the study divided residences into urban and rural categories. The study split ages into seven groups: 15–19, 20–24, 25–29, 30–34, 35–39, 40–44, and 45–49 years. Education level, the last certificate owned, was categorized into four categories: no education, primary, secondary, and higher. Marital status had three categories: never in marriage, married/living with a partner, and divorced/widowed.

Meanwhile, this study divided employment status into two categories: unemployed and employed. Parity, the acknowledgment of the number of live babies ever born, was studied in three categories, namely primiparous (<2), multiparous (2–4), and grand multiparous (>4).

The 2017 IDHS described wealth status as the socioeconomic quintile in a household. It was related to household income as seen from the styles and prices of furniture, such as televisions, bicycles, motorcycles, and household products, including drinking water supplies, bathroom amenities, and flooring materials. The 2017 IDHS determined the value of this variable by using the principal component analysis in the report. Every household's score for the national wealth quintiles was then grouped into the same five classes, accounting for 20% of the population [10]. They were the poorest (quintile 1), poorer (quintile 2), middle (quintile 3), richer (quintile 4), and richest (quintile 5) [19,20].

2.3. Data Analysis

The study employed the bivariate test to analyze all data investigated at the first stage. In the bivariate analysis, the researchers used the chi-square test. Before conducting a binary logistic regression test, the study performed a co-linearity test to ensure no multicollinearity symptoms existed between the independent variables. The study used binary logistic regression to determine the policy's target at the final stage. The author processed all statistical analyses in SPSS 26 software.

The study used ArcGIS 10.3 to map the percentage distribution of LBW cases by provinces in Indonesia (ESRI Inc., Redlands, CA, USA). The study collected the shapefile of administrative boundary polygons from the Indonesian Bureau of Statistics.

2.4. Ethical Approval

The study used secondary data from the 2017 IDHS for a materials analysis. The survey removed the identities of all respondents from the 2017 IDHS dataset. Participants in this study signed written consent forms, and the children's parents or guardians gave their consent (under 16 years). The author has obtained permission to use data for this study through the website <https://dhsprogram.com> (accessed on 1 November 2020).

The 2017 IDHS adheres to the Standard D.H.S. survey protocol under the Demographic and Health Surveys Program (DHS-7), which has been approved by ICF International's Institutional Review Board and was previously reviewed and approved by the ORC Macro IRB in 2002. DHS surveys that adhere to the Standard are classified as DHS-7 program-approved, and the approval document is attached. ICF International's Institutional Review Board followed the US Department of Health and Human Services requirements for the "Protection of Human Subjects" (45 CFR 46).

3. Results

The national average percentage of LBW was 6.981% (95% CI 6.607–7.355%). At least 18 out of the 34 provinces in Indonesia have LBW rates above the national average. Figure 1 shows the percentage distribution of LBW cases by the provinces in Indonesia. The areas with a high percentage of LBW cases tended to be in the Northern Central parts of Indonesia.




Figure 1. Distribution of LBW percentage by the province in Indonesia, 2017.

Table 1 presents the descriptive analysis of the LBW cases in Indonesia. The percentage is in the column percentage. Based on residence, women in urban areas tended to give birth to LBW babies more, and women in the 25–29 age group broadly delivered LBW babies more. In addition to these variables, married women or women who lived with partners tended to give birth to LBW babies more.

In terms of education, women with secondary education were more likely to deliver LBW babies. Unemployed women and multiparous women tended to give birth to LBW babies. Moreover, wealthier women were more likely to have LBW babies.

The collinearity test result indicated no symptoms of multicollinearity between the independent variables tested. The tolerance value of all variables was greater than 0.10. Simultaneously, the variance inflation factor (VIF) value for all independent variables was less than 10.00. Then, referring to the basis of decision-making in the multicollinearity test, the study concluded that there was no indication of a strong relationship between the independent variables in the regression model.

According to maternal education level, mothers with only primary school education and less ruled in both nutritional status categories. Based on maternal age, stunted children under the age of five have mothers with an average age slightly older than normal children under five years.

Table 1. The descriptive analysis of LBW in Indonesia, 2017.

Variables	Low Birth Weight		p-Value
	No (n = 16,602)	Yes (n = 1246)	
Type of residence			<0.001
Urban	48.3%	50.9%	
Rural	51.7%	49.1%	
Age group			<0.001
15–19	2.3%	2.9%	
20–24	16.3%	17.7%	
25–29	25.7%	25.2%	
30–34	26.4%	23.1%	
35–39	19.6%	21.5%	
40–44	8.0%	7.6%	
45–49	1.7%	2.0%	
Marital status			<0.001
Never in union	0.1%	0.2%	
Married/Living with a partner	97.2%	96.0%	
Divorced/Widowed	2.7%	3.9%	
Education Level			<0.001
No education	1.2%	1.0%	
Primary	25.6%	29.4%	
Secondary	57.6%	56.8%	
Higher	15.6%	12.8%	
Employment status			<0.001
Unemployed	55.0%	56.6%	
Employed	45.0%	43.4%	
Parity			<0.001
Primiparous	29.3%	31.5%	
Multiparous	70.7%	68.5%	
Wealth status			<0.001
Poorest	20.6%	21.1%	
Poorer	20.0%	21.8%	
Middle	20.1%	20.0%	
Richer	20.2%	20.1%	
Richest	19.1%	17.0%	

Table 2 displays the binary logistic regression results for LBW cases. The study referenced the category “LBW = No” at this final stage. This analysis indicated that women in rural areas are less likely to give birth to LBW babies in Indonesia. Women living in urban areas are 1.200 times more likely to provide LBW babies than those in rural areas (AOR: 1.200; 95% CI: 1.200–1.200).

While women in all age group categories, except the 45–49 age group, are less likely to give birth to LBW babies. Moreover, pregnant women in all marital status categories, except those who have never been married, are less likely to give birth to LBW babies.

Women in all education level categories showed a higher probability of giving birth to LBW babies than women with higher education levels. Concerning education level, unemployed women are 1.033 times more likely to deliver LBW babies than those employed (AOR: 1.033; 95% CI: 1.032–1.033). This analysis showed that unemployment is a risk factor for women giving birth to LBW babies in Indonesia.

Regarding parity, primiparous women are 1.132 times more likely to labor LBW babies than women with many children (AOR: 1.132; 95% CI: 1.132–1.132). The information indicated that being primiparous is a risk factor for pregnant women giving birth to LBW babies in Indonesia.

Table 2. The result of binary logistic regression on LBW in Indonesia, 2017.

Predictor	Low Birth Weight			
	p-Value	AOR	95% CI	
			Lower Bound	Upper Bound
Residence: Urban	<0.001	1.200	1.200	1.200
Residence: Rural (ref.)	-	-	-	-
Age: 15–19	<0.001	0.932	0.932	0.933
Age: 20–24	<0.001	0.870	0.870	0.871
Age: 25–29	<0.001	0.837	0.836	0.837
Age: 30–34	<0.001	0.774	0.774	0.774
Age: 35–39	<0.001	0.958	0.957	0.958
Age: 40–44	<0.001	0.813	0.813	0.813
Age: 45–49 (ref.)	-	-	-	-
Marital: Never in union (ref.)	-	-	-	-
Marital: Married/Living with a partner	<0.001	0.399	0.399	0.400
Marital: Divorced/Widowed	<0.001	0.550	0.550	0.551
Education: No education	<0.001	1.018	1.018	1.019
Education: Primary	<0.001	1.361	1.360	1.361
Education: Secondary	<0.001	1.146	1.145	1.146
Education: Higher (ref.)	-	-	-	-
Employment: Unemployed	<0.001	1.033	1.032	1.033
Employment: Employed (ref.)	-	-	-	-
Parity: Primiparous	<0.001	1.132	1.132	1.132
Parity: Multiparous (ref.)	-	-	-	-
Wealth: Poorest	<0.001	1.111	1.111	1.111
Wealth: Poorer	<0.001	1.181	1.181	1.181
Wealth: Middle	<0.001	1.067	1.067	1.067
Wealth: Richer	<0.001	1.079	1.079	1.079
Wealth: Richest (ref.)	-	-	-	-

AOR: adjusted odds ratio; LBW: low birth weight.

Finally, Table 2 informs us that the poorest women are 1.111 times more likely to give birth to LBW babies than the most prosperous women (AOR: 1.111; 95% CI: 1.111–1.111). Women of poorer wealth status are 1.181 times more likely to deliver LBW babies than the most affluent women (AOR: 1.181; 95% CI: 1.181–1.181). Women in the middle wealth status group have 1.067 times more chances of giving birth to LBW babies than the most affluent women (AOR: 1.067; 95% CI: 1.067–1.067). Besides, wealthier women are 1.079 times more likely to give birth to LBW babies than the most prosperous women (AOR: 1.079; 95% CI: 1.079–1.079). This analysis implied that women in all wealth status categories except the most decadent have a higher probability of giving birth to LBW babies.

4. Discussion

Nationally, the average percentage of LBW cases, at 6.981% in 2017, was below the target of the National Medium-Term Development Plan in 2015–2019. However, at least 10 out of 34 provinces in Indonesia did not experience a lower percentage of LBW cases below this target. Besides, the Indonesian government recorded 18 regions with a prevalence of LBW cases above the national average [10]. This study informs us that women living in urban areas were more likely to give birth to LBW babies in Indonesia. This finding aligns with the previous research in Afghanistan, reporting that mothers who lived in rural residences were 0.3 times less likely to give birth to LBW babies than those in urban areas [21]. This condition is likely related to higher pollution in urban than rural areas of Indonesia. A previous study found that higher paternal prenatal concentrations of mono-benzyl phthalate and mono-carboxyisononyl phthalate were linked to a 40% and 53% increase in the incidence of LBW in infants who were born spontaneously [22]. Moreover,

previous studies have indicated that a higher pollution level was associated with a higher incidence of LBW [23,24].

This finding contradicts several previous studies. Studies in Ethiopia and Iran provided different conclusions. Mothers in rural Ethiopia and Iran were 1.39 and 1.4 times more likely to give birth to LBW babies than those in urban areas [25]. Another study in Sidama Zone, South Ethiopia, used public hospitals as the research setting for newborns. It found that mothers in rural areas were 3.51 times more likely to give birth to LBW patients than those in urban areas [26].

Based on the age group, the result showed women in all age group categories except the 45–49 age group had a lower likelihood of giving birth to LBW babies. Previous studies in Jordan and Ethiopia also indicated that age influenced LBW incidence [25,27]. A study in Germany reported different results, finding that teenage mothers had higher rates of depression during pregnancy than mothers of older generations. The research indicated that this factor puts adolescent mothers at risk of giving birth to LBW babies [28].

Women in all marital status categories, except those who had never married, had a lower chance of giving birth to LBW babies. An earlier study in Sub-Saharan Africa also reported marital status was associated with LBW babies. Divorced or widowed mothers were more likely to deliver LBW babies than married mothers [29]. Having a partner is a protective factor for women, and wives become a place to share psychological and financial burdens [30]. Besides, mothers in all education level categories had a higher probability of giving birth to LBW babies than women with higher education levels. A study in Jordan and Ethiopia also found similar results: a higher education level was a protective factor for mothers not giving birth to LBW babies [25]. Meanwhile, previous studies in Indonesia showed that pregnant women with primary school education were 2.154 times more likely to give birth to LBW babies (AOR 2.154, 95% CI 1.399–3.315; $p = 0.004$) [1]. Because teaching is a motivating opportunity to live healthily, education is good for your health. Women who receive an education will have more opportunities to develop their abilities and lead healthy lives [31]. Higher-educated women were more likely to experience a better pregnancy and to experience a lower probability of delivering LBW kids [16,32]. A better education level allows mothers to better understand actions needed to provide the best output for themselves and their children [16,33,34]. Many studies have reported that better education is a strong determinant of better production in the health sector [35–38]. Instead, poor education is a barrier to producing quality performance in the health sector [8,32,39]. Unemployment was deemed a risk factor for women giving birth to LBW babies. A previous study in hospital settings in the Gambata-Tembaro zone, Southern Ethiopia, found that unemployed mothers could be 4 times more likely to deliver LBW babies than employed mothers [40]. Another study at a large European maternity hospital found similar effects of a multivariable analysis: unemployed women or homemakers increased adjusted odds ratios for LBW [41]. This condition was possibly related to employment and wealth status. Unemployed women are among low-income families [42].

Primiparous women had a higher probability of giving birth to LBW babies than women with many children. A study in Brazil also reported that primiparous mothers had up to 1.62 times the chance of delivering LBW babies than mothers with many children [43]. Moreover, studies in Sub-Saharan Africa and China also provided similar results, indicating that multiparity was associated with reduced cases of LBW [29]. Research in Southern Africa informed different findings that mothers with more than three births were 1.5 times more likely to give birth to LBW babies [40].

Moreover, this study revealed that women in all wealth status categories, compared to the wealthiest, had a higher probability of giving birth to LBW babies. An earlier study analyzing secondary data from the 2016 Ethiopia Demographic and Health Survey in Ethiopia also reported similar results with the incidence of LBW being dominant among the poor [44]. Poor households have limitations in providing food for all family members, including pregnant women. For pregnant women, the mother's nutritional status is an indicator of the adequacy of food and nutrition. Better wealth status is related to the

availability of food in the family and thus is closely related to food intake during pregnancy. Lack of food availability in terms of quantity and quality. This is related to the increasing need for macronutrients and micronutrients in pregnant women and their importance for the mother and fetus. Therefore, a better wealth status can reduce the incidence of LBW [45]. In terms of food insecurity and consumption, a previous study described acute and chronic malnutrition. Reducing malnutrition in children could be done through prenatal measures and supplementary food for children during food insecurity [45].

Study Limitation

The 2017 IDHS derived the data from a stratified multi-stage sampling procedure. The sampling method allows for an unequal probability of selection.

This current study analyzed secondary data limited to variables reported in the IDHS. It excluded several variables known as determinants of LBW incidence in previous studies. Some were antenatal care during pregnancy [15,45], smoke pollution [46], wanting a later child [25], pregnancy interval and gestational age of <37 weeks at birth [47], hypertension, mid-upper arm circumference [26], maternal height [48], hemoglobin levels [49], and intimate partner violence during pregnancy [50].

Additionally, this study employed a quantitative approach and thus did not uncover the social and cultural phenomena related to LBW regarding family and children value [51–53], pregnancy value, and dietary restrictions in pregnant women [54], and a patriarchal social structure in Indonesia that places women subordinate to men [55].

5. Conclusions

Based on the results, policymakers should target pregnant women who live in urban areas, are old, have never been in a marriage, have low education, and are unemployed, primiparous, and poor to decrease LBW cases in Indonesia. The policymaker should target prospective mothers with a higher risk of having LBW babies.

Author Contributions: Conceptualization, R.D.W. and A.D.L.; methodology, R.D.W.; software, 15 D.L.; validation, R.D.W. and A.D.L.; formal analysis, A.D.L.; investigation, R.M.; resources, R.M.; data curation, R.M.; writing—original draft, 43 eparation, A.D.L.; writing—review and editing, R.D.W.; 3; sualization, R.M.; supervision, R.D.W.; project administration, R.M.; funding acquisition, R.D.W. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: The 2017 IDHS adheres to the Standard D.H.S. survey protocol under the Demographic and Health Surveys Program (DHS-7), which has been approved by ICF International's Institutional Review Board and was previously reviewed and approved by the ORC Macro IRB in 2002. DHS surveys that adhere to the Standard are classified as DHS-7 program-approved, and 17 e approval document is attached. ICF International's Institutional Review Board followed the US Department of Health and Human Services requirements for the "Protection of Human Subjects" (45 CFR 46).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: The author cannot publicly share the data because a third party and the authors who own the data do not have permission to share the data. The 2017 IDHS data set requested from the ICF ('data set of childbearing age women') is available from the ICF contact via <https://dhsprogram.com> (accessed on 1 November 2020) for researchers who meet the criteria for access to confidential data.

Acknowledgments: The author would like to thank ICF International, which has agreed to allow the author to analyze the 2017 IDHS data in this article.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Siramaneerat, I.; Agushybana, F.; Meebunmak, Y. Maternal risk factors associated with low birth weight in Indonesia. *Open Public Health J.* **2018**, *11*, 376–383. [CrossRef]
2. Mardiaturrmah, M.; Anjarwati, A. Relationship between the nutritional status of pregnant women and the incident of Low Birth Weight infant. *Int. J. Health Sci. Technol.* **2020**, *1*, 58–62. [CrossRef]
3. Mediani, H.S. Predictors of Stunting Among Children Under Five Year of Age in Indonesia: A Scoping Review. *Glob. J. Health Sci.* **2020**, *12*, 83. [CrossRef]
4. Aryastami, N.K.; Shankar, A.; Kusumawardani, N.; Besral, B.; Jahari, A.B.; Achadi, E. Low birth weight was the most dominant predictor associated with stunting among children aged 12–23 months in Indonesia. *BMC Nutr.* **2017**, *3*, 16. [CrossRef]
5. Shrestha, S.; Shrestha, S.; Shakya Shrestha, U.; Gyawali, K. Predictors of Low Birth Weight at Lumbini Provincial Hospital, Nepal: A Hospital-Based Unmatched Case Control Study. *Adv. Prev. Med.* **2020**, *2020*, 8459694. [CrossRef]
6. United Nations Children’s Fund; World Health Organization. *UNICEF-WHO Low Birthweight Estimates: Levels and Trends 2000–2015*; World Health Organization: Geneva, Switzerland, 2019. Available online: <https://apps.who.int/iris/bitstream/handle/10665/324783/WHO-NMH-NHD-19.21-eng.pdf?ua=1> (accessed on 22 December 2020).
7. Utami, S.M.; Wulandari, R.D. Factors affecting the occurrence of low birth weight. *Eur. Asian J. Biosci.* **2020**, *14*, 2657–2661.
8. Masruroh Yusuf, A.; Rohmah, N.; Pakki, I.B.; Sujoso, A.D.P.; Andayani, Q.; Laksono, A.D. Neonatal Death Incidence in Healthcare Facility in Indonesia: Does Antenatal Care Matter? *Indian J. Forensic. Med. Toxicol.* **2021**, *15*, 1265–1271.
9. Sutan, R.; Berkati, S. Does cultural practice affects neonatal survival- a case control study among low birth weight babies in Aceh Province, Indonesia. *BMC Pregnancy Childbirth* **2014**, *14*, 342. [CrossRef]
10. National Population and Family Planning Board; Statistics Indonesia, Ministry of Health. *Indonesia Demographic and Health Survey, 2017: Final Report*; BKKBN, BPS, Kemenkes, and ICF: Jakarta, Indonesia, 2017. Available online: <https://www.dhsprogram.com/publications/publication-FR342-DHS-Final-Reports.cfm> (accessed on 17 December 2022).
11. BAPPENAS; UNICEF. *SDG Baseline Report on Children in Indonesia (Laporan Baseline SDG tentang Anak-Anak di Indonesia)*; Kementerian Perencanaan Pembangunan Nasional (Bappenas) dan United Nations Children’s Fund: Jakarta, Indonesia, 2017. Available online: <https://www.unicef.org/indonesia/media/1471/file/SDG%20Baseline%20report%20Indonesian.pdf> (accessed on 22 December 2020).
12. Ministry of Health Republic of Indonesia. *Basic Health Survey (RISKESDAS) 2018. Main Results of Basic Health Survey (RISKESDAS) 2018*; Ministry of Health Republic of Indonesia: Jakarta, Indonesia, 2018; Volume 44. Available online: <http://repository.bkpk.kemkes.go.id/3514/1/Laporan%20Riskasdas%202018%20Nasional.pdf> (accessed on 30 December 2022).
13. Aryastami, N.K. Policy Studies and Management of Stunting Nutritional Problems in Indonesia (Kajian Kebijakan dan Penganggulangan Masalah Gizi Stunting di Indonesia). *Bul. Penelit. Kesehat* **2017**, *45*, 233–240.
14. Indonesian Ministry of Health. *My Peaceful Healthy Family, My City’s Healthy City (Keluarga Sehat Idamanku, Kota Sehat Kotaku)*; Indonesian Ministry of Health: Jakarta, Indonesia, 2020; Volume 5. Available online: <https://promkes.kemkes.go.id/?p=7212> (accessed on 27 December 2022).
15. Hill, P.S.; Goeman, L.; Sofiarini, R.; Djara, M.M. “Desa SIAGA”, the “Alert Village”: The evolution of an iconic brand in Indonesian public health strategies. *Health Policy Plan.* **2014**, *29*, 409–420. [CrossRef]
16. Kusriani, I.; Fuada, N.; Supadmi, S.; Laksono, A.D. Education as Predictor of Low Birth Weight among Female Worker in Indonesia. *Medico-Legal Updat* **2021**, *21*, 360–365.
17. Laksono, A.D.; Wulandari, R.D.; Kusriani, I.; Ibad, M. The effects of mother’s education on achieving exclusive breastfeeding in Indonesia. *BMC Public Health* **2021**, *21*, 14. [CrossRef]
18. Wulandari, R.D.; Laksono, A.D.; Prasetyo, Y.B.; Nandini, N. Socioeconomic Disparities in Hospital Utilization Among Female Workers in Indonesia: A Cross-Sectional Study. *J. Prim. Care Community Health* **2022**, *13*, 21501319211072679. [CrossRef]
19. Wulandari, R.D.; Qomarrudin, M.B.; Supriyanto, S.; Laksono, A.D. Socioeconomic Disparities in Hospital Utilization among Elderly People in Indonesia. *Indian J. Public Health Res. Dev.* **2019**, *10*, 1800–1804.
20. National Development Planning Agency of Indonesia. *Indonesian National Medium Term Development Plan in 2015–2019*; National Development Planning Agency of Indonesia: Jakarta, Indonesia, 2015. Available online: <https://www.bappenas.go.id/id/data-dan-informasi-utama/dokumen-perencanaan-dan-pelaksanaan/dokumen-rencana-pembangunan-nasional/rpjm-2005-2025/rpjm-2015-2019/> (accessed on 3 January 2020).
21. Gupta, R.D.; Swasey, K.; Burrows, V.; Hashan, M.R.; Al Kibria, G.M. Factors associated with low birth weight in Afghanistan: A cross-sectional analysis of the demographic and health survey 2015. *BMJ Open* **2019**, *9*, e025715. [CrossRef]
22. Zang, L.; Lv, H.; Du, J.; Pan, Y.; Lin, Y.; Dai, J. Association of phthalate exposure with low birth weight in couples conceiving naturally or via assisted reproductive technology in a prospective birth cohort. *Sci. Total Environ.* **2023**, *855*, 158852. [CrossRef]
23. Strickland, M.J.; Lin, Y.; Darrow, L.A.; Warren, J.L.; Mulholland, J.A.; Chang, H.H. Associations Between Ambient Air Pollutant Concentrations and Birth Weight: A Quantile Regression Analysis. *Epidemiology* **2019**, *30*, 624–632. [CrossRef]
24. Trerotoli, P.; Bartolomeo, N.; Leogrande, S.; Triggiani, S.; Mincuzzi, A.; Serio, G.; Minerba, A.S. Survey of Low Birthweight and Extremely Low Birthweight Events in a High Environmental Risk Area of Apulia, Italy. *Int. J. Environ. Res.* **2021**, *15*, 11–17. [CrossRef]

25. Kassaw, M.W.; Abebe, A.M.; Kassie, A.M.; Abate, B.B.; Masresha, S.A. Trends of proximate low birth weight and associations among children under-five years of age: Evidence from the 2016 Ethiopian demographic and health survey data. *PLoS ONE* **2021**, *16*, e0246587. [[CrossRef](#)]
26. Bekela, M.B.; Shimbire, M.S.; Gebabo, T.F.; Geta, M.B.; Tonga, A.T.; Zeleke, E.A.; Sidemo, N.B.; Getnet, A.B. Determinants of Low Birth Weight among Newborns Delivered at Public Hospitals in Sidama Zone, South Ethiopia: Unmatched Case-Control Study. *J. Pregnancy* **2020**, *2020*, 4675701. [[CrossRef](#)]
27. Islam, M.M.; Ababneh, F.; Akter, T.; Khan, H.R. Prevalence and risk factors for low birth weight in Jordan and its association with under-five mortality: A population-based analysis. *East. Mediterr. Health J.* **2020**, *26*, 1273–1284. [[CrossRef](#)] [[PubMed](#)]
28. Wong, S.P.W.; Twynstra, J.; Gilliland, J.A.; Cook, J.L.; Seabrook, J.A. Risk Factors and Birth Outcomes Associated with Teenage Pregnancy: A Canadian Sample. *J. Pediatr. Adolesc. Gynecol.* **2020**, *33*, 153–159. [[CrossRef](#)] [[PubMed](#)]
29. Tessema, Z.T.; Tamirat, K.S.; Teshale, A.B.; Tesema, G.A. Prevalence of low birth weight and its associated factor at birth in Sub-Saharan Africa: A generalized linear mixed model. *PLoS ONE* **2021**, *16*, e0248417. [[CrossRef](#)] [[PubMed](#)]
30. Megatsari, H.; Laksono, A.D.; Herwanto, Y.T.; Sarweni, K.P.; Geno, R.A.P.; Nugraheni, E.; Ibad, M. Does husband/partner matter in reduce women's risk of worries?: Study of psychosocial burden of COVID-19 in Indonesia. *Indian J. Forensic. Med. Toxicol.* **2021**, *15*, 1101–1106.
31. Pakaya, N.; Laksono, A.D.; Masruroh Sujoso, A.D.P.; Ibrahim, I.; Marasabessy, N.B.; Rohmah, N.; Seran, A.A.; Wulansari, I. Are education level of women related to contraceptive use? Analysis of the 2017 Indonesia demographic and health survey. *Int. J. Health Sci. (Qassim)* **2022**, *6*, 5561–5570.
32. Rohmah, N.; Yusuf, A.; Hargono, R.; Laksono, A.D.; Masruroh Ibrahim, I.; Walid, S. Determinants of teenage pregnancy in Indonesia. *Indian J. Forensic. Med. Toxicol.* **2020**, *14*, 2080–2085.
33. Wulandari, R.D.; Laksono, A.D. Determinants of knowledge of pregnancy danger signs in Indonesia. *PLoS ONE* **2020**, *15*, e0232550. [[CrossRef](#)]
34. Laksono, A.D.; Ibad, M.; Mursita, A.; Kusriani, I.; Wulandari, R.D. Characteristics of mother as predictors of stunting in toddler. *Pakistan J. Nutr.* **2019**, *18*, 1101–1106. [[CrossRef](#)]
35. Ipa, M.; Widawati, M.; Laksono, A.D.; Kusriani, I.; Dhewantara, P.W. Variation of preventive practices and its association with malaria infection in eastern Indonesia: Findings from community-based survey. *PLoS ONE* **2020**, *15*, e0232909. [[CrossRef](#)]
36. Megatsari, H.; Laksono, A.D.; Ibad, M.; Herwanto, Y.T.; Sarweni, K.P.; Geno, R.A.P.; Nugraheni, E. The community psychosocial burden during the COVID-19 pandemic in Indonesia. *Heliyon* **2020**, *6*, e05136. [[CrossRef](#)]
37. Seran, A.A.; Antaria, M.D.; Haksama, S.; Setijaningrum, E.; Laksono, A.D.; Prahastuti Sujoso, A.D. Disparities of the use of hormonal and non-hormonal contraceptive drugs in urban and rural areas in Indonesia and the world. *Syst. Rev. Pharm.* **2020**, *11*, 66–73.
38. Laksono, A.D.; Wulandari, R.D.; Ibad, M.; Herwanto, Y.T.; Sarweni, K.P.; Geno, R.A.P.; Nugraheni, E. Predictors of Healthy Lifestyle in The COVID-19 Pandemic Period in East Java, Indonesia. *J. Crit. Rev.* **2020**, *7*, 1515–1521.
39. Laksono, A.D.; Wulandari, R.D. The Barrier to Maternity Care in Rural Indonesia. *J. Public Health* **2022**, *30*, 135–140. [[CrossRef](#)]
40. Alemu, A.; Abageda, M.; Assefa, B.; Melaku, G. Low birth weight: Prevalence and associated factors among newborns at hospitals in kambata-tembaro zone, southern Ethiopia 2018. *Pan Afr. Med. J.* **2019**, *34*, 68. [[CrossRef](#)]
41. Reynolds, C.M.E.; McMahon, L.E.; O'Malley, E.G.; O'Brien, O.; Sheehan, S.R.; Turner, M.J. Maternal employment and pregnancy outcomes in a large European maternity hospital. *Eur. J. Obstet. Gynecol. Reprod Biol.* **2020**, *250*, 86–92. [[CrossRef](#)]
42. Wulandari, R.D.; Laksono, A.D. The association of socioeconomic status on early marriage among women in Rural Indonesia (Hubungan status ekonomi terhadap pernikahan dini pada perempuan di perdesaan Indonesia). *J. Kesehatan Reproduksi.* **2020**, *11*, 115–124. [[CrossRef](#)]
43. Falcão, I.R.I.R.; Ribeiro-Silva, R.D.C.R.D.C.; De Almeida, M.F.M.F.; Fiaccone, R.L.R.L.; Dos Srocha, A.; Ortelan, N.; Silva, N.J.N.J.; Paixao, E.S.E.S.; Ichihara, M.Y.M.Y.; Rodrigues, L.C.L.C.; et al. Factors associated with low birth weight at term: A population-based linkage study of the 100 million Brazilian cohort. *BMC Pregnancy Childbirth* **2020**, *20*, 536. [[CrossRef](#)]
44. Shibre, G.; Tamire, M. Prevalence of and socioeconomic gradient in low birth weight in Ethiopia: Further analysis of the 2016 demographic and health survey data. *BMC Pregnancy Childbirth* **2020**, *20*, 608. [[CrossRef](#)]
45. Supadmi, S.; Kusriani, I.; Fuada, N.; Laksono, A.D. The Low Birth Weight in Indonesia: Does Antenatal Care Matter? *Int. J. Innov. Creat Chang.* **2020**, *14*, 490–500.
46. Islam, S.; Mohanty, S.K. Maternal exposure to cooking smoke and risk of low birth weight in India. *Sci. Total Environ.* **2021**, *774*, 145717. [[CrossRef](#)]
47. Endalamaw, A.; Engeda, E.H.; Ekubagewargies, D.T.; Belay, G.M.; Tefera, M.A. Low birth weight and its associated factors in Ethiopia: A systematic review and meta-analysis. *Ital. J. Pediatr.* **2018**, *44*, 141. [[CrossRef](#)] [[PubMed](#)]
48. Desta, S.A.; Damte, A.; Hailu, T. Maternal factors associated with low birth weight in public hospitals of Mekelle city, Ethiopia: A case-control study. *Ital. J. Pediatr.* **2020**, *46*, 124. [[CrossRef](#)] [[PubMed](#)]
49. Mekie, M.; Taklual, W. Magnitude of low birth weight and maternal risk factors among women who delivered in Debre Tabor Hospital, Amhara Region, Ethiopia: A facility based cross-sectional study. *Ital. J. Pediatr.* **2019**, *45*, 86. [[CrossRef](#)]
50. Musa, A.; Chojenta, C.; Loxton, D. The association between intimate partner violence and low birth weight and preterm delivery in eastern Ethiopia: Findings from a facility-based study. *Midwifery* **2021**, *92*, 102869. [[CrossRef](#)]
51. Pratita, I.; Laksono, A.D. "If this child eats, whatever she/he is asked . . .": Exploration of Value of Children and Parenting Patterns in Javanese in the Besowo village, Kediri, East Java. *Amerta Nutr.* **2020**, *4*, 147. [[CrossRef](#)]

52. Maghfiroh, M.S.; Laksono, A.D. "Given sugar water . . . at first the cry became silent, because it was full, not limp, its endurance increased"; Study of Patterns of Infant Intake ("Diberi air gula . . . awalnya nangis menjadi diam, karena kenyang, gak lemas, daya tahan tubuhnya meningkat". *S. Amerta Nutr.* **2020**, *4*, 116–122. [[CrossRef](#)]
53. Kusriani, I.; Ipa, M.; Laksono, A.D. "Is It true that the child is king? Qualitative Study of Factors Related to Nutritional Status of Children in West Lombok, Indonesia. *Indian J. Public Health Res. Dev.* **2019**, *10*, 1729–1733. [[CrossRef](#)]
54. Sholihah, A. *Ethnobiology Study of Taboo Food in Madura, Javanese, and Pendalungan Tribe Communities in Jember Regency and Its Use as Popular Scientific Books (Kajian Etnobiologi Makanan Tabu pada Masyarakat Suku Madura, Jawa, dan Pendalungan di Kabupaten Jember serta*; UNIVERSITAS JEMBER: Jember, Indonesia, 2019; pp. 1–42.
55. Laksono, A.D.; Soedirham, O.; Saptandari, P.; Wulandari, R.D. Study of family size among tenggerese in Indonesia. *Int. J. Innov. Creat Chang.* **2020**, *13*, 964–978.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

Policy to Decrease Low Birth Weight in Indonesia: Who Should Be the Target?

ORIGINALITY REPORT

16%

SIMILARITY INDEX

9%

INTERNET SOURCES

10%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Agung Dwi Laksono, Ratna Dwi Wulandari. "Relationship between Environment, Smoking Behavior, Education, Poverty, and Prevalence of Stunted Toddler in Indonesia: An Ecological Analysis", Research Square, 2020
Publication 1%
- 2 Submitted to Argosy University
Student Paper 1%
- 3 Submitted to Chung Shan Medical University
Student Paper 1%
- 4 Submitted to Poltekkes Kemenkes Banjarmasin
Student Paper 1%
- 5 Sharma, Sudesh Raj, Smith Giri, Utsav Timalina, Sanjiv Sudarshan Bhandari, Bikash Basyal, Kusum Wagle, and Laxman Shrestha. "Low Birth Weight at Term and Its Determinants in a Tertiary Hospital of Nepal: A Case-Control Study", PLoS ONE, 2015.
Publication 1%

6	www.coursehero.com Internet Source	<1 %
7	www.revistaespacios.com Internet Source	<1 %
8	Kidanemaryam Berhe, Lemlem Weldegerima, Freweini Gebrearegay, Amaha Kahsay et al. "Effect of under-nutrition during pregnancy on low birth weight in Tigray regional state, Ethiopia; a prospective cohort study", BMC Nutrition, 2021 Publication	<1 %
9	jknrh.com Internet Source	<1 %
10	pesquisa.bvsalud.org Internet Source	<1 %
11	www.ajol.info Internet Source	<1 %
12	dash.harvard.edu Internet Source	<1 %
13	jurnal.unw.ac.id Internet Source	<1 %
14	Desi Suantari. "Misconceptions and stigma against people living with HIV/AIDS: a cross-sectional study from the 2017 Indonesia	<1 %

Demographic and Health Survey",
Epidemiology and Health, 2021

Publication

15

Rivers Ingersoll, David Lentink. "How the hummingbird wingbeat is tuned for efficient hovering", The Journal of Experimental Biology, 2018

Publication

<1 %

16

Yuki Yamamoto, Ken'ichi Matsumoto. "Choice of contraceptive methods by women's status: Evidence from large-scale microdata in Nepal", Sexual & Reproductive Healthcare, 2017

Publication

<1 %

17

aphrc.org
Internet Source

<1 %

18

www.iipsindia.ac.in
Internet Source

<1 %

19

Submitted to University of Teesside
Student Paper

<1 %

20

journal.ipb.ac.id
Internet Source

<1 %

21

Submitted to Resurrection University
Student Paper

<1 %

22

www.kemhan.go.id
Internet Source

<1 %

23 Amene Ranjbar, Maliheh Shirzadfard Jahromi, Mitra Shekari, Fatemeh Darsareh, Vahid Mehrnoush. "The prevalence and associated risk factors of low birth weight in singleton term pregnancies", Research Square Platform LLC, 2022
Publication

24 Ciara M.E. Reynolds, Léan E. McMahon, Eimer G. O'Malley, Oliver O'Brien, Sharon R. Sheehan, Michael J. Turner. "Maternal employment and pregnancy outcomes in a large European maternity hospital", European Journal of Obstetrics & Gynecology and Reproductive Biology, 2020
Publication

25 Submitted to Gyeongsang National University
Student Paper

26 Submitted to University of Southampton
Student Paper

27 jcreview.com
Internet Source

28 Submitted to Middlesex University
Student Paper

29 Submitted to University of Westminster
Student Paper

30 www.intechopen.com

Internet Source

<1 %

31

ijcrr.com

Internet Source

<1 %

32

Lailatul Muniroh, Triska Susila Nindya, Santi Martini, Rondius Solfaine. "Effect of Plectranthus amboinicus Extract on BUN and Creatinine Levels and Cellular Response Proinflammatory Factors TNF- α and IL-1 β on Gout Arthritis Patients", Makara Journal of Health Research, 2014

Publication

<1 %

33

Sherly Ardi Vantono, Saptawati Bardosono, Luh Ade Ari Wiradnyani. "Mother's Coping strategies toward Food Insecurity during COVID-19 Pandemic: A Review Article", World Nutrition Journal, 2022

Publication

<1 %

34

internationalbreastfeedingjournal.biomedcentral.com

Internet Source

<1 %

35

reliefweb.int

Internet Source

<1 %

36

Umi Farida, Ansar ., Gunawan Bata Ilyas, Jamaludin ., Chuduriah Sahabudin. "Effectiveness of Desa Siaga-Alert Village Programs for Empowerment of Self-Independent Communities Case Study: Desa

<1 %

Siaga Rabies-Rabies Alert Villag of Martajaya Village, Pasangkayu District", International Journal of Engineering & Technology, 2018

Publication

37

dspace.bracu.ac.bd

Internet Source

<1 %

38

ejournal2.litbang.kemkes.go.id

Internet Source

<1 %

39

oaji.net

Internet Source

<1 %

40

scholarworks.waldenu.edu

Internet Source

<1 %

41

www.najms.org

Internet Source

<1 %

42

Peter Onubiwe Nkwo, Lucky Osaheni Lawani, Euzebus Chinonye Ezugwu, Chukwuemeka Anthony Iyoke et al. "Correlates of poor perinatal outcomes in non-hospital births in the context of weak health system: the Nigerian experience", BMC Pregnancy and Childbirth, 2014

Publication

<1 %

43

Rani Moran, Peter Dayan, Raymond J. Dolan. "Efficiency and prioritization of inference-based credit assignment", Current Biology, 2021

Publication

<1 %

44 Sri Nuryani Wahyuningrum, Ika Puspita Asturiningtyas, Cati Martiyana, Anggita Mirzautika. "Low birth weight and low mother education as dominant risk factors of stunting children in Magelang Regency, Central Java", AcTion: Aceh Nutrition Journal, 2023
Publication <1 %

45 Trias Mahmudiono, Triska Susila Nindya, Dini Ririn Andrias, Hario Megatsari, Richard R. Rosenkranz. "The effectiveness of nutrition education for overweight/obese mothers with stunted children (NEO-MOM) in reducing the double burden of malnutrition in Indonesia: study protocol for a randomized controlled trial", BMC Public Health, 2016
Publication <1 %

46 cmsproject.com
Internet Source <1 %

47 dprexternal3.dpr.go.id
Internet Source <1 %

48 journal2.uad.ac.id
Internet Source <1 %

49 www.jphres.org
Internet Source <1 %

50 www.nature.com
Internet Source <1 %

51 www.refworld.org

Internet Source

<1 %

52 Abebe Alemu, Mulatu Abageda, Biruk Assefa, Getnet Melaku. "Low birth weight: prevalence and associated factors among newborns at hospitals in Kambata-Tembaro zone, southern Ethiopia 2018", Pan African Medical Journal, 2019

Publication

<1 %

53 Agani Afaya, Richard Adongo Afaya, Thomas Bavo Azongo, Vida Nyagre Yakong et al. "Maternal risk factors and neonatal outcomes associated with low birth weight in a secondary referral hospital in Ghana", Heliyon, 2021

Publication

<1 %

54 Gebretsadik Shibre, Mulugeta Tamire. "Prevalence of and socioeconomic gradient in low birth weight in Ethiopia: further analysis of the 2016 demographic and health survey data", BMC Pregnancy and Childbirth, 2020

Publication

<1 %

55 Jason Phowira, Felicitas Tania Elvina, Igor Ian Wiguna, Fathurohman Ramadhan Hanif Bari Wahyudi, Bernie Endyarni Medise. "The Association Between Tobacco Exposure During Pregnancy and Newborns' Birth

<1 %

Weight in DKI Jakarta Community Members", Cold Spring Harbor Laboratory, 2020

Publication

56

Melaku Tadege Engidaw, Tahir Eyayu,
Tegenaw Tiruneh. "The effect of maternal
anaemia on low birth weight among
newborns in Northwest Ethiopia", Scientific
Reports, 2022

Publication

<1 %

57

Santosh Kumar Sharma, Devaki Nambiar,
Jaison Joseph. "Hidden educational
inequalities in high blood pressure and high
blood glucose levels in Kerala: evidence from
the National Family Health Survey (2019–
2021)", BMJ Open, 2023

Publication

<1 %

Exclude quotes Off

Exclude matches < 5 words

Exclude bibliography On

Policy to Decrease Low Birth Weight in Indonesia: Who Should Be the Target?

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11
