

About Journal

Unnes Journal of Public Health (UJPH) is an open access and peer-reviewed journal. It publishes original papers, reviews and short reports on all aspects of the science, philosophy, and practice of public health, especially in low-income and middle-income countries which have many problems in technology, human resources, economics, social welfare, etc impacting public health.

Unnes Journal of Public Health is Nationally Accredited (SINTA 2) based on the Decree of The Minister of Research, Technology and Higher Education of Republic Indonesia, Number 10 / E / KPT / 2019, starts from Volume 7 Issue 2 (July 2018). Unnes Journal of Public Health is published by Universitas Negeri Semarang (UNNES) in cooperation with Association of Indonesian Public Health Experts (Ikatan Ahli Kesehatan Masyarakat Indonesia (IAKMI)).

It is aimed at all public health practitioners and researchers and those who manage and deliver public health services and systems. It will also be of interest to anyone involved in provision of public health programmes, the care of populations or communities and those who contribute to public health systems in any way.



Unnes Journal of Public Health is published by Universitas Negeri Semarang (UNNES) in cooperation with the Association of Indonesian Public Health Experts (Ikatan Ahli Kesehatan Masyarakat Indonesia (IAKMI)). Unnes Journal of Public Health contains articles about health epidemiology, biostatistics, health administration, public health nutrition, environmental health, occupational health and safety, health promotion, reproductive health, maternal and child health, health management, health economics, traditional medicine/treatment for preventive or first aid purposes, public health law/ethics and other related health articles, especially in low-income and middle-income countries. The journal can be used by health practitioners, health caregivers, teachers, students, and people who interested in health issues.

UJPH considers submissions on any aspect of public health across age groups and settings.

These include:

- Public health practice and impact
- Epidemiology and Biostatistics
- Applied Epidemiology
- Need or impact assessments
- Health service effectiveness, management, and re-design
- Health Protection including control of communicable diseases
- Health promotion and disease prevention
- Evaluation of public health programs or interventions
- Public health governance, audit and quality
- Public health law and ethics
- Health policy and administration
- Capacity in public health systems and workforce
- Public health nutrition
- Environmental health
- Occupational health and safety
- Reproductive health
- Maternal and child health

This is not an exhaustive list and the Editors will consider articles on any issue relating to public health.

UJPH also publishes invited articles, reviews, and supplements from leading experts on topical issues.

 Scopus®

(<https://suggestor.step.scopus.com/progressTracker/?>

trackingID=8C1168FE0F638FE6)



(<https://scholar.google.co.id/citations?user=1arcYkEAAA&hl=id>)



Dimensions

(<https://app.dimensions.ai/discover/publication?>

search_mode=content&and_facet_source_title=jour.1279460)



(<https://sinta.kemdikbud.go.id/journals/profile/401>)



(<https://garuda.kemdikbud.go.id/journal/view/14754>)



(<https://doaj.org/toc/2252-6781>)

About Journal

Unnes Journal of Public Health (UJPH) is an open access and peer-reviewed journal. It publishes original papers, reviews and short reports on all aspects of the science, philosophy, and practice of public health, especially in low-income and middle-income countries which have many problems in technology, human resources, economics, social welfare, etc impacting public health.

Unnes Journal of Public Health is Nationally Accredited (SINTA 2) based on the Decree of The Minister of Research, Technology and Higher Education of Republic Indonesia, Number 10 / E / KPT / 2019, starts from Volume 7 Issue 2 (July 2018). Unnes Journal of Public Health is published by Universitas Negeri Semarang (UNNES) in cooperation with Association of Indonesian Public Health Experts (Ikatan Ahli Kesehatan Masyarakat Indonesia (IAKMI)).

It is aimed at all public health practitioners and researchers and those who manage and deliver public health services and systems. It will also be of interest to anyone involved in provision of public health programmes, the care of populations or communities and those who contribute to public health systems in any way.

HOME ([HTTPS://JOURNAL.UNNES.AC.ID/SJU/INDEX.PHP/UJPH/INDEX](https://journal.unnes.ac.id/sju/index.php/ujph/index))
/ ARCHIVES ([HTTPS://JOURNAL.UNNES.AC.ID/SJU/INDEX.PHP/UJPH/ISSUE/ARCHIVE](https://journal.unnes.ac.id/sju/index.php/ujph/issue/archive))
/ VOL 9 NO 2 (2020): UNNES JOURNAL OF PUBLIC HEALTH

DOI: <https://doi.org/10.15294/ujph.v0i0> (<https://doi.org/10.15294/ujph.v0i0>)

Published: 2020-07-31

Articles

Service Quality and Unmet Need for Family Planning in Childbearing-Age Women at Jambesari Darus Sholah Sub-District, Bondowoso Regency

(<https://journal.unnes.ac.id/sju/index.php/ujph/article/view/28815>)

Dinda Masitha Aulia, Andrei Ramani, Ni'mal Baroya
78-85

PDF (<https://journal.unnes.ac.id/sju/index.php/ujph/article/view/28815/16708>)

The Effectiveness of Recreation Program on Self-Esteem of Aging in the Elderly Club of Danmeakhmmon Subdistrict, Uttaradit Province, Thailand

(<https://journal.unnes.ac.id/sju/index.php/ujph/article/view/36286>)

Teerawat Piwkom, Sumonratee Nimnatipun
86-90

PDF (<https://journal.unnes.ac.id/sju/index.php/ujph/article/view/36286/16709>)

Anxiety of Health Workers in the Prevention and Management of Covid-19 in Sidrap Regency (<https://journal.unnes.ac.id/sju/index.php/ujph/article/view/38436>)

Fadli Fadli, Andi Sastria Ahmad, Safruddin Safruddin, Sumbara Sumbara, Rohandi Baharuddin
91-97

PDF (<https://journal.unnes.ac.id/sju/index.php/ujph/article/view/38436/16710>)

Literature Review: Cause Factor Analysis and an Effort to prevent Medication Administration Error (MAE) at Hospital (<https://journal.unnes.ac.id/sju/index.php/ujph/article/view/36470>)

Innes Rizma Brigitta, Inge Dhamanti
98-107

PDF (<https://journal.unnes.ac.id/sju/index.php/ujph/article/view/36470/16711>)

An Approach to Measure the Death Impact of Covid-19 in Jakarta using Autoregressive Integrated Moving Average (ARIMA) (<https://journal.unnes.ac.id/sju/index.php/ujph/article/view/38460>)

Ferdian Fadly, Erika Sari
108-116

PDF (<https://journal.unnes.ac.id/sju/index.php/ujph/article/view/38460/16712>)

Smoking and Comorbidities in Covid-19: A Systematic Review (<https://journal.unnes.ac.id/sju/index.php/ujph/article/view/38181>)

Andri Nugraha, Ernawati Ernawati, Tuti Anggriani Utama, Santi Rinjani
117-125

PDF (<https://journal.unnes.ac.id/sju/index.php/ujph/article/view/38181/16716>)

Decision of Implementing Uzlah and Gerbat Techniques in Islamic Boarding School as Preparedness Response for Covid-19 Pandemic (<https://journal.unnes.ac.id/sju/index.php/ujph/article/view/38107>)

Samsul Arifin, Akhmad Zaini
126-134

PDF (<https://journal.unnes.ac.id/sju/index.php/ujph/article/view/38107/16721>)

Efectivity Evaluation among Dengue Control Programs in Semarang City, Indonesia

About Journal

Unnes Journal of Public Health (UJPH) is an open access and peer-reviewed journal. It publishes original papers, reviews and short reports on all aspects of the science, philosophy, and practice of public health, especially in low-income and middle-income countries which have many problems in technology, human resources, economics, social welfare, etc impacting public health.

Unnes Journal of Public Health is Nationally Accredited (SINTA 2) based on the Decree of The Minister of Research, Technology and Higher Education of Republic Indonesia, Number 10 / E / KPT / 2019, starts from Volume 7 Issue 2 (July 2018). Unnes Journal of Public Health is published by Universitas Negeri Semarang (UNNES) in cooperation with Association of Indonesian Public Health Experts (Ikatan Ahli Kesehatan Masyarakat Indonesia (IAKMI)).

It is aimed at all public health practitioners and researchers and those who manage and deliver public health services and systems. It will also be of interest to anyone involved in provision of public health programmes, the care of populations or communities and those who contribute to public health systems in any way.

Editorial Team

Chief Editor

Fitri Indrawati (Scopus ID : 57194183122 (<https://www.scopus.com/authid/detail.uri?authorId=57194183122>))

Universitas Negeri Semarang, Indonesia

Section Editors

Dyah Mahendrasari Sukendra (Scopus ID: 57614897400 (<https://www.scopus.com/authid/detail.uri?authorId=57614897400>))

Universitas Negeri Semarang, Indonesia

Efa Nugroho (Scopus ID : 57192436111 (<https://www.scopus.com/authid/detail.uri?authorId=57192436111>))

Universitas Negeri Semarang, Indonesia

Chatila Maharani (Scopus ID: 57208663133 (<https://www.scopus.com/authid/detail.uri?authorId=57208663133>))

Universitas Negeri Semarang, Indonesia

Adi Nugroho (Scopus ID : 57189875212 (<https://www.scopus.com/authid/detail.uri?authorId=57189875212>))

Universitas Lambung Mangkurat, Indonesia

Christina Rony Nayoan (Scopus ID: 57194601944 (<https://www.scopus.com/authid/detail.uri?authorId=57194601944>))

University of Nusa Cendana, Indonesia

Editorial Advisory Board

Amar Sharaf Eldin Khair (Scopus ID: 57201584803 (<https://www.scopus.com/authid/detail.uri?authorId=57201584803>))

Omdurman Islamic University, Sudan

Bilkis Banu (Scopus ID: 55382310400 (<https://www.scopus.com/authid/detail.uri?authorId=55382310400>))

Northern University Bangladesh

Soong Yuen Quah (Scopus ID: 7003418736 (<https://www.scopus.com/authid/detail.uri?authorId=7003418736>))

Hospital Likas, Malaysia

Madhusudan Subedi (Scopus ID: 56274284400 (<https://www.scopus.com/authid/detail.uri?authorId=56274284400>))

Patan Academy of Health Sciences, Nepal

Mohammad Abdi (Scopus ID: 57202238954 (<https://www.scopus.com/authid/detail.uri?authorId=57218551725>))

Zanjan University, Iran

Nathratul Ayesah Zulkifli (Scopus ID: 57221777730 (<https://www.scopus.com/authid/detail.uri?authorId=57221777730>))

University Sains Islam Malaysia

Budi Aji (Scopus ID: 55911737200 (<https://www.scopus.com/authid/detail.uri?authorId=55911737200>))

Universitas Jenderal Soedirman, Indonesia

Arif Susanto (Scopus ID: 57090334100 (<https://www.scopus.com/authid/detail.uri?authorId=57090334100>))

Universitas Kebangsaan, Indonesia

Editorial Board

Irwan Budiono (Scopus ID : 57194195131 (<https://www.scopus.com/authid/detail.uri?authorId=57194195131>))

Universitas Negeri Semarang, Indonesia

Sri Ratna Rahayu (Scopus ID : 55624888300 (<https://www.scopus.com/authid/detail.uri?authorId=55624888300#>))

Universitas Negeri Semarang, Indonesia

Oktia Woro Kasmini Handayani (Scopus ID : 57192428885 (<https://www.scopus.com/authid/detail.uri?authorId=57192428885#>))

Universitas Negeri Semarang, Indonesia

Lintang Dian Saraswati (Scopus ID : 57193449366 (<https://www.scopus.com/authid/detail.uri?authorId=57193449366#>))

Universitas Diponegoro, Indonesia

Sri Sumartiningsih (Scopus ID : 41662392800 (<https://www.scopus.com/authid/detail.uri?authorId=41662392800#>))
Universitas Negeri Semarang, Indonesia

Adang Bachtiar (Scopus ID : 56683183900 (<https://www.scopus.com/authid/detail.uri?authorId=56683183900#>))
Universitas Indonesia, Indonesia
Ikatan Ahli Kesehatan Masyarakat Indonesia (IAKMI)

Mahalul Azam (Scopus ID : 57194196255 (<https://www.scopus.com/authid/detail.uri?authorId=57194196255#>))
Universitas Negeri Semarang, Indonesia

Sartiah Yusran (Google Scholar ID (<https://scholar.google.com/citations?user=9iUPPvoAAA&hl=en>))
Universitas Halu Oleo, Indonesia

Agustin Kusumayati (Scopus ID : 6507269847 (<https://www.scopus.com/authid/detail.uri?authorId=6507269847&eid=2-s2.0-0032438849>))
Universitas Indonesia, Indonesia
Ikatan Ahli Kesehatan Masyarakat Indonesia (IAKMI)

Bayu Anggileo Pramesona (Scopus ID : 57201555135
(<https://www.scopus.com/authid/detail.uri?authorId=57201555135>))
RSD Mayjend. HM. Ryacudu Kotabumi, Lampung Utara, Indonesia

Marniati (Scopus ID : 57202099589 (<https://www.scopus.com/authid/detail.uri?authorId=57202099589>))
Universitas Ubudiyah Indonesia, Indonesia

Sri Nurlaela (Scopus ID : 57216416092 (<https://www.scopus.com/authid/detail.uri?authorId=57216416092>))
Universitas Jenderal Soedirman, Indonesia

Yanelza Supranelfy (Scopus ID : 57221327196 (<https://www.scopus.com/authid/detail.uri?authorId=57221327196>))
Institute of Health Research and Development (Balai Litbangkes) Baturaja, Health Ministry, Indonesia

Siti Alfiah (Scopus ID : 57203408996 (<https://www.scopus.com/authid/detail.uri?authorId=57203408996>))
Health Research and Development Agency (Badan Litbangkes), Health Ministry, Indonesia

Lulus Susanti (Scopus ID : 57205363850 (<https://www.scopus.com/authid/detail.uri?authorId=57205363850>))
Institute for Vector and Reservoir Control, Research, and Development (B2P2VRP), Health Ministry, Indonesia

Imelda G. Purba (Scopus ID : 57205022590 (<https://www.scopus.com/authid/detail.uri?authorId=57205022590>))
Universitas Sriwijaya, Indonesia

Arum Kartikadewi (Scopus ID : 57208864205 (<https://www.scopus.com/authid/detail?authorId=57208864205>))

Universitas Muhammadiyah Semarang, Indonesia

Elanda Fikri (Scopus ID : 57189573562 (<https://www.scopus.com/authid/detail?authorId=57189573562>))

Bandung Polytechnic of Health, Health Ministry, Indonesia

Vita Nur Latif (Scopus ID : 57208281949 (<https://www.scopus.com/authid/detail?authorId=57208281949>))

Universitas Pekalongan, Indonesia

Evi Widowati (Scopus ID : 57210980764 (<https://www.scopus.com/authid/detail?authorId=57210980764>))

Universitas Negeri Semarang, Indonesia

Novie Susanto (Scopus ID: 56023348100 (<https://www.scopus.com/authid/detail?authorId=56023348100>))

Universitas Diponegoro, Indonesia

ISSN

p-ISSN 2252-6781 (<http://issn.pdii.lipi.go.id/issn.cgi?daftar&1333557581&1&&>)

e-ISSN 2548-7604 (<http://issn.pdii.lipi.go.id/issn.cgi?daftar&1471319300&1&&>)

Sertifikat



Literature Review: Cause Factor Analysis and an Effort to prevent Medication Administration Error (MAE) at Hospital

Innes Rizma Brigitta^{1✉}, Inge Dhamanti^{1,2,3}

¹Department of Health Policy and Administration, Faculty of Public Health, Universitas Airlangga

²School of Public Health, La Trobe University, Victoria, Australia

³Center for Patient Safety Research, Universitas Airlangga, Surabaya, Indonesia

Article Info

Article History:

Submitted January 2020

Accepted April 2020

Published July 2020

Keywords:

Patient safety incidents; medication administration errors; determinants; preventive actions

DOI

<https://doi.org/10.15294/ujph.v9i2.36470>

Abstract

The high number of medical errors, especially in medication administration errors (MAE) in the last few decades that have occurred in hospitals in developed and developing countries makes patient safety an important issue. This requires the hospital to take steps that prioritize patient safety by focusing on preventive measures, so as to reduce the risk of an MAE. The writing of this article of literature review aims to explain the determinants of MAE in hospitals and their prevention efforts through preventive measures, so that patient safety standards in hospitals (zero defects) can be achieved. The design of the article search in this literature study was carried out through Google scholar, JAKI, and PSNet with the keywords of patient safety incident, medication administration error, contributing factor of MAEs, and determinant factor of MAEs. Based on 13 articles that have been obtained in accordance with the criteria, there are 38 determinant factors in MAE which are grouped into three categories, namely ineffective communication factors, work environment factors, and human factors. Various preventive efforts that can be done to prevent MAEs include: implementing crew resource management, clarifying the chain of command, using the communication form of SBAR, designing an ergonomic workplace, implementing Patient Advocacy Reporting System (PARS), providing training and education for health workers, and setting work schedules that do not exceed workload. Implementation of effective MAE preventive measures can reduce the number of MAEs in the hospital directly.

INTRODUCTION

The hospital is a health service institution for the community by providing safe, quality, anti-discrimination and effective health services by prioritizing the interests of patients. Implementation of health services must be carried out in accordance with hospital service standards so that the highest degree of public health can be achieved (Law of the Republic of Indonesia, 2009). The implementation of health services in hospitals has a variety of very complex characteristics, ranging from the availability of various types of health workers to the rapid development of science and technology. This complexity, if not mana-

ged and coordinated properly, will lead to incidents that threaten patient safety (Department of Health of the Republic of Indonesia, 2008).

Patient safety is a discipline in the health service sector that applies the method of safety science towards a goal which is to achieve a reliable health service delivery system. Patient safety is also an attribute of the health care system that minimizes events and impacts, and maximizes recovery from side effects (Emanuel et al., 2008). Patient safety then becomes an important issue because of the large number of medical error cases that occur in various countries. Patient safety incident reports in northern England show that

✉ Correspondence Address:

Faculty of Public Health, Campus C, Mulyorejo, District Mulyorejo, City of Surabaya, East Java 60115
E-mail: innes.rizma.brigitta-2016@fkm.unair.ac.id

1 in 10 patients experience a patient safety incident with a type of medication error (Hara et al., 2018).

Medication errors are the incidents that most often endanger the patient, although not all of them can cause serious failure in patients (Hines et al., 2018). The number of reports of medication errors has continued to increase in the last few decades, so it is a concern in terms of patient safety in hospitals (Saghafi & Zargarzadeh, 2014). In the United States, medication errors account for 28% of all patient safety incidents and can have serious patient complications (Patel & Balkrishnan, 2010). According to a review study in 2013, the prevalence of medication error in Middle Eastern countries including Iran is 7 to 90% (Leber, et al., 2018).

Medication errors can cause losses in terms of treatment, so patients must spend more time to stay in the hospital and increase the burden of care, and costs. The existence of medication errors can threaten the quality of the hospital because the patient or family has the right to report incidents that occur through legal action (Saghafi & Zargarzadeh, 2014) especially in medical centers has become a growing concern for patient safety in recent decades. Patient safety and in particular, medication safety is a major concern and challenge for health care professionals around the world. Our prospective study was designed to detect prescribing, transcribing, dispensing, and administering medication errors in two major university hospitals. Materials and Methods: After choosing 20 similar hospital wards in two large teaching hospitals in the city of Isfahan, Iran, the sequence was randomly selected. Diagrams for drug distribution were drawn by the help of pharmacy directors. Direct observation technique was chosen as the method for detecting the errors. A total of 50 doses were studied in each ward to detect prescribing, transcribing and administering errors in each ward. The dispensing error was studied on 1000 doses dispensed in each hospital pharmacy. Results: A total of 8162 number of doses of medications were studied during the four stages, of which 8000 were complete data to be analyzed. 73% of prescribing orders were incomplete and did not have all six parameters (name, dosage form, dose and measuring unit, administration route, and intervals of administration. Medication errors contribute to morbidity, mortality and an increase in health care costs (Kumar, K.S.A. & Venkateswarlu, K., 2011).

The national coordinating council for medication errors and prevention states that medication errors are preventable events, which lead

to drug abuse or patient error in the treatment process under the control of health workers (Saghafi and Zargarzadeh, 2014) especially in medical centers has become a growing concern for patient safety in recent decades. Patient safety and in particular, medication safety is a major concern and challenge for health care professionals around the world. Our prospective study was designed to detect prescribing, transcribing, dispensing, and administering medication errors in two major university hospitals. Materials and Methods: After choosing 20 similar hospital wards in two large teaching hospitals in the city of Isfahan, Iran, the sequence was randomly selected. Diagrams for drug distribution were drawn by the help of pharmacy directors. Direct observation technique was chosen as the method for detecting the errors. A total of 50 doses were studied in each ward to detect prescribing, transcribing and administering errors in each ward. The dispensing error was studied on 1000 doses dispensed in each hospital pharmacy. Results: A total of 8162 number of doses of medications were studied during the four stages, of which 8000 were complete data to be analyzed. 73% of prescribing orders were incomplete and did not have all six parameters (name, dosage form, dose and measuring unit, administration route, and intervals of administration. The high prevalence of medication error has a detrimental impact on health services. This has the potential to threaten patient safety, making it a challenge for hospitals to prevent the incident from recurring through preventive action (Saghafi & Zargarzadeh, 2014; Mekonnen et al., 2018; Leber et al., 2018) especially in medical centers has become a growing concern for patient safety in recent decades. Patient safety and in particular, medication safety is a major concern and challenge for health care professionals around the world. Our prospective study was designed to detect prescribing, transcribing, dispensing, and administering medication errors in two major university hospitals. Materials and Methods: After choosing 20 similar hospital wards in two large teaching hospitals in the city of Isfahan, Iran, the sequence was randomly selected. Diagrams for drug distribution were drawn by the help of pharmacy directors. Direct observation technique was chosen as the method for detecting the errors. A total of 50 doses were studied in each ward to detect prescribing, transcribing and administering errors in each ward. The dispensing error was studied on 1000 doses dispensed in each hospital pharmacy. Results: A total of 8162 number of doses of medications were studied during the four stages, of which

8000 were complete data to be analyzed. 73% of prescribing orders were incomplete and did not have all six parameters (name, dosage form, dose and measuring unit, administration route, and intervals of administration).

Medication errors can occur at any stage, namely prescribing, documentation, expenditure, preparation, or giving (Kumar K.S.A. & Venkateswarlu, K., 2011). In 2007, the National Patient Safety Agency statistics (NPSA) showed that 59.3% of medication errors occurred during the administration phase (Kumar, K.S.A. & Venkateswarlu, K., 2011). The administrative stage of the treatment process is the most common point where errors occur, with an incidence of 50% of all medication error events in the United Kingdom (Hines, Kynoch & Khalil, 2018). 1 in 3 patients experience medication administration error (Saghafi & Zargarzadeh, 2014).

A review article published in 2013 in Iran reported that administration error was the most common type of medication error with a prevalence range of 14.3-70% (Saghafi & Zargarzadeh, 2014). Levinson, 2011 in his research conducted in Colombia stated that the occurrence of patient safety incidents in hospitals related to medication administration (medical administration error) occurred by 31%.

The higher MAE incidence volume was found in a study from Girma & Feleke (2010), which managed to identify the MAE incidence as much as 89.9% of the 218 observations that had been made. In a study conducted by Feleke, et al. (2015) at one hospital in Ethiopia, showed the results of 360 treatment administration interventions, 98.1% of them experienced MAE events. The results of this study are also in line with studies conducted by Westbrook et al., (2011) in Sydney, Australia.

The reported number of patient safety incidents in Indonesia in 2007 was 145 incidents. DKI Jakarta was ranked first with a number of incidents of 37.9% (KKPRS, 2008). Indonesia has a tendency towards high incidence of patient safety incidents, given the low incidence of incident reporting that is carried out in accordance with procedures. Law of the Republic of Indonesia No. 44 of 2009 concerning Hospitals explains that hospitals have an obligation to implement patient safety standards in order to reduce the number of unexpected events. The magnitude of patient safety incident cases, including MAE, requires hospitals to take steps to prioritize patient safety by focusing on preventive measures.

Based on the number of MAE problems that occur in hospitals in developed and develop-

ing countries that have been mentioned, the objectives of this article are formulated. The writing of this literature review article aims to explain the determinants that influence the medication administration error (MAE) in hospitals and their prevention efforts through preventive measures, so as to prevent the occurrence of patient safety incidents and the achievement of patient safety standards at the hospital (zero defect).

METHOD

The collection of journal articles in this literature study was carried out through searches from Google scholar, JAKI (Indonesian Health Administration Journal), and PSNet (Patient Safety Network) with keywords of patient safety incidents, medication administration errors, contributing factors of MAEs, and determinant factors of MAEs. The inclusion criteria in this literature study are 1) the article has a scope of research on medication administration error; 2) articles were published in 2014-2020; 3) the article is available in full text. Articles discussing medication administration error theoretically are the exclusion criteria in this literature study. The incident factors found were then grouped and analyzed narratively. The next step is to provide recommendations as a follow-up effort to prevent the occurrence of patient safety incidents in the treatment administration process.

RESULTS AND DISCUSSION

Based on data searches using keywords and criteria in the database above, obtained 13 relevant journal articles. Found 38 factors that contributed to the occurrence of medication administration error. All these factors are then described in Table 1.

Of all the factors that have been identified based on research findings, 38 existing factors can be classified into three categories seen in Figure 1. These categories include ineffective communication factors, work environment factors, and human factors. The distribution of these three factors can be seen in Figure 1.

Based on Figure 1, it can be seen that the most dominant factor influencing the incidence of incidents in treatment administration comes from the individual factors of health workers. The distribution and amount of each factor that affects medication administration error can be seen in Table 2.

Based on Table 2, it appears that the most dominant factor in the incidence of MAE is human factor, which amounts to 21 of the total 38

Table 1. Summary of selected studies

| References | Design | Sample | Data collection method | Factors contributing to MAE |
|-----------------------------|---|---|--|--|
| Mansah et al., 2014 | A retrospective (case control) audit | 1 large tertiary metropolitan hospital in New South Wales, Australia | The information from Incident Information Management System (IIMS) | Violation procedures |
| Parry et al., 2015 | A thematic analysis and narrative synthesis | Six electronic databases | Cochrane, MedLine, CINAHL, BNI, Embase, PsycINFO | Workload Work setting Nurse characteristics Experience of work |
| Tehewy et al., 2016 | Descriptive direct-observational study | Aim Shams University hospital | A standardized observational checklist, and a medical record audit form | Workload Illiteracy and elderly patient |
| Dietrich & Norman, 2017 | A cross-sectional study | Six agency CNOs (Chief Nurse Officers) at the hospital | CALNOC direct observation methodology | Perceived skill of nurses |
| Thomas et al., 2017 | Hierarchical | 79 RN from 9 hospital in Newyork | The ISRN Coordinating Center | Workload Human factor |
| Budihardjo, 2017 | Descriptive observational | 56 nurses in 7 rooms at the Haji General Hospital in Surabaya | Questionnaire, interview | Skill of nurses Nurses' knowledge Communication |
| Handayani, 2017 | Qualitative, explanative | Employees of the Antapura General Hospital of Palu City | Observation, interview, document review | Health worker Patient factors Work environment |
| Baraki et al., 2018 | Prospective observational based cross-sectional study | 1251 medication administration in Jimma University Specialized Hospital | Pre-tested structured questionnaire and blind observation checklist | Patient factor Nurses' knowledge Work environment Experience of work |
| Keers et al., 2015 | Qualitative interviewing | 20 nurses in two NHS in the North West of England | Face-to-face semistructured interviews | Violation procedures Team's support Workload Shift changes |
| Felekeet et al., 2015 | A prospective, observation-based, cross-sectional study | 82 nurses in the Felege Hiwot Referral Hospital inpatient department | Pre-tested structured questionnaire | Nurse's age Experience of work Patient factor Interruption Shifts taken |
| Elasrag & Abu-Snieneh, 2020 | Descriptive exploratory cross-sectional design | 146 nurses from two regional hospitals in Egypt | Medication Administration Error Reporting Scale | Nursing Staffing Communication |
| Alemu et al., 2017 | A quantitative and observational approach, cross-sectional design | 141 nurses working in two hospitals in Southern X | A structured, pretested, self-administered questionnaire and a semi-structured, pretested, observational checklist | Lack of sufficient training Inadequate staffing Distraction Like-looking like drugs |
| Ayorinde & Alabi, 2019 | A descriptive cross-sectional study | 300 nurses in University College Hospital, West Africa | A structured pre-tested questionnaire | Nurse Staffing Confusion about drug Illegible writing |




Figure 1. Contributing factors to MAE

factors identified.

Factors that contribute to MAE
Ineffective communication

Ineffective communication is one of the factors associated with the occurrence of medication administration errors in hospitals. Communication failures in medication administration are caused by several things, namely the delivery of ambiguous verbal instructions to nurses, the weakness of giving direct instructions to nurses, and nurses' lack of understanding in interpreting communication that is being established among health workers (Delgado et al., 2015).

The flow of communication in health services is divided into three, namely communication of health workers to patients, communication among health workers, and communication of patients to health workers (Mansah, et al., 2014; Hara et al., 2018). Most of the communication error is caused by health workers, namely the lack of availability of health workers to respond to patient questions in administering treatment. Therefore, this will lead to miscommunication between patients and health workers so that the risk of developing MAEs will increase (Hara et al., 2018). Health workers who do not establish good communication among staff, especially in terms of treatment schedules will trigger MAE (Elasrag & Abu-Snieneh, 2020). Communication failures among nurses and nurses with doctors contribute

to 64% of medication error events (Mansah et al., 2014).

A health professional needs high concentration during the treatment process, from preparation to administration of the drug. Nurses who face interruptions, including communication that takes place during drug administration, are twice as likely to develop medical administration errors compared to nurses who work without disturbances (Feleke et al., 2015). One source of interruption can be done by medical personnel, namely doctors. Examples of such interruptions include discussing patient care plans while other medical personnel are doing other works (Johnson et al., 2017) interruptions during medication rounds have been associated with medication errors. Method: A non-participant observational study was undertaken of nurses conducting medication rounds. Results: Fifty-six medication events (including 101 interruptions).

A health service must be able to develop its strategy in order to improve the effectiveness of communication between nurses and doctors. The better communication is established among health workers, the smaller the number of medication errors in a service unit occurs (Budihardjo, 2017).

Work environment

One factor related to the occurrence of medication errors is the work environment. Work

Table 2. Contributing factors to MAE

| No | Contributing Factor to MAE | Total |
|-------|----------------------------|-------|
| 1 | Ineffective communication | 3 |
| 2 | Work environment | 14 |
| 3 | Human factor | 21 |
| Total | | 38 |

environment is a composition of three main sub-environments namely technical environment, human environment, and organizational environment (Oludeyi, 2015).

Technical environment consists of equipment, infrastructure, and other physical environmental factors in the workplace (Oludeyi, 2015). Health facilities, availability of drug preparation rooms, number of drug prescriptions per patient and availability of drug administration guidelines were found to be significant independent factors related to MAE. In the administration of treatment that is prepared without the availability of drug preparation space 13.5 times more risk of developing MAE (Baraki et al., 2018). Likewise, during the treatment administration process, namely the provision of drugs which are prepared in places where there are no drug administration guidelines 4 times more at risk of developing MAE (Baraki et al., 2018). If viewed from the physical work environment, the ones that contribute to the occurrence of medication error are the noise level, lighting level, and work-space layout (Hara et al., 2018).

Human environment is related to interpersonal relations of staff with staff or staff with their superiors (Oludeyi, 2015). Errors at the administrative stage are more often caused by busy workers in the related service units (Handayani, 2017). Medication administration error occurs when nurses are not supported by the work team in their work unit, and / or when the workload is increasing due to shifts in work, or because of an emergency (Keers et al., 2015) where they were asked to discuss perceived causes of intravenous MAEs that they had been directly involved with. Transcribed interviews were analysed using the Framework approach and emerging themes were categorised according to Reason's model of accident causation. Results: In total, 21 intravenous MAEs were discussed containing 23 individual active failures which included slips and lapses (n=11).

Interpersonal relationships among doctors and nurses with other health professionals can be at risk for MAE. This includes inefficiencies in the division of roles during examinations in patients, resulting in overlapping work. Examples of errors that occur are health workers not doing double checking during the treatment administration process (Keers et al., 2015). So adaptive coordination is needed in establishing interpersonal relationships among health workers, including dynamic allocation of tasks and responsibilities (Manser, 2009).

Organizational environment is related to

the structure of hospital organization, assignment of duties, leadership of department heads, communication in teamwork, and patient safety programs in hospitals (Oludeyi, 2015; Parry, Barriball & While, 2015). Delgado, et al (2015) states that excessive workload is a major factor in workplace conditions that triggers medication errors, followed by work time pressure and the addition of new members in the work unit.

An increase in workload results in work stress experienced by nurses so that it affects the incidence of medication administration errors (Parry, et al., 2015). The number of work shift taking is the implication of the nurse's workload. Workload consisting of taking night shift work and work on holidays is a significant independent variable on the occurrence of MAE. The workload can be represented through the high number of nurse services to patients and the number of shifts taken per month. The severity of the nurse's workload will have an impact on patient safety (Tehewy et al., 2016).

Nurses who work during the night shift can experience circadian disorders that cause fatigue, disturbed hours of sleep, so that the impact on performance is not optimal. The process of treatment to patients at night is twice as likely to experience to MAE compared to treatment during the day (Feleke, Mulatu and Yesmaw, 2015). The complexity of the nurse's work, interruption and interruption of work, and high workload demands are the cause of the MAE. The higher the workload that must be done, especially in the treatment administration process, the risk of a MAE will be higher too (Thomas et al., 2017).

Elasrag & Abu-Snieneh (2020), in his research stated that the inadequate number of nursing staff affected the MAE incidence. The limited number of staff and the presence of distraction factors in the organizational environment are proven to affect the MAE (Alemu et al., 2017). The number of staff who do not meet the needs will have an impact on work fatigue due to high workload so that MAE can occur (Ayorinde & Alabi, 2019).

Human factor

Human factor is a scientific discipline that deals with understanding interactions between humans and other elements of a system, and professions that apply theories, principles, data, and methods to optimize overall system performance (Hayden et al., 2018).

Medication errors (ME) caused by human factors include various aspects of triggers therein. Violations of Operational Standards The appli-

cable procedures, writing errors, lack of work experience, and work stress are examples of several aspects involved in the human error (health care practitioners) factor to the incidence of medication errors (Mansah, et al., 2014). Human error contributes to medication errors by 19.4% which consists of the level of knowledge of health workers, work stress, and lack of work motivation of health workers in doing their work optimally (Delgado et al., 2015).

Human error factor by health workers is the main factor causing medication administration error. The majority of incidents (88%) are caused by health workers who do not comply with policies and procedures that apply to the treatment administration process (Mansah, et al., 2014). A health worker can violate SOPs when administering drugs, when working with work teams, to violations using irrelevant medical or non-medical equipment (Keers et al., 2015).

Parry et al. (2015) in his study also stated that individual factors of health workers included factors that contributed to MAE. This factor is divided into two, namely the characteristics of nurses and experience of working in the health field. Health workers who work in service units in a period of less than 12 months have a 63% potential to experience MAE, compared to health workers who have experienced more than 24 months working in the health sector (Baraki et al., 2018).

In addition, the skills of nurses have a significant relationship to MAE (Budihardjo, 2017). The higher the level of nurse skills, the higher the level of accuracy in the treatment administration process to patients (Dietrich & Norman, 2017).

Various factors that cause MAE in the human error category have several elements in it. Age of nurses also has a significant relationship with the risk of MAE. The more a nurse ages, the higher the risk of MAE. The lower the knowledge of health workers and the level of nurse compliance with policies along with existing Standard Operating Procedures, the risk of MAE will be higher (Thomas et al., 2017). The better the level of knowledge of nurses, the smaller the number of medication error events in a service unit (Budihardjo, 2017).

Lack of adequate training of health workers contributes to the occurrence of MAEs (Alemu et al., 2017). Nurse accuracy also influences MAE, such as errors in reading doctor's writing about prescribed drugs, and errors in administering drugs that look similar (look alike sound alike) (Ayorinde & Alabi, 2019)

Types of violations that occur can be classified based on the source, namely violations by

patients and violations by health workers. Research conducted by (Handayani, 2017) explains the human error factor in more detail from the two parties. Judging from the factors of health workers, the cause of administration error is the work culture that is applied in the related unit. Judging from the patient's factors, the cause of administration error is the family of the patient who is not cooperative and the lack of understanding of the patient's family about the treatment procedure.

Preventive Efforts to MAE

1. Implementing Crew Resource Management (CRM) is a technique used by a team to reduce performance errors and create a safety culture. CRM can improve communication between team members in terms of delivering briefings, a shared mental model, and encourage each team member to be able to express their opinions (McKoin et al., 2010). The use of communication guidelines is recommended in a team regarding work processes, standards for the transfer of patient information, team training in communication, and the use of standardized tools (Link, 2018). The MedTeams project has implemented the CRM principle in the delivery of health services and obtained results that decreased clinical error by 30% (Lindquist, 2009).
2. Clarifying the chain of command in health services for the patient care process and appropriate feedback to ensure that patients receive quality care. Command chain policies also play a role in reducing the risk of incidents and increasing patient safety (McKoin et al., 2010). Studies at the RSUI of Malang show that a well-structured organization can affect the flow of the chain of command that runs well as well, so that health workers can prevent medication errors (Hastuti et al., 2014). Using simple and structured communication forms to efficiently convey information to patients thereby limiting the possibility of miscommunication. The skills of health workers in communicating effectively can be obtained through the SBAR method (McKoin et al., 2010).
3. Redesign an ergonomic workplace so that it can increase work productivity. This is done through workspace settings, lighting, temperature, and noise levels that do not exceed excessive values. Good work design can produce good service processes and improve patient safety (Carayon, Alvarado and Schoofs Hundt, 2007).

4. Implementing a surveillance system, namely Patient Advocacy Reporting Systems (PARS), to recognize the behavior patterns of health professionals who are unprofessional and endanger patient safety. This is done based on patient or family complaints and the perception of coworkers. In its application, it is necessary to ensure that staffs have the opportunity to fully participate in risk identification and mitigation activities that will prevent patient safety incidents (Mckoin et al., 2010). PARS can improve supervision of the entire system of health personnel behavior that deviates from safety culture and seeks to improve patient safety culture (Battles et al., 2017).
5. Teamwork training through application *Team Strategies and Tools to Enhance Performance and Patient Safety™ (TeamSTEPPS™)* which consists of leadership, team member support, condition monitoring, and communication (Mckoin et al., 2010). TeamSTEPPS enables a team to adapt to changing situations, share understanding of patient care, develop positive teamwork attitudes, provide efficient and reliable patient care, and most importantly, achieve safer treatment outcomes (Hunt, 2010).
6. Encouraging the involvement of communication leaders in a service unit in the direction of the delivery of safety values, and provide feedback to members to improve patient safety (Mattson & Hellgren, 2015)
7. Providing education and training for health workers and enforcing policies or Standard Operating Procedures at work (Scanlon & Karsh, 2010). Simulation-based training program with scenario design and debriefing in accordance with the health service situation (Sarfati et al., 2019). Providing training of health personnel on patient safety that is relevant to the state of the hospital. Providing skills-based training and coaching is also needed for all unit leaders to provide feedback on the chain of command, unprofessional staff behavior, and problem resolution.

Work policies and procedures need to be developed and implemented in accordance with the principle of “zero tolerance”, which means that staff who behave unprofessionally will receive disciplinary action (Timmons et al., 2014).
8. Striving for work schedule retrieval in accordance with work standards and capabilities, accompanied by giving instructions on patient handover and work procedures (Weinger & Gaba, 2014).
9. Creating a hospital environment that supports the process of care for patients to improve the level of service security by allocating special resources to deal with problems in the hospital environment (Sahlström, 2018)

CONCLUSION

Based on the literature review conducted on 13 journal articles, the results obtained are that there are three groups of factors that influence Medication Administration Error (MAE). These three factors include ineffective communication, work environment, and human factors.

There are various preventive efforts that can be done to prevent MAE in hospitals, namely implementing crew resource management, clarifying the chain of command, using SBAR communication forms, designing an ergonomic workplace, implementing a monitoring system namely Patient Advocacy Reporting System (PARS), providing training and education for health workers, along with setting work schedules that do not exceed workload.

A hospital can consider various preventive measures based on the level of importance and available resources to deal with patient safety incidents, especially in the case of medication administration errors. The implementation of MAE preventive programs will directly affect the decline in MAE cases that occur in hospitals.

REFERENCES

- Alemu, W., Belachew, T. & Yimam, I. 2017. Medication administration errors and contributing factors: A cross sectional study in two public hospitals in Southern Ethiopia. *International Journal of Africa Nursing Sciences*, 7(September): 68–74. doi: 10.1016/j.ijans.2017.09.001.
- Ayorinde, M. O. and Alabi, P. I. 2019. Perception and contributing factors to medication administration errors among nurses in Nigeria. *International Journal of Africa Nursing Sciences*, 11(May): 100153. doi: 10.1016/j.ijans.2019.100153.
- Baraki, Z., et al. 2018. Medication administration error and contributing factors among pediatric inpatient in public hospitals of Tigray, northern Ethiopia. *BMC Pediatrics*, 18(1): 1–8. doi: 10.1186/s12887-018-1294-5.
- Battles, J. et al. 2017. *Advances in Patient Safety and Medical Liability*. AHRQ Publication No. 17-0017. Rocville, MD: Agency for Healthcare Research and Quality
- Budihardjo, V. S. 2017. Faktor Perawat Terhadap Kejadian Medication Administration Error di Instalasi Rawat Inap. *Jurnal Administrasi Kesehatan Indonesia*, 5(1): 52–61. doi: 10.20473/jaki.v5i1.2017.52-61
- Carayon, P., Alvarado, C. J. & Hundt, A.S. 2007.

- Work design and patient safety. *Theoretical Issues in Ergonomics Science*, 8(5): 395–428. doi: 10.1080/14639220701193157.
- Delgado, M.C.M., et al. 2015. Analysis of contributing factors associated to related patients safety incidents in Intensive Care Medicine. *Medicina Intensiva Journal (English Edition)*, 39(5): 263–271. doi: 10.1016/j.medine.2015.05.001.
- Department of Health of the Republic of Indonesia. 2008. *Panduan Nasional Keselamatan Pasien Rumah Sakit*. Departemen Kesehatan Republik Indonesia. Jakarta: Department of Health of the Republic of Indonesia.
- Dietrich, M. & Norman, L. 2017. Nurses' Perceived Skills and Attitudes About Updated Safety Concepts: Impact on Medication Administration Errors and Practices. *Journal of Nursing Care Quality*, 32(3): 226–233. doi: 10.1097/NCQ.0000000000000226.
- Elasrag, G.A.E. & Abu-Snieneh, H.M. 2020. Nurses' perception of factors contributing to medication administration errors. *International Journal of Research in Pharmaceutical Sciences*, 11(1): 44–56. doi: 10.26452/ijrps.v11i1.1781.
- Emanuel, L. et al. 2008. What Exactly Is Patient Safety?. *Advances in Patient Safety: New Directions and Alternative Approaches*, 1(Assessment): 1–18. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21249863>.
- Feleke, S.A., Mulatu, M.A. & Yesmaw, Y.S. 2015. Medication administration error: Magnitude and associated factors among nurses in Ethiopia. *BMC Nursing*, 14(1): 1–8. doi: 10.1186/s12912-015-0099-1.
- Girma, B. & Feleke, Y. 2010. Medication administration errors involving paediatric in-patients in a hospital in Ethiopia. *Tropical Journal of Pharmaceutical Research*, 9(4): 401–407. doi: 10.4314/tjpr.v9i4.58942.
- Handayani, T.W. 2017 Faktor Penyebab Medication Error di RSUD Anutapura. *Jurnal Pengembangan Sumber Daya Insani*, 02(2): 224–229. doi: 10.26618/perspektif.v2i2.1285
- Hara, J.K.O., et al. 2018. What can patients tell us about the quality and safety of hospital care? Findings from a UK multicentre survey study. *BMJ Quality and Safety*, 27: 673–682. doi: 10.1136/bmjqs-2017-006974.
- Hastuti, A.P., Nursalam & Triharini, M. 2014. Preventing Medication Error Based on Knowledge Management Against Adverse Event. *Jurnal Ners*, 12(1): 133–141. doi: 10.20473/jn.v12i1.2297
- Hayden, E.M., Wong, A.H., Ackerman, J., Sande, M.K., Lei, C., Kobayashi, L., et al. 2018. Human factors and simulation in emergency medicine. *Academic Emergency Medicine*, 25(2): 221–229. <https://doi.org/10.1111/acem.13315>.
- Hines, S., Kynoch, K. & Khalil, H. 2018. Effectiveness of interventions to prevent medication errors. *JBI Database of Systematic Reviews and Implementation Reports*, 16(2): 291–296. doi: 10.11124/jbisrir-2017-003481.
- Hunt, C. M. 2010. Patient Safety is Enhanced by Teamwork. *Pennsylvania Patient Safety Advisory*, 7(2): 14–17. available at: <http://www.patient-safetyauthority.org>
- Johnson, M. et al. 2017. The impact of interruptions on medication errors in hospitals: an observational study of nurses. *Journal of Nursing Management*, 25(7): 498–507. doi: 10.1111/jonm.12486.
- Keers, R.N. et al. 2015. Understanding the causes of intravenous medication administration errors in hospitals: A qualitative critical incident study. *BMJ Open*, 5(3): 1–10. doi: 10.1136/bmjopen-2014-005948.
- Kumar K.S.A. & Venkateswarlu, K.R.A. 2011. A Study of medication administration errors in a tertiary care hospital. *Indian Journal of Pharmacy Practice*, 4(2): 37–42. doi:10.5530/ijopp
- Leber, M., Carillo, C.J.D., Cassano, A.T., et al. 2018. ASHP guidelines on the prevention of medication errors in hospitals. *Am J Health Syst Pharm*, 75: 1493-1517. doi: 10.2146/ajhp170811
- Levinson, D.R. 2011. *Adverse Events in Hospitals: National Incidence Among Medicare Beneficiaries*. Washington DC: Department of Health & Human Services USA.
- Lindquist, D.G. 2009. Improving Patient Safety Using Crew Resource Management Principles Taught Via Medical Simulation. *Prescriptions for excellence in health care newsletter supplement*, 1(7): 5. Available at: <http://jdc.jefferson.edu/pehc/vol1/iss7/5>.
- Link, Terri. 2018. Guideline Implementation: Team Communication. *AORN Journal*, 102(2): 165–174. doi: 10.1002/aorn.12300.
- Mansah, M., Hons, B.N., et al. 2014. Older Folks in Hospitals: The Contributing Factors and Recommendations for Incident Prevention. *Journal of Patient Safety*, 10(3): 146–153. doi: 10.1097/PTS.0b013e31829954fd
- Manser, T. 2009. Teamwork and patient safety in dynamic domains of healthcare: A review of the literature. *Acta Anaesthesiologica Scandinavica*, 53(2): 143–151. doi: 10.1111/j.1399-6576.2008.01717.x.
- Mattson, M. & Hellgren, J. 2015. Leader communication approaches and patient safety: An integrated model. *Journal of Safety Research*, 53: 53–62. doi: 10.1016/j.jsr.2015.03.008.
- Mckoin, S. et al. 2010. Chain of Command: When Disruptive Behavior Affects Communication and Teamwork. *Pennsylvania Patient Safety Advisory*, 7(2): 4–14. available at: <http://www.patientsafetyauthority.org>
- McKoin, S., et al. 2010. Building a culture of operating room safety using crew resource management. *Pennsylvania Patient Safety Advisory*, 7(2): 1–3. Available at: <http://www.patientsafetyauthority.org>.
- Mekonnen, A.B., et al. 2018. Adverse Drug Events and Medication Errors in African Hospitals: A Systematic Review. *Drugs - Real World Outcomes*,

- 5(1): 1–24. doi: 10.1007/s40801-017-0125-6.
- Oludeyi, O. S. 2015. A review of literature on work environment and work commitment: Implication for future research in citadels of learning. *Journal of Human Resource Management*, 18: 32–46.
- Parry, A.M., Barriball, K.L. & While, A.E. 2015. International Journal of Nursing Studies Factors contributing to Registered Nurse medication administration error: A narrative review. *International Journal of Nursing Studies*, 52(1): 403–420. doi: 10.1016/j.ijnurstu.2014.07.003.
- Patel, I. & Balkrishnan, R. 2010. 'Medication error management around the globe: An overview. *Indian Journal of Pharmaceutical Sciences*, 72(5): 539–545. doi: 10.4103/0250-474X.78518.
- Saghafi, F. & Zargarzadeh, A.H. 2014. Medication error detection in two major teaching hospitals: What are the types of errors? *Journal of Research in Medical Sciences*, 19(7): 617–623.
- Sahlström, M. 2018. Patient participation in patient safety — An exploration of promoting factors. *Journal Nursing Management*, 27: 1–9. doi: 10.1111/jonm.12651.
- Sarfati, L., et al. 2019. Human-simulation-based learning to prevent medication error: A systematic review. *Journal of Evaluation in Clinical Practice*, 25(1): 11–20. doi: 10.1111/jep.12883.
- Scanlon, M.C. & Karsh, B. 2010. Value of human factors to medication and patient safety in the intensive care unit. *Critical Care Med*, 38(6): 90–96. doi: 10.1097/CCM.0b013e3181dd8de2.
- Tehewy, M. et al. 2016. Medication Administration Errors in a University Hospital. *Journal of Patient Safety*, 12(1): 34–39.
- Thomas, L. et al. 2017. Impact of Interruptions, Distractions, and Cognitive Load on Procedure Failures and Medication Administration Errors. *Journal of Nursing Care Quality*, 32(4):309–317. doi: 10.1097/NCQ.000000000000256.
- Timmons, S., et al. 2014. Implementing human factors in clinical practice. *Emergency Medicine Journal*, 32(5):368–72. doi: 10.1136/emmermed-2013-203203.
- Undang-Undang Republik Indonesia no 44 tahun 2009, Tentang Rumah Sakit.* [Law of the Republic of Indonesia No. 44 of 2009]
- Weinger, M.B. & Gaba, D.M. 2014. Human Factors Engineering in Patient Safety. *Anesthesiology*, Vol. 120(4): 801–806. doi: 10.1097/ALN.000000000000144
- Westbrook, J.I., et al. 2011. Errors in the administration of intravenous medications in hospital and the role of correct procedures and nurse experience. *BMJ Quality and Safety*, 20(12): 1027–1034. doi: 10.1136/bmjqs-2011-000089.



UNIVERSITAS AIRLANGGA

FAKULTAS KESEHATAN MASYARAKAT

Kampus C Mulyorejo Surabaya 601115 Telp.031-5920948, 5920949 Fax 031-5924618

Laman : <http://www.fkm.unair.ac.id> : E-mail: info@fkm.unair.ac.id

SURAT KETERANGAN

Nomor : 3022/UN3.1.10/KP/2023

Yang bertanda tangan di bawah ini :

Nama : Dr. Santi Martini, dr. M.Kes
NIP : 196609271997022001
Pangkat/Golongan : Pembina/ Gol (IV/a)
Jabatan : Dekan

Dengan ini menerangkan bahwa :

Nama : Inge Dhamanti, SKM. M.Kes, MPH., Ph.D
NIP : 19801224 200501 2 002
Pangkat/Golongan : Penata (Gol. III/c)
Jabatan : Lektor

Telah melaksanakan penelitian dengan judul sebagai berikut :

| No. | Judul Karya Ilmiah | Tahun Pelaksanaan Penelitian |
|-----|---|------------------------------|
| 1. | Implementation of Computerized Physician Order Entry in Primary Care: A Scoping Review | 2021 |
| 2. | Adverse Reactions of COVID-19 Vaccines: A Scoping Review of Observational Studies | 2023 |
| 3. | Literature Review: Cause Factor Analysis and an Effort to prevent Medication Administration Error (MAE) at Hospital | 2020 |
| 4. | A Literature review on the Identification of Variables for Measuring Hospital Efficiency in the Data Envelopment Analysis (DEA) | 2021 |
| 5. | Telemedicine Use In Health Facility During Covid-19 Pandemic: Literature Review | 2022 |
| 6. | Faktor yang Menghambat Pelaporan Insiden Keselamatan Pasien di Rumah Sakit: Literature Review | 2021 |
| 7. | Comparison of Four Methods To Detect Adverse Events in Hospital | 2015 |
| 8. | Infections Prevention and Control (IPC) Programs in Hospitals | 2021 |
| 9. | Studi Komparatif Pengembangan Contact Tracing Applications Di Singapura dan Indonesia (Studi Kasus: TraceTogether dan PeduliLindungi) | 2022 |
| 10. | Faktor Penghambat Pelayanan Kesehatan Rutin di Rumah Sakit saat Pandemi COVID-19 | 2021 |
| 11. | Governmental Policies in Managing COVID-19 Pandemic: Comparative Study Between Indonesia and Vietnam, Period of January – March 2020 | 2021 |

| | | |
|-----|---|------|
| 12. | Akses Pelayanan Kesehatan Ibu dan Anak di Puskesmas Selama Pandemi Covid-19 | 2022 |
| 13. | Comparison of Dental Care Policies Before and During The COVID-19 Pandemic: A Literature Review | 2022 |
| 14. | Analysis of Implementation of Patient Identification In Hospitals to Improve Patient Safety in Indonesia | 2022 |
| 15. | Literature Review: Implementation Of Patient Safety Goals In Hospitals In Indonesia | 2021 |
| 16. | Literature Review: Hospital Service Quality During The COVID-19 Pandemic | 2022 |
| 17. | Comparison of Hospital Emergency Room Management Regulations in Indonesia Before and During The COVID-19 Pandemic: Literature Study | 2022 |
| 18. | Analisis Pelaksanaan Pelayanan Kesehatan Perorangan (Ukp) Di Puskesmas Sebelum Dan Selama Pandemi Covid-19: Literature Review | 2022 |
| 19. | Perbandingan Kebijakan Pelaksanaan Imunisasi Rutin pada Anak sebelum dan selama Pandemi | 2022 |
| 20. | Recommendation Analysis Of Mental Health Services For Health Workers During Pandemic Covid-19 | 2021 |
| 21. | Impact Of Implementing A Surgical Safety Checklist In Hospital: Literature Review | 2023 |
| 22. | Quality Improvement For Maternal And Child Health In Primary Health Care: A Scoping Review | 2023 |
| 23. | Implementation Of Root Cause Analysis On Patient Safety Iincidence In Hospital: Literature Review | 2022 |
| 24. | Analisis Peran Stakeholder dalam Kapasitas Rumah Sakit akibat COVID-19: Literature Review | 2022 |
| 25. | Lessons from Indonesia, a country with highest COVID-19 mortality rate in the world: dissecting multiple aspects | 2022 |

Adapun penelitian tersebut layak dilakukan dan menghasilkan output yang sangat baik, meskipun belum ada **Uji Etical Clearence** karena menggunakan metode litteratur review . Demikian surat keterangan ini kami buat untuk dapat dipergunakan sebagai persyaratan pengusulan Jabatan Fungsional Lektor Kepala.

Surabaya, 13 April 2023



Gana Martini, dr. M.Kes
NIP. 196609271997022001