

Health and behavioral outcomes of a telemedicine strategy for COVID-19 and other COVID-19 related diseases and the impact of the COVID-19 pandemic on the health of the community: a cross-sectional study in the city of Surabaya, Indonesia

The relationship between level of a telemedicine and mortality in diabetes foot ulcers patients admitted to Dr. H. Abdul Wahab General Hospital, Lumajang  
Jurnal Kedokteran Universitas Muhammadiyah Surabaya

Epidemiology of pelvic fractures in the emergency room at Dr. Soetomo General Hospital between 2018-2019  
Jurnal Kedokteran Universitas Surabaya

Comparison of subjective and functional results after operative and nonoperative management of distal radius fractures with the proximal ulna: a retrospective study

Kliniscomparative comparison of midshaft clavicle plate fixation between two screws and three screws on each side of the fracture  
Jurnal Kedokteran Universitas Surabaya

The efficacy of physiotherapy in diabetes mellitus and non-diabetes mellitus  
Jurnal Kedokteran Universitas Muhammadiyah Surabaya

Effect of low energy CO<sub>2</sub> laser on facial wrinkles: histological and clinical study in Surabaya  
Jurnal Kedokteran Universitas Muhammadiyah Surabaya

Correlation between obesity and endothelial dysfunction: evidence with Doppler flow study  
Jurnal Kedokteran Universitas Muhammadiyah Surabaya

The difference of physical fitness and lung capacity in individuals with obstructive pulmonary dysfunction (COPD) and acute respiratory distress syndrome (ARDS) in the intensive care unit (ICU) of RSUD Haji Surabaya  
Jurnal Kedokteran Universitas Muhammadiyah Surabaya

Comparison of using an individualized program and family-based program with a family approach  
Jurnal Kedokteran Universitas Muhammadiyah Surabaya

Analysis of the relationship between the personal protective equipment (PPE) usage and the incidence of respiratory symptoms disorders of online respondents that works in Malang  
Jurnal Kedokteran Universitas Muhammadiyah Surabaya

The effect of ginger (Zingiber officinale) extract on the respiratory tract and CBT (COPD Assessment Test) scores in smokers with COPD on level moderate  
Jurnal Kedokteran Universitas Muhammadiyah Surabaya

Microscopic, histological, and immunohistochemical (IHC) study of RT viral bodies in case report of a patient with EBV-EBV seronegative nasopharyngeal carcinoma  
Jurnal Kedokteran Universitas Muhammadiyah Surabaya

Management and quality of life of the elderly with non-typhoidal salmonellosis of birds  
Jurnal Kedokteran Universitas Muhammadiyah Surabaya

Management and quality of life of the elderly with dengue fever: longitudinal study  
Jurnal Kedokteran Universitas Muhammadiyah Surabaya

diterbitkan oleh:  
Fakultas Kedokteran  
Universitas Muhammadiyah Surabaya **2022**

**Person in Charge**

Dr.H.M.Yusuf Wibisono,Sp.P(K),FCCP  
 Dean of Faculty of Medicine, Muhammadiyah University of Surabaya

**Editor in Chief**

dr.Yelvi Levani.,M.Sc

**Member of Editorial Board**

Abdullah Al-Tarique.,Ph.D (Queensland University, Australia)  
 J.L Nouwen, MD, Ph.D (Erasmus University Rotterdam, the Netherlands)  
 Prof. Murat Coskun, Md, Ph.D (IstanbulUniversitesi, Turkey)  
 Prof.Dr.dr.Suhartono Taat Putera.,M.S (Airlangga University, Surabaya)  
 Prof. Takashi Yashiro (JICHI Medical School, Japan)  
 Aziz Alimul Hidayat (Muhammadiyah University of Surabaya,)

**Section Editor**

dr.Syafarinah Nur Hidayah Akil  
 dr.Ayu Lidya Paramita

**Layout Editor**

Dede Nasrullah,S.Kep.Ns.M.Kep

**Reviewer**

Dr.Reny I'tishom, M.Si  
 Dr. dr. Muhammad Anas, Sp.OG  
 dr. Findra Setyaningrum, M.Sc., Ph.D  
 dr.Uning Marlina.,MHSM.,Sp.OG  
 dr.Yudith Annisa Ayu Rezkitha.,Sp.PD  
 dr.Kukuh Dwi Putera Hemugrahanto.,Sp.OT  
 dr. Afrita Amalia Laitupa, Sp.P  
 dr. Musa Ghuffron, MMR  
 dr. Muhammad Reza Utama, MHPE  
 dr.Hafid Algristian.,Sp.KJ  
 dr. Mohammad Subkhan, Sp.P  
 dr. Fadhool Romdhoni, M.Si

**FOREWORD**

Alhamdulillah, praised to Allah, Journal *Qanun Medika: Fakultas Kedokteran Universitas Muhammadiyah Surabaya* vol 06 no 01 has been published. It consists of 15 articles including 1 literature review, 3 case report and 11 research articles in the medical field. In addition, there is 1 international article from India. We would like to thanks our reviewers and editorial board members who helped us in this publication. In order to be internationalized, we only published articles written in English since July 2019. We hope that these articles can be read widely both by domestic and foreign readers.

Thank you,

Yelvi Levani, MD.,M.Sc

Editor in Chief

**Address**

Faculty of Medicine  
 Muhammadiyah University of Surabaya  
 Jl.Sutorejo 59 Surabaya 60113  
 Telp.031-3811966 fax.031-3813096Email:  
 qanunmedika@um-surabaya.ac.id

**Bank account**

Qanun Medika  
 Bank Jatim cabang Dr.Soetomo  
 Bank account number 0323055441





p-ISSN. 2541-2272

e-ISSN. 2548-9526

**Mental and behavioral disorders due to substances during the COVID-19 Pandemic: Prevalence, diagnosis, and management**

Era Catur Prasetya<sup>1</sup>, Kharisma Sukma Nanda<sup>2</sup>, Khofifah Dwi Oktaviana<sup>2</sup>, Lens Hanin Herda Zhafirah<sup>2</sup>, Lia Lisa Arista<sup>2</sup>, M. Dedi Dermawan<sup>2</sup>

**The relationship between bacterial characteristics and mortality in diabetic foot ulcers' patients admitted to Dr. H. Abdul Moeloek General Hospital, Lampung**

Iswandi Darwis<sup>1</sup>, Gusti Ngurah P Pradnya Wisnu<sup>2</sup>, Sekar Mentari<sup>2</sup>

**Epidemiology of pelvic fracture in the emergency room at Dr. Soetomo General Hospital between 2016-2018**

Yuga Rahmadana<sup>1</sup>, M. Zaim Chilmi<sup>1</sup>

**Comparison of subjective and functional results on the operative and non-operative application of clavicle fractures**

Haris Dwi Khoirur Rofiq<sup>1</sup>, Erwin Ramawan<sup>1</sup>

**A biomechanical comparison of midshaft clavicle plate fixation between two screws and three screws on each side of the fractures**

Naufal Ranadi Firas<sup>1</sup>, Erwin Ramawan<sup>2</sup>

**The success of glaucoma therapy in diabetes mellitus and non-diabetes mellitus**

Nur Shani Meida<sup>1</sup>, Ahmad Ikhliluddin<sup>2</sup>, Yunani Setyandriana<sup>3</sup>, Ilma Naafisa Anthrasita<sup>4</sup>

**Shallot (*Allium cepa* L.) skin ethanol extract neutralizes liver oxidative stress in diazinon-induced Wistar rats**

Shahifa Audy Rahima<sup>1</sup>, Rosita Dewi<sup>2</sup>, Nindya Shinta Rusmatika<sup>3</sup>, Dina Helianti<sup>2</sup>

**Correlation between obesity and successful ovulation induction with Clomiphene citrate**

Rida Eka Setiani<sup>1</sup>, Sri Ratna Dwiningsih<sup>2\*</sup>, Gadis Meinari Sari<sup>3</sup>

**The differences of parasitemia and lungs size in malaria-associated acute respiratory distress syndrome (MA-ARDS) and non-MA-ARDS in mice infected with *Plasmodium berghei* ANKA**

Eka Noviya Fuzianingsih<sup>1</sup>, Cyuzuzo Callixte<sup>1</sup>, Marselaonety La'lang<sup>1</sup>, Dinda Eka Putri<sup>1</sup>, Heny Arwati<sup>2</sup>, Lucia Tri Suwanti<sup>3</sup>

**Compliance in taking medication for hypertension patients listed at Healthy Indonesia Program with a Family Approach**

Merry Tiyas Anggraini<sup>1</sup>, Aisyah Lahdji<sup>2</sup>, Hema Dewi Anggraheny<sup>3</sup>

**Analysis of the relationship between using personal protective equipment (PPE) masks on the incidence of respiratory symptoms disorders of online motorcycle taxi drivers in Malang**

Jeremi Setiawan<sup>1</sup>, Hanna Cakrawati<sup>2</sup>, Anung Putri Illahika<sup>3</sup>

**The effect of ginger (*Zingiber officinale*) extract on the neutrophil level and CAT (COPD Assessment Test) scores in workers with COPD due to dust exposure**

Susilo Budi Pratama<sup>1,2</sup>, Yuliani Setyaningsih<sup>2</sup>, Daru Lestyanto<sup>2</sup>

**Bradycardia, renal failure, shock, and hyperkalemia (BRASH) caused by AV nodal blocker: a case report of a patient with BRASH syndrome resistant to calcium administration**

Sidhi Laksono<sup>1,2\*</sup>, Ananta Siddhi Prawara<sup>3</sup>

**Management and quality of life extranodal non hodgkin lymphoma of testis**

Bagus Aulia Mahdi<sup>1</sup>, Pradana Zaky Romadhon<sup>1</sup>, Ugroseno Yudho Bintoro<sup>1</sup>, Merlyna Savitri<sup>1</sup>, Putu Niken Ayu Amrita<sup>1</sup>, Muhammad Noor Diansyah<sup>1</sup>, Ami Ashriati Prayoga<sup>1</sup>, Kartika Prahasanti<sup>2</sup>

**Management and quality of life extranodal non hodgkin lymphoma of testis**

Bagus Aulia Mahdi<sup>1</sup>, Pradana Zaky Romadhon<sup>1</sup>, Ugroseno Yudho Bintoro<sup>1</sup>, Merlyna Savitri<sup>1</sup>, Putu Niken Ayu Amrita<sup>1</sup>, Muhammad Noor Diansyah<sup>1</sup>, Ami Ashriati Prayoga<sup>1</sup>, Kartika Prahasanti<sup>2</sup>



# QANUN MEDIKA

## JURNAL KEDOKTERAN FK UM SURABAYA

<http://journal.um-surabaya.ac.id/index.php/qanunmedika>



### Research Article

## Epidemiology of pelvic fracture in the emergency room at Dr. Soetomo General Hospital between 2016-2018

Yuga Rahmadana<sup>1\*</sup>, M. Zaim Chilmi<sup>1</sup>

1) Department of Orthopaedic and Traumatology, Faculty of Medicine Universitas Airlangga- Dr. Soetomo General Hospital, Surabaya

### ARTICLE INFO

*Submitted* : 24<sup>th</sup> February 2021

*Accepted* : 2<sup>nd</sup> August 2021

*Published* : 25<sup>th</sup> January 2022

#### *Keywords:*

Epidemiology Pelvic Trauma,  
Prevention Pelvic Trauma,  
Management Pelvic Trauma

#### *\*Correspondence:*

yugarahmadana@gmail.com

*This is an Open acces article  
under the CC-BY license*



### ABSTRACT

Pelvic fracture is a rare case. The incidence is about 3% to 8% of the whole kind of fractures. The most common cause is high energy trauma and is associated with multi-trauma. The amount of pelvic trauma patients is 63,000 and is associated with increased mortality rates. The scientific data is needed because it is helpful for trauma prevention and management. This research is a descriptive observational study that examines the epidemiology of pelvic trauma. The population was patients with pelvic trauma who came to the ER RSUD Dr. Soetomo in 2016-2018. The total sample was 229 patients. A total of 149 patients (65.1%) were male, and 80 patients (34.9%) were female. Traffic accidents caused 173 patients (76%) cases. Tile A classification was most frequent in 117 patients (51.1%). There were 178 patients (77.7%) who were multi-trauma patients. The most common treatment was pelvic bandage with 95 patients (41.5%). There were 178 multi-trauma patients, 109 patients (61.2%) who underwent emergency surgery for trauma, and 69 patients (38.8%) who did not. The most common cause of pelvic fracture in Dr. Soetomo general hospital during 2016-2018 was trauma with high energy due to traffic accidents. The most common type of pelvic fracture was Tile A. The more severe the pelvic trauma, the treatment should be more aggressive.



# QANUN MEDIKA

## JURNAL KEDOKTERAN FK UM SURABAYA

<http://journal.um-surabaya.ac.id/index.php/qanunmedika>



### INTRODUCTION

Fracture pelvic is one of the main results of low energy trauma, such as falls, especially in older people. Fracture pelvic is closely associated with morbidity and mortality, with an 8-27% mortality rate of all cases (Andrich *et al.*, 2015). Fracture pelvic has been reported to occur in 3% to 8% of all bone injuries and is often associated with high-energy trauma such as high-speed motor vehicle accidents or falls from heights in younger people. From Epidemiological studies of fracture pelvic, all incidence is 28-37 per 100,000 people for a year (Verbeek *et al.*, 2018).

The main goal of managing fracture pelvic is to reduce the risk of mortality. Severe pelvic trauma can cause significant complications. A recent analysis of more than 63,000 trauma patients showed that fracture pelvic is associated with a high mortality rate (Sathy *et al.*, 2009). There is a significant variability related to risk factors that increase mortality in pelvic trauma. In 15–30%, patients with a high-energy pelvic fracture are commonly unstable hemodynamics. It is directly associated with blood loss due to pelvic trauma. Extensive bleeding is a major cause of death in fracture pelvic (Schulman *et al.*, 2010). The high-power mode of injury is closely related to the mortality rate in fracture pelvic. Several factors influence the prognosis, such as characteristics, diagnosis, and management of fracture pelvic. Therefore, it should be determined immediately to decrease the mortality of these patients. (Breuil, Roux and Carle, 2016). However, there is no data on the incidence rate and characteristics of Dr. Soetomo Hospital and how to handle it in emergency cases.

Therefore, it is important to carry out epidemiological research in orthopedic trauma because epidemiological data can accelerate

diagnosis and management that improve the prognosis. This research aimed to describe the distribution of gender, age, the motion of injury (MOI), hemodynamic condition, classification of fracture pelvic, type of injury, and treatment of pelvic trauma patients in Dr. Soetomo general hospital during 2016-2018. In addition, this study aimed to produce scientific data that can be useful for preventing and managing trauma in the population.

### METHODS

This research was a descriptive observational study that examines the epidemiology of pelvic trauma. This study population was patients with pelvic trauma who came to the ER RSUD Dr. Soetomo in 2016-2018. This study's sample was a total sampling, all patients with pelvic trauma who went to the emergency room Dr. Soetomo in 2016-2018. The patient characteristics such as gender, age, MOI, hemodynamic condition, classification of fracture pelvic, type of injury and treatment were included in this study. Data of this study was taken from the recapitulation of the patient's medical records. Data was manually collected, and the data were analyzed descriptively; it was first selected, edited, coded, and tabulated. Descriptive data were displayed in tables and graphs and analyzed variables (gender, age, MOI, hemodynamic condition, classification of fracture pelvic, type of injury, and treatment). This study had received Ethical Clearance Information No. 0175/LOE/301.4.2/X/2020 issued by the Research Commission of the Dr. Soetomo Hospital Surabaya.

### RESULTS

This retrospective study was conducted with a descriptive design of pelvic trauma profiles in the Emergency Room of Dr. Soetomo Hospital Surabaya in 2016-2018 with a total sample of 229 patients.



# QANUN MEDIKA

## JURNAL KEDOKTERAN FK UM SURABAYA

<http://journal.um-surabaya.ac.id/index.php/qanunmedika>



**Table 1.** Distribution of sample frequencies by gender and age

No.	Characteristic	Amount (%)
1.	<b>Gender</b>	
	Male	149 (65.1%)
	Women	80 (34.9%)
	Total	229 (100%)
2.	<b>Age</b>	
	0-10	5 (2%)
	11-20	55 (24%)
	21-30	45 (20%)
	31-40	44 (19%)
	41-50	45 (20%)
	51-60	23 (10%)
	61-70	9 (4%)
	>70	3 (1%)
	Total	229 (100%)
3.	<b>MOI</b>	
	Traffic Accident	173 (76%)
	Fallen	51 (22%)
	Work accident	5 (2%)
	<b>Total</b>	229 (100%)

According to table 1, out of 229 patients in this study, 149 patients (65.1%) were male, while the remaining 80 patients (34.9%) were female with a ratio of 1.8: 1. The mean patient age was 33.5 years, with the youngest patient was eight years, and the oldest age was 74 years. Traffic accidents are the leading cause of pelvic fractures (76%).

As shown in Table 2, based on pelvic trauma diagnosis classification, most pelvic trauma cases were patients with the Tile A classification, namely 117 patients (51.1%), followed by the APC 2 classification of 26 patients (11.4%). The

third most classification of pelvic trauma is the APC 1 classification of 23 patients (10%) and followed by LC 2 classification of 19 patients (8.3%), LC 1 as many as 15 patients (6.6%), Vertical Shear as many as 14 patients (6.1%), APC 3 was ten patients (4.4%) and LC 3 was five patients (2.2%).

Table 3 shows the distribution of the type of therapy based on hemodynamic conditions. Mostly, the patients were 182 patients with stable hemodynamic conditions, 85 patients were subjected to conservative therapy, 83



**QANUN MEDIKA**  
*JURNAL KEDOKTERAN FK UM SURABAYA*  
<http://journal.um-surabaya.ac.id/index.php/qanunmedika>



patients had a pelvic bandage, 4 patients had C-Clamp insertion, 10 patients had external fixation. 47 patients with unstable hemodynamic conditions, 8 patients received conservative therapy, 12 received pelvic bandage therapy, 17 received C-Clamp insertion, and 10 received external fixation

Based on the type of trauma, from 229 patients, 178 patients, or 77.7% were multi-trauma patients. In contrast, the remaining 22.3% or 51 patients were single-trauma patients. The most common type of comorbid trauma to pelvic trauma was another extremity trauma with 68 cases (38.2%). This study found that

the highest number of other trauma surgeries performed by C-clamp installation was in other orthopedic trauma cases, as many as 5 patients out of a total of 14 patients. Meanwhile, the most other trauma surgery that performed external fixation on the pelvis was urogenital trauma surgery with abdominal trauma surgery as many as 4 out of 17 patients. The pelvic bandage was mainly placed on the patient’s pelvis, who underwent surgery on other extremities in 19 out of 35 patients. Conservative therapy was mostly performed on patients who underwent surgery in other orthopedic fields, in 16 patients from 35.

**Table 2** Distribution of sample frequencies based on the classification of pelvic trauma diagnoses

No	Pelvic Trauma Classification	Amount (n)	(%)
1	<b>Tile A</b>	117	51,1%
2	<b>APC 1</b>	23	10,0%
3	<b>APC 2</b>	26	11,4%
4	<b>APC 3</b>	10	4,4%
5	<b>LC 1</b>	15	6,6%
6	<b>LC 2</b>	19	8,3%
7	<b>LC 3</b>	5	2,2%
8	<b>Vertical Shear</b>	14	6,1%
	<b>Total</b>	229	100%

**Table 3.** Distribution of sample frequencies based on the classification of pelvic trauma diagnoses

		Therapy				Total
		Conservative	Pelvic Bandage	C-Clamp	External Fixation	
<b>Hemodynamic Condition</b>	<b>Stable</b>	85	83	4	10	182
	<b>Unstable</b>	8	12	17	10	47
	<b>Total</b>	93	95	21	20	229



# QANUN MEDIKA

## JURNAL KEDOKTERAN FK UM SURABAYA

<http://journal.um-surabaya.ac.id/index.php/qanunmedika>



		Treatment				Total
		C-Clamp	External Fixation	Pelvic Bandage	Conservative	
Operations with other trauma	A	5	2	19	16	42
	B	0	0	3	3	6
	C	2	1	7	2	12
	D	0	0	2	1	3
	E	1	1	6	2	10
	A+B	0	0	1	3	4
	A+C	1	0	0	0	1
	A+E	3	3	1	4	10
	B+C	0	0	0	1	1
	B+E	0	0	1	0	1
	C+E	1	4	2	3	10
	A+B+E	0	1	0	0	1
	A+C+E	1	2	0	0	3
	A+D+E	0	1	0	0	1
	B+C+E	0	0	1	0	1
	A+B+C+E	0	2	0	0	2
	<b>Total</b>		14	17	43	35

\*Note

- A: Other extremity operation
- B: Thorax operation
- C: Urogenital operation
- D: Head and neck operation
- E: Abdomen operation

## DISCUSSION

The result of this study is consistent with fracture pelvic epidemiology data in other countries, where most patients are male, and the most common age range is 15-25 years. (Wedel and Galloway, 2013). Another study also obtained similar results where pelvic trauma in men was three times more than in women. This is due to their high participation in physical activities such as construction work, motorbike driving, and other activities. In another study by Eluwa et al., it was also shown that the number of men who experienced traumatic events was greater than that of women. When driving, this might be due to men being often impatient, aggressive, and losing concentration (Eluwa et al., 2010). Also, men have a higher risk of traffic accidents (Young, Birrell, and Stanton, 2011).

Traffic accidents are the most common mode of injury of pelvic trauma in this study. This is also consistent with other studies, which further confirm that pelvic trauma is closely related to high energy trauma (Breuil, Roux and Carle, 2016). This study's results are also in line with research conducted by Ghosh et al., which showed that traffic accidents cause as many as 77% of pelvic trauma. Research conducted by Yang et al. 2014 in Taiwan from 2000 - 2011 also showed that 62% of pelvic trauma causes were traffic accidents. (Yang et al., 2014; Ghosh et al., 2019). Data from the World Report on Traffic Injury Prevention, the death rate due to traffic accidents reaches 1.2 million people each year. The causes of the high number of traffic accidents in developing countries are the steady increase of motorized vehicles, poor enforcement of traffic safety regulations, and





# QANUN MEDIKA

## JURNAL KEDOKTERAN FK UM SURABAYA

<http://journal.um-surabaya.ac.id/index.php/qanunmedika>



poor road infrastructure. (Young, Birrell and Stanton, 2011; Organization, 2015)

From the data obtained, 82,1% of patients came with multi-trauma, with about 79,5% stable hemodynamic conditions. The most complicated cases of pelvic trauma in this study were other limb trauma (multiple trauma), in about 38.2%, followed by trauma to other organ systems, which could include trauma to the head and neck, trauma to the thorax, trauma to the urogenital area, trauma to the gastrointestinal area. This result is in line with a study conducted by Yang et al. with 36,594 patients, 23,888 patients experienced multi-trauma, where the most common accompanying trauma was trauma to other extremities in about 60.6%, followed by head injury for 17.6%, and the rest was trauma to thoracic, abdominal and urogenital. Another study in India also obtained similar results where trauma to the other extremities is the most common comorbid trauma to pelvic trauma (Yang *et al.*, 2014; Vidyarthi and Nayak, 2018). An isolated trauma to the pelvis is extremely rare because of pelvic trauma caused by a high-energy injury. High energy trauma transmits the force to the bones and other organs, so pelvic trauma is often accompanied by other accompanying trauma. (Ghosh *et al.*, 2019).

The Tile and Young and Burgess classifications were used to classify each of the pelvic injuries in this study, where these two classifications can be related. Tile A is a stable pelvic trauma condition. Tile B is a partially unstable pelvic trauma condition, where Tile B1 is the same as Young and Burgess APC I and APC II, Tile B2 is the same as Young and Burgess LC I and LC II, Tile B3 is the same as Young and Burgess LC III. Furthermore, Tile C is the same unstable pelvic trauma condition as the Young and Burgess Vertical Shear (VS) and APC III. (Tosounidis *et al.*, 2015). Based on the type of trauma, Tile A (Stable) trauma

was the most common type, accounting for 51.1% of the total patients. Whereas for type B Tile, a total of 40.7% of the total patients were obtained, with the distribution for Tile-type B1 as much as 21.4% (APC 1 10% and APC 2 11.4%), Tile-type B2 as much as 14.9% (LC I was 6.6% and LC II was 8.3%), Tile-type B3 was 2.2% (LC III was 2.2%).

Moreover, for Tile Type C, 10.5% of the total patients were obtained (APC III 4.4% and VS 6.1%). This result is in line with a study conducted by Pereira et al. in Brazil where Tile A pelvic trauma was the largest classification of pelvic trauma at 54.5%, followed by Tile B and Tile C pelvic trauma. (Pereira *et al.*, 2017).

In this study, 20 patients (8.7%) had pelvic trauma who underwent external fixation and 21 patients (9.2%) with C-clamp. These results are similar to that of a study conducted by Pereira et al., wherein 9.1% of patients had external fixation, and 7.6% of patients had C-clamp insertion. External fixation provides temporary pelvic ring stability that is rigid and serves as an adjunct to control early bleeding in hemodynamically unstable pelvic ring disorders. External fixation of the anterior via the iliac crest or the supra-acetabular route provides adequate transient pelvic stability in the APC-II / -III and LC-II / -III injury patterns. A posterior pelvic C-clamp may be indicated to control bleeding in vertical shear injuries with sacroiliac joint disorders. Pelvic C-clamp application is contraindicated in communitive and transformational sacral trauma, iliac wing trauma, and type LC pelvic trauma. (Magnone *et al.*, 2014; Weaver and Heng, 2015).

There were some limitations to this study. Firstly, the number of samples was relatively small. Future studies should involve a larger sample size. Secondly, the year span of operation was too short. The length of data is more representative of the real data about the case. Thirdly, this research is a descriptive study



# QANUN MEDIKA

## JURNAL KEDOKTERAN FK UM SURABAYA

<http://journal.um-surabaya.ac.id/index.php/qanunmedika>



that is needed to comparison study to improve the study.

### CONCLUSION

The most common cause of trauma to the pelvis is trauma with high energy due to traffic accidents. Men had 1.8 times more pelvic trauma than women. Most pelvic trauma types are Tile A. Pelvic trauma patients who experience trauma elsewhere (multi-trauma) are more likely to experience more than just pelvic trauma (single trauma). The more severe the pelvic trauma suffered (see classification), the more aggressive the therapy was. The more hemodynamically unstable the patient, the more aggressive the pelvic trauma therapy was. If an invasive treatment is performed on accompanying trauma in the abdominal and urogenital area, then therapy for pelvic trauma will be more aggressive.

### REFERENCES

- Andrich, S. *et al.* (2015). Epidemiology of pelvic fractures in Germany: considerably high incidence rates among older people, *PLoS One*, 10(9), p. e0139078.
- Breuil, V., Roux, C. H. and Carle, G. F. (2016). Pelvic fractures: epidemiology, consequences, and medical management, *Current opinion in rheumatology*, 28(4), pp. 442–447.
- Eluwa, M. A. *et al.* (2010). Disposition of fractures and dislocations among road accident victims in rivers and bayelsa states of Nigeria from 1992–2005, *IJE*, 8(1).
- Ghosh, S. *et al.* (2019). Epidemiology of pelvic fractures in adults: Our experience at a tertiary hospital, *Chinese journal of traumatology*, 22(3), pp. 138–141.
- Magnone, S. *et al.* (2014). Management of hemodynamically unstable pelvic trauma : results of the first Italian consensus conference ( cooperative guidelines of the Italian Society of Surgery , the Italian Association of Hospital Surgeons , the Multi-specialist Italian Society of Young Surgeons , the Italian Society of Emergency Surgery and Trauma , the Italian Society of Anesthesia , Analgesia , Resuscitation and Intensive Care , the Italian Society of Orthopaedics and Traumatology , the Italian Society of Emergency Medicine , the Italian Society of Medical Radiology -Section of Vascular and Interventional Radiology- and the World Society of Emergency Surgery ), pp. 1–8.
- Organization, W. H. (2015). *Global status report on road safety 2015*. World Health Organization.
- Pereira, G. J. C. *et al.* (2017). Epidemiology of pelvic ring fractures and injuries, *Revista Brasileira de Ortopedia*, 52(3), pp. 260–269.
- Sathy, A. K. *et al.* (2009). The effect of pelvic fracture on mortality after trauma: an analysis of 63,000 trauma patients, *JBJS*, 91(12), pp. 2803–2810.
- Schulman, J. E. *et al.* (2010). Pelvic ring fractures are an independent risk factor for death after blunt trauma, *Journal of Trauma and Acute Care Surgery*, 68(4), pp. 930–934.
- Tosounidis, T. H. *et al.* (2015). The use of buttress plates in the management of acetabular fractures with quadrilateral plate involvement: is it still a valid option?, *International orthopaedics*, 39(11), pp. 2219–2226.
- Verbeek, D. O. *et al.* (2018). Pelvic fractures in the Netherlands: epidemiology, characteristics and risk factors for in-hospital mortality in the older and



# QANUN MEDIKA

## JURNAL KEDOKTERAN FK UM SURABAYA

<http://journal.um-surabaya.ac.id/index.php/qanunmedika>



- younger population, *European Journal of Orthopaedic Surgery & Traumatology*, 28(2), pp. 197–205.
- Vidyarthi, D. A. and Nayak, D. (2018). Study of incidence and prevalence of pelvic fractures in Central India, *J Med Sci Clin Res*.
- Weaver, M. J. and Heng, M. (2015). Orthopedic Approach to the Early Management of Pelvic Injuries Orthopedic Approach to the Early Management of Pelvic Injuries, (March). doi: 10.1007/s40719-014-0005-4.
- Wedel, V. L. and Galloway, A. (2013). *Broken bones: anthropological analysis of blunt force trauma*. Charles C Thomas Publisher.
- Yang, N.-P. *et al.* (2014). Epidemiology of hospitalized traumatic pelvic fractures and their combined injuries in Taiwan: 2000–2011 National Health Insurance data surveillance, *BioMed research international*, 2014.
- Young, M. S., Birrell, S. A. and Stanton, N. A. (2011). Safe driving in a green world: A review of driver performance benchmarks and technologies to support ‘smart’ driving, *Applied ergonomics*, 42(4), pp. 533–539.