

Hospital Utilization among Urban Poor in Indonesia in 2018: is Government-run Insurance Effective?

by Ratna Dwi Wulandari

Submission date: 07-Apr-2023 02:48PM (UTC+0800)

Submission ID: 2058232079

File name: r_in_Indonesia_in_2018_is_Government-run_Insurance_Effective.pdf (753.35K)

Word count: 6470

Character count: 34881

RESEARCH

Open Access

Hospital utilization among urban poor in Indonesia in 2018: is government-run insurance effective?



Ratna Dwi Wulandari^{1*}, Agung Dwi Laksono², Rofingatul Mubasyiroh², Rika Rachmalina², Mara Ipa² and Nikmatur Rohmah³

Abstract

Background An urban poor is a vulnerable group that needs government financing support to access health services. Once they are sick, they will fall deeper into poverty. The study aims to analyze the effectiveness of government-run insurance in hospital utilization in urban poor in Indonesia.

Methods The research analyzed the 2018 Indonesian Basic Health Survey data. This cross-sectional survey collected 75,970 participants through stratification and multistage random sampling. Meanwhile, the study employed hospital utilization as an outcome variable and health insurance ownership as an exposure variable. Moreover, the study looked at age, gender, marital status, education, and occupation as control factors. The research employed a binary logistic regression to evaluate the data in the final step.

Results The results show that someone with government-run insurance is 4.261 times more likely than the uninsured to utilize the hospital (95% CI 4.238–4.285). Someone with private-run insurance is 4.866 times more likely than the uninsured to use the hospital (95% CI 4.802–4.931). Moreover, someone with government-run and private-run insurance has 11.974 times more likely than the uninsured to utilize the hospital (95% CI 11.752–12.200).

Conclusion The study concluded that government-run insurance is more effective than the uninsured in improving hospital utilization among the urban poor in Indonesia. Meanwhile, private-run is more effective than government-run and uninsured in improving hospital utilization among the urban poor in Indonesia. Moreover, the most effective is to combine the kind of health insurance ownership (government-run and private-run).

Keywords Government-run insurance, Health insurance, Hospital utilization, Healthcare evaluation, Health policy, Public health

Introduction

An urban area is an area that has main non-agricultural activities with the structure of the function of the site as a place for urban settlements, concentration and distribution of government services, social services, and economic activities [1]. Vulnerability in urban communities is often caused by having a high population density and diverse livelihoods [2]. The main problem for urban residents is slum settlements resulting from poverty. Poverty is deprivation, namely the lack of everything needed for

* Correspondence:

Ratna Dwi Wulandari

a-d-w@fkm.unair.ac.id

¹ Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia

² National Research and Innovation Agency, the Republic of Indonesia,

Jakarta, Indonesia

³ Faculty of Health Science, Muhammadiyah University of Jember, East Java, Indonesia



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

well-being (experience of a good quality of life). Poverty has many dimensions: physical, social, economic, political, psychological, and spiritual, including the vulnerabilities suffered by the poor, such as physical weakness, isolation, vulnerability, and powerlessness [3]. As a result, poor people in urban areas are more susceptible to illness. This situation is a double burden due to poor environmental quality and poverty [4].

About 7% of the urban population in Indonesia is poor, almost above the poverty line [5]. In congested conditions, the imperfect urban lives near open sewers and standing water are constantly exposed to harmful waste. Inadequate sanitation can lead to the transmission of worms and other intestinal parasites. The urban poor has limited financial resources and higher food costs, so they often experience a lack of nutritious food. Environmental pollution and food often prepared under unhygienic conditions cause disease [6, 7]. In illness, the urban poor must face shocks that impact their limited assets, livelihoods, and savings [5].

The main urban problems in developing countries show two interacting subsystems, namely: 1) Social subsystems such as unemployment, scarcity of access to clean water and sanitation, health services, and; 2) Environmental subsystem (water, air, and soil pollution, waste management, scarcity of clean water, settlements). The needs of the urban poor to survive in good health are often unmet because of the difficulty of accessing public services [25]. The majority of the poor population in urban areas live in temporary buildings. They made it of temporary materials such as bamboo, wood, corrugated zinc, and straw. Vulnerability in these settlements is further exacerbated by the lack of essential services and the disaster hazard stemming from natural disasters [8, 9].

The research results on poor people in Jakarta's slum areas show that they choose to move to Jakarta because they have lost their work relationships and hope Jakarta will provide a better life than their original area. Most work as laborers or porters or porters. Because their income was insufficient to rent a house, the situation forced them to move under the bridge. Environmental sanitation problems become expensive to obtain, and all places around the house can be trash cans [10]. Previous studies reported that urban residents living in slums said sick household members for two weeks [2]. Meanwhile, other studies state that the most vulnerable groups in informal urban settlements are toddlers, children, the elderly, and people with low immune systems [11, 12].

Apart from being susceptible to illness, the condition of urban poor is also vulnerable to becoming poorer. Previous studies have concluded that living in urban areas is associated with a higher likelihood of being poor [13]. Living in urban areas comes with a high price tag, making

people more vulnerable to poverty. Prices of goods in urban areas are more expensive and susceptible to fluctuations. Therefore, they put vulnerable people at risk of falling into poverty if their wages are insufficient to cover the cost of living. Moreover, urban needs are more complex than those in rural areas [14]. So to survive, the poor in urban areas must continue working, even when sick.

People with low economic status often cannot meet their health needs [37], therefore, the Government of Indonesia launched the National Health Insurance (NHI) in 2014. The government has expanded the contribution subsidy for underprivileged families [15]. This National Health Insurance Policy aims to improve the quality of health services and be able to reach all groups, especially the lower middle class. As the implementer of the policy, Health Social Security Administrator (SSA) [15] made maximum efforts to provide socialization and increase public awareness of the importance of health insurance. Still, guaranteeing national health insurance takes a process and time [16].

A previous study has reported that pro-poor interventions such as the national social health insurance scheme have been shown to have contributed to the quality of health services [17]. If successful, the NHI will become the most extensive single-payer system globally. Indonesia has made steady progress, but about a third of its population remains without protection and direct payments for health, even among NHI members. To help close this gap, particularly among the poor, the Indonesian government [22] implementing a series of Universal Health Coverage (UHC) policy reforms that include the integration of the government's remaining insurance schemes into the NHI [18]. UHC achievement is realized through participation in NHI by all Indonesian people, without exception. The local government registered the poor and underprivileged and paid as a participant in the Contribution Assistance Recipient (CAR) [19]. Based on the background narration, the study aims to analyze the effectiveness of government-run insurance in hospital utilization in urban poor in Indonesia.

Materials and methods

Data source

The study looked at secondary data from the Indonesian Basic Health Survey from 2018, and the Ministry of Health of the Republic of Indonesia conducted a national-scale cross-sectional survey. Furthermore, the study collected data from May to July 2018 through interviews with Household Instruments and Individual Instruments [20].

The 2018 Indonesian Basic Health Survey's population included all Indonesian households. The survey took the sample structure from the 2018 National

Socio-Economic Survey, which was performed in March 2018. Furthermore, the survey visited 300,000 families from 30,000 census blocks in the 2018 Socio-Economic Survey (run by the Central Statistics Agency) [20].

The survey used the PPS (probability proportional to size) approach, employing systematic linear sampling in two stages: Stage 1: Implicit stratification based on welfare strata of all census blocks resulting from the 2010 Population Census. PPS chose the sample survey as the sampling frame for selecting census blocks from a master frame of 720,000 from the 2010 Population Census, of which 180,000 were chosen (25%). The survey used the PPS method to determine numerous census blocks in each urban/rural strata per regency/city to create a Census Block Sample List. There are 30,000 Census Blocks in total that have been chosen. Stage 2: Using systematic sampling, select ten homes in each Census Block with the highest implicit stratification of education completed by the Head of Household to preserve the representation of the diversity value of household characteristics. All household members in the selected household will be questioned as part of the 2018 Indonesian Basic Health Survey [20].

The participants were all adults (aged 15 and up) living in Indonesia's impoverished urban societies. The study defined 75,970 people as a weighted sample based on the sampling procedures.

Setting

The study setting was the urban poor in Indonesia at the national level. The study used the Indonesian Central Statistics Agency's provisions for urban categorization in the survey. The survey employed the wealth index formula to determine wealth status. The survey calculated the wealth index by taking a weighted average of a family's overall spending. Meanwhile, the poll used household expenditures such as health insurance, food, and accommodation to calculate the wealth index. Moreover, the survey divided the income index into five categories: quintile 1 (poorest), quintile 2 (poorer), quintile 3 (middle), quintile 4 (richer), and quintile 5 (richest) [21, 22]—the study categories quintile 1 and 2 as the poor.

Outcome variable

The study's outcome variable was hospital utilization, which included adults' access to outpatient or inpatient hospitals. We had unutilized and utilized hospital beds in hospital utilization (outpatient and inpatient). Outpatient was limited to the preceding month, while inpatient was limited to the previous year. The survey asked respondents to recall outpatient and inpatient episodes appropriately depending on the time reference [20].

Exposure variable

The study included health insurance ownership as an exposure variable. The study categorized health insurance ownership into four categories: uninsured, government-run insurance, private-run insurance, and a combination of both insurances (government-run and private-run insurance) [20].

Control variables

The study employed five elements as control variables as part of such factors. The five criteria were age, gender, marital status, educational level, and occupation type.

The study calculated the respondent's age using the respondent's last birthday. However, we separated gender into two categories: male and female. In addition, the study divided marital status into three categories: never married, married/living with a partner, and divorced/widowed.

The acknowledgment of the respondent's most recent diploma is their education. The survey covers four levels of education: none, primary, secondary, and higher. Meanwhile, there are six occupations: no job, government employee/army/police, private sector, entrepreneur, farmer/fisherman/laborer, and others.

Data analysis

The Chi-Square test was used in the early phases of the sample to create a bivariate comparison for the dichotomous variable. We also used the T-test for the continuous variable in the study (age). The study also used a collinearity test to check that the independent variables in the final regression model did not have a strong relationship. The analysis used a binary logistic regression at the study's last point. We employed the previous test to examine the multivariate link between all independent factors and hospital utilization. Throughout the statistical analysis portion of the investigation, we used the IBM SPSS 26 application.

Ethics approval and consent to participate

The National Ethics Committee in the National Institute of Health Research and Development granted the 2018 Indonesian Basic Health Survey (LB.02.01/2/KE.024/2018). The survey deleted all respondents' identities from the dataset. Respondents have provided written approval for their involvement in the study. The author has obtained permission to use data for this study through the website: <http://www.litbang.kemkes.go.id/layanan-permintaan-data-riset/>.

Results

The analysis found that Indonesia's national average hospital utilization among the urban poor in 2018 was 4.2%. Meanwhile, the distribution of health insurance ownership among urban-poor communities was uninsured at 34.9%, government-run insurance at 63.1%, private-run insurance at 1.6%, and simultaneous government-run and private-run insurance at 0.4%.

Table 1 shows descriptive statistics of the participants. Based on hospital utilization, unutilized people are mainly in all types of health insurance ownership. Meanwhile, those with government-run insurance have an older average age than those with other types of health insurance ownership. Moreover, based on gender, females lead in uninsured and government-run insurance; males lead in private-run and combined government-run and private-run insurance.

Regarding marital status, they are married or living with a partner dominated in all kinds of health insurance ownership. On the other hand, based on education level, primary education leads to uninsured, government-run insurance, and combined government-run and

private-run insurance; meanwhile, secondary education leads to private-run insurance. Furthermore, no job leads to all types of health insurance ownership according to occupation type.

The study performed a collinearity test for the next stage. According to the outcome, each factor's variance inflation factor (VIF) value is less than 10.0, and the tolerance value for each variable is more significant than 0.10. The research suggests the test's decision-making basis since the regression model exhibited no multicollinearity.

Table 2 explains the binary logistic regression of hospital utilization among the urban poor in Indonesia. Table 2 shows, based on health insurance ownership, someone with government-run insurance is 4.261 times more likely than the uninsured to utilize the hospital (AOR 4.261; 95% CI 4.238–4.285). Someone with private-run insurance is 4.866 times more likely than the uninsured to use the hospital (AOR 4.866; 95% CI 4.802–4.931). Moreover, someone with government-run and private-run insurance is 11.974 times more likely than the uninsured to utilize the hospital (AOR 11.974; 95% CI 11.752–12.200).

Table 1 Descriptive statistic of participants (n = 75,970)

Participants Characteristics	Health Insurance Ownership				P-value
	Uninsured (n = 24,534)	Government-run (n = 50,233)	Private-run (n = 938)	Government-run and Private-run (n = 265)	
Hospital utilization					< 0.001
Unutilized	98.6%	94.3%	93.9%	87.3%	
Utilized	1.4%	5.7%	6.1%	12.7%	
Age (mean)	(39.47)	(40.71)	(35.55)	(37.75)	< 0.001
Gender					
Male	49.2%	48.4%	51.5%	52.7%	
Female	50.8%	51.6%	48.5%	47.3%	
Marital status					< 0.001
Never in union	22.9%	23.6%	20.9%	23.0%	
Married/living with a partner	67.6%	67.1%	75.2%	67.2%	
Divorced/Widowed	9.5%	9.2%	3.9%	9.8%	
Education level					< 0.001
No education	6.0%	6.5%	1.0%	4.6%	
Primary	65.9%	66.6%	40.3%	48.6%	
Secondary	25.5%	24.4%	52.8%	43.9%	
Higher	2.6%	2.6%	5.8%	2.9%	
Work type					< 0.001
No job	42.6%	42.6%	32.9%	26.0%	
Civil servant/army/police	0.3%	0.7%	1.6%	2.1%	
Private sector	6.2%	7.7%	29.8%	32.4%	
Entrepreneur	16.9%	13.8%	11.7%	11.3%	
Farmer/fisherman/labor	28.1%	29.8%	18.9%	23.7%	
Others	5.8%	5.4%	5.0%	4.4%	

20

Table 2 The result of binary logistic regression of hospital utilization among the urban poor in Indonesia in 2018 (n = 75,970)

Predictor	P-value	Hospital Utilization		
		AOR	95% Confidence Interval	
			Lower Bound	Upper Bound
Insurance: Uninsured	-	-	-	-
Insurance: Government-run	< 0.001	4.261	4.238	4.285
Insurance: Private-run	< 0.001	4.866	4.802	4.931
Insurance: Government-run & Private-run	< 0.001	11.974	11.752	12.200
Age	< 0.001	1.018	1.018	1.018
Gender: Male	-	-	-	-
Gender: Female	< 0.001	1.109	1.105	1.114
Marital: Never in union	-	-	-	-
Marital: Married/Living with partner	< 0.001	1.860	1.848	1.872
Marital: Divorced/Widowed	< 0.001	1.431	1.417	1.444
Education: No Education	-	-	-	-
Education: Primary	< 0.001	1.249	1.239	1.258
Education: Secondary	< 0.001	1.633	1.620	1.647
Education: Higher	< 0.001	1.475	1.455	1.496
Occupation: no job	-	-	-	-
Occupation: civil servant/army/police	< 0.001	0.485	0.474	0.497
Occupation: private sector	< 0.001	0.483	0.478	0.487
Occupation: entrepreneur	< 0.001	0.619	0.615	0.623
Occupation: farmer/fisherman/labor	< 0.001	0.471	0.469	0.474
Occupation: Others	< 0.001	0.784	0.778	0.790

AOR Adjusted Odds Ratio

In addition to health insurance ownership, the study found five control variables significant to hospital utilization among the urban poor in Indonesia: Age, gender, marital status, education level, and occupation type.

Based on gender, female is 1.109 times more likely to utilize the hospital than males (AOR 1.109; 95% CI 1.105–1.114). Regarding marital status, those married or living with a partner are 1.860 times more likely than those who never married to use the hospital (AOR 1.860; 95% CI 1.848–1.872). Moreover, those divorced or widowed are 1.431 times more likely than those who never married to utilize the hospital (AOR 1.431; 95% CI 1.417–1.444).

According to education level, all kinds of education have more possibilities to utilize the hospital than no education. On the other hand, based on occupation type, all occupations have a lower chance than no job to use the hospital.

Discussion

The policy for implementing NHI in Indonesia has stated in Presidential Regulation of the Republic of Indonesia Number 82 of 2018 that health protection guarantees target the entire population without exception (universal

coverage). The implementation of universal health protection coverage is given to every person (participant) who contributes, further divided into two, namely Contribution Assistance Recipient (CAR), which includes the poor and the needy non-CAR participants [23–25].

The results show someone with government-run insurance is more likely than the uninsured to utilize the hospital. Meanwhile, someone with private-run insurance is more likely to use the hospital than the uninsured. Moreover, someone with government-run and private-run insurance is more likely than the uninsured to utilize the hospital. In addition to guaranteed financing from participation in health insurance, hospitals' comprehensive services and facilities (referral services, inpatient care, emergency services) are driving factors for individuals choosing the type of health service facility. Private insurance usually plays a role by providing excess or additional coverage for services that are not included in government-run insurance [26, 27]. The study results align with previous studies, which show that the urban poor with insurance is more likely to take advantage of hospital services [28, 29]. This pattern is the same in the general population who are non-urban poor, both in Indonesia and other countries such as India and Ghana [4, 30, 31].

In general, poor people are less likely to have private insurance, although this does not rule them out, especially for those with chronic illnesses. Even though the Indonesian government has issued a CAR policy to cover the poor in NHI, there are several crucial issues that the government must address. For example regarding the quality of service that is considered lower, service that takes longer, or procedures that are difficult to understand. Resolving these issues can potentially increase NHI utilization [27, 32, 33].

The study found age is related to hospital utilization among the urban poor in Indonesia. People in urban, poor, and old age groups have more excellent opportunities to use hospitals as healthcare facilities. These results can explain that the catastrophic disease, which increases with age, often causes a person to require referral care. Furthermore, old age with low-income characteristics and unhealthy lifestyles will potentially have more diseases and body disorders. This condition can increase the risk of needing care compared to the elderly in social groups with better economic conditions [28, 34].

The result indicates female is more likely to utilize the hospital than males. This study's results align with the previous research [35, 36]. One thing that might be explained is related to the perception that is followed by health behavior from the side of the individual woman. Women who live less healthy lives than men are more likely to take advantage of the hospital, although biologically, women generally live longer than men [1, 37].

The study shows all marital statuses have more likely than those never in a union to use the hospital. This result supports the statement that marital status is an essential predictor of hospital utilization [38]. A study on colon cancer patients in Norway showed a similar effect that revealed more married or ever-married patients used health care in hospitals than those who never married [39]. In addition, other evidence from health insurance beneficiaries in the US and Puerto Rico. The studies reported that married people increase the likelihood of using health services efficiently compared to currently not-married people because they tend to choose outpatient care or skilled nursing facility [38]. The concept of married couples' protective role may explain this result. Spouses can provide physical and emotional support, including taking or accompanying their partner to a health facility for consultation or treatment [40, 41].

Moreover, marriage facilitates health budget-sharing engagement, allowing them to seek health care at health facilities [38, 42]. Those who are not married tend to be more prone to health problems [41]. Policymakers must understand health facilities utilization according to marriage status because the marriage number in Indonesia

has declined in the last decade, from 2.3 million in 2011 to 1.7 million in 2021 [43]. It may affect health-seeking behavior, health status, and health budget.

The study informs all kinds of education have more possibilities to utilize the hospital than no education. Education is strongly associated with better income and wealth [44], and it can lead to better access to information, health insurance, and health. The Wisconsin cohort study also confirms that adults with higher educational attainment have more access to health care, particularly for preventive care, such as physical and dental check-ups, flu vaccinations, and cholesterol tests [45, 46]. This behavior can accelerate positive health outcomes and increase healthcare costs as a short-term impact [46]. Therefore, policies related to health investment should align with intervention strategies to improve education, prioritizing the dropped out-of-school group for better health literacy.

Moreover, based on occupation type, all occupations have a lower chance than no job to use the hospital. A study in Sweden may explain that employed adults have a higher health-related quality of life scores than those unemployed [47]. This condition may prevent them from using hospitals or other health facilities frequently. Contrarily, being unemployed harms health, such as mental health [48]. A review study concluded that the unemployed was associated with increased healthcare utilization because universal health coverage was available and accessible for this group [49]. The Indonesian government has launched a national health insurance program, Healthy Indonesia Card (HIC), and poor people receive a beneficiary contribution scheme [50]. In 2016, this program protected 14.5 million poor people from severe poverty due to health care costs [51]. The program should run parallel with other social protection programs to achieve a sustainable impact targeting the poor or unemployed, such as business assistance for the working-age or small business loans.

Strength and limitation

The study examines a large amount of data to represent information nationally. On the other hand, the study examines secondary data; therefore, the variables analyzed are limited to acceptable ones. Other variables linked to hospital utilization reported in previous studies, such as trip time, hospital travel costs, and disease type, cannot be investigated [4, 52, 53].

16 Conclusion

Based on the results, the study concluded that government-run insurance is more effective than the uninsured in improving hospital utilization in urban poor in

Indonesia. On the other hand, private-run is more effective than government-run and uninsured in improving hospital utilization in urban-poor communities in Indonesia. Furthermore, the most effective is to combine the kind of health insurance ownership (government-run and private-run).

Abbreviations

6	National Health Insurance
UHC	Universal Health Coverage
SDGs	Sustainable Development Goals
PPS	Probability proportional to size
VIF	Variance inflation factor
AOR	Adjusted odds ratio

Acknowledgements

The author would like to thank the National Institute of Health Research and Development, which has agreed to the author to analyze the 2018 Indonesian Basic Health Survey data.

Authors' contributions

RDW and ADL developed the proposal, analyzed the data, and interpreted the results. RM and RR were significant contributors in conducting the study, interpreting the data and writing the manuscript. MI and NR were substantial contributors to conducting the research and writing the manuscript. All authors read and approved the final manuscript.

Funding

Not applicable.

Availability of data and materials

The author cannot publicly share the data because a third party and the Ministry of Health of the Republic of Indonesia, who owns the data, do not have permission to share it. The 2018 Indonesian Basic Health Survey data set is available on the web <http://www.litbang.kemkes.go.id/layanan-permintaan-data-riset/> for researchers who meet the criteria for access to confidential data.

Declarations

Ethics approval and consent to participate

The National Ethics Committee in the National Institute of Health Research and Development granted the 2018 Indonesian Basic Health Survey (LB.02.01/2/KE.024/2018). The survey deleted all respondents' identities from the dataset. Respondents have provided written approval for their involvement in the study. The author has obtained permission to use data for this study through the website: <http://www.litbang.kemkes.go.id/layanan-permintaan-data-riset/>. The author carried out all methods following relevant guidelines and regulations.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Received: 18 August 2022 Accepted: 10 January 2023

Published online: 12 January 2023

References

- Wulandari RD, Laksono AD, Nantabah ZK, Rohmah N, Zuardin Z. Hospital utilization in Indonesia in 2018: do urban-rural disparities exist? *BMC Health Serv Res.* 2022;22(22):491.
- Mpanje D, Gibbons P, McDermott R. Social capital in vulnerable urban settings: an analytical framework. *J Int Humanit Action.* 2018;3(4):1–14.
- Suprijadi B. Poverty and environmental degradation problems are main obstacles in urban development (Permasalahan Kemiskinan dan Kerusakan Lingkungan Merupakan Hambatan Utama dalam Pembangunan Perkotaan). *MADANI Kaji Ilm Bid Sos Polit Komun dan Inf.* 2016;17(1):1.
- Laksono AD, Wulandari RD, Efendi F. Determinants of hospital utilisation among urban poor societies in Indonesia. *Int J Innov Creat Chang.* 2020;12(9):375–87.
- Asian Development Bank. Building resilience of the urban poor in Indonesia. 2022. p. 120. <https://doi.org/10.22617/TCS210404-2>.
- Kuddus MA, Tynan E, McBryde E. Urbanization: a problem for the rich and the poor? *Public Health Rev.* 2020;41(1):1–4.
- Wulandari RD, Laksono AD, Nantabah ZK, Rohmah N, Zuardin Z. Hospital utilization in Indonesia in 2018: do urban-rural disparities exist? *BMC Health Serv Res.* 2022;22(1):1–11.
- Haque AN. A 'whole systems' view of vulnerability to climatic risks: the case of the urban poor in Dhaka. *Bangladesh Prog Dev Stud.* 2020;20(2):101–18.
- Wulandari RD, Laksono AD, Rohmah N. Urban-rural disparities of antenatal care in South East Asia: a case study in the Philippines and Indonesia. *BMC Public Health.* 2021;21(1):1221.
- Yandri P, Juanda B. Understanding the character of urban poverty using an observational approach (Memahami Karakter Kemiskinan Perkotaan Dengan Pendekatan Observasional). *J Ekon Stud Pembang.* 2018;19(1):75–84.
- Zerbo A, Delgado RC, González PA. Vulnerability and everyday health risks of urban informal settlements in Sub-Saharan Africa. *Glob Heal J.* 2020;4(2):46–50.
- Weimann A, Oni T. A systematised review of the health impact of urban informal settlements and implications for upgrading interventions in South Africa, a rapidly urbanising middle-income country. *Int J Environ Res Public Health.* 2019;16(19):1–17.
- Hadijah Z, Sadali MI. The effect of urbanization on poverty reduction in Indonesia (Pengaruh Urbanisasi Terhadap Penurunan Kemiskinan di Indonesia). *J Wil dan Lingkung.* 2020;8(3):290–306.
- Suryahadi A, Marlina C. Understanding metropolitan poverty: the profile of poverty in Jabodetabek Area. 2018.
- Soewondo P, Johar M, Pujisubekti R. Access to health services for low-economy families in the JKN Era (Akses Pelayanan Kesehatan Keluarga Berstatus Ekonomi Rendah di Era JKN). *J Ekon dan Pembang Indones.* 2021;21(1):108–24.
- Basuki EW, Sulistyowati, Herawati NR. Implementation of National Health Insurance Policy by BPJS Health in Semarang City (Implementasi Kebijakan Jaminan Kesehatan Nasional Oleh BPJS Kesehatan Di Kota Semarang). *Diponegoro J Soc Polit Sci Tahun.* 2016;5(04):1–14.
- Liyanto E, Nuryana D, Cahyani RA, Utomo B, Magnani R. How well are Indonesia's urban poor being provided access to quality reproductive health services? *PLoS ONE.* 2022;17(4):e0265843.
- Wiseman V, Thabrany H, Asante A, Haemmerli M, Kosen S, Gilson L, et al. An evaluation of health systems equity in Indonesia: Study protocol. *Int J Equity Health.* 2018;17(138):1–9.
- Annida. Health financing policy for the poor in achieving universal health coverage in Banjar District (Kebijakan Pembiayaan Kesehatan terhadap Masyarakat Miskin dalam Pencapaian Universal Health Coverage di Kabupaten Banjar). *J Kebijak Pembang.* 2020;15(2):219–29.
- National Institute of Health Research and Development of the Ministry of Health of the Republic of Indonesia. The 2018 Indonesian Basic Health Survey. Jakarta; 2018. p. 99. Available from: http://labdata.litbang.kemkes.go.id/images/download/laporan/RKD/2018/Laporan_Nasional_RKD2018_FINAL.pdf. Cited 2021 Oct 7.
- Wulandari RD, Qomarrudin MB, Supriyanto S, Laksono AD. Socioeconomic disparities in hospital utilization among elderly people in Indonesia. *Indian J Public Heal Res Dev.* 2019;10(11):1800–4.
- Wulandari RD, Laksono AD, Prasetyo YB, Nandini N. Socioeconomic disparities in hospital utilization among female workers in Indonesia: a cross-sectional study. *J Prim Care Community Health.* 2022;13(2):1–7.
- Laksono AD, Wulandari RD, Zuardin Z, Nopianto N. The disparities in health insurance ownership of hospital-based birth deliveries in eastern Indonesia. *BMC Health Serv Res.* 2021;21(1):1261.

24. Nasution SK, Mahendradhata Y, Trisnantoro L. Can a national health insurance policy increase equity in the utilization of skilled birth attendants in Indonesia? A Secondary analysis of the 2012 to 2016 national socio-economic survey of Indonesia. *Asia-Pacific J Public Heal.* 2020;32(1):19–26.
25. Laksono AD, Wulandari RD, Matahari R. The determinant of health insurance ownership among pregnant women in Indonesia. *BMC Public Health.* 2021;281(1):1538.
26. Jiang Y, Ni W. Impact of supplementary private health insurance on hospitalization and physical examination in China. *China Econ Rev.* 2020;63:101514.
27. Laksono AD, Nantabah ZK, Wulandari RD, Khoiri A, Tahangnacca M. Barriers to expanding the national health insurance membership in Indonesia: who should the target? *J Prim Care Community Heal.* 2022;13:21501319221111110.
28. Singh LM, Siddhanta A, Singh AK, et al. Potential Impact of the Insurance on Catastrophic Health Expenditures Among the Urban Poor Population in India. *J Health Manag.* 2022;0(0). <https://doi.org/10.1177/09720634221088425>.
29. Laksono AD, Wulandari RD, Soedirham O. Urban and rural disparities in hospital utilization among Indonesian adults. *Iran J Public Health.* 2019;48(2):247–55.
30. Nandi S, Schneider H, Dixit P. Hospital utilization and out of pocket expenditure in public and private sectors under the universal government health insurance scheme in Chhattisgarh State, India: Lessons for universal health coverage. *PLoS ONE.* 2017;12(11):e0187904.
31. Dalinjong PA, Welaga P, Akazili J, Kwarteng A, Bangha M, Oduro A, et al. The association between health insurance status and utilization of health services in rural Northern Ghana: evidence from the introduction of the National Health Insurance Scheme. *J Heal Popul Nutr.* 2017;36(42):1–10.
32. Laksono AD, Rukmini R, Tumaji T, Ipa M, Wulandari RD. National health insurance membership among urban poor societies in Indonesia in 2019: are they protected? *Economies.* 2022;10(8):196.
33. Sumadi AF, Mardiyoko I, Pratama YY. Differences in the levels of satisfaction of JKN patients and general patients on the quality of inpatient unit services: literature review (Perbedaan Tingkat Kepuasan Pasien JKN Dan Pasien Umum Terhadap Mutu Pelayanan Unit Rawat Inap: Literatur Review). *J Manaj Kesehat.* 2022;8(1):86–97.
34. Wulandari RD, Laksono AD. Urban-rural disparity: the utilization of primary health care center among elderly in East Java. *Indonesia J Adm Kesehat Indones.* 2019;7(2):147–54.
35. Zhao X, Liu Y, Zhang X, Bärnighausen T, Chen S. The impact of retirement on inpatient healthcare utilization in Guangzhou, China: a regression discontinuity analysis of 189,031 health insurance claims. *BMC Geriatr.* 2022;22(1):380.
36. Mahmudiono T, Laksono AD. Disparity in the hospitals utilization among regions in Indonesia. *Open Access Maced J Med Sci.* 2021;9(E):1461–6.
37. Zorrilla-Muñoz V, Agulló-Tomás MS, Rodríguez-Blázquez C, Ayala A, Fernandez-Mayoralas G, Forjaz MJ. Ageing perception as a key predictor of self-rated health by rural older people—a study with gender and inclusive perspectives. *Land.* 2022;11(3):323.
38. Pandey KR, Yang F, Cagney KA, Smieliauskas F, Meltzer DO, Ruhnke GW. The impact of marital status on health care utilization among Medicare beneficiaries. *Medicine (Baltimore).* 2019;98(12-p e14871):1–8.
39. Bjørnelv GMW, Edwin B, Fretland AA, Deb P, Aas E. Till death do us part: the effect of marital status on health care utilization and costs at end-of-life. A register study on all colorectal cancer decedents in Norway between 2009 and 2013. *BMC Health Serv Res.* 2020;20(115):1–13.
40. Wulandari RD, Laksono AD, Matahari R. Does husband's education level matter to antenatal care visits? A study on poor households in Indonesia. *Indian J Community Med.* 2022;47(2):192–5.
41. Megatsari H, Laksono AD, Herwanto YT, Sarweni KP, Geno RAP, Nugraheni E, et al. Does husband/partner matter in reduce women's risk of worries?: Study of psychosocial burden of covid-19 in Indonesia. *Indian J Forensic Med Toxicol.* 2021;15(1):1101–6.
42. Laksono AD, Wulandari RD, Sukoco NEW, Suharmiati S. Husband's involvement in wife's antenatal care visits in Indonesia: What factors are related? *J Public Health Res.* 2022;11(2):1–8.
43. Central Bureau of Statistic Indonesia. Jumlah Nikah, Talak dan Cerai, serta Rujuk (Pasangan Nikah), 2011–2021. *Data Sensus.* 2022.
44. Scott A, Wolla and Jessica Sullivan. Education, Income, and Wealth. 2017: 1–7. Source <https://research.stlouisfed.org/publications/page1-econ/2017/01/03/education-income-and-wealth/>. Accessed 22 Sept 2022.
45. Fletcher JM, Frisvold DE. Higher education and health investments: does more schooling affect preventive health care use? *J Hum Cap.* 2009;3(2):144–76.
46. Raghupathi V, Raghupathi W. The influence of education on health: an empirical assessment of OECD countries for the period 1995–2015. *Arch Public Health.* 2020;78(1):1–18.
47. Norström F, Waenerlund A-K, Lindholm L, Nygren R, Sahlén K-G, Brydsten A. Does unemployment contribute to poorer health-related quality of life among Swedish adults? *BMC Public Health.* 2019;19:457.
48. Norström F, Virtanen P, Hammarström A, Gustafsson PE, Janlert U. How does unemployment affect self-assessed health? A systematic review focusing on subgroup effects. *BMC Public Health.* 2014;14(1):1310.
49. Jin RL, Shah CP, Svoboda TJ. The impact of unemployment on health: a review of the evidence. *Can Med Assoc J.* 1995;153(5):529–666.
50. President of the Republic of Indonesia RI. Presidential Instruction Number 8 of 2017 concerning Optimizing the Implementation of the National Health Insurance Program. Jakarta: Republic of Indonesia; 2017. p. 1–9.
51. BPJS Kesehatan. Gathering of the President and participants beneficiaries of National Health Insurance - Healthy Indonesia Card. 2018.
52. Laksono AD, Wulandari RD. Predictors of hospital utilization among papuans in Indonesia. *Indian J Forensic Med Toxicol.* 2020;14(2):2319–24.
53. Wei Y, Yu H, Geng J, Wu B, Guo Z, He L, et al. Hospital efficiency and utilization of high-technology medical equipment: a panel data analysis. *Heal Policy Technol.* 2018;7(1):65–72.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions



Hospital Utilization among Urban Poor in Indonesia in 2018: is Government-run Insurance Effective?

ORIGINALITY REPORT

16%

SIMILARITY INDEX

12%

INTERNET SOURCES

10%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

1	tropmedhealth.biomedcentral.com Internet Source	2%
2	Submitted to Politeknik Kesehatan Kemenkes Semarang Student Paper	1%
3	jurnal.unej.ac.id Internet Source	1%
4	Submitted to Cypress Fairbanks Independent School District Student Paper	1%
5	pubmed.ncbi.nlm.nih.gov Internet Source	1%
6	www.hindawi.com Internet Source	1%
7	Agung Dwi Laksono, Ratna Dwi Wulandari, Noor Edi Widya Sukoco, Suharmiati Suharmiati. "Husband's involvement in wife's antenatal care visits in Indonesia: What	1%

factors are related?", Journal of Public Health Research, 2022

Publication

8	e-journal.unair.ac.id Internet Source	1 %
9	eprints.uad.ac.id Internet Source	<1 %
10	Submitted to Darton State College Student Paper	<1 %
11	Submitted to Badan PPSDM Kesehatan Kementerian Kesehatan Student Paper	<1 %
12	Andrias Feri Sumadi, Muhammad Syamsu Hidayat, Sulistyawati Sulistyawati. "A Narative Review: The Difference of Satisfaction Level in BPJS Patient And General Patient Toward The Quality of Health Service In Inpatient's Unit", Jurnal Aisyah : Jurnal Ilmu Kesehatan, 2021 Publication	<1 %
13	translational-medicine.biomedcentral.com Internet Source	<1 %
14	www.frontiersin.org Internet Source	<1 %
15	www.iiste.org Internet Source	<1 %

www.medicolegalupdate.org

16

Internet Source

<1 %

17

media.neliti.com

Internet Source

<1 %

18

pubs.sciepub.com

Internet Source

<1 %

19

www.intechopen.com

Internet Source

<1 %

20

Hanifa M. Denny, Agung D. Laksono, Ratu Matahari, Bina Kurniawan. "The Determinants of Four or More Antenatal Care Visits Among Working Women in Indonesia", Asia Pacific Journal of Public Health, 2021

Publication

<1 %

21

Ratna Dwi Wulandari, Agung Dwi Laksono. "Determinants of knowledge of pregnancy danger signs in Indonesia", PLOS ONE, 2020

Publication

<1 %

22

Submitted to Southern New Hampshire University - Continuing Education

Student Paper

<1 %

23

ijop.net

Internet Source

<1 %

24

Elvira Liyanto, Dewi Nuryana, Restu Adya Cahyani, Budi Utomo, Robert Magnani. "How well are Indonesia's urban poor being

<1 %

provided access to quality reproductive health services?", PLOS ONE, 2022

Publication

25 journals.sagepub.com <1 %
Internet Source

26 jphres.org <1 %
Internet Source

27 openaccess.city.ac.uk <1 %
Internet Source

28 pdfs.semanticscholar.org <1 %
Internet Source

29 worldwidescience.org <1 %
Internet Source

30 www.medicopublication.com <1 %
Internet Source

31 Agung Dwi Laksono, Rukmini Rukmini, Ratna Dwi Wulandari. "Regional disparities in antenatal care utilization in Indonesia", PLOS ONE, 2020 <1 %
Publication

32 Anne Lykkeskov, Ezio Di Nucci. "Chapter 4 COVID-19 and Intergenerational Justice: The Case of Denmark", Springer Science and Business Media LLC, 2022 <1 %
Publication

apps.who.int

33

Internet Source

<1 %

34

es.scribd.com

Internet Source

<1 %

35

www.grafiati.com

Internet Source

<1 %

36

Anika Nasra Haque. "A 'Whole Systems' View of Vulnerability to Climatic Risks: The Case of the Urban Poor in Dhaka, Bangladesh", Progress in Development Studies, 2020

Publication

<1 %

37

Budiarsih Budiarsih. "Analysis of the Impact of Corruption in the Health System and Some Recommendations Concepts", Research Square Platform LLC, 2021

Publication

<1 %

Exclude quotes Off

Exclude matches < 5 words

Exclude bibliography On

Hospital Utilization among Urban Poor in Indonesia in 2018: is Government-run Insurance Effective?

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8
