

[ABOUT THE JOURNAL](#)[ANNOUNCEMENTS](#)

### ISSN International Centre

p-ISSN : 2580-0140

e-ISSN : 2597-7571

Journal of Public Health Research and Community Health Development , well known as JPH RECODE, is a journal published by Public Health Study Program Sekolah Ilmu Kesehatan Dan Ilmu Alam (SIKIA) Universitas Airlangga. Our journal is published biannually in October and March.

JPH RECODE is on public health as discipline and practices related to preventive and promotion measures to enhance health of the public through scientific approach applying variety of technique. This focus includes area and scope such as biostatistics and health population, epidemiology, health education and promotion, health policy and administration, environmental health, public health nutrition, sexual and reproductive health, and occupational health and safety.

We accept manuscripts in the following categories: 1) Original research papers, 2) Critical review articles, 3) Short communications, 4) Literature Review, 5) Case Study, on applied or scientific research relevant to public health and or Community Health Development. JPH RECODE has been indexed in [DOAJ](#), [Sinta 4](#), [Dimensions](#), [EBSCO Essentials](#) and [others indexing](#). JPH RECODE have a collaboration with [Association of Indonesian Public Health Scholars and Professionals \(PERSAKMI\)](#), [The Indonesian Epidemiology Association \(PAEI\)](#), [Indonesian Association of Construction Safety Experts \(PAKKI\)](#), [Indonesian Association of Behavior Change Intervention Practitioners \(IPIPIPI\)](#), [Indonesian Hyperhealth and Work Safety Association \(AHKKI\)](#), [Association of Indonesian Public Health Experts \(IAKMI\)](#), [Association of Indonesian Environmental Health Experts \(HAKLI\)](#).

JPH RECODE was first published since October 2017 (Volume 1 Issue 1) with 6 manuscripts in 60 pages, then from March 2018 until now our journal has been publishing 8 manuscripts in every edition. Every edition now consists of 65-75 pages. Manuscripts are written in full Indonesian (Bahasa Indonesia) or full English except on abstracts used both languages are Indonesian and English.

The journal employs peer-review mechanism where each submitted article should be anonymously reviewed by expert peers appointed by the editor. Articles published in this journal could be in the

form of original article.

Please note that Volume 1 Issue 1 2017 - Volume 2 Issue 1 2019 was previously processed offline. Volume 1 Issue 1 2017 can be accessed at <http://journal.unair.ac.id/JPHRCHD>, while Volume 1 Issue 2 2018 -Volume 2 Issue 1 2019 are available in archive (<https://e-journal.unair.ac.id/JPHRECODE/issue/archive>)

## Current Issue



Vol. 4 No. 2 (2021): March

Current Issue



Vol. 4 No. 2 (2021): March

Published: 2021-03-04

## Front Matter

Front Matter Vol 4 No 2, 2021

 DOI : 10.20473/jphrecode.v4i2.25944

 Abstract : 77

 PDF : 106

 PDF

## Back Matter

Back Matter Vol 4 No 2, 2021

 DOI : 10.20473/jphrecode.v4i2.25945

 Abstract : 99

 PDF : 158


 PDF

## Research Articles

Correlation Between Knowledge And Attitude Toward Personal Menstrual Hygiene Practice Among Adolescents

 DOI : 10.20473/jphrecode.v4i2.14460

 Pinesa Rachma Andani

 88-98


 Abstract : 1057


 PDF : 1318

 PDF

Factors Associated With Pediatric Pulmonary Tuberculosis Incidence In Surakarta City Health Office Work Area

 DOI : 10.20473/jphrecode.v4i2.18568

 Rezania Asyfiradayati, S.KM., M.PH.

 99-106


 Abstract : 913

 PDF : 723

 PDF

Utilization Of Electronic Reminder Method On Enhancement Hazard Reporting Cards In Oil & Gas Company X

 DOI : 10.20473/jphrecode.v4i2.21559

 Hannisa Yanuar Utama

 107-121

 Abstract : 371


 PDF : 597

 PDF

The Correlation Between Safety Climate With Unsafe Act In Apartment Building Construction Workers Pt. Multikon 2020

 DOI : 10.20473/jphrecode.v4i2.21545

 Rahmah Amalina , Rahmah Hida Nurriszka , Fandita Tonyka Maharani

 122-129


 Abstract : 973


 PDF : 985

 PDF

Relationship Between The Use Of Iodine Salt And The Growth And Development Of Babies

 DOI : 10.20473/jphrecode.v4i2.15418

 weni tri purnani

 130-136


 Abstract : 395


 PDF : 567

 PDF

Risk Factors That Associated With Hypertension In Population Of Tlatah Village, Purwosari, Bojonegoro

 DOI : 10.20473/jphrecode.v4i2.22862

 Ayu Sekar Pawening , Dani Nasirul Haqi

 137-143


 Abstract : 1612

 PDF : 1078

 PDF

Recommendation Analysis Of Mental Health Services For Health Workers During Pandemic Covid-19

 DOI : 10.20473/jphrecode.v4i2.23230

 Anisa Nur Kholipah , Inge Dhamanti

 144-156

 Abstract : 1110


 PDF : 930


 PDF

## Risk Factors Associated With The Incidence Of Ari In Infants Ages 0-2 Years In One Of The Highest Contributing Areas For Ari, East Java

 DOI : 10.20473/jphrecode.v4i2.23394

 Eti Vera Asmaningrum , Dani Nasirul Haqi

 157-165

 Abstract : 504

 PDF : 603

 PDF

### National Accreditation



Accreditation

Number: 1429/E5.3/HM.01.01/2022

### Journal Policy

Focus and Scope	Publication Ethics
Privacy Statement	Peer Reviewers
Editorial Team	Peer Reviewers Process
Open Access Statement	Plagiarism
Copyright Transfer Agreement	Copyright Notice
Archiving	Article Processing Charge
Indexing	Old Website
Contact	

### Instruction for Author

Guide for Authors	Online Submission
-------------------	-------------------




## Editorial Team


### Editor in Chief



**Susy Katikana Sebayang, Ph.D**

Public Health Study Program, Universitas Airlangga-Banyuwangi  
Campus, Indonesia

 0000-0003-0470-8308  
y\_tZ-9IAAAAJ&hl  
24068188900


 ABI-1686-2022


### Managing Editor



**Ayik Mirayanti Mandagi, S.KM., M.Kes**

Public Health Study Program, Universitas Airlangga-Banyuwangi  
Campus, East Java, Indonesia

 0000-0002-6566-2182  
l3zVXxMAAAAJ&hl  
57223457661


 AEM-6430-2022


### Editorial Boards



**Ni Wayan Septarini, dr., M.PH**

Curtin University, Perth, Australia


 0000-0002-9998-0947  
PtL7XDYAAAAJ&hl  
57156581600


 AFA-9126-2022



**Dr. Dian Kusuma, ScD. MPH**

City University of London, London, United Kingdom


 0000-0002-1909-9341  
QuXXWqkAAAAJ&hl  
57190047530

 ACM-5692-2022




**Setho Hadisuyatmana,**  
**S.Kep.Ns.,M.NS(CommHlth&PC)**

Honorary Fellow at the La Trobe University, Victoria, Australia

 0000-0002-3560-7596


RxPPDCEAAAAJ&hl  
57209738095

 ABC-7537-2020




**Assoc. Prof. Dr Satvinder Kaur**

Faculty of Applied Sciences UCSI University Kuala Lumpur, Malaysia

 0000-0003-0808-9612


p5deu7IAAAAAJ&hl  
56211436300

 AAE-7782-2019




**Dr. Normala Binti Ibrahim**

Hospital Pengajar Universiti Putra Malaysia, Malaysia

 0000-0003-2868-9834


Xk8FqC4AAAAJ&hl  
54414227000

 AFN-3092-2022



**Ssekalembe Geofrey**

FELizabeth Glaser Pediatric AIDS Foundation, Uganda

 0000-0001-6757-7275

VUUK6pAAAAAJ




**Stefania Widya Setyaningtas**

Interdepartmental Nutrition Program, Purdue University, United State  
of America

 0000-0002-2331-2037

cSF3RrAAAAAJ  
57208284644

 AGE-8185-2022




**Dr. dr. Daru Lestantyo, MSi**

Public Health Study Program, Diponogoro University, Indonesia

 0000-0002-6614-773X


mNMGqiMAAAAAJ&hl  
57194585807

 GQQ-0368-2022




**Luh Putu Lila Wulandari, dr., M.PH**

Department of Public Health and Preventive Medicine, Medical Faculty, Universitas Udayana, Indonesia

 0000-0002-3397-3648


biyMoywAAAAJ&hl

54394664900

 H-8156-2019


**Syifa'ul Lailiyah, S.KM., M.Kes**

Public Health Study Program, Universitas Airlangga-Banyuwangi Campus, East Java, Indonesia

 0000-0002-1237-3213


lcZsuOgAAAAJ&hl

57213354857

 AAU-2716-2020


**Novia Handayani, S.KM., M.A., M.Kes**

Health Promotion and Behavioural Sciences Departemen, Faculty of Public Health, Universitas Diponegoro, Indonesia

 0000-0002-5701-7263

Z\_tKYyKAAAAJ&hl

57194601772

 GOP-1684-2022


**Feranita Utama, S.KM., M.Kes**

Departemen of Epidemiology, Faculty of Public Health, Universitas Sriwijaya, Indonesia

 0000-0003-3505-7106


yNkoOgwAAAAJ&hl

57211550400

 AFF-1058-2022


**Septa Indra Puspikawati, SKM.,MPH**

Public Health Study Program, Universitas Airlangga-Banyuwangi Campus, East Java, Indonesia

 0000-0003-0289-9745

\_aldm3cAAAAJ&hl

57214820573

 ABE-4655-2020






**Erni Astutik, S.K.M., M.Epid**

Epidemiology Department, Faculty of Public Health, Universitas Airlangga, Indonesia





 0000-0003-2934-1290

YSY-LPUAAAAJ&hl

57208131686

 ABI-2046-2020**Kurnia Ardiansyah Akbar, S.KM., M.KKK**Occupational Health and Safety Department, Faculty of Public Health,  
Universitas Jember, Indonesia 0000-0001-6265-3064  
WTM88jAAAAAJ&hl  
57214728526 AAC-2926-2021**Septa Katmawanti, S.Gz., M.Kes**Public Health Program Study, Faculty of Sport and Science,  
Universitas Negeri Malang, Indonesia 0000-0002-5115-0311  
wrCHArAAAAAJ&hl  
57219311239 AAV-8485-2021**Rahmi Susanti, SKM., M.Kes**

Faculty of Public Health, Universitas Mulawarman, Indonesia

 0000-0001-9921-7717  
ua3u3AMAAAAAJ&hl  
57210159858 AIE-8295-2022**Desak Made Sintha Kurnia Dewi, S.KM., M.Kes**Public Health Study Program, Universitas Airlangga-Banyuwangi  
Campus, Indonesia 0000-0003-3662-0060  
\_OkcZtcAAAAAJ&hl  
57204352395 AAV-3927-2020

## Administration Staff

**Rika Yunita S.Tr.Par**Sekolah Ilmu Kesehatan Dan Ilmu Alam (SIKIA), Universitas  
Airlangga, Indonesia

National Accreditation



**RECOMMENDATION ANALYSIS OF MENTAL HEALTH SERVICES FOR HEALTH WORKERS DURING PANDEMIC COVID-19**  
*Analisis Rekomendasi Layanan Kesehatan Mental Bagi Tenaga Kesehatan Pada Masa Pandemi Covid-19*

Anisa Nur Kholipah<sup>1</sup>, Inge Dhamanti<sup>1</sup>

<sup>1</sup>Public Health Faculty, Universitas Airlangga, Indonesia  
Anisa.nur.kholipah-2017@fkm.unair.ac.id

---

**ARTICLE INFO**

*Article History:*

Received:  
November, 16<sup>th</sup>, 2020

Revised:  
From November, 27<sup>th</sup>,  
2020

Accepted:  
December, 24<sup>th</sup>, 2020

Published online:  
March, 4<sup>th</sup>, 2021

---

**ABSTRACT**

**Background:** Health workers have a higher risk of experiencing mental health problems during the Covid-19 pandemic. This causes the need for the provision of mental health services given the important role of health workers in dealing with Covid-19 and the high psychological burden of health workers while serving patients. **Research objectives:** This study aims to analyze recommendations of mental health services for health workers during the Covid-19 pandemic. **Method:** The method used in writing this article is a literature review. Articles were obtained through the Google Scholar, Science Direct and Pubmed databases published in 2020. Articles reviewed were eight articles. **Results:** Based on the disaster management cycle which consists of response, recovery, and mitigation stages. The response stage requires identifying risks of mental health problems; the recovery stage can be done by providing psychological support and reward; the mitigation stages can be done by compiling a health care system preparedness and improving the skills of health workers. **Conclusion:** Mental health problems in health workers during the Covid-19 pandemic can be overcome through the provision of health service based on disaster risk management which consists of a response, recovery, and mitigation stages.

**Keywords:** Mental Healthcare, Health Workers, Covid-19

**ABSTRAK**

**Latar belakang:** Tenaga kesehatan memiliki risiko lebih tinggi mengalami masalah kesehatan mental selama pandemi Covid-19. Hal ini menyebabkan perlunya penyediaan pelayanan kesehatan mental mengingat pentingnya peran tenaga kesehatan dalam menangani Covid-19 serta tingginya beban psikologis tenaga kesehatan selama melayani pasien. **Tujuan penelitian:** Penelitian ini bertujuan untuk menganalisis rekomendasi pelayanan kesehatan mental bagi tenaga kesehatan pada masa pandemi Covid-19. **Metode:** Metode yang digunakan dalam penulisan artikel ini adalah *literature review*. Artikel didapatkan melalui database *Google Scholar*, *Science Direct* dan *Pubmed* yang dipublikasikan pada tahun 2020. Artikel yang direview berjumlah delapan artikel. **Hasil penelitian:** Terdapat berbagai masalah kesehatan mental yang didasarkan pada tahapan *disaster risk management* yang terdiri dari tahap *response*, *recovery*, dan *mitigation*. Upaya yang dapat dilakukan pada tahap *response* adalah identifikasi risiko masalah kesehatan mental; upaya yang dapat dilakukan pada tahap *recovery* adalah pemberian dukungan psikologis, dan *reward*; upaya yang dapat dilakukan pada tahap *mitigation* adalah menyusun *healthcare system preparedness* dan meningkatkan keterampilan tenaga kesehatan. **Kesimpulan:** Masalah kesehatan mental pada tenaga kesehatan selama pandemi Covid-19 dapat diatasi melalui penyediaan layanan kesehatan mental berdasarkan *disaster risk management* yang terdiri dari tahap *response*, *recovery*, dan *mitigation*.

**Kata Kunci:** Pelayanan Kesehatan Mental, Tenaga Kesehatan, Covid-19

---

## INTRODUCTION

The case of Corona Virus 2019 disease (Covid-19) was first detected at the end of December 2019 in Wuhan City, Hubei Province, China. This case has increased and has began to spread outside Hubei province in the middle of January 2020. The high level of community mobilization on the Chinese New Year holiday becomes the cause why this case is quickly spreading to various regions. Until January 23, 2020, the outbreak had reached other countries with attractive spread throughout the world. On March 11, 2020, WHO declared Covid-19 a global pandemic based on the high spread of cases in various countries (Zu, 2020). Based on data on the development of Covid-19, until 7 November 2020 there were 433,836 positive cases (+4,262 cases from the previous day), 364,417 recovered (83.9% of confirmed), 54,879 in treatment (12.65% of confirmed) and 14,540 died ( 3,351% of confirmed). Accumulatively, confirmed cases continue to experience a higher increase than the accumulated recovery of Covid-19 patients (Covid-19 Management Task Force, 2020).

The study of the Global Burden of Disease conducted by IMHE (The Institute for Health Metrics and Evaluation) in 2015 revealed convincing data on the map of burden of disease around the world. The years lost due to disability (YLD) data from the study states that six of the 20 types of diseases that are considered the most responsible for causing disability are mental disorders. Contrary to popular belief, the impact of a poor mental health condition is no better than that of infectious diseases and is getting worse every year. (Ridlo, and Zein, 2018)

Several studies have indicated mental health problems for health workers during the Covid-19 pandemic. Lai, J., et al in 2020 conducted a study on anxiety, depression, and insomnia symptoms on 1,830 health workers consisting of 702 doctors and 1,128 nurses. Based on the results of this study, more than 70% of respondents reported experiencing psychological distress, in which female nurses suffered the majority of symptoms of depression, anxiety, and distress. This psychological response occurs since 71.5% of nurses are junior nurses who do not have much work experience. Walton, et al (2020) also explained that the Covid-19 pandemic caused various mental health problems such as acute

stress reactions, moral injury, and posttraumatic stress disorder.

According to Wen, et al (2020), healthcare professionals have a higher risk of experiencing insomnia, anxiety, and depression during the pandemic, which are caused by difficulties in getting work safety. The non-fulfillment of the right to work safety includes a lack of knowledge about the prevention and control of Covid-19 infection, limited personal protective equipment, and an increased workload. Meanwhile, according to Walton (2020) mental health problems in healthcare professionals occur because of concerns regarding to limited contact with family, and dealing with patients with conditions that worsen easily. The provision of mental health services is imperative considering the high risk of health workers in dealing with Covid-19 and the high psychological burden of healthcare professionals while serving patients.

According to BNPB (2020), a disaster is an event or series of events that threatens and disrupts people's lives and livelihoods, which are caused by either natural factors and / or non-natural factors as well as human factors resulting in human casualties, environmental damage, property loss, and psychological impacts. Non-natural disasters are caused by non-natural events such as pandemics and disease outbreaks. Based on this definition, it can be concluded that the Covid-19 pandemic is a form of non-natural disaster. According to APCICT (2011), disasters can be handled using a disaster risk management approach, such as a systematic process through various strategies and policies to reduce the adverse effects of hazards and the possibility of disasters. Disaster risk management consists of a prevention stage to prevent a disaster, response to deal with bad impacts when a disaster occurs, recovery to recover people from the effects of a disaster, and mitigation to reduce disaster risk (APCICT, 2011). The Covid-19 pandemic as a non-natural disaster has a psychological impact on healthcare professionals, which can be overcome using a disaster risk management approach (APCICT, 2011). Thus, this article aims to discuss mental health efforts for health workers during the Covid-19 pandemic, which are based on disaster risk management at the response, recovery and mitigation stages. This issue is considered as important, considering that there are limited studies that discuss efforts to

address mental health problems for healthcare professionals based on the disaster risk management stage.

## METHOD

The design employed in this study was a literature review of articles published in 2020. The literature review design was used to obtain information about the object of research being studied in various countries, thus creating comprehensive information. Articles obtained through the Google Scholar database, Science Direct and Pubmed in August 2020 with the keywords "mental health care" or "mental health" and "health worker" and "Covid-19". The selection of articles was based on inclusion criteria, including

Indonesian or English articles, open access articles, healthcare professionals research targets, and cross-sectional or online surveys research design. The researcher does not limit the country of origin of the articles used as references.

The results of the search for articles on the Pubmed database found 3 articles, 44 articles for Science Direct and 98 articles on Google Scholar, thus the total articles were 145 articles. The results of the assessment based on conformity with the inclusion criteria obtained 25 relevant articles. After going through the review process of full text, eight relevant articles were obtained. Below is an outline of the article search results :




Figure 1. Article Search Results Framework

Based on the search results, all reference articles were published online in 2020. The articles came from China, Italy, Saudi Arabia, the Dominican Republic, Iran, and Nepal. Each

article contains the main information analyzed, which are mental health issues and recommendations for mental health services for healthcare professionals.

## RESULTS

The following is *literature review* Matrix:

Table 1. *Literature Review* Matrix

No	Authors, Country	Research Design	Respondents	Instruments	Research Findings	Recommendation for Mental Health Services
1.	Xiao, X. et al., China	<i>Cross sectional</i>	958 health workers from 26 provinces in China. 39.5% of the respondents were doctors, 37.5% were nurses, and 54.8% were young graduates.	<ol style="list-style-type: none"> <li><i>Questionnaires regarding demographic and exposure factors</i></li> <li><i>Perceived Stress Scale (PSS-14)</i></li> <li><i>Hospital anxiety and depression scale (HAD-scale)</i></li> </ol>	55.1% of respondents experienced higher psychological stress than health workers who faced SARS. 54.2% had symptoms of anxiety, and 58% had symptoms of depression. Stress is influenced by position and length of work. Anxiety is influenced by gender, position, level of protection and contact history. Depression is influenced by the level of protection and contact history.	<ol style="list-style-type: none"> <li>Grouping new health workers and female health workers based on work experience to improve practical skills.</li> <li>Psychological education to reduce pressure and develop psychological abilities.</li> <li>Promote Balint Group to develop the ability to face emotional challenges, and improve the quality of medical care.</li> <li>Perform job promotions and provide financial incentives</li> </ol>
2.	Temsah, M. H. et al., Saudi Arabia	<i>Survei online</i>	582 health worker in all departments and clinical units of King Khalid University Hospital. 75% of respondents were female with a nurse percentage of 62.4%. 44.8% of respondents from acute care areas (ICU and emergency), 28% of <i>outpatient clinics, and</i>	<ol style="list-style-type: none"> <li><i>Generalized Anxiety Disorder (GAD-7)</i></li> <li><i>Anxiety Severity screening tool</i></li> </ol>	41.1% of health workers are more anxious about Covid-19 than MERS-CoV, 41.4% have the same level of anxiety in dealing with MERS-CoV and Covid-19, 17.5% are more stressful in facing MERS-CoV. The frequency of anxiety in transmitting the virus to family and friends was	<ol style="list-style-type: none"> <li>Provide control assessment and preventive measures to ensure the safety of health workers</li> <li>Communicate policy effectively</li> <li>Deliver the right information</li> <li>Manage the workers</li> <li>Manage workload</li> <li>Provide emotional support</li> <li>Provide education and training to prepare health</li> </ol>

Table 1. <i>Liteature Review Matrix</i> (continue)						
No	Authors, Country	Research Design	Respondents	Instruments	Research Findings	Recomendation for Mental Health Services
			19.4% of general hospital wards.		greater (2.71 / 5) than in transmitting the virus to himself (2.57 / 5).	workers for pandemic conditions
3.	Tella, M. D. e al., Italia	Survei <i>online</i>	145 health workers practicing at Piedmont (72 doctors and 73 nurses). 72% are female respondents, and 43% work in the Covid-19 handling unit.	<ol style="list-style-type: none"> <li>1 <i>Quality of life and health-related Visual Analogue Scales (VAS)</i></li> <li>2 <i>State-Trait Anxiety Inventory-Form Y1 (STAI Y1)</i></li> <li>3 <i>Beck Depression Inventory (BDI-II)</i></li> <li>4 <i>PTSD Checklist for DSM-5 (PCL-5) to investigate the presence of PTSS.</i></li> </ol>	Health workers who treat Covid-19 patients experience higher symptoms of depression and PTSS than health workers in other units. Single or unmarried female health workers have a higher risk of developing symptoms of depression. Older female health workers have a higher risk of experiencing PTSS.	<ol style="list-style-type: none"> <li>1. Provide psychological care such as cognitive therapy</li> <li>2. Provide the needs of health workers</li> <li>3. Provide psychosocial support</li> <li>4. Identify mental health risks to health workers.</li> </ol>
4.	Giusti, E. M. et al., Italia	<i>Cross sectional</i>	330 health professionals working at Istituto Auxologico Italiano Hospital.	<ol style="list-style-type: none"> <li>1. <i>State-Trait Anxiety Inventory—State form Psychological distress:Depression, Anxiety and Stress Scale-21 (DASS)</i></li> <li>2. <i>Post-traumatic</i></li> </ol>	235 health workers (71.2%) had anxiety levels above the clinical limit, 88 health workers (26.8%) had clinical level depression, 103 (31.3%) experienced anxiety, 113 (34.3%) experienced stress, and 121 (36.7%) experienced	<ol style="list-style-type: none"> <li>1. Setting working hours</li> <li>2. Giving awards</li> <li>3. Providing psychological support programs</li> <li>4. Development of organizational work</li> <li>5. Skills training</li> <li>6. Online cognitive attitude therapy</li> <li>7. Timely monitoring and</li> </ol>

Tabel 1. Matriks <i>Liteature Review</i> (Lanjutan)						
No	Authors, Country	Research Design	Respondents	Instruments	Research Findings	Recomendation for Mental Health Services
				<i>symptoms:</i> <i>Impact of Event Scale Revised–6 items version (IES-6)</i> 3. <i>Maslach Burnout Inventory (MBI)</i>	post-traumatic stress. 107 (35.7%) experienced moderate level of burnout, 105 (31.9%) experienced severe emotional fatigue, 46 (14%) had moderate level. 40 (12.1%) had severe depersonalization level, 132 (40.1%) moderate level, and 113 (34.3%) decreased self-achievement.	maintenance to improve healthcare system preparation in the medium and long term.
5.	Batista, Z. E. et al., Republik Dominika	Tes psikologi	115 doctors from the Dominican Republic aged 23 to 66 years, of which 67.9% were female and 32.1% male.	1. <i>Perceived Stress Scale (PSS-14)</i> 2. <i>Emotion Regulation Questionnaire (ERQ)</i>	Contact with Covid-19 patients can increase stress on health care providers during the Covid-19 pandemic. This leads to the need for emotional suppression strategies, although this is not related to the use of cognitive re-evaluation.	Create interventions that focus on promoting the flexibility of using Emotion Regulation (ER) strategies to reduce stress levels and possibly develop Post-Traumatic Symptoms.
6.	Shoja, E. et al., Iran	Survei online	495 health workers in Iran	1. <i>Demographic Questionnaire</i> 2. <i>NASA-TLX Questionnaire</i> 3. <i>General Health Questionnaire (GHQ-12)</i>	Health workers handling Covid-19 patients significantly had a greater workload and mental health problems than other units ( $p < 0.001$ ). Nurses are	1. Providing psychological interventions for health workers. 2. Planning work shifts based on the psychological characteristics of health workers to reduce workload



No	Authors, Country	Research Design	Respondents	Instruments	Research Findings	Recomendation for Mental Health Services
					health workers with the highest workload ( $71 \pm 16.13$ , $p < 0.001$ ).	3. Attention to macro ergonomic factors to increase motivation, job involvement, job enrichment and work commitment 4. Providing financial and social support to increase working spirit.
7.	Khanal, P. et al, Nepal	<i>Cross sectional</i>	475 health workers working in Covid-19 response management units in Nepalese health facilities. These health workers include doctors, nurses, pharmacists, diagnostic personnel, paramedics and public health practitioners.	1. <i>Hospital anxiety and depression scale (HADS: 0-21)</i> 2. <i>Insomnia severity index (ISI:0-28)</i>	41, 9% of health workers experienced symptoms of anxiety, 37.5% had symptoms of depression, 33.9% had symptoms of insomnia.	1. Reducing stigma 2. Ensure an adequate support system such as PPE and family support.
8.	Zhang, W. et al, China	<i>Survei online</i>	2182 Chinese who are more than 16 years old.	1. <i>The Insomnia Severity Index (ISI)</i> 2. <i>The Symptom Check List-revised (SCL-90-R)</i> 3. <i>The Patient Health Questionnaire-4 (PHQ-4)</i> 4. <i>The Patient Health</i>	Health worker had a prevalence of insomnia (38.4 vs. 30.5%, $p < 0.01$ ), anxiety (13.0 vs. 8.5%, $p < 0.01$ ), depression (12.2 vs. 9.5%; $p < 0.04$ ), somatic symptoms (1.6 vs. 0.4 %; $p < 0.01$ ), and obsessive <i>Compulsive symptoms</i> (5.3 vs. 2.2%; $p < 0.01$ ) were greater than that of non-health worker.	1. Providing the adequacy work condition 2. Providing the recovery program purposes to support medical practitioners adjusting with the work environment and keeping the mental and physical balance 3. Reducing workload 4. Increasing job control and reward 5. Provision of PPE

Table 1. *Literature Review Matrix (Continue)*

No	Authors, Country	Research Design	Respondents	Instruments	Research Findings	Recomendation for Mental Health Services
				<i>Questionnaire-2 (PHQ-2)</i>	Health worker have higher total ISI, GAD-2, PHQ-2, and SCL-90-R obsessive-compulsive symptoms ( $p \leq 0.01$ ) scores than non-health worker.	6. Providing intervention via smartphone.

Table 2. Matrix of mental health service recommendations for health workers:

No	Recomendation	Mental Health Efforts	Disaster Risk Management
1.	<ul style="list-style-type: none"> <li>a. Grouping of new health workers and female health workers based on work experience to improve practical skills. (Xiao, X. et al)</li> <li>b. Identify mental health risks to health workers. (Tella, M. D. e al.)</li> </ul>	Identification of Risk of Mental Health Problems in Health Workers.	<i>Response</i>
2.	<ul style="list-style-type: none"> <li>a. Managing workloads (Temsah, M. H. et al)</li> <li>b. Setting working hours (Giusti, E. M. et al)</li> <li>c. Planning work shifts based on the psychological characteristics of health workers to reduce the workload, especially in the operation unit, ICU, handling Covid-19 and emergencies (Shoja, E. et al)</li> <li>d. Reducing workload (Zhang, W. et al)</li> </ul>	Manage work Shift	
3.	<ul style="list-style-type: none"> <li>a. Providing the needs of health workers (Tella, M. D. e al.)</li> <li>b. Ensuring adequate support systems such as PPE (Khanal, P. et al)</li> <li>c. Provision of adequate working conditions (Zhang, W. et al)</li> <li>d. Provision of PPE (Zhang, W. et al)</li> <li>e. Delivering the right information (Temsah, M. H. et al)</li> </ul>	Pemberian Dukungan <i>Tangible</i>	
4.	<ul style="list-style-type: none"> <li>a. Psychological education to reduce stress and develop</li> <li>b. psychological abilities (Xiao, X. et al)</li> <li>c. Provides psychological care such as cognitive therapy (Tella, M. D. e al.)</li> <li>d. Providing psychological support programs (Giusti, E. M. et al)</li> </ul>	Giving psychological support	

Table 2. Matrix of mental health service recommendations for health workers (Continue)			
No	Recomendation	Mental Health Efforts	Disaster Risk Management
	<ul style="list-style-type: none"> <li>e. Providing psychosocial support (Tella, M. D. e al.)</li> <li>f. Providing emotional support (Temsah, M. H. et al)</li> <li>g. Online cognitive attitude therapy (Giusti, E. M. et al)</li> <li>h. Create interventions that focus on promoting the flexibility of using emotion regulation (ER) strategies to reduce stress levels and the possibility of developing post-traumatic symptoms (Batista, Z. E. et al).</li> <li>i. Providing psychological interventions for health workers (Shoja, E. et al)</li> <li>j. Reducing stigma (Khanal, P. et al)</li> <li>k. Providing intervention via smartphone (Zhang, W. et al)</li> <li>l. Promote Balint Group to develop the ability to face emotional challenges, and improve the quality of medical care (Xiao, X. et al)</li> <li>m. Provision of a recovery program to support medical personnel to adapt to the work environment and maintain a balance of mental and physical health (Zhang, W. et al).</li> </ul>		<i>Recovery</i>
5.	<ul style="list-style-type: none"> <li>a. Doing job promotions and providing financial incentives (Xiao, X. et al)</li> <li>b. Awarding (Giusti, E. M. et al)</li> <li>c. Providing financial and social support to increase morale (Shoja, E. et al)</li> <li>d. Giving attention to macro ergonomic factors to increase motivation, job involvement, job enrichment, and Work comitment. (Shoja, E. et al)</li> </ul>	Giving Reward	
6.	<ul style="list-style-type: none"> <li>a. Timely monitoring and care to improve healthcare system preparation in the medium and long term (Giusti, E. M. et al)</li> <li>b. Communicating policies effectively (Temsah, M. H. et al)</li> <li>c. Providing control assessment and preventive measures to ensure</li> <li>d. the safety of health workers (Temsah, M. H. et al)</li> <li>e. Improve job control (Zhang, W. et al)</li> <li>f. Development of organizational work (Giusti, E. M. et al)</li> <li>g. Managing the workforce (Temsah, M. H. et al)</li> </ul>	Creating Healthcare System Preparedness	<i>Mitigation</i>
7.	<ul style="list-style-type: none"> <li>a. Providing education and training to prepare health workers for pandemic conditions (Temsah, M. H. et al)</li> <li>b. Skills training (Giusti, E. M. et al)</li> </ul>	Increase the Skills of Health Workers	

Based on the results of the instrument analysis, the most mental health problems which were examined was anxiety and depression. There were seven instruments to measure anxiety (HAD Scale, GAD-7, Anxiety Severity screening tool, STAI Y1, STAI-S, DASS, HADS: 0 -21) and five instruments to measure depression (HAD scale, BDI III, DASS, HADS: 0-21, PHQ-4, PHQ-2). Other instruments were PCL-5, IES-6, PSS-14, DASS, NASA-TLX, MBI, ISI 0-28, ERQ, GHQ-12, and SCL-90-R. Based on the analysis of the article, the most dominant psychological response that occurred in health workers were anxiety, depression, and stress. Other mental health problems experienced by health professionals were depersonalization, PTSD, emotional exhaustion, trauma, moral injury, somatic symptoms, and obsessive-compulsive disorder. According to the results, it can be concluded that during a pandemic, health workers did experience mental health problems, especially anxiety, depression, and stress.

Regarding the psychological responses experienced by health workers, there were various recommendations for mental health services that were important to be implemented. These recommendations can be classified based on the disaster risk management stage which consisted of preparedness, response, recovery, and mitigation (APCICT, 2011). Recommendations at the preparedness stage were not found since the recommendations were given after the Covid-19 pandemic. Recommendations that can be suggested at the response stage were identification of the risk of mental health problems in health workers, work shift arrangements, and provision of tangible support. Recovery efforts that can be performed provided psychological support and gave rewards. Efforts at the mitigation stage were the preparation of healthcare system preparedness and improved the skills of health workers.

## DISCUSSION

Law Number 24 of 2007 stated that a disaster is an event or series of events that threatens and disrupts people's lives and livelihoods. These events can be caused by natural factors, non-natural factors, and human factors that result in human casualties, environmental damage, property losses, and

psychological impacts. Thus, it can be concluded that the Covid-19 pandemic is a non-natural disaster that has various impacts, one of them is a psychological impact on health workers. The following are recommendations for mental health efforts for health workers based on disaster risk management:

### Response

Response is a series of activities performed immediately at the time of a disaster to prevent bad impacts (APCICT, 2011). Based on the results of the article analysis, efforts that can be made at the response stage are the identification of risks to mental health problems in health workers, work shift arrangements, and provision of tangible support, such as provision of PPE and the needs of health workers. These various efforts can ensure the availability of an adequate work environment. Those are performed to minimize the risk of mental health problems for health workers at the beginning of the Covid-19 pandemic.

An increased workload which is followed by pressure to be in traumatic situations such as seeing a patient's death, can increase the risk of mental health problems for health professionals. This condition is often exacerbated by the existence of a work environment that does not guarantee the safety of health workers in carrying out their responsibilities. Hence, the hospital needs to identify the risk factors that can cause mental health problems in health workers.

Work shift arrangements are an important recommendation to maximize productivity and provide adequate rest time for health workers. This arrangement can be based on the psychological characteristics and work experience of health workers (Shoja, E., et al. 2020). Health workers also need tangible support. Tangible support includes listening to complaints from health workers, creating a healthy work environment, job promotion, providing financial support, and rewards according to the needs of health workers. The creation of a healthy work environment is one of the recommendations of ISQua (2020) to ensure the psychological safety of health workers during a pandemic. The various supports are expected to maintain health and increase the morale of health workers.

## Recovery

Based on the results of the analysis of the article, efforts that can be made to recover mental health problems in health workers are psychological support and reward. Based on ISQua recommendations, organizations should prepare their staff to deal with future trauma through PFA (Psychological First Aid), which is a response to staff who are suffering or need psychological support. This support includes providing education about the cumulative effects of trauma, stress management training, and psychotherapy (ISQua, 2020).

Psychosocial intervention recommendations during a pandemic must be tailored to the needs without violating physical distancing. The use of online media or hotlines will increase the required psychological encounter. Tele-mental health uses useful information and communication to provide remote mental health services including consultation, evaluation, medical management, and psychotherapy. It cannot be denied that in the future, tele-mental health is a method for providing mental health services that are safe, effective, comfortable, and measurable (Whaibeh, Mahmoud & Naal, 2020). This telemedicine is expected to be able to provide psychological assistance and stress relief efforts both individually and in groups.

According to the analysis of resource needs, training of doctors at public health centers, specialists in other medical services, psychologists and paramedical teams regarding psychological interventions are crucial (De Sousa, et al, 2020). This education and training can help health workers develop experiences, skills, and mental wellbeing during a pandemic (Temsah, M. H et al, 2020). This psychological support needs to be accompanied by the provision of rewards such as incentives, job promotions, and other efforts according to the needs of health workers to increase motivation and work commitment.

## Mitigation

Mitigation is a series of efforts to reduce disaster risk (APCICT, 2011). Based on the results of the article analysis, efforts to reduce the risk of mental health problems for health workers are the preparation of a Healthcare System Preparedness and improvement of the skills of health workers. Psychosocial support in disaster management is more important than infection control during an outbreak.

Guidelines for psychosocial services during a pandemic must be issued quickly to facilitate the implementation of interventions as well as an essential implication for disaster preparedness in general during an outbreak (Windarwati, H. D. et al, 2020).

Intervention plans in healthcare system preparedness need to be tailored to the needs of health workers and implemented at the organizational, team, and individual levels. This can support the creation of adequate working conditions, thus the health workers can adapt to the work environment quickly and are able to better maintain a balance of physical and mental conditions (Zhang, W. R, 2020).

## RESEARCH LIMITATIONS

This research only reviews articles in English and Indonesian. Future researchers are expected to be able to review articles from more diverse languages with a more comprehensive discussion.

## CONCLUSION

Based on the results of the article analysis, mental health problems experienced by health workers are anxiety, depression, stress, insomnia, depersonalization, PTSD, emotional exhaustion, trauma, moral injury, somatic symptoms, and obsessive-compulsive disorder. This problem can be overcome through various recommendations for mental health services at each stage of disaster risk management. The response phase requires identifying risks of mental health problems, arranging work shifts, and providing tangible support to prevent the adverse effects of the pandemic on health workers. Recovery efforts can be performed by providing psychological support and rewards. Mitigation efforts to reduce the risk of health problems can be conducted by preparing healthcare system preparedness and improving the skills of health workers.

## SUGGESTION

Policy makers, both from the central government to health services, are expected to work together in creating policies that support the fulfilment of mental health rights for health workers. The policy is expected to be well implemented as well as monitoring, and conducting the evaluation, thus it can resolve

mental health problems for health workers and prevent these problems in the future.

## REFERENCE

- APCICT. 2011. Module 9 Introduction to *Disaster risk management*, Training of Trainers Workshop on ICT for *Disaster risk management*.
- Barry, M. & Al Amri M, M. Z., 2020. COVID-19 in the shadows of MERS-CoV in the Kingdom of Saudi Arabia. *Journal Epidemiol Glob Health*, 10(1), pp. 1-3.
- Batista, Z. E. G., et al. 2020. COVID-19 pandemic and health worker stres: The mediating effect of emotional regulation. 20 June.pp. 1-23.
- BNPB. 2020. Definisi Bencana. [Online] Available at: <https://bnpb.go.id/definisi-bencana> [Accessed 9 Desember 2020].
- De Sousa, A. & Mohandas, E. & J. A., 2020. Psychological interventions during COVID-19: Challenges for low and middle income countries. *Asian Journal of*, April.51(102128).
- Giusti, E. M., et al. 2020. The Psychological Impact of the COVID-19 Outbreak on Health Professionals: A Cross-Sectional Study. *Frontiers in Psychology*, 10 July, Volume 11, pp. 1-9.
- ISQua. 2020. *PATIENT SAFETY RECOMMENDATIONS FOR COVID-19 EPIDEMIC OUTBREAK*. Italia
- Khanal, P., et al. 2020. Mental health impacts among health workers during COVID-19 in a low resource setting: a crosssectional survey from Nepal. *Research Square*, pp. 1-27.
- Lai, J., et al. 2020. Factors associated with mental health outcomes among health careworkers exposed to coronavirus disease 2019. *Jama Network*. 23 March. 3(3), pp. 1-12.
- Neto, M., et al. 2020. When health professionals look death in the eye: the mental health of professionals who deal daily with the 2019 coronavirus outbreak. *Psychiatry Research*, 13 April, Volume 288, pp. 1-3.
- Percudani, M., et al. 2020. Mental Health Services in Lombardy during COVID-19 outbreak. *Psychiatry Research*, 12 April, 288(112980), pp. 1-3.
- Presti, G., et al. 2020. Mental health support to staff in a major hospital in Milan (Italy) during the COVID-19 pandemic: a framework of actions. *General Psychiatry*, 1 July, Volume 33, pp. 1-4.
- Rajkumar, R. P. 2020. COVID-19 and mental health: A review of the existing literature. *Asian Journal of Psychiatry*. 31 March. Volume 52. 1-5.
- Ridlo, I. A. & Zein, R. A., 2018. Arah Kebijakan Kesehatan Mental: Tren Global dan Nasional Serta Tantangan Aktual. *Buletin Penelitian Kesehatan*, 1 March, 46(1), pp. 45-52.
- Satgas Penanganan Covid-19. 2020. *Peta Sebaran Covid-19*. [Online] Available at: <https://covid19.go.id/peta-sebaran-covid19> [Accessed 11 November 2020].
- Shoja, E., et al. 2020. Covid-19 effects on the workload and mental health of Iranian healthcare workers. *BMC Public Health*, pp. 1-16.
- Tella, M. D., et al. 2020. Mental health of healthcare workers during the COVID-19 pandemic in Italy. *Jorunal of Evaluation in Clinical Practice*, 28 June.pp. 1-5.
- Temet M. & Mc Michael. 2020. Covid-19 in a long-term care facility King County, Washington, February 27 - March 9, 2020. *US Department of Health and Human Services/Centers for Disease Control and Prevention*. 69(12), pp. 339-343.
- Temsah, M. H. et al. 2020. The psychological impact of COVID-19 pandemic on health care workers in a MERS-CoV endemic country. *Journal of Infection and Public Health*, 21 May, Volume 13, pp. 877-882.
- Walton, M., et al. 2020. Mental health care for medical staff and affiliated healthcare workers during the COVID-19 pandemic. *European Heart Journal: Acute Cardiovascular Care*, 8 April, 0(0), pp. 1-7.
- Wen, L., et al. 2020. progression of mental health services during the COVID-19 outbreak in China. *International Journal of Biological Sciences*. 16(10), pp. 1732-1738.

- Whaibeh, E. & Mahmoud, H. & N. H., 2020. Telemental Health in the Context of a Pandemic: the COVID-19 Experience. *Current Treatment Options in Psychiatry*, 7(2), pp. 198-202.
- Windarwati, H. D., et al. 2020. In the Middle of the COVID-19 Outbreak: Early Practical Guidelines for Psychosocial Aspects of COVID-19 in East Java, Indonesia. *Journal Pre-proof*, 17 August.
- Xiao, X., et al. 2020. Psychological impact of healthcare workers in China during COVID-19 pneumonia epidemic: A multi-center cross-sectional survey investigation. *Journal of Affective Disorders*, 19 May, Volume 274, pp. 405-410.
- Zhang, W. R. et al. 2020. Mental Health and Psychosocial Problems of Medical Health Workers during the COVID-19 Epidemic in China. *Psychotherapy Psychosomatics*, 9 April, pp. 1-9
- Zu, Z. Y. et al. 2020. Coronavirus Disease 2019 (COVID-19): A Perspective from China. In Press



# UNIVERSITAS AIRLANGGA

FAKULTAS KESEHATAN MASYARAKAT

Kampus C Mulyorejo Surabaya 601115 Telp.031-5920948, 5920949 Fax 031-5924618

Laman : <http://www.fkm.unair.ac.id> : E-mail: info@fkm.unair.ac.id

## SURAT KETERANGAN

Nomor : 3022/UN3.1.10/KP/2023

Yang bertanda tangan di bawah ini :

Nama : Dr. Santi Martini, dr. M.Kes  
NIP : 196609271997022001  
Pangkat/Golongan : Pembina/ Gol (IV/a)  
Jabatan : Dekan

Dengan ini menerangkan bahwa :

Nama : Inge Dhamanti, SKM. M.Kes, MPH., Ph.D  
NIP : 19801224 200501 2 002  
Pangkat/Golongan : Penata (Gol. III/c)  
Jabatan : Lektor

Telah melaksanakan penelitian dengan judul sebagai berikut :

No.	Judul Karya Ilmiah	Tahun Pelaksanaan Penelitian
1.	Implementation of Computerized Physician Order Entry in Primary Care: A Scoping Review	2021
2.	Adverse Reactions of COVID-19 Vaccines: A Scoping Review of Observational Studies	2023
3.	Literature Review: Cause Factor Analysis and an Effort to prevent Medication Administration Error (MAE) at Hospital	2020
4.	A Literature review on the Identification of Variables for Measuring Hospital Efficiency in the Data Envelopment Analysis (DEA)	2021
5.	Telemedicine Use In Health Facility During Covid-19 Pandemic: Literature Review	2022
6.	Faktor yang Menghambat Pelaporan Insiden Keselamatan Pasien di Rumah Sakit: Literature Review	2021
7.	Comparison of Four Methods To Detect Adverse Events in Hospital	2015
8.	Infections Prevention and Control (IPC) Programs in Hospitals	2021
9.	Studi Komparatif Pengembangan Contact Tracing Applications Di Singapura dan Indonesia (Studi Kasus: TraceTogether dan PeduliLindungi)	2022
10.	Faktor Penghambat Pelayanan Kesehatan Rutin di Rumah Sakit saat Pandemi COVID-19	2021
11.	Governmental Policies in Managing COVID-19 Pandemic: Comparative Study Between Indonesia and Vietnam, Period of January – March 2020	2021



12.	Akses Pelayanan Kesehatan Ibu dan Anak di Puskesmas Selama Pandemi Covid-19	2022
13.	Comparison of Dental Care Policies Before and During The COVID-19 Pandemic: A Literature Review	2022
14.	Analysis of Implementation of Patient Identification In Hospitals to Improve Patient Safety in Indonesia	2022
15.	Literature Review: Implementation Of Patient Safety Goals In Hospitals In Indonesia	2021
16.	Literature Review: Hospital Service Quality During The COVID-19 Pandemic	2022
17.	Comparison of Hospital Emergency Room Management Regulations in Indonesia Before and During The COVID-19 Pandemic: Literature Study	2022
18.	Analisis Pelaksanaan Pelayanan Kesehatan Perorangan (Ukp) Di Puskesmas Sebelum Dan Selama Pandemi Covid-19: Literature Review	2022
19.	Perbandingan Kebijakan Pelaksanaan Imunisasi Rutin pada Anak sebelum dan selama Pandemi	2022
20.	Recommendation Analysis Of Mental Health Services For Health Workers During Pandemic Covid-19	2021
21.	Impact Of Implementing A Surgical Safety Checklist In Hospital: Literature Review	2023
22.	Quality Improvement For Maternal And Child Health In Primary Health Care: A Scoping Review	2023
23.	Implementation Of Root Cause Analysis On Patient Safety Iincidence In Hospital: Literature Review	2022
24.	Analisis Peran Stakeholder dalam Kapasitas Rumah Sakit akibat COVID-19: Literature Review	2022
25.	Lessons from Indonesia, a country with highest COVID-19 mortality rate in the world: dissecting multiple aspects	2022

Adapun penelitian tersebut layak dilakukan dan menghasilkan output yang sangat baik, meskipun belum ada **Uji Etical Clearence** karena menggunakan metode litteratur review . Demikian surat keterangan ini kami buat untuk dapat dipergunakan sebagai persyaratan pengusulan Jabatan Fungsional Lektor Kepala.

Surabaya, 13 April 2023



Gana Martini, dr. M.Kes  
NIP. 196609271997022001