


[Skip to main content](#)

Advertisement



Search

- [Explore journals](#)
- [Get published](#)
- [About BMC](#)
- [Login](#)

Menu

- [Explore journals](#)
- [Get published](#)
- [About BMC](#)
- [Login](#)

## BMC Public Health

- [Home](#)
- [About](#)
- [Articles](#)
- [Submission Guidelines](#)
- [Press Releases](#)

Search articles within this journal

[Submit manuscript](#)

## Collections open to submissions



### [The COVID-19 pandemic and intimate partner violence](#)

Guest Edited by Jhumka Gupta, Lindsay Stark, Tiara Willie and Paul Bukuluki



**LG** **BTQ+ Health**

Guest Edited by Monica Malta and Wangnan Cao



**Air po** **llution and health**

Guest Edited by Hualiang Lin, Om Kurmi and Bing-Fang Hwang



**Food security and food systems**

Guest Edited by Sherly Hendriks, Hirotugu Aiga and Tarra Penney

[Click here for all journal collections](#)

**Editors' picks**



How issues surrounding food security, including shortages, disruption and stockpiling, were represented in media reporting on Brexit.



Testing three common correction formats finds that COVID-19 myths can be effectively corrected using materials and formats typical of health campaigns.

## Articles

- [Recent](#)
- [Most accessed](#)

### 1. [Post-traumatic stress disorder and mental health assessment of seafarers working on ocean-going vessels during the COVID-19 pandemic](#)

Authors: Fereshteh Baygi, Christine Blome, Andrew Smith, Nami Mohammadian Khonsari, Arash Agoushi, Arman Maghoul, Mohammad Esmacili-Abdar, Armita Mahdavi Gorabi and Mostafa Qorbani

Content type: Research

6 February 2022

### 2. [Development of an intervention for the social reintegration of adolescents and young adults affected by cancer](#)

Authors: Marie Broholm-Jørgensen, Tine Tjørnhøj-Thomsen and Pia Vivian Pedersen

Content type: Research article

5 February 2022

### 3. [Health-seeking behavior of COVID-19 cases during the first eight weeks of the outbreak in Singapore: differences between local community and imported cases and having visits to single or multiple healthcare providers](#)

Authors: Min Zhi Tay, Li Wei Ang, Wycliffe Enli Wei, Vernon J. M. Lee, Yee-Sin Leo and Matthias Paul H. S. Toh

Content type: Research article

5 February 2022

[Skip to main content](#)

Advertisement



Search

- [Explore journals](#)
- [Get published](#)
- [About BMC](#)
- [Login](#)

Menu

- [Explore journals](#)
- [Get published](#)
- [About BMC](#)
- [Login](#)

[BMC Public Health](#)

- [Home](#)
- [About](#)
- [Articles](#)
- [Submission Guidelines](#)
- [Press Releases](#)

[Articles](#)

- [Supplements](#)
- [Collections](#)
- [Reviewer acknowledgments](#)

## Articles

- [Search by keyword](#)
- [Search by citation](#)

Search BMC Public Health for results from

Show results from

### 2338 result(s) within Volume 21 of BMC Public Health

Page 1 of 47

Sort by

1. [Factors influencing variation in implementation outcomes of the redesigned community health fund in the Dodoma region of Tanzania: a mixed-methods study](#)

Micro-health insurance (MHI) has been identified as a possible interim solution to foster progress towards Universal Health Coverage (UHC) in low- and middle- income countries (LMICs). Still, MHI schemes suffe...

Authors: Albino Kalolo, Lara Gautier, Ralf Radermacher, Siddharth Srivastava, Menoris Meshack and Manuela De Allegri

Citation: *BMC Public Health* 2021 21:1

Content type: Research article

Content type: Research article

Published on: 4 January 2021

- > [View Full Text](#)
- > [View PDF](#)

#### 16. [Conceptualising changes to tobacco and alcohol policy as affecting a single interlinked system](#)

To support a move towards a coordinated non-communicable disease approach in public health policy, it is important to conceptualise changes to policy on tobacco and alcohol as affecting a single interlinked sy...

Authors: Duncan Gillespie, Jenny Hatchard, Hazel Squires, Anna Gilmore and Alan Brennan

Citation: *BMC Public Health* 2021 21:17

Content type: Research article

Published on: 4 January 2021

- > [View Full Text](#)
- > [View PDF](#)

#### 17. [Less than one-fifth of the mothers practised exclusive breastfeeding in the emerging regions of Ethiopia: a multilevel analysis of the 2016 Ethiopian demographic and health survey](#)

The burden of low coverage of exclusive breastfeeding (EBF) has a significant impact on the health of a newborn and also on the family and social economy in the long term. Even though the prevalence of EBF pra...

Authors: Tsegaye Gebremedhin, Demiss Mulatu Geberu and Asmamaw Atnafu

Citation: *BMC Public Health* 2021 21:18

Content type: Research article

Published on: 4 January 2021

- > [View Full Text](#)
- > [View PDF](#)

#### 18. [The potential role of local pharmacies to assess asthma control: an Italian cross-sectional study](#)

Asthma control and monitoring still represents a challenge worldwide. Although the international guidelines suggest the interplay between secondary and primary care services as an effective strategy to control...

Authors: M. Caminati, L. Cegolon, M. Bacchini, N. Segala, A. Dama, C. Bovo, B. Olivieri, F. Furci and G. Senna

Citation: *BMC Public Health* 2021 21:19

Content type: Research article

Published on: 5 January 2021

- > [View Full Text](#)
- > [View PDF](#)

#### 19. [Constructing a malaria-related health service readiness index and assessing its association with child malaria mortality: an analysis of the Burkina Faso 2014 SARA data](#)

The Service Availability and Readiness Assessment surveys generate data on the readiness of health facility services. We constructed a readiness index related to malaria services and determined the association...

Authors: Ouhouiré Millogo, Jean E. O. Doamba, Ali Sié, Jürg Utzinger and Penelope Vounatsou

Citation: *BMC Public Health* 2021 21:20

Content type: Research article

Published on: 5 January 2021

- > [View Full Text](#)
- > [View PDF](#)

#### 20. [Knowledge, attitude and perception of Pakistanis towards COVID-19; a large cross-sectional survey](#)

The Novel Coronavirus Disease (COVID-19) has created havoc globally as countries worldwide struggle to combat this pandemic. Since prevention and social isolation are known to be the only ways to prevent the s...

Authors: Zoya Fatima Rizwan Ladiwala, Rubaid Azhar Dhillon, Ibrahim Zahid, Omar Irfan, Muhammad Sharjeel Khan, Safia Awan and Javaid Ahmad Khan

Citation: *BMC Public Health* 2021 21:21

Content type: Research article

Published on: 5 January 2021

- > [View Full Text](#)
- > [View PDF](#)

21. [\*\*Temporal trends and regional disparities in cancer screening utilization: an observational Swiss claims-based study\*\*](#)

We examined colorectal, breast, and prostate cancer screening utilization in eligible populations within three data cross-sections, and identified factors potentially modifying cancer screening utilization in ...

Authors: Caroline Bähler, Beat Brüngger, Agne Ulyte, Matthias Schwenkglenks, Viktor von Wyl, Holger Dressel, Oliver Gruebner, Wenjia Wei and Eva Blozik

Citation: *BMC Public Health* 2021 21:23

Content type: Research article

Published on: 5 January 2021

- > [View Full Text](#)
- > [View PDF](#)

22. [\*\*Estimation of losses of quality-adjusted life expectancy attributed to the combination of cognitive impairment and multimorbidity among Chinese adults aged 45 years and older\*\*](#)

This study aims to estimate the losses of quality-adjusted life expectancy (QALE) due to the joint effects of cognitive impairment and multimorbidity, and to further confirm additional losses attributable to t...

Authors: Suting Xiong, Siyuan Liu, Yanan Qiao, Dingliu He, Chaofu Ke and Yueping Shen

Citation: *BMC Public Health* 2021 21:24

Content type: Research article

Published on: 5 January 2021

- > [View Full Text](#)
- > [View PDF](#)

23. [\*\*The potential role of network-oriented interventions for survivors of sexual and gender-based violence among asylum seekers in Belgium\*\*](#)

Social support and social network members have been identified as an important factor in mitigating the effects of sexual and gender-based violence (SGBV) and improving the coping process for many survivors. N...

Authors: Emilomo Ogbe, Alaa Jbour, Ladan Rahbari, Maya Unnithan and Olivier Degomme

Citation: *BMC Public Health* 2021 21:25

Content type: Research article

Published on: 5 January 2021

- > [View Full Text](#)
- > [View PDF](#)

24. [\*\*The effects of mother's education on achieving exclusive breastfeeding in Indonesia\*\*](#)

Even though the Indonesian government have set regulations for maintaining exclusive breastfeeding practices, the coverage remains low. The study aims to analyze the effects of mother's education level on the ...

Authors: Agung Dwi Laksono, Ratna Dwi Wulandari, Mursyidul Ibad and Ina Kusriani

Citation: *BMC Public Health* 2021 21:14

Content type: Research article


Published on: 6 January 2021

- > [View Full Text](#)
- > [View PDF](#)

[Skip to main content](#)

Advertisement

---



Search

- [Explore journals](#)
- [Get published](#)
- [About BMC](#)
- [Login](#)

Menu

- [Explore journals](#)
- [Get published](#)
- [About BMC](#)
- [Login](#)

[BMC Public Health](#)

- [Home](#)
- [About](#)
- [Articles](#)
- [Submission Guidelines](#)
- [Press Releases](#)

[About](#)

- [Contact](#)
- [Editorial board](#)
- [In Review](#)

## Editorial Board

**Natalie Pafitis, Senior Editor**

[Back to top](#)



Natalie joined BMC after having completed an MSc in Environmental Health at the Cyprus International Institute, in Association with the Harvard School of Public Health, USA where she completed courses including global climate change, environmental epidemiology, sustainable development and exposure assessment. Prior to this, she gained her BSc in Environmental Science at the University of Indianapolis, USA. Natalie is a Senior Editor for the BMC-series and has been Editor of *BMC Public Health* since 2007, also having worked on several biology and medical journals since this time. She has a keen interest in the use of research evidence to underpin public health policy and to advance the Sustainable Development Agenda. She is an Editor of the [BMC Series blog](#) and an ambassador of the Sustainable Development Goals for the BMC Series journals.

## Ana Donnelly, Editor

[Back to top](#)



Ana joined the BMC Series as a Manuscript Editor in 2018 and has been an Editor of *BMC Public Health* since 2021. Prior to joining BMC Series, Ana was a post-doctoral researcher with the School of Public Health at University College Cork and the National Suicide Research Foundation developing a program to improve mental health services in Ireland. Ana completed her Masters and PhD in Pharmacology with emphasis in Neuroscience at the Federal University of Santa Catarina (Brazil) exploring animal models of anxiety and depression. As a scientist from a developing country, Ana has also developed a special interest in social determinants of health and evidence based medicine.

### Senior Editorial Board Members

Christopher Ardern, *York University, Canada*  
 Noriko Cable, *University College London, UK*  
 Jennifer Emond, *Dartmouth College Geisel School of Medicine, USA*  
 Jianguang Ji, *Lund University, Sweden*  
 Youcheng Liu, *Wayne State University School of Medicine, USA*  
 Carol Maher, *University of South Australia, Australia*  
 Isabelle Niedhammer, *INSERM, France*  
 David Rehkopf, *Stanford University School of Medicine, USA*  
 Richard Rosenkranz, *Kansas State University, USA*  
 Jose Tuells, *University of Alicante, Spain*  
 Shankar Viswanathan, *Albert Einstein College of Medicine, USA*

### Editorial Board Members

#### *Biostatistics and methods*

Shakeel Ahmed, *Wah Medical College, Pakistan*



Dawit Getnet Ayele, *John Hopkins University, USA*  
Ashish Awasthi, *Public Health Foundation of India, India*  
Ram Bajpai, *Keele University, United Kingdom*  
Jeannette Beasley, *NYU School of Medicine, USA*  
Tzu-An Ann Chen, *University of Houston, USA*  
Ka Chun Chong, *The Chinese University of Hong Kong, Hong Kong*  
Patrizio Pezzotti, *Lazio Sanità - Agenzia di Sanità Pubblica, Italy*  
Ezra Gayawan, *Federal University of Technology, Akure, Nigeria*  
Jenine Harris, *Washington University in St. Louis, USA*  
Md. Golam Hossain, *University of Rajshahi, Bangladesh*  
Mehwish Hussain, *Dow University of Health Sciences, Pakistan*  
Kim Kiely, *School of Psychology, University of NSW, Australia*  
Ryung Kim, *Albert Einstein College of Medicine, USA*  
Peter Kolarčík, *Safarik University in Kosice, Slovakia*  
Karen Lamb, *University of Melbourne, Australia*  
Leah Li, *University College London, UK*  
Mahdi Mahdizadeh, *Hakim Sabzevari University, Iran*  
Samuel Manda, *South Africa Medical Research Council, South Africa*  
Justin Manjourides, *Northeastern University, USA*  
Haider Mannan, *Western Sydney University, Australia*  
Zindoga Mukandavire, *Coventry University, UK*  
Fredrik Norström, *Umeå University, Sweden*  
Jasmina Panovska-Griffiths, *University College London, UK*  
Mariamma Philip, *National Institute of Mental Health and NeuroSciences, India*  
Jinma Ren, *Pfizer Inc., USA*  
Marc Saez, *Universitat de Girona, Spain*  
Odile Sauzet, *Bielefeld University, Germany*  
Manik Sharma, *DAV University, India*  
Chuen Sang Tan, *National University of Singapore, Singapore*  
Tinku Thomas, *St John's Medical College, India*  
Fenicia Vescio, *Istituto Superiore di Sanità, Italy*  
Tzai-Hung Wen, *National Taiwan University, Taiwan*  
Darren Wraith, *Queensland University of Technology, Australia*  
Chengdong Xu, *Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, China*  
Paola Zaninotto, *University College London, UK*

### ***Chronic disease epidemiology***

Aderonke Akinkugbe, *Virginia Commonwealth University, USA*  
Arsham Alamian, *University of Miami, USA*  
Amir Almasi-Hashiani, *Arak University of Medical Sciences, Iran*  
Leopold Aminde, *School of Medicine, Griffith University, Australia*  
Christopher Ardern, *Bethune College, Canada*  
Hisatomi Arima, *Fukuoka University, Japan*  
Kate Bartlem, *University of Newcastle, Australia*  
Estifanos Baye, *Harvard Medical School, USA*  
Leonelo Bautista, *University of Wisconsin-Madison, United States*  
Danielle Berkovic, *Monash University, Australia*  
Boris Bikbov, *Istituto di Ricerche Farmacologiche Mario Negri IRCCS, Italy*  
Henrik Boggild, *Aalborg University, Denmark*  
Genny Carrillo, *Texas A&M School of Public Health, USA*  
Tawanda Chivese, *Qatar University, Qatar*  
Enayet Chowdhury, *Monash University, Australia*  
Lara Dugas, *Loyola University Chicago, USA*  
Jennifer Emond, *Geisel School of Medicine at Dartmouth, USA*  
Kaniz Fatema, *Centre for Health Equity Training, Research & Evaluation, Australia*  
Saruna Ghimire, *University of Miami, USA*  
Sameer Gopalani, *University of Oklahoma Health Sciences Center, USA*

Bhawna Gupta, *Griffith University, Australia*  
Hassan Haghparast Bidgoli, *University College London, UK*  
Chisato Hamashima, *Teikyo University, Japan*  
Gang Hu, *Pennington Biomedical Research Center, USA*  
Md Rafiqul Islam, *Goulburn Valley Health, Australia*  
Seow Wei Jie, *National University of Singapore, Singapore*  
Deanna Kepka, *University Of Utah, United States*  
Malcolm Koo, *Tzu Chi University of Science and Technology, Taiwan*  
Julia Kravchenko, *Duke University School of Medicine, USA*  
Yu Heng Kwan, *Duke-NUS Medical School, Singapore*  
Michael Laxy, *Technical University of Munich, Germany*  
Yo Han Lee, *Ajou University, South Korea*  
Yan Li, *Cleveland Clinic, USA*  
Gang Liu, *Huazhong University of Science and Technology, China*  
Maria Maynard, *Leeds Beckett University, UK*  
Christa Meisinger, *Ludwig-Maximilians-University Munchen, Chair of Epidemiology at UNKA-T Augsburg, Germany*  
Mustafa Mümtaz Mazicioglu, *Erciyes University, Turkey*  
Roberto Monastero, *University of Palermo, Italy*  
Keiko Murakami, *Tohoku University, Japan*  
Giesje Nefs, *Radboud University Medical Center, Netherlands*  
Leila Nikniaz, *Golgasht St. Tabriz University of Medical Sciences, Tabriz, Iran*  
Sisse Njor, *Randers Regional Hospital, Denmark*  
Victor Oguoma, *Health Research Institute, University of Canberra, Australia*  
Chen-Wei Pan, *Medical College of Soochow Univeristy, China*  
Feng Pan, *Menzies Institute for Medical Research, University of Tasmania, Australia*  
Ngoc Minh Pham, *Curtin University, Australia*  
Francesco Pistelli, *Pisa University Hospital, Italy*  
Muhammad Aziz Rahman, *Federation University, Australia*  
Venkataraghavan Ramamoorthy, *University of Central Missouri, USA*  
Zengliang Ruan, *Southeast University, China, People's Republic*  
Jennifer Salinas, *Texas Tech University Health Sciences Center El Paso, USA*  
János Sándor, *University of Debrecen, Hungary*  
Anshul Saxena, *Florida International University, USA*  
Kashif Shafique, *Dow University of Health Sciences, Karachi, Pakistan*  
David Shoham, *East Tennessee State University, USA*  
Anabela Silva, *University of Aveiro, Portugal*  
Fuschia M. Sirois, *University of Sheffield, UK*  
Sherri Stewart, *Centers for Disease Control and Prevention, USA*  
Saverio Stranges, *Western University, Canada*  
Jing Tian, *University of Tasmania, Australia*  
JS Thakur, *Ghent University, Belgium*  
Stephen Ojiambo Wandera, *Makerere University, Uganda*  
Peizhong Peter Wang, *Memorial University of Newfoundland, Canada*  
ShaoMing Wang, *Peking Union Medical College, China*  
Youxin Wang, *Capital Medical University, China*  
Yuan-Pang Wang, *University of São Paulo, Brazil*  
James White, *Cardiff University, UK*  
Tsu-Yin Wu, *Eastern Michigan University, USA*  
Fei Xu, *Nanjing Municipal Center for Disease Control and Prevention, China*  
Chin Kok Yong, *Universiti Kebangsaan, Malaysia*  
Xiaotao Zhang, *Baylor College of Medicine, USA*  
Melissa Zullo, *Kent State University, USA*

### ***Dental public health***

Juliana Balbinot Hilgert, *Universidade Federal do Rio Grande do Sul, Brazil*  
Ratilal Laloo, *University of Queensland, Australia*  
Ajesh George, *Western Sydney University, Australia*

RESEARCH ARTICLE

Open Access



# The effects of mother's education on achieving exclusive breastfeeding in Indonesia

Agung Dwi Laksono<sup>1,2</sup>, Ratna Dwi Wulandari<sup>3\*</sup> , Mursyidul Ibad<sup>4</sup> and Ina Kusri<sup>5</sup>

## Abstract

**Background:** Even though the Indonesian government have set regulations for maintaining exclusive breastfeeding practices, the coverage remains low. The study aims to analyze the effects of mother's education level on the coverage of exclusive breastfeeding in Indonesia.

**Methods:** This study used data from the 2017 Nutrition Status Monitoring Survey. It covered data of 53,528 children under 5 years old (7–59 months) as the samples. Variables included exclusive breastfeeding status, mother's education level, mother's age, marital status, employment status, gender, residence, under five's age and gender. A binary logistics regression was performed in the final test.

**Results:** Mothers who graduated from elementary school were 1.167 times more likely to perform exclusive breastfeeding compared to mothers who never attended schools. Additionally, those who graduated from junior high school had 1.203 times possibilities to give exclusive breastfeeding compared to mothers without educational records. While, mothers who graduated from high school were 1.177 times more likely to perform exclusive breastfeeding compared to those without educational records. Mothers who graduated from tertiary education had 1.203 times more possibilities to perform exclusive breastfeeding compared to mothers who were never enrolled to schools. Other variables also became affecting predictors on exclusive breastfeeding, such as mother's age, mother's employment status, child's age, and residence.

**Conclusions:** The mother's education level positively affects exclusive breastfeeding practice in Indonesia.

**Keywords:** Breastfeeding, Exclusive breastfeeding, Education level, Nutrition education, Health education

## Background

Breastfeeding is a common method of providing breast milk as food supply for infants and young children. It is the cheapest and simplest method to meet the baby's nutritional needs. Breast milk will improve sensory and cognitive abilities and protect children from infectious and chronic diseases. Poor infant feeding practices may impact on children's growth and development [1, 2]. A meta-analysis study used 17 previous studies that

discover an increase in IQ scores of 3.44 points in breastfed children. Meanwhile, in the meta-regression, none of the studies explained the heterogeneity [3].

Exclusive breastfeeding (EBF) is a method of giving breast milk merely for infants to provide complete nutrition in the first 6 months of life [4]. Exclusive breastfeeding has many benefits for mothers and babies. For babies, EBF can reduce infant mortality due to common infectious diseases, help recovery, and increase children immunity. Regarding its benefits for mothers, EBF is a safe feeding method that can protect mothers from the risk of ovarian and breast cancer and reduce obesity [5–7].

\* Correspondence: [ratna-d-w@fkm.unair.ac.id](mailto:ratna-d-w@fkm.unair.ac.id)

<sup>3</sup>Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia  
Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Although EBF has been proven to have a positive effect [3, 8, 9], the coverage of EBF is still very low. Globally, the coverage of EBF was 30–50% [1], but was around 35.7% in Indonesia [10]. Previous studies have identified intrinsic and extrinsic factors that affect EBF. Despite having regulations to maintain EBF practices (Law Number 36/2009 concerning Health), the coverage of EBF in Indonesia remains low, especially for uneducated and employed mothers [11].

The low coverage has something to do with traditional breastfeeding practices in native Indonesian tribes. Some native tribes introduce food or drinks to babies who are recently a few days old. The Gayo, Javanese, and Muyu tribes commonly feed their babies honey, sugar water, and sago solution [12–15].

Based on the background, this study aimed to analyze the effects of mother's education on exclusive breastfeeding practice in Indonesia. The results of this study vitally become basic references for policymakers to formulate policies in improving the coverage of EBF in Indonesia.

## Methods

### Data source

This study utilized raw data of the 2017 Nutrition Status Monitoring Survey. This survey was a national survey using a multi-stage cluster random sampling method and was conducted by the Directorate of Community Nutrition of the Indonesian Ministry of Health [10]. The survey population was all children under 5 years old or 7–59 months in Indonesia, and there were 53,528 babies as the samples.

### Data analysis

Mother's education level was to which degree a mother has ever completed an education. It has five categories, such as not attending school, primary school, junior high school, senior high school, and tertiary education. Another variable was mother's employment status, which defines a type of mother's work or employment.

The dependent variable in this study was exclusive breastfeeding practice. Seven independent variables were divided into 3 groups. The first variable was the characteristics of mothers (education level, age, marital status, employment status). The second variable was the characteristics of the children under five (age and gender). Finally, the third was a residence (urban-rural).

The Chi-Square test was utilized to test the dichotomous variables, while continuous variables were tested using the T-test. This statistical test assessed whether there was a statistically significant relationship between mother's education level variable and another variable. Besides, a binary logistic regression test was performed at the final stage to identify disparities in the

contribution of mother's education levels to EBF practices in Indonesia. All data analyses were performed in SPSS 22 version software.

## Results

Figure 1 shows the distribution of breastfed children under five by Indonesian provinces. It indicates the largest coverage of EBF in Java.

### Descriptive statistics

Prior to a binary logistic regression test, a co-linearity test was carried out. The results of co-linearity test show that there was no co-linearity between dependent and independent variables. Table 1 shows that the tolerance value of all variables was greater than 0.10, and the VIF value for all variables was less than 10.00. Based on the result of multicollinearity test, there were no symptoms of multicollinearity in the regression model.

Table 2 explains the statistical description of mother's education level. The table points out mothers in all categories of education level mostly did not breastfeed their children. It means the practice of EBF in all education categories was relatively low.

In Table 2, mothers who had primary education record are slightly older than mothers with other education levels. The majority of the mothers were married, but most mothers in four education levels were unemployed. Only mothers with tertiary education record were dominantly employed (65.52%).


Table 2 informs that there was a significant relationship between education levels and babies' age. Based on babies' gender, male babies were dominant in all education categories. However, the test results stated that the relationship between mother's education level and babies' gender was not statistically significant. This study also reveals most mothers lived in rural areas. This variable was proven statistically significant to affect the practice of EBF.

### Multivariate regression analysis

Table 3 shows the results of the binary logistic regression test that used "No EBF" as a reference.

It informs that all three groups of variables significantly affected the practice of EBF. Those affecting variables were mother's age, mother's education level, mother's employment status, babies' age, and residence.

This study finds mothers who graduated from primary school were 1.167 times more likely to perform EBF compared to mothers without education records (OR 1.167; 95% CI 1.026–1.328). Those who graduated from junior high school had 1.203 times possibilities to perform EBF compared to those who never attended schools (OR 1.203; 95% CI 1.057–1.368). Mothers who are high school graduates were 1.177 times more likely



to perform EBF than mothers without formal education experience (OR 1.177; 95% CI 1.037–1.335). Additionally, mothers with tertiary education record had 1.203 times possibilities to give EBF than mothers without formal education record (OR 1.231; 95% CI 1.073–1.411).

Children under five whose mothers were employed had 0.954 times more likely to be breastfed those whose mothers were unemployed (OR 0.954; 95% CI 0.913–0.997). Table 3 also shows that children under five in urban areas had 1.138 times possibilities to perform EBF than those in rural areas (OR 1.138; 95% CI 1.092–1.186).

**Table 1** Results for a co-linearity test of EBF in Indonesia (n = 53,528)

Variables	Collinearity Statistics	
	Tolerance	VIF
Mother's Characteristics		
Education level	0.958	1.044
Age (in years; mean)	0.978	1.022
Marital status	0.997	1.003
Employment Status	0.980	1.020
Under five's Characteristic		
Under five's Age (in months; mean)	0.997	1.003
Under five's Gender	1.000	1.000
Residence	0.969	1.032

\*Dependent Variable: EBF status

### Discussion

This study discovers that mother's education affected the practice of EBF in Indonesia. A better education tends to give mothers more possibilities of EBF. This finding is in line with findings of other studies [16, 17]. The surveys to postnatal mothers in Nigeria and China show better education positively contributed to the breastfeeding process and the rate level of EBF [16, 17]. Some studies in America added self-efficacy as a variable to find the correlation of maternal education and the practice of EBF. Higher education levels were correlated with better self-efficacy scores. Mother's education level had a positive relationship with the practice of EBF [8, 9].

Similarly, another study in European multiregions finds that mothers who were younger and less educated were more likely to stop breastfeeding before their babies were aged 6 months. In other words, they did not perform complete EBF. Education level, parity, and socioeconomic factors could indicate whether mothers breastfeed or not their babies [18]. Some studies in Chile also found psychosocial factors, such as maternal IQ and low-risk prenatal behavior at birth, could affect the duration of breastfeeding [19, 20]. Another study conducted in Eastern Indonesia, which analyzed data of the 2012 Indonesian Family Life Survey, concluded the same thing. This study took a smaller sample size (1138 under five) and found that mother's education levels had a positive effect on the success rates of EBF [21].

Moreover, this study uncovers unemployed mothers had better EBF practice. This result is similar to the

**Table 2** Descriptive Statistics of Mother's Education Level and Related Variables (n = 53,528)

Variables	Mother's Education level					All	P
	No School	Primary School	Junior High School	Senior High School	College		
EBF status							0.013*
• EBF	367 (29.84%)	4063 (34.12%)	4066 (34.49%)	8180 (34.15%)	1632 (35.13%)	18,308 (34.20%)	
• Not EBF (Ref.)	863 (70.16%)	7846 (65.88%)	7723 (65.51%)	15,774 (65.85%)	3014 (64.87%)	35,220 (65.80%)	
Mother's Characteristics							
Age (in years; mean)	1230 (28.48)	11,909 (31.07)	11,789 (29.07)	23,594 (28.50)	4646 (30.06)	53,528 (29.33)	≤ 0.001*
Marital status							≤ 0.001*
• Single (Ref.)	5 (0.41%)	53 (0.45%)	46 (0.39%)	82 (0.34%)	18 (0.39%)	204 (0.38%)	
• Married	1208 (98.21%)	11,719 (98.40%)	11,622 (98.58%)	23,731 (99.07%)	4606 (99.14%)	52,886 (98.80%)	
• Divorce	17 (1.38%)	137 (1.16%)	121 (1.03%)	141 (0.59%)	22 (0.47%)	438 (0.82%)	
Employment Status							≤ 0.001*
• Unemployed (Ref.)	617 (50.16%)	9077 (76.22%)	9323 (79.08%)	19,867 (82.94%)	1602 (34.48%)	40,486 (75.64%)	
• Employed	613 (49.84%)	2832 (23.78%)	2466 (20.92%)	4087 (17.06%)	3044 (65.52%)	13,042 (24.36%)	
Under five's Characteristic							
Age (in months; mean)	1230 (14.66)	11,909 (14.81)	11,789 (14.76)	23,594 (14.53)	4646 (14.69)	53,528 (14.66)	≤ 0.001*
Gender							0.605
• Male	641 (52.11%)	5987 (50.27%)	5993 (50.84%)	12,180 (50.85%)	2367 (50.95%)	27,168 (50.75%)	
• Female (Ref.)	589 (47.89%)	5922 (49.73%)	5793 (49.14%)	11,771 (49.14%)	2279 (49.05%)	26,354 (49.23%)	
Residence							≤ 0.001*
• Urban	167 (13.58%)	1989 (16.70%)	2392 (20.29%)	7559 (31.56%)	1583 (34.07%)	13,690 (25.58%)	
• Rural (Ref.)	1063 (86.42%)	9920 (83.30%)	9397 (79.71%)	16,395 (68.44%)	3063 (65.93%)	39,838 (74.42%)	

Note: Chi-Square used for dichotomous variables and the T-test used for continuous variables. \*Significant at the 95% level

**Table 3** Results of Binary Logistic Regression (n = 53,528)

Predictors	Sig.	EBF		
		OR	Lower Bound	Upper Bound
Mother's Characteristics				
Education level: no school	–	–	–	–
Education level: primary school	0.019 <sup>a</sup>	1.167	1.026	1.328
Education level: junior high school	0.005 <sup>a</sup>	1.203	1.057	1.368
Education level: senior high school	0.011 <sup>a</sup>	1.177	1.037	1.335
Education level: college	0.003 <sup>a</sup>	1.231	1.073	1.411
Age	≤ 0.001 <sup>a</sup>	1.009	1.006	1.012
Marital Status: single	–	–	–	–
Marital Status: married	0.561	1.092	0.812	1.468
Marital Status: divorced	0.817	0.959	0.670	1.371
Employment status: unemployed	–	–	–	–
Employment status: employed	0.038 <sup>a</sup>	0.954	0.913	0.997
Toddler's Characteristic				
Age	≤ 0.001 <sup>a</sup>	1.022	1.018	1.026
Residence				
Area: Urban	≤ 0.001 <sup>a</sup>	1.138	1.092	1.186
Area: Rural	–	–	–	–

Note: The reference EBF status category was "Not EBF"; confidence interval of 95% for OR; <sup>a</sup>significant at 95% level

results of other studies indicating employed mothers were less likely to perform EBF. They tend to have less time and opportunity to interact with children, including giving breastfeed to their children [21, 22]. Studies in Qatar and Ethiopia find that the employment status was one of the barriers to the success of EBF. Plausibly mothers must return to work immediately after the maternity leave period runs out [23, 24]. Having flexible work schedules and workplace proximity to home can assist the sustainability of breastfeeding [25].

Another study carried out in Vietnam indicates that teamwork parenting could increase the success of EBF. In this study, husband's role was also observed to be one of the predictors of EBF success [26]. Supporting this fact, RCT research in Canada finds collaboration between parents influenced knowledge, perceptions, practices, and duration of EBF [27]. In India, social support became one of the determinants of EBF success [1].

This present study, furthermore, shows that children under five in urban areas had a better EBF intake. This is probably due to better information exposure about the benefits of EBF in urban areas than in rural areas. The effect of information exposure that may improve mother's knowledge about EBF was proven as one determinant of EBF success by several studies [17, 21, 28, 29]. In Indonesia, urban areas tend to have better access to health services and information than rural areas, and this leads to higher coverage of health programs [30, 31].

To end this discussion, this study also suggests the government to formulate policies focusing on clear targets to be achieved by referring to these study findings. It can target mothers who have poor education, are employed, and live in rural areas. Policies focusing on these targets need to be implemented for a wider coverage of EBF.

### Limitation

The study utilized big data of the 2017 IDHS survey. The data are not presenting facts deeply so that the findings in this study have not captured different phenomena in the number of indigenous Indonesians, who have their local wisdom of breastfeeding practices. Mother's cultural background can be an adequate differentiator [12, 14, 32]. Besides, the study also did not involve the participation of parents, parents-in-law, and family to indicate the practice of EBF. While, previous studies find this variable could affect EBF [32, 33]. It is necessary to conduct further qualitative studies to capture or deepen some phenomena in question.

### Conclusions

It could be concluded that there was a positive effect of mother's education on the practice of EBF in Indonesia. Other variables, such as mother's age, mother's

employment status, babies' age, and residence were some predictors of EBF success rates in Indonesia.

The government needs to issue policies that focus on specific targets as identified in this study. It can target mothers who have poor education, are employed, and live in rural areas to expand the coverage of EBF in Indonesia.

### Acknowledgments

The author would like to thank the Directorate of Community Nutrition of the Ministry of Health of the Republic of Indonesia for allowing to processing of data of the 2017 Indonesia Nutritional Status Monitoring.

### Authors' contributions

ADL developed the proposal, analyzed, and interpreted the patient data. RDW, MI, and IK was a major contributor in conducting the study, interpreting the data, and writing the manuscript. The authors read and approved the final manuscript.

### Funding

Not applicable.

### Availability of data and materials

The 2017 Nutrition Status Monitoring Survey data used to support these findings of this study were supplied by the Directorate of Community Nutrition of the Indonesian Ministry of Health under license and so can not be made freely available. Requests for access to these data should be made to the Directorate of Community Nutrition of the Indonesian Ministry of Health.

### Ethics approval and consent to participate

The Nutrition Status Monitoring Survey in 2017 has an ethical license approved by the Health Research Ethics Commission at the Indonesian Ministry of Health (ethics number: LB.02.01/2/KE.244/2017). In this survey, informed consent was used during data collection, which considers aspects of procedures for data collection, voluntary, and confidentiality. Respondents gave written consent.

### Consent for publication

Not applicable.

### Competing interests

The authors declare that they have no competing interests.

### Author details

<sup>1</sup>National Institute of Health Research and Development, the Ministry of Health of the Republic of Indonesia, Jakarta, Indonesia. <sup>2</sup>Doctoral Program, Faculty of Public Health, University of Airlangga, Surabaya, Indonesia. <sup>3</sup>Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia. <sup>4</sup>Faculty of Health, Nadlatul Ulama University, Surabaya, Indonesia. <sup>5</sup>Unit of Health Research and Development Magelang, Ministry of Health, Center Java, Java, Indonesia.

Received: 1 July 2020 Accepted: 8 December 2020

Published online: 06 January 2021

### References

- Pareek S. Exclusive breastfeeding in India: an ultimate need of infants. *Nurs Pract Today*. 2019;6(1):4–6.
- Binda V, Figueroa-Leigh F, Olhaberry M. Low quality of mother-child interaction in infants at psychosocial risk is associated with risk of developmental delay. *Rev Chil Pediatr*. 2019;90(3):260–6.
- Horta BL, de Mola CL, Victora CG. Breastfeeding and intelligence: a systematic review and meta-analysis. *Acta Paediatr*. 2015;104(467):14–9.
- Charlick SJ, McKellar L, Gordon AL, Pincombe J. The private journey: an interpretative phenomenological analysis of exclusive breastfeeding. *Women Birth*. 2019;32(1):e34–42.
- Arikawa S, Rollins N, Jourdain G, Humphrey J, Kourtis AP, Hoffman I, et al. Contribution of maternal antiretroviral therapy and breastfeeding to 24-

- month survival in human immunodeficiency virus-exposed uninfected children: an individual pooled analysis of African and Asian studies. *Clin Infect Dis*. 2018;66(11):1668–77.
6. MacVicar S, Humphrey T, Forbes-McKay KE. Breastfeeding and the substance-exposed mother and baby. *Birth*. 2018;45(4):450–8.
  7. Fan HSL, Wong JYH, Fong DYT, Lok KYW, Tarrant M. Breastfeeding outcomes among early-term and full-term infants. *Midwifery*. 2019;71:71–6.
  8. Glassman ME, McKearney K, Saslaw M, Sirota DR. Impact of breastfeeding self-efficacy and sociocultural factors on early breastfeeding in an urban, predominantly Dominican community. *Breastfeed Med*. 2014;9(6):301–7.
  9. Bahorski JS, Childs GD, Loan LA, Azuero A, Morrison SA, Chandler-Laney PC, et al. Self-efficacy, infant feeding practices, and infant weight gain: an integrative review. *J Child Health Care*. 2019;23(2):286–310.
  10. Directorate of Community Nutrition of The Ministry of Health of The Republic of Indonesia. The 2017 Indonesia Nutritional Status Monitoring (Pemantauan Status Gizi 2017) [Internet]. Jakarta; 2017. Available from: [http://www.kesmas.kemkes.go.id/assets/upload/dir\\_519d41d8cd98f00/files/Buku-Saku-Nasional-PSG-2017\\_975.pdf](http://www.kesmas.kemkes.go.id/assets/upload/dir_519d41d8cd98f00/files/Buku-Saku-Nasional-PSG-2017_975.pdf).
  11. Ratnasari D, Paramashanti BA, Hadi H, Yugistyowati A, Astiti D, Nurhayati E. Family support and exclusive breastfeeding among Yogyakarta mothers in employment. *Asia Pac J Clin Nutr*. 2017;26:531–5.
  12. Pratiwi NL, Fitrianti Y, Nuraini S, Rachmawati T, Laksono AD, Afreni M, et al. Concealed pregnant women or Kemel of Gayo ethnic in Blang Pegayon District, Gayo Lues District, Aceh. *Bull Heal Syst Res*. 2019;22(2):81–90.
  13. Laksono AD, Faizin K, Raunsay EM, Soerachman R. Muyu women in exile (Perempuan Muyu dalam Pengasingan) [internet]. Jakarta: Lembaga Penerbitan Balitbangkes; 2014. Available from: <https://www.scribd.com/doc/261673624/Perempuan-Muyu-dalam-Pengasingan-Riset-Etnografi-Kesehatan-2014-Boven-Digoel>.
  14. Laksono AD, Soerachman R, Angkasawati TJ. Case study of Muyu Ethnic's maternal health in Mindiptara District-Boven Digoel (Studi Kasus Kesehatan maternal Suku Muyu di Distrik Mindiptana, Kabupaten Boven Digoel). *J Reprod Heal*. 2016;07/03:145–55.
  15. Maghfiroh MS, Laksono AD. Given sugar water ... "at first the cry became silent, because it was full, not limp, its endurance increased"; study of patterns of infant intake ("Diberi air gula ... awalnya nangis menjadi diam, karena kenyang, gak lemas, daya tahan tubuhnya meningkat"). *S Amerta Nutr*. 2020;4(2):116–22.
  16. Agho KE, Ogeleka P, Ogbo FA, Ezeh OK, Eastwood J, Page A. Trends and predictors of prelacteal feeding practices in Nigeria (2003–2013). *Nutrients*. 2016;8(8):462.
  17. Hamze L, Mao J, Reifsnider E. Knowledge and attitudes towards breastfeeding practices: a cross-sectional survey of postnatal mothers in China. *Midwifery*. 2019;74:68–75.
  18. Bonnet C, Blondel B, Piedvache A, Wilson E, Bonamy A-KE, Gortner L, et al. Low breastfeeding continuation to 6 months for very preterm infants: a European multiregional cohort study. *Matern Child Nutr*. 2019;15(1):e12657.
  19. Farkas C, Girard L-C. Breastfeeding initiation and duration in Chile: understanding the social and health determinants. *J Epidemiol Community Health*. 2019;73(7):637.
  20. Girard L-C, Farkas C. Breastfeeding and behavioural problems: propensity score matching with a national cohort of infants in Chile. *BMJ Open*. 2019; 9(2):e025058.
  21. Bue MCL, Priebe J. Revisiting the socioeconomic determinants of exclusive breastfeeding practices: evidence from eastern Indonesia. *Oxford Dev Stud*. 2018;46(3):398–410.
  22. Laksono AD, Ibad M, Mursita A, Kusriani I, Wulandari RD. Characteristics of mother as predictors of stunting in toddler. *Pak J Nutr*. 2019;18(12):1101–6.
  23. Hendaus MA, Alhammadi AH, Khan S, Osman S, Hamad A. Breastfeeding rates and barriers: a report from the state of Qatar. *Int J Womens Health*. 2018;10:467–75.
  24. Tadesse F, Alemayehu Y, Shine S, Asresahegn H, Tadesse T. Exclusive breastfeeding and maternal employment among mothers of infants from three to five months old in the Fafan zone, Somali regional state of Ethiopia: a comparative cross-sectional study. *BMC Public Health*. 2019;19(1): 1015.
  25. Chen J, Xin T, Gaoshan J, Li Q, Zou K, Tan S, et al. The association between work related factors and breastfeeding practices among Chinese working mothers: a mixed-method approach. *Int Breastfeed J*. 2019;14(1):28.
  26. Rempel JK, Rempel LA, Hoa DTP, Vui LT, Long TK. Parenting teamwork: the impact of a fathering intervention on mothers and infants in Vietnam. *Child Dev*. 2019;91(2):e345–e364. <https://doi.org/10.1111/cdev.13244>.
  27. Abbass-Dick J, Stern SB, Nelson LE, Watson W, Dennis C-L. Coparenting breastfeeding support and exclusive breastfeeding: a randomized controlled trial. *Pediatrics*. 2015;135(1):102–10.
  28. Buss IM. Knowledge, attitudes and behaviours towards exclusive breastfeeding amongst mothers in Sarawak, Malaysia: a qualitative study. *Int Med J Malaysia*. 2019;18(1):45–54.
  29. Wulandari RD, Laksono AD. Does the place of residence affect the achievement of exclusive breastfeeding? A study in eastern Indonesia. *Syst Rev Pharm*. 2020;11(9):872–6.
  30. Laksono AD, Wulandari RD, Soedirham O. Urban and rural disparities in hospital utilization among Indonesian adults. *Iran J Public Health*. 2019;48(2): 247–55.
  31. Laksono AD, Wulandari RD. Urban-rural disparities of facility-based childbirth in Indonesia. In: 4th international symposium on Health Research (ISHR 2019) [internet]. Denpasar: Atlantis Press; 2020. p. 33–9. Available from: <https://www.atlantis-press.com/proceedings/ishr-19/125935017>.
  32. Sutan R, Berkat S. Does cultural practice affects neonatal survival- a case control study among low birth weight babies in Aceh Province, Indonesia. *BMC Pregnancy Childbirth*. 2014;14(1):1–13.
  33. Alianmoghaddam N, Phibbs S, Benn C. The impact of family culture on six months exclusive breastfeeding: a qualitative study in New Zealand. *Breastfeed Rev*. 2018;26(1):23–36.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

