

Chlorogenic Acid Fractionation in Robusta Green Bean Extract as a Combination Agent of Dental Pulp Stem Cells in Periodontal Tissue Engineering

by Retno Widyowati

Submission date: 14-Apr-2023 03:54AM (UTC+0800)

Submission ID: 2063761304

File name: C-18_artikel.pdf (300.71K)

Word count: 4171

Character count: 22005

RESEARCH ARTICLE

Chlorogenic Acid Fractionation in Robusta Green Bean Extract as a Combination Agent of Dental Pulp Stem Cells in Periodontal Tissue Engineering

Desi Sandra Sari¹, Neira Sakinah¹, Nuri², Enny Suswati³, Retno Widayati⁴, Ernie Maduratna⁵

¹Department of Periodontic, Faculty of Dentistry, University of Jember, Jember, Indonesia.

²Department of Pharmaceutical Biology, Faculty of Pharmacy, University of Jember, Jember, Indonesia.

³Department of Microbiology, Faculty of Medicine, University of Jember, Jember, Indonesia.

⁴Department Fitokimia & Farmakognosi, Faculty of Pharmacy, University of Airlangga, Surabaya, Indonesia.

⁵Department of Periodontic, Faculty of Dental Medicine, University of Airlangga, Surabaya, Indonesia.

*Corresponding Author E-mail: desi_sari.fkg@unej.ac.id

ABSTRACT:

Background: Robusta coffee beans contain very high chlorogenic acid. In the last decade, chlorogenic acid was developed as an adjunct in stem cells to enhance the anti-inflammatory and antioxidant properties of stem cells when used as a therapeutic agent. Chlorogenic acid can increase proliferation and migration and inhibit the production of pro-inflammatory cytokines in stem cells thereby increasing the ability of stem cells to regenerate tissue. **Purpose:** To analyze the levels of chlorogenic acid in robusta coffee bean extract which can be used as a combination agent for Dental Pulp Stem Cells (DPSC) in periodontal tissue engineering therapy. **Materials and Methods:** Robusta coffee bean extract was obtained from the Coffee and Cocoa Research Center, Jember Regency which was processed and processed using the fractionation method. This study used DPSC with the extraction of premolar teeth of orthodontic patients. The toxicity test was performed on the coffee extract 0.0625%; 0.125%; 0.25%; 0.5% to determine the biocompatible concentration of DPSC. **Results:** The highest fractionation and measurement of chlorogenic acid content obtained was 30.49%. Robusta coffee bean extract with a chlorogenic acid content of 30.49% at concentrations of 0.125% and 0.0625% did not cause toxicity to DPSC. **Conclusion:** Chlorogenic acid in robusta coffee bean extract concentrations of 0.125% and 0.0625% are relatively biocompatible as a combination agent for DPSC in periodontal tissue engineering therapy.

KEYWORDS: Robusta coffee beans, DPSC, Periodontal engineering, Periodontitis, Regeneration.

INTRODUCTION:

Robusta coffee beans contain several anti-inflammatory and antioxidant compounds whose benefits have been widely studied, including caffeine, caffeic acid, chlorogenic acid, and trigonelline. These chemical components are a good source of anti-inflammatory and antioxidants for the body.¹ However, of these compounds, chlorogenic acid is the highest content of polyphenolic compounds in coffee beans compared to other compounds.²⁻⁴

Chlorogenic acid is well known for its various biological properties such as anti-inflammatory, antioxidant, antimicrobial, antifungal, and antiviral activities.^{5,6} In addition, several studies have revealed that chlorogenic acid is an effective anti-inflammatory, antipyretic, and analgesic agent in both in vitro and in vivo experiments.⁷⁻¹⁰ The highest content of chlorogenic acid can be found in robusta coffee beans, which is 6.1-11.3%. This amount is the largest content compared to other coffee beans.³

In addition to the high anti-inflammatory content in robusta coffee beans, robusta coffee is the main and largest commodity in Indonesia. The production and export value of Indonesia's robusta coffee globally ranks second after Vietnam. High exports in several countries

Received on 05.11.2021

Modified on 04.02.2022

Accepted on 13.03.2022

© RJPT All right reserved

Research J. Pharm. and Tech 2022; 15(11):5005-5010.

DOI: 10.52711/0974-360X.2022.00841

make coffee a relatively large foreign exchange contributor. Its availability is very abundant in Indonesia, making this type of coffee easy to obtain.¹¹ However, its use in the medical field is still not known.

Several researchers have proven the potential of chlorogenic acid as an additive in stem cells to increase the anti-inflammatory and antioxidant power of stem cells when used as a therapeutic agent. One of them is chlorogenic acid which can increase proliferation and migration and inhibit the production of pro-inflammatory cytokines in stem cells, thereby increasing the ability of stem cells to regenerate tissue.^{9,12,13}

In the periodontal field, stem cells are used as materials for tissue engineering therapy. In vivo studies reported that Dental Pulp Stem Cells (DPSC) have the potential to repair periodontal bone defects by providing optimal results of bone repair, cementum formation, and periodontal ligament around the bone defect. DPSC is reported to be effective in increasing bone regeneration in the treatment of socket preservation and intrabony defects.^{14,15} However, the presence of bacterial toxins, especially LPS *P. gingivalis* in the periodontal tissue can affect the success of DPSC in regenerating periodontal tissue. Several studies reported that *P. gingivalis* LPS can stimulate the secretion of proinflammatory cytokines such as IL-1 β and TNF- α in excess, thereby inhibiting the proliferation and differentiation of osteogenic stem cells. This has an impact on delaying treatment success.^{16,17}

Besides the advantages of stem cells as a regenerative therapy for periodontal tissue, this therapy must also be able to control the possibility of inflammation that occurs and prevent disease progression. Because basically, stem cell therapy will be used in patients with periodontitis, which is a chronic inflammatory disease. Therefore, the administration of chlorogenic acid to increase the anti-inflammatory and antioxidant power of DPSC needs to be explored further so that it can increase the success of periodontal regeneration treatment.^{18,19}

The utilization of robusta coffee bean extract to increase the anti-inflammatory and antioxidant power of DPSC itself is still not well understood until now. Therefore, this study aims to analyze the optimal levels of chlorogenic acid and the concentration of robusta coffee bean extract to be used as a combination of stem cells in periodontal tissue engineering therapy.

MATERIALS AND METHODS:

Materials:

Robusta coffee bean extract was obtained from the Coffee and Cocoa Research Center, Jember Regency

which was processed and processed using the fractionation method so that it had a chlorogenic acid content of 30.49%. The processed coffee bean extract was then made at a concentration of 0.5%; 0.25%; 0.125%; 0.0625%. Dental pulp stem cells (DPSCs) were obtained from the pulp of the 1st or 2nd premolars that were extracted due to indications for orthodontic treatment. The criteria for teeth that can be used are healthy teeth, no caries, and perfectly closed tooth roots in young patient donors aged between 19-29 years, healthy, free from infectious and infectious diseases.

Extraction and Isolation of Chlorogenic Acid in Robusta Coffee Beans:

Extraction and isolation of chlorogenic acid were carried out using the fractionation method. Fractionation was carried out using the elution liquid in a suitable TLC as the mobile phase and silica gel as the stationary phase. A total of 10 mg of robusta coffee bean extract was put into a glass column that already contained silica gel. The elution liquid was added in a gradient using n-hexane : ethyl acetate (10:1 - 1:1) followed by chloroform : methanol (5:1 - 1:1) and allowed to flow through the column. The presence of compounds in these fractions was detected by TLC, fractions having the same pattern were then combined into one so that fractions with almost the same properties were obtained.

Analysis of Chlorogenic Acid Levels in Robusta Coffee Bean Extract:

Analysis of chlorogenic acid levels using the Thin Layer Chromatography (TLC) method. TLC was carried out to determine the chromatogram pattern resulting from the fractionation process of chlorogenic acid compounds in the sample. Determination of chlorogenic acid levels in robusta coffee bean extract begins with making a standard curve with a range according to linearity. Furthermore, the sample solution was prepared by weighing a certain number of samples of robusta coffee bean extract as much as 3 replications then dissolved in methanol. Then the standard solution and sample were put in a vial and spotted on a Silica Gel 60 F254 TLC plate to be analyzed under analytical conditions (eluent composition, wavelength, and test concentration) according to the optimization results. The spots formed are scanned at the optimized wavelength. Then do the calculation of % w/w levels.

DPSC Culture and Isolation:

The pulp tissue was chopped and given 3 ml of collagenase type 1, then incubated in a 5% CO₂ incubator, at 37°C for 30 minutes. After incubation, 3 ml of 25% MEM solution was added and incubated again. The pulp and media were transferred to a 15 ml conical tube, then centrifuged at 3500 rpm for 5 minutes at 28°C. After being centrifuged, the supernatant was

discarded and the pellet was planted in a petri dish containing 3 ml of 25% MEM solution media. Incubate and observe cell growth periodically on day 3 and day 7 to ensure cells can grow and there is no contamination. If the cell culture is 80% confluent, the culture is harvested up to 2-3 times (passage). Cell culture used in the experiment was cell culture 4 times (passages).²⁰

DPSC characteristic analysis:

DPSC characteristics test includes identification of morphology and cell surface proteins. Morphological identification of DPSC was carried out using an inverted light microscope to confirm the shape of the stem cells used in the study. While the identification of cell surface proteins using flow cytometry with antibody markers CD105, CD90, CD45, CD34.

Robusta Coffee Bean Extract Toxicity Analysis on DPSC:

Cells were cultured on 96 well plates and incubated for 24 hours. After 24 hours, all well plates were given robusta coffee bean extract in various concentrations. Giving Robusta coffee bean extract each concentration of 0.5%; 0.25%; 0.125%; 0.0625% as much as 200 µl/well plate. Each well plate was incubated for 24, 48, and 72 hours. After incubation, 25 µl of MTT (3-4-5-dimethylthiazol-2-yl-2,5-dibromodiphenyltetrazolium) was added to each well. Then incubated for 4 hours at 37°C. The color change was observed with Elisa Reader at a wavelength of 595 nm.²¹

RESULT:

Analysis of Chlorogenic Acid Levels in Robusta Coffee Bean Extract:

The results of the standard calculation of chlorogenic

acid obtained a calibration curve of chlorogenic acid levels with absorbance obtained with a linearity value of $R^2 = 0.9992$. The calibration curve is a graph that forms a straight line (linear) which states the relationship between the concentration of the working solution and the proportional response of the instrument used. Measurements at the lowest level of 15 µg/ml and the highest level of 90 µg/ml with each concentration were repeated 3 times to calculate the standard deviation. The results showed a concentration of 90 µg/ml was the peak of a linear calibration curve (Figure 1).




Figure 1. Chlorogenic acid calibration curve

Fractionation of chlorogenic acid levels in robusta coffee bean extract obtained the highest chlorogenic acid fraction content of 30.49%. This concentration is the largest concentration that has been obtained from 13.5 mg of robusta coffee bean extract. From the sample weight, the chlorogenic acid content of 4.1 mg of chlorogenic acid was calculated (Table 1, Figure 2).

Table 1. Table of recapitulation of chlorogenic acid sample calculations. Sample P2 can be fractionated with the highest chlorogenic acid content.

Sample	Sample weight (g)	Area	Measured concentration (µg/ml)	Dilution factor	Weirght (µg)	Calculated concentration (µg/g)	Level (%)
P1	0,0135	393.209,99	201,73	20	4.034,60	298.859,49	29,89
P2	0,0135	401.065,93	205,80	20	4.116,09	304.895,40	30,49
P3	0,0135	396.455,22	203,41	20	4.068,26	301.352,88	30,14
C1	0,0189	385.278,35	197,62	20	3.952,33	209.118,16	20,91
C2	0,0189	414.555,50	212,80	20	4.256,01	225.185,54	22,52
C3	0,0189	411.400,59	211,16	20	4.223,28	223.454,12	22,35
A1	0,0137	165.694,43	83,74	20	1.674,73	122.242,93	12,22
A2	0,0137	159.062,51	80,30	20	1.605,94	117.221,85	11,72
A3	0,0137	167.342,96	84,59	20	1.691,83	123.491,04	12,35




Figure 2. Chromatogram of sample P2 with a concentration of chlorogenic acid found at 30.49%

DPSC Characteristics Analysis

On observation, the morphology of DPSC was fusiform in shape with a tapered tip, a cell nucleus in the middle, and a large cytoplasm, which can be seen in Figure 3 below.




Figure 3. Morphology of DPSC (section 1). A fibroblast-like morphology appears, which is a fusiform shape with a tapered tip (blue arrow), a cell nucleus in the middle, and a large cytoplasm (red arrow). Observations under a microscope with a magnification of 100x.

Observation of DPSC surface expression markers was carried out using a flow cytometry (FCM) test to measure and analyze the surface characteristics of stem cells used in this study. The markers used as positive controls were CD90 and CD105 as markers of

mesenchymal stem cell lineage. Meanwhile, the markers used as negative controls were CD34 and CD45, which are markers of hematopoietic stem cell lineage. The FCM test on the DPSC of this study shows the results according to Figure 4 below.




Figure 4. DPSC FCM test histogram graph. (A, B) DPSCs express the cell surface protein markers CD90 and CD105 which are shown in the green and red diagrams on the right side of the histogram. (C, D) While DPSC did not express the CD34 and CD45 cell surface protein markers which were indicated by the absence of green and red diagrams on the right side of the histogram.

Robusta Coffee Bean Extract Toxicity Analysis on DPSC:

The results of the toxicity test of robusta coffee bean extract at various concentrations on DPSC showed results as shown in Figure 5. Robusta coffee bean extract at concentrations of 0.5% and 0.25% caused toxicity to DPSC at 24, 48, and 72 hours of observation. This was indicated by the percentage of DPSC living cells <80% after the concentration was given. Meanwhile, robusta coffee bean extract concentrations of 0.125% and 0.0625% were not toxic to DPSC.




Figure 5. Graph of Robusta coffee bean extract toxicity test against DPSC. Concentrations that do not cause toxicity are 0.125% and 0.0625%.

DISCUSSION:

The chlorogenic acid content of robusta coffee bean extract in this study was 30.49%. This fractionation result is the highest concentration of chlorogenic acid that has been successfully fractionated. The most chlorogenic acid is contained in robusta coffee beans compared to other herbal sources. This content is mostly contained in robusta coffee beans without the roasting process.^{22,23} Farhaty's research (2016) states that the average chlorogenic acid content in robusta coffee beans is around 6.1-11.3%.³

Chlorogenic acid has biological properties such as anti-inflammatory, antioxidant, antimicrobial, antifungal, and antiviral activities.^{5,6} In addition, several studies have revealed that chlorogenic acid is an effective anti-inflammatory, antipyretic, and analgesic agent in both *in vitro* and *in vivo* experiments.⁷⁻¹⁰ The utilization of chlorogenic acid in stem cells as an agent to increase the anti-inflammatory potential of cells is currently the focus of researchers to develop the success rate of stem cell treatments. In addition, the application of chlorogenic acid to stem cells aims to increase cell proliferation and regeneration, so that cells are susceptible to premature aging. However, the use of chlorogenic acid in periodontal tissue engineering therapy using DPSC still requires further research.

Based on observations of morphological characteristics carried out in this study, it was confirmed that DPSCs are fusiform in shape with a tapered tip, a cell nucleus in the middle, and a large cytoplasm, which is often referred to as a fibroblast-like spindle formation. These results are following the previous theory, which confirmed that DPSCs, which are mesenchymal stem cells, have a morphology similar to fibroblast cells.²⁴⁻²⁶ In this study, DPSCs can also grow and attach to plastic substrates and can colonize. These results are in line with the research of Martens et al. (2010) who investigated the characteristics and surface expression of DPSC *in vitro*. His research explains that in the first 24 hours of culture, cells are able to grow and migrate out of the cultured pulp tissue and will stick to the bottom of the plastic media container which will then form colonies.²⁷

Observation of other characteristics by analyzing DPSC surface expression markers was performed using flow cytometry (FCM) test. The results of the FCM test showed the expression of CD90 and CD105, while CD34 and CD45 were not expressed. CD90 and CD105 specific antibodies are markers of mesenchymal stem cell lineage. While the specific antibodies CD34 and CD45 are markers of hematopoietic stem cell lineage. The results of this study are following pre-existing theories.^{24,25,28} Martens et al. (2010) in their research in

vitro also showed the same results.²⁷

A cell has the criteria for mesenchymal stem cells if it has the following criteria: Able to stick to plastic medium containers when cultured, Expressing CD105, CD90, CD73 and CD44, and lacking expression of CD45, CD34, CD14 or CD11b, CD79 or CD19 and HLA-DR, Can differentiate into osteoblasts, adipocytes, and chondroblasts *in vitro*²⁹. Although this study did not test the ability of cells to differentiate into osteoblasts, adipocytes, or chondroblasts, the other two criteria for mesenchymal stem cells were met.

Toxicity test of robusta coffee bean extract against DPSC showed that 30.49% robusta coffee bean extract at a concentration of 0.125% and 0.0625% was not toxic to DPSC, with the percentage of DPSC living cells >80%. Meanwhile, robusta coffee bean extract concentrations of 0.5% and 0.25% caused toxicity to DPSC. The cytotoxicity level of each concentration of antibacterial agent was calculated as a percentage of cell viability including absorbance values obtained for each treatment. Based on ISO 10993-5, there are cytotoxicity criteria in the viability test, including: The percentage of cell viability > 80% is considered as non-cytotoxic; The percentage of cell viability 80%-60% is considered as weak cytotoxicity; Cell viability percentage 60%-40% is considered as moderate cytotoxicity; The percentage of cell viability < 40% is considered as strong cytotoxicity.³⁰

CONCLUSION:

Robusta coffee bean extract with a chlorogenic acid content of 30.49% at concentrations of 0.125% and 0.0625% did not cause toxicity to DPSC. This concentration is relatively safe to be used as a stem cell combination in periodontal tissue engineering to increase the anti-inflammatory and antioxidant power of stem cells.

CONFLICT OF INTEREST:

The authors have no conflicts of interest regarding this investigation.

ACKNOWLEDGMENTS:

The author would like to thank the Islamic Development Bank through LP2M University of Jember who has funded this research

REFERENCES:

1. Kiattisin K, Nantarat T, Leelapornpisid P. Evaluation of antioxidant and anti-tyrosinase activities as well as stability of green and roasted coffee bean extracts from *Coffea arabica* and *Coffea canephora* grown in Thailand. *J Pharmacogn Phyther*. 2016. doi:10.5897/JPP2016.0413
2. Budiman A, Lutfi A, Mughtaridi M. The Stability of Chlorogenic Acid in Syrup *Coffea arabica* (*Coffea arabica* L.) Extract With

- Deffaceination Process . Asian J Pharm Clin Res. 2017.Special Issue (May):1-5.doi: 10.22159/ajpcr.2017.v10s2.19469
3. Farhady N. Tinjauan Kimia dan Aspek Farmakologi Senyawa Asam Klorogenat pada Biji Kopi : Review. Farmaka Suplemen. 2018.
 4. Liang N, Kitts DD. Role of chlorogenic acids in controlling oxidative and inflammatory stress conditions. *Nutrients*. 2015;8(1):1-20. doi:10.3390/nu8010016
 5. Farah A, dePaula Lima J. Consumption of Chlorogenic Acids through Coffee and Health Implications. *Beverages*. 2019. 5(11):1-29. doi:10.3390/beverages5010011
 6. Naveed M, Hejazi V, Abbas M, et al. Chlorogenic acid (CGA): A pharmacological review and call for further research. *Biomed Pharmacother*. 2018 Jan;97:67-74. doi: 10.1016/j.biopha.2017.10.064.
 7. Gao R, Yang H, Jing S, et al. Protective effect of chlorogenic acid on lipopolysaccharide-induced inflammatory response in dairy mammary epithelial cells. *Microb Pathog*. 2018;124:178-182. doi:10.1016/j.micpath.2018.07.030
 8. Hwang SJ, Kim YW, Park Y, Lee HJ, Kim KW. Anti-inflammatory effects of chlorogenic acid in lipopolysaccharide-stimulated RAW 264.7 cells. *Inflamm Res*. 2014;63(1):81-90. doi:10.1007/s00011-013-0674-4
 9. Liu CC, Zhang Y, Dai BL, et al. Chlorogenic acid prevents inflammatory responses in IL-1 β -stimulated human SW-1353 chondrocytes, a model for osteoarthritis. *Mol Med Rep*. 2017;16(2):1369-1375. doi:10.3892/mmr.2017.6698
 10. Zhou RP, Deng MT, Chen LY, et al. Shp2 regulates chlorogenic acid-induced proliferation and adipogenic differentiation of bone marrow-derived mesenchymal stem cells in adipogenesis. *Mol Med Rep*. 2015;11(6):4489-4495. doi:10.3892/mmr.2015.3285
 11. Suryanendra A, Suryani E. The analysis of coffee productivity and production improvement strategies in Indonesia: A system a. thinking approach. AIP Conference Proceedings. 2021. 2329(1):1-7.doi.org/10.1063/5.0042157
 12. Bin HS, Jeong JH, Choi UK. Chlorogenic acid promotes osteoblastogenesis in human adipose tissue-derived mesenchymal stem cells. *Food Sci Biotechnol*. 2013;22 (suppl.1):107-112. doi:10.1007/s10068-013-0055-3
 13. Liu CC, Zhang Y, Dai BL, et al. Chlorogenic acid prevents inflammatory responses in IL 1 β stimulated human SW 1353 chondrocytes, a model for osteoarthritis. *Mol Med Rep*. 2017 Aug;16(2):1369-1375. doi: 10.3892/mmr.2017.6698. .
 14. Citterio F, Gualini G, Fierravanti L, Aimetti M. Stem cells and periodontal regeneration: present and future. *Plast Aesthetic Res*. 2020. doi:10.20517/2347-9264.2020.29
 15. Hu J, Cao Y, Xie Y, et al. Periodontal regeneration in swine after cell injection and cell sheet transplantation of human dental pulp stem cells following good manufacturing practice. *Stem Cell Res Ther*. 2016. doi:10.1186/s13287-016-0362-8
 16. Kato H, Taguchi Y, Tominaga K, Umeda M, Tanaka A. Porphyromonas gingivalis LPS inhibits osteoblastic differentiation and promotes pro-inflammatory cytokine production in human periodontal ligament stem cells. *Arch Oral Biol*. 2014;59(2):167-175. doi:10.1016/j.archoralbio.2013.11.008
 17. Tang J, Wu T, Xiong J, et al. Porphyromonas gingivalis lipopolysaccharides regulate functions of bone marrow mesenchymal stem cells. *Cell Prolif*. 2015;48(2):239-248. doi:10.1111/cpr.12173
 18. Kizil C, Kyritsis N, Brand M. Effects of inflammation on stem cells: together they strive? *EMBO Rep*. 2015;16(4):416-426. doi:10.15252/embr.201439702
 19. Michael S, Achilleos C, Panayiotou T, Strati K. Inflammation shapes stem cells and stemness during infection and beyond. *Front Cell Dev Biol*. 2016;4(NOV):1-7. doi:10.3389/fcell.2016.00118
 20. Gronthos S, Mankani M, Brahimi J, Robey PG, Shi S. Postnatal human dental pulp stem cells (DPSCs) in vitro and in vivo. *Proc Natl Acad Sci U S A*. 2000. doi:10.1073/pnas.240309797
 21. Bahuguna A, Khan I, Bajpai VK, Kang SC. MTT assay to evaluate the cytotoxic potential of a drug. *Bangladesh J Pharmacol*. 2017. doi:10.3329/bjp.v12i2.30892
 22. Vinson JA, Chen X, Garver DD. Determination of Total Chlorogenic Acids in Commercial Green Coffee Extracts. *J Med Food*. 2019. doi:10.1089/jmf.2018.0039
 23. Farah A, De Paulis T, Trugo LC, Martin PR. Effect of roasting on the formation of chlorogenic acid lactones in coffee. *J Agric Food Chem*. 2005;53(5):1505-1513. doi:10.1021/jf048701t
 24. Nuti N, Corallo C, Chan BMF, Ferrari M, Gerami-Naini B. Multipotent Differentiation of Human Dental Pulp Stem Cells: a Literature Review. *Stem Cell Rev Reports*. 2016;12(5):511-523. doi:10.1007/s12015-016-9661-9
 25. Yamada Y, Nakamura-Yamada S, Kusano K, Baba S. Clinical potential and current progress of dental pulp stem cells for various systemic diseases in regenerative medicine: A concise review. *Int J Mol Sci*. 2019;20(5). doi:10.3390/ijms20051132
 26. Yang X, Li L, Xiao L, Zhang D. Recycle the dental fairy's package: Overview of dental pulp stem cells. *Stem Cell Res Ther*. 2018;9(1):1-8. doi:10.1186/s13287-018-1094-8
 27. Martens W, Wolfs E, Struys T, Politis C, Bronckaers A, Lambrechts I. Expression pattern of basal markers in human dental pulp stem cells and tissue. *Cells Tissues Organs*. 2012;196(6):490-500. doi:10.1159/000338654
 28. Nakashima M, Iohara K, Murakami M. Dental pulp stem cells and regeneration. *Endod Top*. 2013;28(1):38-50. doi:10.1111/etp.12027
 29. Kobolak J, Dinnyes A, Memic A, Khademhosseini A, Mobasher A. Mesenchymal stem cells: Identification, phenotypic characterization, biological properties and potential for regenerative medicine through biomaterial micro-engineering of their niche. *Methods*. 2016;99:62-68. doi:10.1016/j.ymeth.2015.09.016
 30. Cheng Zhang, Ning Wang, Yu Xu et al. Identification of Key Contributive Compounds in a Herbal Medicine: A Novel Mathematic—Biological Evaluation Approach. *Adv. Theory Simul*. 2021: 1-15 DOI: 10.1002/adts.202000279

Chlorogenic Acid Fractionation in Robusta Green Bean Extract as a Combination Agent of Dental Pulp Stem Cells in Periodontal Tissue Engineering

ORIGINALITY REPORT

17%

SIMILARITY INDEX

11%

INTERNET SOURCES

11%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	garuda.kemdikbud.go.id Internet Source	2%
2	balimedicaljournal.org Internet Source	1%
3	eprints.bmsu.ac.ir Internet Source	1%
4	Kobolak, Julianna, Andras Dinnyes, Adnan Memic, Ali Khademhosseini, and Ali Mobasheri. "Mesenchymal stem cells: Identification, phenotypic characterization, biological properties and potential for regenerative medicine through biomaterial micro-engineering of their niche", Methods, 2015. Publication	1%
5	www.ncbi.nlm.nih.gov Internet Source	1%
6	interstellarsuperherbs.com Internet Source	

1 %

7

Sri Wulandari, Makhmudun Ainuri, Anggoro Cahyo Sukartiko. "Biochemical content of Robusta coffees under fully-wash, honey, and natural processing methods", IOP Conference Series: Earth and Environmental Science, 2021
Publication

1 %

8

www.entomologyjournals.com
Internet Source

1 %

9

N L E Wahyuni, R Rispiandi, T Hariyadi. "Effect of bean maturity and roasting temperature on chemical content of robusta coffee", IOP Conference Series: Materials Science and Engineering, 2020
Publication

1 %

10

www.wjgnet.com
Internet Source

1 %

11

rjptonline.org
Internet Source

1 %

12

Rebecca A. Flaherty, Deborah L. Donahue, Katelyn E. Carothers, Jessica N. Ross et al. "Neutralization of Streptolysin S-Dependent and Independent Inflammatory Cytokine IL-1 β Activity Reduces Pathology During Early Group A Streptococcal Skin Infection",

1 %

13

Pravin D. Potdar, Stephanie B. D'souza.
"Isolation of Oct4+, Nanog+ and SOX2-
mesenchymal cells from peripheral blood of a
diabetes mellitus patient", Human Cell, 2011

Publication

<1 %

14

Haibo Jiang, Kewei He, Xueping Luo, Muzi
Zhang, Jian Shao, Lei Gan, Yanhong Lin,
Chuanjie Qin, Hui Zhang, Qiwei Wei.
"Chlorogenic acid attenuates inflammation,
oxidative stress, apoptosis and protects head
kidney macrophage of yellow catfish from
ammonia toxicity", Aquaculture Research,
2021

Publication

<1 %

15

Jeroen S. Werij. "Unravelling enzymatic
discoloration in potato through a combined
approach of candidate genes, QTL, and
expression analysis", Theoretical and Applied
Genetics, 06/26/2007

Publication

<1 %

16

edoc.ub.uni-muenchen.de

Internet Source

<1 %

17

tigerprints.clemson.edu

Internet Source

<1 %

18 Juciane Prois Fortes, Fernanda Wouters Franco, Julia Baranzelli, Gustavo Andrade Ugalde et al. "Enhancement of the Functional Properties of Mead Aged with Oak (Quercus) Chips at Different Toasting Levels", *Molecules*, 2022
Publication

19 www.mdpi.com
Internet Source

20 Ailone Tichon. "Telomerase activity and expression in adult human mesenchymal stem cells derived from amyotrophic lateral sclerosis individuals", *Cytotherapy*, 2009
Publication

21 Dianri Wang, Yuhao Wang, Weidong Tian, Jian Pan. "Advances of tooth - derived stem cells in neural diseases treatments and nerve tissue regeneration", *Cell Proliferation*, 2019
Publication

22 Heba Khalil Alyahya, Pandurangan Subash-Babu, Ahmad Mohammad Salamatullah, Khizar Hayat et al. "Quantification of Chlorogenic Acid and Vanillin from Coffee Peel Extract and its Effect on α -Amylase Activity, Immunoregulation, Mitochondrial Oxidative Stress, and Tumor Suppressor Gene Expression Levels in H₂O₂-Induced Human

Mesenchymal Stem Cells", Frontiers in Pharmacology, 2021

Publication

23

bsdwebstorage.blob.core.windows.net

Internet Source

<1 %

24

docksci.com

Internet Source

<1 %

25

link.springer.com

Internet Source

<1 %

26

www.biorxiv.org

Internet Source

<1 %

27

A. Mukkundur Vasudevaiah, A. Chaturvedi, R. Kulathooran, I. Dasappa. "Effect of green coffee extract on rheological, physico-sensory and antioxidant properties of bread", Journal of Food Science and Technology, 2017

Publication

<1 %

28

Hariyanti Hariyanti, Arry Yanuar, Kusmardi Kusmardi, Hayun Hayun. "Synthesis and in vitro cytotoxic activity of novel indazole analogues of curcumin against MCF-7, HeLa, WiDr, and vero cell lines", Journal of Applied Pharmaceutical Science, 2022

Publication

<1 %

29

parjournal.net

Internet Source

<1 %

30	pubmed.ncbi.nlm.nih.gov Internet Source	<1 %
31	www.researchgate.net Internet Source	<1 %
32	"Bio-orthopaedics", Springer Science and Business Media LLC, 2017 Publication	<1 %
33	"MSCs and Innovative Biomaterials in Dentistry", Springer Science and Business Media LLC, 2017 Publication	<1 %
34	Karola Vorauer-Uhl, Dominik Jeschek, Gabriele Lhota, Andreas Wagner, Stefanie Strobach, Hermann Katinger. "Simultaneous Quantification of Complex Phospholipid Compositions Containing Monophosphoryl Lipid-A by RP-HPLC", Journal of Liquid Chromatography & Related Technologies, 2009 Publication	<1 %
35	"Clinical Regenerative Medicine in Urology", Springer Science and Business Media LLC, 2018 Publication	<1 %
36	"Stem Cells", Springer Science and Business Media LLC, 2019 Publication	<1 %

37

Ashraf Al Madhoun, Sardar Sindhu, Dania Haddad, Maher Atari, Rasheed Ahmad, Fahd Al-Mulla. "Dental Pulp Stem Cells Derived From Adult Human Third Molar Tooth: A Brief Review", *Frontiers in Cell and Developmental Biology*, 2021

Publication

<1 %

38

Jian-Guo Xu, Qing-Ping Hu, Yu Liu. "Antioxidant and DNA-Protective Activities of Chlorogenic Acid Isomers", *Journal of Agricultural and Food Chemistry*, 2012

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Chlorogenic Acid Fractionation in Robusta Green Bean Extract as a Combination Agent of Dental Pulp Stem Cells in Periodontal Tissue Engineering

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
