

March 28, 2023

PLOS ONE

An inclusive journal community working together to advance science by making all rigorous research accessible without barriers

Read the latest COVID-19 research

This Collection highlights content published across the PLOS journals relating to the COVID-19 pandemic.

[READ MORE](#)


EDUCATION

03/08/2023

Neighbourhood effects on educational attainment. What matters more: exposure to poverty or exposure to affluence?

Troost and colleagues show that, in the Netherlands, neighbourhood affluence has a stronger effect on educational attainment than neighbourhood poverty, and that children with higher educated parents are not affected by neighbourhood poverty.

Image credit: [Amsterdam](#) by PublicDomainPictures/ 17901 images, [Pixabay](#)



CANCER RESEARCH

03/16/2023

Validation of a 3D perfused cell culture platform as a tool for humanised preclinical drug testing in breast cancer using established cell lines and patient-derived tissues

Liu and colleagues demonstrate how 3D cell culture can be used to investigate drug effects on breast cancer cells. The approach presents a potential future alternative to animal models and holds promise for personalized medicine.

Image credit: [Fig 1c](#) by Liu et al., [CC BY 4.0](#)



MENTAL HEALTH

03/15/2023

Network analysis reveals abnormal functional brain circuitry in anxious dogs

Xu and colleagues investigated abnormalities in function brain networks in anxious dogs and find that network topological changes at the node level are correlated with anxiety symptoms.

Image credit: [Fig 1](#) by Xu et al., [CC BY 4.0](#)



PLASTIC POLLUTION

03/08/2023

A growing plastic smog, now estimated to be over 170 trillion plastic particles afloat in the world's oceans – urgent solutions required.

Eriksen and colleagues provide an update on their seminal 2014 analysis of plastics floating in the ocean ([pone.0111913](#)), using new data from the peer-reviewed literature and 5 Gyres Institute expeditions. They observe a trend of increasing numbers of plastic particles since 2006, to about 82–358 trillion today.

Image credit: [Fig 1](#) by Eriksen et al., [CC BY 4.0](#)



EVERYONE

03/13/2023

Editor Spotlight: Haroldo V. Ribeiro

In this interview, PLOS ONE Academic Editor Haroldo V. Ribeiro talks about his editorial experience, complex systems, and sharing data and code.

Image credit: Haroldo V. Ribeiro

PUBLISH WITH PLOS ONE

[SUBMISSION INSTRUCTIONS](#)
[SUBMIT YOUR MANUSCRIPT](#)


CALL FOR PAPERS

03/28/2023

Open Calls for Papers

We make it our priority to communicate and promote expert research to the widest possible audience. Built by and for researchers actively working in the field, Calls for Papers surface the latest research in emerging topics and explore important issues. These collections are open for active submissions.

Image credit: [Tourists](#) by Jonathan Ford, [Unsplash License](#)

EVERYONE

03/01/2023

EveryONE

Posted March 13

Editor Spotlight: Haroldo V. Ribeiro

SHEIN EI CHO

Posted March 1

An Update to the Human Subjects Research Policy on PLOS ONE

PLOS ONE EDITORS

Posted February 24



An Update to the Human Subjects Research Policy on PLOS ONE
 Safeguarding the integrity of, and trust in, research involving human subjects.

Image credit: PLOS

Editor Spotlight: Branislav Šiler

SHEIN EI CHO

Posted February 16

Facilitated Preprint Posting is now available for Lab Protocols at PLOS ONE

PLOS

[SEE ALL BLOGS](#)




COMPUTATIONAL BIOLOGY 03/15/2023

B-cell epitope discovery: the first protein flexibility-based algorithm—Zika virus conserved epitope demonstration

Ortoleva and colleagues present a very complete analysis of the epitopes on Zika virus surface.

Image credit: [Eg.2](#) by Biner et al., [CC BY 4.0](#)



COMPUTER SCIENCE 03/15/2023

Identification of social groups and waiting pedestrians at railway platforms using trajectory data

Küpper and Seyfried study non-walking social group dynamics, using data from crowds waiting on train platforms.

Image credit: [Eg.7](#) by Küpper et al., [CC BY 4.0](#)

CONNECT WITH US

[✉](#) [RSS](#) [Twitter](#) [Facebook](#) [PLOS Blogs](#)

[Tweets by @plosone](#)



RESEARCH DESIGN 03/09/2023

Shedding light on participant selection bias in Ecological Momentary Assessment (EMA) studies: Findings from an Internet panel study

Stone and colleagues examine person selection bias in momentary data collection studies.

Image credit: [Work desk](#) by Pixabay, [Pixabay Licence](#)



SPORTS SCIENCE 03/22/2023

Surface effects on kinematics, kinetics and stiffness of habitual rearfoot strikers during running


Wang and colleagues find that rearfoot runners are at a higher risk of injuries when running on harder surfaces.

Image credit: [Jogging](#) by Pixabay, [Pixabay Licence](#)

Call for Papers

Reproducibility and Replicability in Neuroscience and Mental Health


To be considered for this call, papers must be submitted by April 28, 2023.



Collections

Urban Transportation

PLOS ONE presents a curated collection of research from 2020-2022 in the field of Urban Transportation.



Get new content from PLOS ONE in your inbox

YOUR EMAIL ADDRESS

PLOS will use your email address to provide content from PLOS ONE. You can find out more about how PLOS processes your data by reading our [Privacy Policy](#). You can unsubscribe at any time by clicking the unsubscribe link in our emails or by contacting us at privacy@plos.org.

SIGN UP

primary healthcare utilization

[Advanced Search](#)475 results for **primary healthcare utilization**Sort By: **Relevance**

SEARCH ALERT

Filters: Mar 15, 2023 TO Apr 7, 2023 PLOS ONE [Clear all filters](#)

Journal

PLOS ONE

PLOS Global Public Health (50)

PLOS Pathogens (25)

PLOS Neglected Tropical Diseases (23)

PLOS Computational Biology (14)

PLOS Genetics (10)

[show more](#)**Regional differences in primary healthcare utilization in Java Region—Indonesia**

Ratna Dwi Wulandari, Agung Dwi Laksono, Nikmatur Rohmah, Hadi Ashar

Research Article | published 27 Mar 2023 PLOS ONE

<https://doi.org/10.1371/journal.pone.0283709>**Views: 207** • **Citations: 0** • **Saves: 0** • **Shares: 0****Primary care provider uptake of intensive behavioral therapy for obesity in Medicare patients, 2013–2019**

Mounira Ozoor, Mark Gritz, Rowena J. Dolor, Jodi Summers Holtrop, Zhehui Luo

Research Article | published 24 Mar 2023 PLOS ONE

<https://doi.org/10.1371/journal.pone.0266217>**Views: 101** • **Citations: 0** • **Saves: 0** • **Shares: 0**

Subject Area

Biology and life sciences (393)

Medicine and health sciences (393)

Research and analysis methods (230)

Social sciences (211)

People and places (204)

[show more](#)**Impact of COVID-19 on healthcare utilization, cases, and deaths of citizens and displaced Venezuelans in Colombia: Complementary comprehensive and safety-net systems under Colombia's constitutional commitment**

Donald S. Shepard, Adelaida Boada, Douglas Newball-Ramirez, Anna G. Sombrio, Carlos William Rincon Perez, Priya Agarwal-Harding, Jamie S. Jason, Arturo Harker Roa, Diana M. Bowser

Research Article | published 28 Mar 2023 PLOS ONE

<https://doi.org/10.1371/journal.pone.0282786>**Views: 153** • **Citations: 0** • **Saves: 0** • **Shares: 0****Understanding the barriers and facilitators of COVID-19 vaccine hesitancy amongst healthcare workers and healthcare students worldwide: An umbrella review protocol**

Jemma McCready, Dania Comparcini, Bethany Nichol, Mary Steen, John Unsworth, Marco Tomietto

Study Protocol | published 31 Mar 2023 PLOS ONE

<https://doi.org/10.1371/journal.pone.0283897>**Views: 295** • **Citations: 0** • **Saves: 0** • **Shares: 14**

Article Type

Research Article (445)

Study Protocol (27)

Lab Protocol (2)

Collection Review (1)

Remote Assessment in healthcare—Technologies, methods, benefits, and challenges

Jakob Eyvind Bardram

Collection Review | published 06 Apr 2023 PLOS ONE

<https://doi.org/10.1371/journal.pone.0283945>**Views: 24** • **Citations: 0** • **Saves: 0** • **Shares: 0**

Author

Atalay Mulu Fentie (2)

Benjamin M Davies (2)

Bradley N Gaynes (2)

Brian W Pence (2)

Elizabeth Thomas (2)

[show more](#)**Predicting a diagnosis of ankylosing spondylitis using primary care health records—A machine learning approach**

Jonathan Kennedy, Natasha Kennedy, Roxanne Cooksey, Ernest Choy, Stefan Siebert, Muhammad Rahman, Sinead Brophy

Research Article | published 31 Mar 2023 PLOS ONE

<https://doi.org/10.1371/journal.pone.0279076>**Views: 197** • **Citations: 0** • **Saves: 0** • **Shares: 11**

Where my keywords appear

Body (372)

Results and Discussion (137)

Materials and Methods (108)

References (97)

Introduction (73)

[show more](#)**Cost-effectiveness of transdiagnostic group cognitive behavioural therapy versus group relaxation therapy for emotional disorders in primary care (PsicAP-Costs2): Protocol for a multicentre randomised controlled trial**

César González-Blanch, Sara Barrio-Martínez, Amador Priede, Sandra Martínez-Gómez, Saioa Pérez-García-Abad, María Miras-Aguilar, José Ruiz-Gutiérrez, Roger Muñoz-Navarro, Paloma Ruiz-Rodríguez, Leonardo A. Medrano, Maider Prieto-Vila, María Carpallo-González, Ángel Aguilera-Martín, Mario Gálvez-Lara, Fátima Cuadrado, Eliana Moreno, Francisco García-Torres, José F. Venceslá, Jorge Corpas, Francisco J. Jurado-González, Juan A. Moriana, Antonio Cano-Vindel

Study Protocol | published 16 Mar 2023 PLOS ONE

<https://doi.org/10.1371/journal.pone.0283104>**Views: 393** • **Citations: 0** • **Saves: 0** • **Shares: 5**

Publication Date

2023-03-15

to

Health care utilizations and costs of *Campylobacter* enteritis in Germany: A claims data analysis

2023-04-07

Apply

Elisabeth Schorling, Sonja Lick, Pablo Steinberg, Dagmar Adeline Brüggemann

Research Article | published 05 Apr 2023 PLOS ONE

<https://doi.org/10.1371/journal.pone.0283865>

Views: 76 • Citations: 0 • Saves: 0 • Shares: 1

Defining avoidable healthcare-associated harm in prisons: A mixed-method development study

Richard N. Keers, Verity Wainwright, Joy McFadzean, Kate Davies, Stephen M. Campbell, Caroline Stevenson, Thomas Purchase, Jennifer Shaw, Andrew Carson-Stevens

Research Article | published 15 Mar 2023 PLOS ONE

<https://doi.org/10.1371/journal.pone.0282021>

Views: 298 • Citations: 0 • Saves: 0 • Shares: 6

Investment case for primary health care in low- and middle-income countries: A case study of Kenya

Daniel Mwai, Salim Hussein, Agatha Olago, Maureen Kimani, David Njuguna, Rose Njiraini, Elizabeth Wangia, Easter Olwanda, Lilian Mwaura, Wesley Rotich

Research Article | published 23 Mar 2023 PLOS ONE

<https://doi.org/10.1371/journal.pone.0283156>

Views: 651 • Citations: 0 • Saves: 2 • Shares: 1

Healthcare system action on employment as a social determinant of health in people living with HIV: A qualitative study

Amy Craig-Neil, Julia Ho, Melissa Perri, Mark Gaspar, Charlotte Hunter, Beth Rachlis, Claire E. Kendall, Sergio Rueda, Ann N. Burchell, Andrew D. Pinto

Research Article | published 06 Apr 2023 PLOS ONE

<https://doi.org/10.1371/journal.pone.0282421>

Views: 45 • Citations: 0 • Saves: 0 • Shares: 4

Mining actionable combined high utility incremental and associated sequential patterns

Min Shi, Yongshun Gong, Tiantian Xu, Long Zhao

Research Article | published 29 Mar 2023 PLOS ONE

<https://doi.org/10.1371/journal.pone.0283365>

Views: 107 • Citations: 0 • Saves: 0 • Shares: 0

Ethnobotanical survey of medicinal plants used by indigenous knowledge holders to manage healthcare needs in children

Peter Tshepiso Ndhlovu, John Awungnjia Asong, Abiodun Olusola Omotayo, Wilfred Otang-Mbeng, Adeyemi Oladapo Aremu

Research Article | published 27 Mar 2023 PLOS ONE

<https://doi.org/10.1371/journal.pone.0282113>

Views: 298 • Citations: 0 • Saves: 0 • Shares: 0

Evaluating the adoption of voice recognition technology for real-time dictation in a rural healthcare system: A retrospective analysis of dragon medical one

Adedayo A. Onitilo, Abdul R. Shour, David S. Puthoff, Yusuf Tanimu, Adedayo Joseph, Michael T. Sheehan

Research Article | published 23 Mar 2023 PLOS ONE

<https://doi.org/10.1371/journal.pone.0272545>

Views: 109 • Citations: 0 • Saves: 0 • Shares: 0

Healthcare resource utilisation and cost of pneumococcal disease from 2003 to 2019 in children ≤ 17 years in England

Salini Mohanty, B el ene Podmore, Ana Cu nado Moral, Ian Matthews, Eric Sarpong, Agueda Azpeitia, Nawab Qizilbash

Research Article | published 05 Apr 2023 PLOS ONE

<https://doi.org/10.1371/journal.pone.0283084>

Views: 48 • Citations: 0 • Saves: 0 • Shares: 0

Results per page 15 ▾

Editorial Board

The *PLOS ONE* Editorial Board is powered by thousands of academic experts from all over the world. Our board comprises working scientists who are established principal investigators/group leaders with extensive publication records.

Academic Editors oversee the peer review process for the journal, including evaluating submissions, selecting reviewers and assessing their comments, and making editorial decisions. Together with fellow Editorial Board Members and internal staff, Academic Editors uphold journal policies and ethics standards and work to promote the *PLOS ONE* mission to provide free public access to scientific research.

Editorial board members affiliated with the Centers for Disease Control and Prevention are serving in a personal capacity. The views expressed are their own and do not necessarily represent the views of the Centers for Disease Control and Prevention or the United States Government.

Editorial board members affiliated with the World Health Organization are serving in a personal capacity. The views expressed are their own and do not necessarily represent the decisions, policy or views of the World Health Organization.

Interested in serving on the Editorial Board?

PLOS welcomes volunteers to the Editorial Board who support our mission, values, and commitment to providing a high-quality experience for our authors. [Explore the Academic Editor role](#) and [apply online](#).

Academic Editors

To find a member, browse the list, search by name, or search by Section or [Keyword](#). To search for multiple sections or keywords, enter a comma-separated list of terms.

The list of Academic Editors is updated weekly. If you are an Editorial Board member and would like to update any of the information below, please contact plosone@plos.org

Search for:

Displaying 1-50 of 10338 Editors.

< 1 2 3 4 5 ... 207 >

Rana Muhammad Aadil

orcid.org/0000-0002-0185-0096

University of Agriculture Faisalabad
PAKISTAN

Sections: Food science and technology, Nutrition

Keywords: Biology and life sciences, Nutrition, Medicine and health sciences, Nutrition, Diet, Food, Pathology and laboratory medicine, Pathogens

Katriina Aalto-Setälä

University of Tampere
FINLAND

Sections: Developmental biology - Cell differentiation; Cell fate determination; Stem cells; Embryology; Fertilization

Keywords: Biology and life sciences, Cellular types, Stem cells, Induced pluripotent stem cells, Embryonic stem cells, Developmental biology, Cell differentiation, Genetics, Genetics of disease, Genetic testing, Human genetics, Genetic association studies, Molecular genetics, Cardiovascular anatomy, Electrophysiology, Medicine and health sciences, Cardiology, Arrhythmia, Clinical genetics, Personalized medicine, Animal models, Mouse models, Model organisms

Mohd Nadhir Ab Wahab

orcid.org/0000-0002-3549-6443

Universiti Sains Malaysia
MALAYSIA

Sections: Computer and information sciences - Artificial intelligence, machine learning and data science, Engineering and technology - Aerospace and automotive engineering, Robotics

Keywords: Computer and information sciences, Artificial intelligence, Real time computing, Engineering and technology, Mechanical engineering, Robotics, Robotics systems, Nanorobotics, Navigation, Optimization

Nurul Hidayah Ab Rahman

Universiti Tun Hussein Onn Malaysia
MALAYSIA

Sections: Computer and information sciences - Cryptography and computer security

Keywords: Computer and information sciences, Computing systems, Digital computing, Information technology, Computer security, Computing methods, Cloud computing

Andrew Max Abaasa

orcid.org/0000-0002-6770-5588

London School of Hygiene and Tropical Medicine Faculty of Epidemiology and Population Health
UGANDA

Sections: Infectious diseases - Epidemiology and prevention, Public health and epidemiology - Biostatistics and methods

Keywords: Medicine and health sciences, Epidemiology, Health care, Clinical trials, Research and analysis methods, Research design

Nidaa Ababneh

orcid.org/0000-0002-2155-3013

University of Jordan
JORDAN

Sections: Genetics - Mutation; Genetics of disease; Heredity, Stem cells and regenerative medicine, Neuroscience - Neurodegenerative diseases and dementia
Keywords: Medicine and health sciences, Neurology

Muktar Abadiga

[id. orcid.org/0000-0002-3006-3155](https://orcid.org/0000-0002-3006-3155)

Wollega University
ETHIOPIA

Sections: Cardiovascular science and medicine - Epidemiology and prevention, Nursing, Health care - General
Keywords: Medicine and health sciences, Health care, Neurology

Ukachukwu Okoroafor Abaraogu

[id. orcid.org/0000-0002-1967-1459](https://orcid.org/0000-0002-1967-1459)

Glasgow Caledonian University
UNITED KINGDOM

Sections: Physiotherapy, Health care - Health services research, Cardiovascular science and medicine - Vascular diseases
Keywords: Medicine and health sciences, Health care, Cardiology, Cardiovascular medicine, Rehabilitation medicine, Vascular medicine, Complementary and alternative medicine

Francisco Martinez-Abarca

Estacion Experimental del Zaidin - CSIC
SPAIN

Sections: Agriculture - General
Keywords: Biology and life sciences, Genome evolution, Comparative genomics, Microbial ecology, Evolutionary biology, Transposable elements, Genomics, Metagenomics, Microbiology, Plant microbiology, Genome sequencing, Biochemistry, Nucleic acids, Biotechnology, Applied microbiology

Behnam Abasht

[id. orcid.org/0000-0003-2374-8145](https://orcid.org/0000-0003-2374-8145)

University of Delaware
UNITED STATES

Sections: Agriculture - Animals, Genetics - Gene function, Genomics
Keywords: Biology and life sciences, Agriculture, Livestock, Genetics

Kalkidan Hassen Abate

Jimma University
ETHIOPIA

Sections: Sport and exercise science, Physiotherapy
Keywords: Nutrition, Physiology, Sports and exercise medicine, Medicine and health sciences

Ayew Mose Abawa

[id. orcid.org/0000-0003-4353-7791](https://orcid.org/0000-0003-4353-7791)

Wolkite University
ETHIOPIA

Sections: Obstetrics and gynecology, Nursing
Keywords: Medicine and health sciences

Eric Fosu Oteng-Abayie

[id. orcid.org/0000-0002-4598-2066](https://orcid.org/0000-0002-4598-2066)

Kwame Nkrumah University of Science and Technology
GHANA

Sections: Economics - Econometrics, Economics - General, Economics - Behavioral; Experimental
Keywords: Social sciences, Economics, Ecology and environmental sciences, Environmental economics

Sani Isah Abba

[id. orcid.org/0000-0001-9356-2798](https://orcid.org/0000-0001-9356-2798)

King Fahd University of Petroleum and Minerals
SAUDI ARABIA

Sections: Computer and information sciences - Artificial intelligence, machine learning and data science, Engineering and technology - General
Keywords: Ecology and environmental sciences, Natural resources, Environmental chemistry, Engineering and technology, Membrane technology, Sanitary engineering, Chemical engineering, Civil engineering, Environmental engineering

Gianmarco Abbadessa

[id. orcid.org/0000-0001-8912-3055](https://orcid.org/0000-0001-8912-3055)

University of Campania Luigi Vanvitelli: Universita degli Studi della Campania Luigi Vanvitelli
ITALY

Sections: Digital health, Neuroscience - Cellular and molecular, Neuroscience - Neurobiology of disease; Neuropathology
Keywords: Immunology, Clinical immunology, Neuroimmunology, Medicine and health sciences, Health care, Medical services, Telemedicine, Immunology, Clinical immunology, Autoimmune diseases, Multiple sclerosis, Neurology, Demyelinating disorders, Multiple sclerosis

Cristiana Abbafati

[id. orcid.org/0000-0003-2811-6251](https://orcid.org/0000-0003-2811-6251)

Sapienza University of Rome
ITALY

Sections: Economics - Health economics, Health care - General, Public health and epidemiology - Health policies, systems and management
Keywords: Social sciences, Economics, Medicine and health sciences, Health care, Health care facilities, Hospitals, Outpatient clinics, Health care policy, Health care providers, Health care quality, Health care utilization, Health economics, Health services administration and management, Health services research, Health statistics, Morbidity, Community based intervention, Quality of life, Socioeconomic aspects of health, Public and occupational health

Faisal Abbas

[id. orcid.org/0000-0002-9312-5659](https://orcid.org/0000-0002-9312-5659)

National University of Sciences and Technology
PAKISTAN

Sections: Economics - Health economics, Public health and epidemiology - Global health, Public health and epidemiology - Health policies, systems and management
Keywords: Nutrition, Malnutrition, Social sciences, Economics, Economic history, Development economics, Economic development, Economic analysis

Mazhar Abbas

University of Ha'il Kingdom of Saudi Arabia
 PAKISTAN

Sections: Management science

Keywords: Social sciences, Economics, Research and analysis methods, Research design

Asad Abbas

[id orcid.org/0000-0003-1395-4009](https://orcid.org/0000-0003-1395-4009)

Tecnologico de Monterrey
 MEXICO

Sections: Education research, Management science

Keywords: Social sciences, Sociology, Education, Political science, Public administration

Qaiser Abbas

Ghazi University
 PAKISTAN

Sections: Agriculture - General, Economics - General, Economics - Health economics

Keywords: Agricultural economics, Ecology, Ecological economics, Neuroeconomics, Behavioral economics, Social sciences, Economics, Economic history, Economic models, Experimental economics, Labor economics, Resource management, Development economics, Economic analysis, Soil science, Ecology and environmental sciences, Environmental impacts, Environmental management, Natural resources, Environmental protection, Environmental economics, Health economics

Qaisar Abbas

[id orcid.org/0000-0001-5176-5685](https://orcid.org/0000-0001-5176-5685)

The Shaikh Ayaz University Shikarpur, Pakistan
 PAKISTAN

Sections: Education research

Keywords: Social sciences, Anthropology, Sociology, Education, Educational attainment, Training (education), Home education, Archaeology, Human geography, Law and legal sciences, Linguistics, Philosophy, Political science, Psychology, People and places, Population groupings, Educational status, Research and analysis methods, Research assessment, Research design, Research facilities

Alhamzah F. Abbas

[id orcid.org/0000-0002-7508-9340](https://orcid.org/0000-0002-7508-9340)

UTM Skudai: Universiti Teknologi Malaysia
 MALAYSIA

Sections: Management science, Economics - General

Keywords: Medicine and health sciences, Health care, Health care quality, Quality of care, Quality of life

Mahdieh Abbasalizad Farhangi

[id orcid.org/0000-0002-7036-6900](https://orcid.org/0000-0002-7036-6900)

Tabriz University of Medical Sciences
 IRAN, ISLAMIC REPUBLIC OF

Sections: Nutrition, Cardiovascular science and medicine - Arrhythmias and cardiac electrophysiology, Health care - Health education

Keywords: Biology and life sciences, Genetics, Immunology, Neuroscience, Nutrition, Diet, Malnutrition, Nutrients, Nutritional deficiencies, Nutritional diseases, Nutritional disorders, Physiology, Medicine and health sciences, Endocrinology, Epidemiology, Cardiovascular medicine, Pharmacology, Drug research and development, Women's health, Clinical trials, Complementary and alternative medicine

Mohsen Abbasi-Kangevari

[id orcid.org/0000-0001-8405-7482](https://orcid.org/0000-0001-8405-7482)

Tehran University of Medical Sciences
 IRAN, ISLAMIC REPUBLIC OF

Sections: Clinical trials, Digital health, Public health and epidemiology - Chronic diseases

Keywords: Social sciences, Political science, Public policy, Medicine and health sciences, Health care, Military medicine, Gulf War syndrome, Public and occupational health, Clinical medicine

Mahdi Abbasi

[id orcid.org/0000-0002-5373-5778](https://orcid.org/0000-0002-5373-5778)

Bu-Ali Sina University: Bu Ali Sina University
 IRAN, ISLAMIC REPUBLIC OF

Sections: Computer and information sciences - Artificial intelligence, machine learning and data science, Computer and information sciences - Computer Hardware, Computer and information sciences - General

Keywords: Computer and information sciences, Artificial intelligence, Artificial neural networks, Expert systems, Genetic programming, Machine learning, Computing systems, Digital computing, Digital imaging, Network analysis, Network theory, Signaling networks, Network control, Neural networks, Feedforward neural networks, Recurrent neural networks, Random number generators, Real time computing, Computer applications, Computer architecture, Computer hardware, Microprocessors, Computer networks, Internet, Internet of Things, Computer vision, Computers, Computing methods

A. M. Abd El-Aty

[id orcid.org/0000-0001-6596-7907](https://orcid.org/0000-0001-6596-7907)

Cairo University
 EGYPT

Sections: Veterinary science, Pharmacology, Food science and technology

Keywords: Antimicrobials, Phytochemicals, Environmental chemistry, Medicine and health sciences, Pharmacology, Drugs, Pharmacodynamics, Pharmacogenetics, Pharmacokinetics, Pharmacologic analysis, Pharmacokinetic analysis, Drug interactions, Drug-food interactions, Physical sciences, Chemistry, Analytical chemistry, Chemical analysis, Water analysis, Liquid chromatography-tandem mass spectrometry, Phytochemistry, Phytopharmacology, Research and analysis methods, Extraction techniques, Supercritical fluid extraction, Liquid-liquid extraction, Solid-phase extraction, Chromatographic techniques, Liquid chromatography

Khaled Abd EL-Hamid Abd EL-Razik

[id orcid.org/0000-0002-8895-2895](https://orcid.org/0000-0002-8895-2895)


National Research Center (NRC)

RESEARCH ARTICLE

Regional differences in primary healthcare utilization in Java Region—Indonesia

Ratna Dwi Wulandari^{1,2*}, Agung Dwi Laksono^{2,3}, Nikmatur Rohmah⁴, Hadi Ashar³

1 Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia, 2 The Airlangga Centre for Health Policy (ACeHAP), Surabaya, Indonesia, 3 National Research and Innovation Agency Republic of Indonesia, Jakarta, Indonesia, 4 Faculty of Health Science, Muhammadiyah University of Jember, East Java, Indonesia

* ratna-d-w@fkm.unair.ac.id

Abstract

Background

Policymakers must understand primary healthcare utilization disparity to minimize the gap because they must seek fair service for every citizen. The study analyzes regional differences in primary healthcare utilization in Java Region-Indonesia.

Methods

The cross-sectional research analyzes secondary data from the 2018 Indonesian Basic Health Survey. The study setting represented Java Region-Indonesia, and the participants were adults 15 years or more. The survey explores 629,370 respondents. The study used primary healthcare utilization as an outcome variable and province as the exposure variable. Moreover, the study employed eight control variables (residence, age, gender, education, marital, employment, wealth, and insurance). The study evaluated data using binary logistic regression in the final step.

Results

People in Jakarta are 1.472 times more likely to utilize primary healthcare than those in Banten (AOR 1.472; 95% CI 1.332–1.627). People in Yogyakarta are 1.267 times more likely to use primary healthcare than those in Banten (AOR 1.267; 95% CI 1.112–1.444). In addition, people in East Java are 15% less likely to utilize primary healthcare than those in Banten (AOR 0.851; 95% CI 0.783–0.924). Meanwhile, direct healthcare utilization was the same between West Java, Central Java, and Banten Province. They are sequentially starting from the minor primary healthcare utilization: East Java, Central Java, Banten, West Java, Yogyakarta, and Jakarta.

Conclusion

Disparities between regions exist in the Java Region-Indonesia. They are sequentially starting from the minor primary healthcare utilization: East Java, Central Java, Banten, West Java, Yogyakarta, and Jakarta.

OPEN ACCESS

Citation: Wulandari RD, Laksono AD, Rohmah N, Ashar H (2023) Regional differences in primary healthcare utilization in Java Region—Indonesia. PLoS ONE 18(3): e0283709. <https://doi.org/10.1371/journal.pone.0283709>

Editor: Retno Asti Werdhani, Universitas Indonesia Fakultas Kedokteran, INDONESIA

Received: October 28, 2022

Accepted: March 14, 2023

Published: March 27, 2023

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0283709>

Copyright: © 2023 Wulandari et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The data supporting this study's findings are available from the National Institute of Health Research and Development of Indonesia's Ministry of Health. Still, restrictions apply to the availability of these data used under

license for the current study, so they are not publicly available. However, data are available from the authors upon reasonable request and with permission of the National Institute of Health Research and Development of Indonesia Ministry of Health via the web page <http://www.litbang.kemkes.go.id/jasa-permintaan-data-riset>.

Funding: The author(s) received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Introduction

According to the World Health Organization, primary healthcare is essential healthcare based on methods and techniques that are practical, scientific, socially acceptable, and universally accessible to individuals, families, and communities [1]. Puskesmas (Pusat Kesehatan Masyarakat/Center of Public Health) is a primary healthcare facility unique to Indonesia. The primary entry point to the government-run healthcare system is Puskesmas. In addition to general practice and primary clinics, Puskesmas is Indonesia's entry point to critical services. Every subdistrict, or 30,000 residents in Indonesia, must have access to primary healthcare [2]. At least 9,993 Puskesmas in Indonesia will serve around 250 million people through 2018 [3].

In Indonesia, Puskesmas, as the embodiment of primary healthcare, is the first-level health service facility in the era of national health insurance (NHI). Puskesmas are required to provide comprehensive health services, covering 144 diseases for participants. Services must have multiple benefits, both medical and non-medical; namely, the services provided are complete, ranging from preventive, promotive, curative, and rehabilitative. Promotive is a process for the community's ability to maintain and improve their health; One of the efforts made by the Puskesmas is health counseling. Meanwhile, prevention is an effort to take various actions to avoid health problems threatening ourselves and others in the future [4].

On the other hand, curative is a health effort to prevent the disease from getting worse through treatment; The Puskesmas provides the most appropriate and quickest possible treatment for each type of disease so that complete and immediate healing is achieved. Puskesmas serve this effort by providing medical personnel. Moreover, curative is an action or a series of activities aimed at ex-sufferers (patients who are no longer suffering from the disease), so they can generally interact in the social environment [4].

There are often some misconceptions about primary healthcare, namely the assumption that primary healthcare provides only basic care when primary healthcare provides essential care that can meet most of a person's health needs throughout life. Primary healthcare serves health for all ages for all diseases, including promotive, preventive, curative, rehabilitation, and palliative [1]. Several factors are related to the utilization of primary healthcare, such as the variety of services, the consultation process, affordability, and convenience. The situation correlates with patient comfort and satisfaction [5]. Patient satisfaction is the most crucial and is one of the main reasons patients visit primary healthcare [6]. Experience in Canada, several factors influence the use of primary healthcare. It includes doctor's practice, various services, incentives, combination payment schemes, electronic medical records, and efforts to improve quality, showing that primary healthcare transformation is achieved pluralistically [7].

Health development objectives are to increase the reach and distribution of quality health services for the community. With Indonesia's existing geographical conditions, it is challenging for the government to solve. So far, comprehensive health services have been centered in Java; some of the best-advanced healthcare facilities (hospitals) are in Java [8]. The Java region is a benchmark for success for other areas in Indonesia, including models for policies addressing gaps in primary healthcare utilization.

Achieving the primary healthcare target requires additional investment in more comprehensive health services. At the United Nations high-level universal health coverage meeting in 2019, countries committed to strengthening primary healthcare. The WHO recommends that each country allocate an additional budget of 1% of GDP [9]. All stakeholders, from the government and doctors to community members, need to be aware of primary healthcare's role and benefits [1]. Policy policymakers should pursue patient satisfaction in direct healthcare services by improving the quality of medical services, especially in rural areas [5]. According to the research background, the study analyzes regional differences in primary healthcare utilization in Java Region in Indonesia.

Methods

Study design and data source

We designed a cross-sectional study using secondary data from the 2018 Indonesian Basic Health Survey. The 2018 Indonesian Basic Health Survey was a national-scale cross-sectional survey conducted by the Ministry of Health of the Republic of Indonesia. The survey collected data from May to July 2018 through Household and Individual Instruments interviews.

The survey included all Indonesian households. This survey employs the sample framework of the 2018 National Socio-Economic Survey (conducted by the Central Statistics Agency in March 2018). Furthermore, in the 2018 Socio-Economic Survey, the 2018 Indonesian Basic Health Survey visited a target sample of 300,000 households from 30,000 census blocks [10].

The PPS (probability proportional to size) method is used in the 2018 Indonesian Basic Health Survey, with systematic linear sampling in two stages: Stage 1: Implicit stratification based on the welfare strata of all census blocks from the 2010 Population Census. PPS selected the sample survey as the sampling frame from a master frame of 720,000 census blocks from the 2010 Population Census, of which 180,000 were chosen (25%). The survey used the PPS method to determine several census blocks in each urban/rural strata per regency/city to create a Census Block Sample List. A total of 30,000 Census Blocks have been selected. Stage 2: Using systematic sampling, choose ten households in each Census Block with the highest implicit education stratification completed by the Head of Household to maintain the representation of the diversity value of household characteristics. As part of the 2018 Indonesian Basic Health Survey, all selected household members will be interviewed [10].

The study population comprised all adults (15 years old) in Indonesia's Java Region. According to the sampling methods, we obtained a weighted sample of 213,140 adults for this study.

Outcome variable

The study's outcome variable was primary healthcare utilization—adults' access to primary healthcare; whatever the reason for the need for such access, whether outpatient or inpatient, was the utilization of primary healthcare. Outpatient care was limited to the previous month, while inpatient care was limited to last year. The survey asked respondents to recall the correct outpatient and inpatient incidences [10]. Primary healthcare utilization comprises unutilized and utilized.

Exposure variable

We employed region or province as an exposure variable in this study. The region comprises all provinces (six) on Java Island, namely Jakarta, West Java, Central Java, Yogyakarta, East Java, and Banten.

Control variables

Moreover, the study employed eight factors as control variables. The eight factors were the type of residence, age group, gender, education level, marital status, employment status, wealth status, and health insurance ownership. The study divided the available residence into two categories: urban and rural. We used the Indonesian Central Statistics Agency's provisions for urban-rural categorization in the survey. Furthermore, the study calculated age based on the most recent birthday, and the age group comprises ≤ 17 , 18–64, and ≥ 65 . Gender includes two categories: male and female.

The study also classified marital status into three categories: single, married/living with a partner, and divorced/widowed. The study defined education as accepting the respondent's most recent diploma. The study has four levels of education: no education, primary, secondary, and higher education. Meanwhile, there are two employment options: unemployed and employed.

The pool used the wealth index formula to determine wealth status in the study. The survey calculated the wealth index using a weighted average of a family's total spending. Meanwhile, the survey calculated the wealth index using primary household expenditures such as health insurance, food, and lodging, among other things. In addition, the poll classified income into five categories: poorest, poorer, middle, wealthier, and most prosperous [11, 12]. According to the survey, there are two types of health insurance owners: uninsured and insured.

Study setting

We are setting the study to represent the Java Region in Indonesia.

Data analysis

The Chi-Square test was used in the early stages of the sample to produce a bivariate comparison between region (province) with other variables. Furthermore, we used a collinearity test to ensure that the independent variables in the final regression model did not have a strong relationship. The study's final point employed binary logistic regression (enter method). The survey used the last test to examine the multivariate relationship between all independent variables and primary healthcare utilization. We declare it's statistically significant at $p < 0.050$. With 95% confidence intervals, adjusted odds ratios (AOR) were presented (95% CI). Throughout the statistical analysis process, we employed the IBM SPSS 26 application.

Ethical approval

The 2018 Indonesian Basic Health Survey received Ethical Clearance from the National Ethics Committee (LB.02.01/2/KE.024/2018). During data collection, the survey used informed consent, which accounted for aspects of the data collection procedure, such as voluntariness and confidentiality. Respondents provided written permission.

Results

The analysis found that the average primary healthcare utilization in Java Region in 2018 was 5.1%. Meanwhile, Table 1 shows descriptive statistics of direct healthcare utilization and respondents' characteristics in Indonesia in 2018. The result indicates that Yogyakarta Province has the highest primary healthcare utilization compared to other provinces in the Java Region. Based on the type of residence, a minor proportion of urban areas is in Central Java Province. According to the age group, 18–64 dominated all provinces. Moreover, regarding gender, males ruled in West Java and Banten Provinces.

Based on marital status, Table 1 indicates marriage dominated all regions. Regarding education level, primary education leads all provinces, except Jakarta Provinces, which are led by secondary education. According to employment status, employed dominated all areas. The most prosperous lead in Jakarta, West Java, Yogyakarta, and Banten Province is based on wealth status. Regarding health insurance ownership, insured people dominated all regions.

Table 2 shows the binary logistic regression of primary healthcare utilization in Java Region-Indonesia. We used "unutilized primary healthcare" as a reference. Based on the province, people in Jakarta are 1.472 times more likely to utilize primary healthcare than those in

Table 1. Descriptive statistic of primary healthcare utilization and respondents' characteristics in Java Region in Indonesia, 2018 (n = 213,140).

Demographic Characteristics	Province						p-value
	Jakarta (n = 9,711)	West Java (n = 48,491)	Central Java (n = 63,118)	Yogyakarta (n = 8,312)	East java (n = 68,747)	Banten (n = 14,761)	
Primary Healthcare Utilization							< 0.001
Unutilized	93.8%	94.7%	94.8%	93.6%	95.4%	95.4%	
Utilized	6.2%	5.3%	5.2%	6.4%	4.6%	4.6%	
Type of residence							< 0.001
Urban	100.0%	74.1%	51.3%	72.9%	52.3%	71.9%	
Rural	-	25.9%	48.7%	27.1%	47.7%	28.1%	
Age Group							< 0.001
≤ 17	5.7%	7.5%	6.8%	5.3%	6.2%	7.0%	
18–64	88.6%	84.8%	82.0%	82.7%	83.5%	88.2%	
≥ 65	5.7%	7.7%	11.2%	12.1%	10.3%	4.8%	
Gender							< 0.001
Male	49.7%	50.4%	49.0%	48.7%	48.8%	50.8%	
Female	50.3%	49.6%	51.0%	51.3%	51.2%	49.2%	
Marital status							< 0.001
Never in union	25.5%	22.5%	21.4%	23.3%	19.6%	24.4%	
Married/Living with a partner	66.9%	69.4%	69.2%	66.9%	69.7%	68.6%	
Divorced/Widowed	7.5%	8.1%	9.5%	9.9%	10.7%	7.0%	
Education level							< 0.001
No education	2.6%	3.5%	7.6%	6.0%	8.4%	5.1%	
Primary	38.9%	62.5%	63.4%	41.6%	59.3%	53.9%	
Secondary	44.0%	26.7%	22.5%	38.9%	25.4%	32.5%	
Higher	14.5%	7.3%	6.5%	13.5%	7.0%	8.5%	
Employment status							< 0.001
Unemployed	39.7%	42.9%	32.7%	32.9%	34.4%	43.5%	
Employed	60.3%	57.1%	67.3%	67.1%	65.6%	56.5%	
Wealth status							< 0.001
Poorest	1.0%	16.3%	25.3%	18.4%	22.0%	8.6%	
Poorer	4.7%	19.3%	20.7%	19.4%	20.5%	14.6%	
Middle	16.3%	18.1%	18.7%	15.4%	18.2%	19.5%	
Richer	25.0%	18.1%	18.3%	19.8%	20.0%	22.9%	
Richest	53.0%	28.2%	17.1%	27.0%	19.3%	34.4%	
Health Insurance							< 0.001
Uninsured	15.5%	34.4%	31.2%	17.5%	37.4%	35.3%	
Insured	84.5%	65.6%	68.8%	82.5%	62.6%	64.7%	

<https://doi.org/10.1371/journal.pone.0283709.t001>

Banten (AOR 1.472; 95% CI 1.332–1.627). People in Yogyakarta are 1.267 times more likely to use primary healthcare than those in Banten (AOR 1.267; 95% CI 1.112–1.444). In addition, people in East Java are 15% less likely to utilize primary healthcare than those in Banten (AOR

Table 2. The result of binary logistic regression of primary healthcare utilization in Java Region in Indonesia, 2018 (n = 213,140).

Predictor	Utilized Primary Healthcare			
	p-value	AOR	95% CI	
			Lower Bound	Upper Bound
Province: Jakarta	* <0.001	1.472	1.332	1.627
Province: West Java	0.492	1.028	0.950	1.114
Province: Central Java	0.123	0.936	0.861	1.018
Province: Yogyakarta	* <0.001	1.267	1.112	1.444
Province: East Java	* <0.001	0.851	0.783	0.924
Province: Banten	-	-	-	-
Residence: Urban	** 0.001	1.080	1.034	1.129
Residence: Rural	-	-	-	-
Age: ≤ 17	-	-	-	-
Age: 18–64	0.292	1.062	0.950	1.187
Age: ≥ 65	* <0.001	1.522	1.342	1.726
Gender: Male	-	-	-	-
Gender: Female	* <0.001	1.326	1.267	1.387
Marital: Never in union	-	-	-	-
Marital: Married/Living with partner	* <0.001	1.716	1.592	1.849
Marital: Divorced/Widowed	* <0.001	2.174	1.977	2.391
Education: No Education	-	-	-	-
Education: Primary	0.102	0.942	0.876	1.012
Education: Secondary	* <0.001	0.560	0.513	0.611
Education: Higher	* <0.001	0.302	0.262	0.348
Employment: Unemployed	-	-	-	-
Employment: Employed	* <0.001	0.805	0.769	0.842
Wealth: Poorest	-	-	-	-
Wealth: Poorer	0.252	0.966	0.910	1.025
Wealth: Middle	0.160	0.957	0.901	1.017
Wealth: Richer	* <0.001	0.866	0.814	0.922
Wealth: Richest	* <0.001	0.617	0.576	0.661
Health insurance: Uninsured	-	-	-	-
Health insurance: Insured	* <0.001	1.841	1.757	1.929

Note: *p<0.001

**p<0.010

<https://doi.org/10.1371/journal.pone.0283709.t002>

0.851; 95% CI 0.783–0.924). Meanwhile, there was no significant difference in direct healthcare utilization between West Java, Central Java, and Banten Province. They are sequentially starting from the minor primary healthcare utilization: East Java, Central Java, Banten, West Java, Yogyakarta, and Jakarta.

Moreover, the study also found eight control variables significantly correlated with primary healthcare utilization. Regarding the type of residence, people in urban areas are 1.080 times more likely to use primary healthcare than those in rural areas (AOR 1.080; 95% CI 1.034–1.129). According to the age group, those ≥ 65 are 1.522 times more likely than those ≤ 17 to utilize primary healthcare (AOR 1.522; 95% CI 1.342–1.726). Based on gender, females are 1.326 times more likely to use primary healthcare than males (AOR 1.326; 95% CI 1.267–1.387).

Married or living with a partner is 1.716 times more likely to use primary healthcare than never in a union (AOR 1.716; 95% CI 1.592–1.849). Furthermore, divorced or widowed are

2.174 times more likely to utilize primary healthcare than those never in a union (AOR 2.174; 95% CI 1.977–2.391). Based on education level, all education levels are less likely than no school to utilize primary healthcare, except primary education, which is not significantly related to no education.

According to employment, the employed are 20% times less likely to use primary healthcare than the unemployed (AOR 0.805; 95% CI 0.769–0.842). All wealth statuses are less likely than the poorest to use primary healthcare, except for the poorer and the middle. There are no significant differences among the poorest in utilizing primary healthcare. Based on health insurance ownership, the insured are 1.841 times more likely than the uninsured to use primary healthcare (AOR 1.841; 95% CI 1.757–1.929).

Discussion

Essential health services are the first level of health services and are the first contact of the population with the healthcare system. The primary healthcare management policy aims to increase the utilization of essential health services in Indonesia. One of these policies is establishing the Regional Public Service Agency (RPSA). This RPSA makes primary healthcare more developed because the funds/financial support issued follow the needs and are not dependent on the health department [13]. The utilization of primary healthcare will be a critical factor in Indonesia's successful use of health infrastructure [14]. The Indonesian government has implemented a series of Universal Health Coverage (UHC) policy reforms that include the integration of the government insurance scheme into the National Health Insurance (NHI), expanding the provider network, and accreditation of all health facilities [15]. However, the study results show that UHC in Indonesia still faces the burden of state finances for healthcare and a lack of healthcare facilities, especially in primary health units [16].

The results show differences in primary healthcare utilization between regions in Java Region in Indonesia. The utilization of essential health services between areas in the Java Region of Indonesia is still not evenly distributed. The results are similar to the geographic inequalities in health services utilization in 497 districts in Indonesia. The study results indicate a significant variation in the utilization of health services at the district level and a minor variation at the provincial level [17, 18]. Studies on the reform of the Indonesian public health system also find indications that post-reform, the gap is widening in the propensity of the poor to file insurance claims depending on their level of access to health centers [19]. The study supports previous studies that reported disparities in the utilization of primary health among regions in rural Indonesia. Regions in western Indonesia tend to have better utilization of Puskesmas in rural areas [8, 20, 21].

The study indicates people in urban areas are more likely to use primary healthcare than those in rural areas. Rural people have more significant barriers to using primary healthcare than people in urban areas. The results support the study in the US, which states that compared to urban residents, rural residents have lower access to primary care providers [22, 23]. However, the survey on the satisfaction level of primary healthcare did not show a significant difference between rural and urban dwellings [24]. Studies in India also reveal rural-urban inequalities that do not benefit the rural population and the utilization of healthcare for the elderly [25]. This disparity also exists in China; a study reports that rural residents in China are still underutilizing healthcare when compared to their urban counterparts [26]. A similar study in Ghana shows a generally low utilization of health services among rural communities [27].

All age groups are more likely than ≤ 17 to utilize primary healthcare, and young adults and above mainly use primary healthcare. These results support an integrative review study

that states that age is the predisposing factor that most influences the use of direct health services by the elderly [28]. The survey in Gomoa confirmed that age was the key factor determining the utilization of health services in the sub-district [29]. Studies in the Chongqing or Guizhou areas also report that age is associated with knowledge, utilization of health education, and satisfaction in the primary healthcare sector. The situation is related to previous treatment experience and degenerative disease factors [30].

All marital statuses are more likely to use primary healthcare than those who are never in a union, and those in marital status have a greater need to utilize direct health services. In line with the results of this study, marital status determines the utilization of primary health services [31]. Studies in China also stated that marital status was associated with knowledge, utilization, and satisfaction with health education in primary healthcare [32, 33]. Compared to unmarried respondents, married respondents tend to make outpatient visits, and married patients use more healthcare services at the secondary and primary healthcare levels [34].

The study shows education level related to primary healthcare utilization in Java Region in Indonesia. Those not educated tend to use more essential health services than those with secondary and higher education. This study supports a study in Ghana, which reported that education had a positive and significant effect on health services utilization in rural communities [27]. The level of education also has a statistically significant relationship with the likelihood of seeking services and the level of services used in dealing with simple ailments in the Iranian Shiraz community [35]. Education level was also significantly associated with using primary healthcare for antenatal care [36]. This result is different from the study in Gomoa, which reported that education level was insignificant as a determinant of primary healthcare utilization [29].

According to employment, the employed are less likely to use primary healthcare than the unemployed. People who work generate income, and employed people may have a higher likelihood of using primary healthcare. Another study states that income level is significantly related to the probability of finding services and the services used in dealing with simple diseases [35]. This result is different from the study in Gomoa, which confirmed that the average monthly income was insignificant as a determinant of primary health service utilization [29].

All wealthy statuses were less likely to use primary healthcare than the poorest, except for the poorer and middle class. There is no significant difference between the poorest communities in direct health services, and people mostly use primary healthcare, with the most impoverished. The situation is in line with the results of a study in Ethiopia which stated that the supporting factor in the form of monthly income determines the utilization of primary health services. Low household and socio-economic status encourage direct health services [31]. Moreover, wealth status is also related to health insurance ownership, and wealthier people tend to have health insurance [37, 38].

The insured is more likely than the uninsured to use primary healthcare based on the possession of health insurance. People who have health insurance make more use of direct health services. This study's results align with reflections on the impact of subsidy policies on the utilization of various types of healthcare facilities in Indonesia. The results show that subsidized NHI premiums for the poor at the first level of healthcare facilities can increase the likelihood of outpatient visits to primary healthcare [39, 40]. The study in Shiraz also stated that insurance and type had a significant relationship with the probability of finding services and the level of service used in dealing with simple illnesses [35]. However, previous research in Indonesia found that general health insurance for the poor and public health insurance had minimal impact on the utility of health facilities as measured by outpatient and inpatient visits by program beneficiaries [41]. The NHI program increases the utilization of outpatient and inpatient care in the contribution group. To a lesser extent, those with subsidized insurance have

increased access to inpatient-only facilities [42]. Expanding health insurance coverage reduces sociodemographic disparities in Indonesia's access to maternal health services. However, significant differences in utilization remain across regions and economic subgroups [43].

The government should minimize inter-regional service disparities in its territory, including differences in health services. To overcome this situation, the government has made several efforts. Among them encourage the expansion of NHI membership by regulating in more detail assistance or premium subsidies for NHI membership for the poor and requiring employers to bear the NHI membership premium for their workers [37, 44]. On the other hand, the government is also expanding the Puskesmas network by establishing *Pondok Kesehatan Desa (Ponkesdes)* or Village Health Cottages for each village [45]. Furthermore, the government is trying to collaborate with more primary clinics and general practices in the cooperation mechanism within NHI [46].

Strength and limitation

The study examines a large amount of data to represent information on a national scale. On the other hand, the study examines secondary data; thus, the accepted variables limit the factors analyzed. Several other factors related to primary healthcare utilization discovered in previous studies, such as travel time, travel cost, and disease type, cannot be studied [17, 47].

Conclusion

Based on the study results, the study concluded that disparities between regions exist in the Java Region in Indonesia. The study listed East Java, Central Java, Banten, West Java, Yogyakarta, and Jakarta in order of minor primary healthcare utilization.

Interventions to increase primary healthcare utilization focusing on regions are insufficient as the primary strategy to address health disparities at the population level. Future policies should concentrate more on interventions targeting social determinants of upstream health, targeting this study's results. Efforts should focus on increasing interventions for those who live in rural areas, are younger, have never been in a union, are better educated, are employed, have better wealth status, and are uninsured.

Acknowledgments

The author would like to thank the National Institute of Health Research and Development, which has agreed to allow the authors to analyze the data set in this article.

Author Contributions

Conceptualization: Ratna Dwi Wulandari, Agung Dwi Laksono.

Data curation: Nikmatur Rohmah, Hadi Ashar.

Formal analysis: Ratna Dwi Wulandari, Agung Dwi Laksono.

Funding acquisition: Ratna Dwi Wulandari.

Investigation: Nikmatur Rohmah, Hadi Ashar.

Methodology: Ratna Dwi Wulandari, Agung Dwi Laksono.

Project administration: Hadi Ashar.

Resources: Ratna Dwi Wulandari.

Software: Agung Dwi Laksono, Nikmatur Rohmah.

Validation: Nikmatur Rohmah, Hadi Ashar.

Visualization: Nikmatur Rohmah, Hadi Ashar.

Writing – original draft: Agung Dwi Laksono, Nikmatur Rohmah, Hadi Ashar.

Writing – review & editing: Ratna Dwi Wulandari.

References

1. World Health Organization. Primary health care. 2022 [cited 15 Aug 2022] p. 1. Available: https://www.who.int/health-topics/primary-health-care#tab=tab_1
2. Ministry of Health of the Republic of Indonesia. Regulation of the Minister of Health of the Republic of Indonesia Number 75 of 2014 concerning Community Health Centers (Peraturan Menteri Kesehatan Republik Indonesia Nomor 75 Tahun 2014 tentang Pusat Kesehatan Masyarakat). Permenkes 75/2014 Jakarta, Indonesia: Kementerian Kesehatan Republik Indonesia; 2014.
3. Ministry of Health of the Republic of Indonesia. Puskesmas Basic Data; Conditions December 31, 2018 (Data Dasar Puskesmas; Kondisi 31 Desember 2018). Jakarta; 2019. Available: [https://www.kemkes.go.id/resources/download/pusdatin/data-dasar-puskesmas/2018/00.Buku Data Dasar Puskesmas kondisi 31 des 2018-Nasional.pdf](https://www.kemkes.go.id/resources/download/pusdatin/data-dasar-puskesmas/2018/00.Buku%20Data%20Dasar%20Puskesmas%20kondisi%2031%20des%202018-Nasional.pdf)
4. Laksono AD, Nantabah ZK, Wulandari RD, Khoiri A, Tahangnacca M. Barriers to Expanding the National Health Insurance Membership in Indonesia: Who Should the Target? *J Prim Care Community Heal.* 2022;13. <https://doi.org/10.1177/21501319221111112> PMID: 35818670
5. Gao Q, Liu M, Peng L, Zhang Y, Shi Y, Teuwen DE, et al. Patient Satisfaction and Its Health Provider--related Determinants in Primary Health Facilities in Rural China. *BMC Health Serv Res.* 2022; 22: 1–12. <https://doi.org/10.1186/s12913-022-08349-9> PMID: 35883080
6. Peng X, Zhang JH. Socio-economic Inequality in Public Satisfaction with The Healthcare System in China. *Arch Public Heal.* 2022;80.
7. Hutchison Brian, Levesque JF, Strumpf E, Coyle N. Primary Health Care in Canada: Systems in Motion. *Milbank Q.* 2016; 62. <https://doi.org/10.1111/j.1468-0009.2011.00628.x>
8. Laksono AD, Wulandari RD, Soedirham O. Regional Disparities of Health Center Utilization in Rural Indonesia. *Malaysian J Public Heal Med.* 2019; 19: 158–166. <https://doi.org/10.37268/mjphm/vol.19/no.1/art.48>
9. Chan Margaret. Primary Health Care: Now More than Ever. *UN Chron.* 2021; 47: 4–7. <https://doi.org/10.18356/cbabf986-en>
10. National Institute of Health Research and Development of the Ministry of Health of the Republic of Indonesia. The 2018 Indonesian Basic Health Survey. Jakarta; 2018 [cited 7 Oct 2021] p. 99. Available: http://labdata.litbang.kemkes.go.id/images/download/laporan/RKD/2018/Laporan_Nasional_RKD2018_FINAL.pdf
11. Wulandari RD, Qomarrudin MB, Supriyanto S, Laksono AD. Socio-economic Disparities in Hospital Utilization among Elderly People in Indonesia. *Indian J Public Heal Res Dev.* 2019; 10: 1800–1804. <https://doi.org/10.5958/0976-5506.2019.03885.3>
12. Wulandari RD, Laksono AD, Prasetyo YB, Nandini N. Socio-economic Disparities in Hospital Utilization Among Female Workers in Indonesia: A Cross-Sectional Study. *J Prim Care Community Health.* 2022; 13: 1–7. <https://doi.org/10.1177/21501319211072679> PMID: 35068256
13. Shubhan MH, Aisyah RHS, . . . Optimization Of Primary Health Care To Improve Health Services For Participants Of Social Insurance Administering Agency Of Health In Indonesia. *J Crit . . .* 2020; 7: 4498–4505.
14. Wenang S, Schaefers J, Afdal A, Gufron A, Geyer S, Dewanto I, et al. Availability and Accessibility of Primary Care for the Remote, Rural, and Poor Population of Indonesia. *Front Public Heal.* 2021; 9: 1–11. <https://doi.org/10.3389/fpubh.2021.721886> PMID: 34621720
15. Wiseman V, Thabrany H, Asante A, Haemmerli M, Kosen S, Gilson L, et al. An evaluation of health systems equity in Indonesia: Study protocol. *Int J Equity Health.* 2018; 17: 1–9. <https://doi.org/10.1186/s12939-018-0822-0> PMID: 30208921
16. Mutiarin D, Suranto, Darumurti A, Tamronglak A, Whangmahaporn P, Sakir. Improving Policy Model of Universal Health Coverage Policy: A Comparison Study between Indonesia and Thailand. *International Conference on Governance Innovation and Social Science.* 2019. pp. 1–17. <https://doi.org/10.32528/pi.v0i0.2530>
17. Laksono AD, Wulandari RD. Predictors of hospital utilization among papuans in Indonesia. *Indian J Forensic Med Toxicol.* 2020; 14: 2319–2324. <https://doi.org/10.37506/ijfmt.v14i2.3374>

18. Mulyanto J, Kunst AE, Kringos DS. Geographical inequalities in healthcare utilisation and the contribution of compositional factors: A multilevel analysis of 497 districts in Indonesia. *Heal Place*. 2019; 60: 102236. <https://doi.org/10.1016/j.healthplace.2019.102236> PMID: 31778844
19. Rinasih R. Has Reform of the System of Public Health Insurance for the Poor in Indonesia Increased Health Inequality? Considering Different Accessibility of Puskesmas. *Info Artha*. 2019; 3: 101–118. <https://doi.org/10.31092/jia.v3i2.584>
20. Mahmudiono T, Laksono AD. Disparity in the hospitals utilization among regions in Indonesia. *Open Access Maced J Med Sci*. 2021; 9: 1461–1466. <https://doi.org/10.3889/oamjms.2021.7304>
21. Laksono AD, Wulandari RD, Rohmah N, Rukmini R, Tumaji T. Regional disparities in hospital utilisation in Indonesia: a cross-sectional analysis data from the 2018 Indonesian Basic Health Survey. *BMJ Open*. 2023; 13: e064532. <https://doi.org/10.1136/bmjopen-2022-064532> PMID: 36596635
22. Chen X, Orom H, Hay JL, Waters EA, Schofield E, Li Y, et al. Differences in Rural and Urban Health Information Access and Use. *J Rural Heal*. 2019; 35: 405–417. <https://doi.org/10.1111/jrh.12335> PMID: 30444935
23. Wulandari RD, Laksono AD, Nantabah ZK, Rohmah N, Zuardin Z. Hospital utilization in Indonesia in 2018: do urban-rural disparities exist? *BMC Health Serv Res*. 2022; 2022: 491. <https://doi.org/10.1186/s12913-022-07896-5> PMID: 35413914
24. Yaya S, Bishwajit G, Ekholuenetale M, Shah V, Kadio B, Udenigwe O. Urban-rural difference in satisfaction with primary healthcare services in Ghana. *BMC Health Serv Res*. 2017; 17: 1–9. <https://doi.org/10.1186/s12913-017-2745-7> PMID: 29178876
25. Banerjee S. Determinants of rural-urban differential in healthcare utilization among the elderly population in India. *BMC Public Health*. 2021; 21: 1–18. <https://doi.org/10.1186/s12889-021-10773-1> PMID: 34001026
26. Li J, Shi L, Liang H, Ding G, Xu L. Urban-rural disparities in health care utilization among Chinese adults from 1993 to 2011. *BMC Health Serv Res*. 2018; 18: 1–9. <https://doi.org/10.1186/s12913-018-2905-4> PMID: 29426313
27. Agyemang S, Asibey BO. Effect Of Education On Health Care Utilization In Rural Ghana: The Case Of Selected Communities In The Bekwai Municipality. *KNUST J Geogr Dev*. 2018; 2: 114–127.
28. Baithesda B, Chen C. Factors Influencing Primary Healthcare Utilization by the Elderly: an Integrative Review of the Literature. *Proceeding 2nd Manad Heal Polytech Int Conf*. 2019; 1–7.
29. Konneh-Archer M. Determinants of Primary Healthcare Utilization in the Gomoa East Sub-District. *Mod-ish Proj*. 2019; July.
30. Zhang R, Chen Y, Liu S, Liang S, Wang G, Li L, et al. Progress of equalizing basic public health services in Southwest China—health education delivery in primary healthcare sectors. *BMC Health Serv Res*. 2020; 20: 1–13. <https://doi.org/10.1186/s12913-020-05120-w> PMID: 32209085
31. Kim D, Birmeta K, Kim C, Sharma B, Dhaka S, Lee H, et al. Determinants of Primary Health Care Utilization in Ethiopia. *Inst Poverty Alleviation Int Dev*. 2016; 25s: 269–293. <https://doi.org/10.18350/ipaid.2016.25s.269>
32. Wulandari RD, Laksono AD, Matahari R. Does Husband's Education Level Matter to Antenatal Care Visits? A Study on Poor Households in Indonesia. *Indian J Community Med*. 2022; 47: 192–195. https://doi.org/10.4103/ijcm.ijcm_981_21 PMID: 36034251
33. Megatsari H, Laksono AD, Herwanto YT, Sarweni KP, Geno RAP, Nugraheni E, et al. Does husband/partner matter in reduce women's risk of worries?: Study of psychosocial burden of covid-19 in indonesia. *Indian J Forensic Med Toxicol*. 2021; 15: 1101–1106. <https://doi.org/10.37506/ijfmt.v15i1.13564>
34. Bjørnelv GMW, Edwin B, Fretland AA, Deb P, Aas E. Till death do us part: The effect of marital status on health care utilization and costs at end-of-life. A register study on all colorectal cancer decedents in Norway between 2009 and 2013. *BMC Health Serv Res*. 2020; 20: 1–13. <https://doi.org/10.1186/s12913-019-4794-6> PMID: 32054492
35. Niyas M, Karimi M, Kavosi Z. Utilization of primary health care services in rural and urban areas in Shiraz. *Shiraz E Med J*. 2018; 19. <https://doi.org/10.5812/semj.66219>
36. Okonofua F, Ntoimo L, Ogungbangbe J, Anjorin S, Imongan W, Yaya S. Predictors of Women's Utilization of Primary Health Care for Skilled Pregnancy Care in Rural Nigeria. *BMC Pregnancy Childbirth*. 2018; 18: 1–15. <https://doi.org/10.1186/s12884-018-1730-4> RESEARCH
37. Laksono AD, Nugraheni WP, Rohmah N, Wulandari RD. Health insurance ownership among female workers in Indonesia: does socio-economic status matter? *BMC Public Health*. 2022; 22: 1798. <https://doi.org/10.1186/s12889-022-14189-3> PMID: 36138387
38. Laksono AD, Rukmini R, Tumaji T, Ipa M, Wulandari RD. National Health Insurance Membership among Urban Poor Societies in Indonesia in 2019: Are They Protected? *Economies*. 2022; 10: Article number 196. <https://doi.org/10.3390/economies10080196>

39. Zamaro RT. The Impact of Health Subsidy on Health Care Utilization in Indonesia. *J Anggar dan Keuang Negara Indones*. 2021; 3: 16–37. <https://doi.org/10.33827/akurasi2021.vol3.iss2.art129>
40. Laksono AD, Wulandari RD, Efendi F. Determinants of hospital utilisation among urban poor societies in Indonesia. *Int J Innov Creat Chang*. 2020; 12: 375–387.
41. Astuti RD, Nugrohowati RNI. Impact evaluation of subsidized health insurance programs on utilization of healthcare facilities: evidence from Indonesia. *Int J Heal Econ Manag*. 2022;Jan: 1. <https://doi.org/10.1007/s10754-021-09321-6> PMID: 35028789
42. Erlangga D, Ali S, Bloor K. The impact of public health insurance on healthcare utilisation in Indonesia: evidence from panel data. *Int J Public Health*. 2019; 64: 603–613. <https://doi.org/10.1007/s00038-019-01215-2> PMID: 30737522
43. Anindya K, Lee JT, McPake B, Wilopo SA, Millett C, Carvalho N. Impact of Indonesia's national health insurance scheme on inequality in access to maternal health services: A propensity score matched analysis. *J Glob Health*. 2020; 10: 1–12. <https://doi.org/10.7189/jogh.10.010429> PMID: 32566167
44. Nasution SK, Mahendradhata Y, Trisnantoro L. Can a National Health Insurance Policy Increase Equity in the Utilization of Skilled Birth Attendants in Indonesia? A Secondary Analysis of the 2012 to 2016 National Socio-Economic Survey of Indonesia. *Asia-Pacific J Public Heal*. 2020; 32: 19–26. <https://doi.org/10.1177/1010539519892394> PMID: 31810376
45. Munir M, Nursalam N, Triyoga RS. Village health post (Ponkesdes) development into community nursing center-based Health Promotion Model, nursing center, and behavioral performance. *Int J Med Sci Public Heal*. 2016; 5: 1. <https://doi.org/10.5455/ijmsph.2016.07092015119>
46. Kurniawati W, Rachmayanti RD. Identification of the Causes of Low JKN Participation in Informal Sector Workers in Rural Areas (Identifikasi Penyebab Rendahnya Kepesertaan JKN pada Pekerja Sektor Informal di Kawasan Pedesaan). *J Adm Kesehat Indones*. 2018; 6: 33. <https://doi.org/10.20473/jaki.v6i1.2018.33-39>
47. Wei Y, Yu H, Geng J, Wu B, Guo Z, He L, et al. Hospital efficiency and utilization of high-technology medical equipment: A panel data analysis. *Heal Policy Technol*. 2018; 7: 65–72. <https://doi.org/10.1016/j.hlpt.2018.01.001>



KEMENTERIAN KESEHATAN REPUBLIK INDONESIA
BADAN PENELITIAN DAN PENGEMBANGAN KESEHATAN

Jalan Percetakan Negara No. 29 Jakarta 10560 Kotak Pos 1226

Telepon (021) 4261088 faksimile (021) 4243933

Laman : www.litbang.depkes.go.id Surat Elektronik : sesban@litbang.depkes.go.id



PERSETUJUAN AMANDEMEN PROTOKOL
PROTOCOL AMENDMENT APPROVAL

No. : LB.02.01/2/KE.024/2018

Ref. : Persetujuan/Approval no : LB.02.01/2/KE.267/2017 tanggal 28 Juli 2017

Komisi Etik Penelitian Kesehatan, Badan Penelitian dan Pengembangan Kesehatan (KEPK-BPPK) dengan berdasarkan Deklarasi Helsinki, telah melakukan telaah, pembahasan dan penilaian melalui proses **expedited**.

memutuskan amandemen protokol penelitian yang berjudul :

Health Research Ethics Committee, National Institute of Health Research and Development (HREC-NIHRD), in accordance with Helsinki Declaration, has conducted a thorough expedited review of research protocol amendment entitled :

"Riset Kesehatan Dasar (RISKESDAS) 2017-2018"

yang akan mengikutsertakan manusia sebagai partisipan/subyek penelitian; dengan Ketua Pelaksana/Peneliti Utama :

in which will involve human participant(s), with Principal Investigator :

dr. Agus Suprpto, M.Kes.

dapat diberikan persetujuan amandemen sesuai surat pengantar no. LB.02.03/1/406/2018 tanggal 16 Januari 2018. Masa berlaku surat persetujuan etik ini adalah :

*has hereby declared the amendment is **approved** for implementation. This letter is valid from/to*

24 Januari 2018 s/d 28 Juli 2018


Jika ada perubahan protokol (amandemen) dan/atau perpanjangan penelitian, Ketua Pelaksana/Peneliti Utama harus mengajukan kembali protokol versi terbaru untuk kaji etik penelitian. Pada akhir penelitian, laporan pelaksanaan penelitian juga harus diserahkan kepada KEPK-BPPK.

Should there be any modification (amendment) and/or extension of the study, the Principal Investigator is required to resubmit the latest version of protocol for approval. The final summary reports should also be submitted to HREC-NIHRD.

Chair of HREC-NIHRD :

Jakarta, 24 Januari 2018

Ketua
Komisi Etik Penelitian Kesehatan
Badan Litbang Kesehatan,



Prof. Dr. M. Sudomo

