



rr retno widyowati <rr-retno-w@ff.unair.ac.id>

[MPI] Submission Acknowledgement

1 message

Kartini <mpi@unit.ubaya.ac.id>

Fri, Jan 21, 2022 at 2:36 PM

To: Retno Widyowati <rr-retno-w@ff.unair.ac.id>

Retno Widyowati:

Thank you for submitting the manuscript, "Review: Quality Control Study of Crude Drug of Justicia gendarussa Burm. f. Leaves as Male Contraceptive" to MPI (Media Pharmaceutica Indonesiana). With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Submission URL: <https://journal.ubaya.ac.id/index.php/MPI/authorDashboard/submission/4838>

Username: retnowidyowati

If you have any questions, please contact me. Thank you for considering this journal as a venue for your work.

Kartini

[MPI \(Media Pharmaceutica Indonesiana\)](#)



rr retno widyowati <rr-retno-w@ff.unair.ac.id>

[MPI] Editor Decision

2 messages

Kartini <kartini@staff.ubaya.ac.id>

Fri, Mar 18, 2022 at 10:04 AM

To: Rokhmatul Ummah <rokhmatul.ummah-2018@ff.unair.ac.id>, Bambang Prajogo <prajogo_ew@yahoo.com>, Retno Widyowati <rr-retno-w@ff.unair.ac.id>

Rokhmatul Ummah, Bambang Prajogo, Retno Widyowati:

We have reached a decision regarding your submission to MPI (Media Pharmaceutica Indonesiana), "Review: Quality Control Study of Crude Drug of Justicia gendarussa Burm. f. Leaves as Male Contraceptive".

Our decision is to: Accept Submission

Kartini

kartini@staff.ubaya.ac.id

[MPI \(Media Pharmaceutica Indonesiana\)](#)

rr retno widyowati <rr-retno-w@ff.unair.ac.id>

Tue, Mar 22, 2022 at 2:22 PM

To: Kartini <kartini@staff.ubaya.ac.id>

Selamat siang,


Terima kasih atas informasi dan kesempatan pada artikel kami yang akan dipublikasi pada MPI. Berikut ini saya kirimkan hasil proofread yang telah kami lakukan dan bukti proofread. Terima kasih.


Salam,

Rr. Retno Widyowati, PhD

[Quoted text hidden]

2 attachments

 **2 4838-Article Text-13054-1-15-20220226 (Proffread).doc**
965K

 **2 4838-Article Text-13054-1-15-20220226-final.doc**
962K

Review: Quality Control Study of Crude Drug of *Justicia gendarussa* Burm. f. Leaves as Male Contraceptive

ASUS 13/2/22 17.49

Comment [1]: Lebih cocok sbg mini review

Abstract

Justicia gendarussa Burm. f. (Gendarussa) leaves contain gendarusin A, which is a potential anti-fertility agent. Preclinical researches (*in vitro*, *in vivo*, & toxicity) and clinical trials have been carried out proving that Gendarussa leaves are safe to be used as a male contraceptive. Quality control of medicine must be carried out from the beginning when selecting crude drugs until the production process. Quality control of the crude drug is needed to maintain the quality, safety, and efficacy of their crude drug. That standardization has been done in Kediri, Mojokerto, and Ponorogo. The research output data used were specific and non-specific parameters (macroscopic and microscopic assay, determination of ash content, compound levels, etc.). The quality of crude drugs is different in each region. Crude drug from Mojokerto is better than crude drug from Kediri and Ponorogo. This is influenced by environmental difference factors in how plants grow. This review describes the differences in the quality control results of crude drug of Gendarussa from several areas. So in the future, the quality of this plant can be guaranteed quality to be used as a contraceptive drug.

Key Words: *Justicia gendarussa* Burm. f.; quality control; crude drug

1. Introduction

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

Justicia gendarussa Burm. f. (Gendarussa) is a plant from the Acanthaceae family spread in Indonesia and several Asian countries such as Sri Lanka, India, and Malaysia [1]. This plant is often used as a folk remedy to cure rheumatic, bronchitis, eczema, fever, etc. [2]. Apigenin and vitexin glycosides in Gendarussa can be used as anti-inflammatory and anti-tumor [3]. However, *in vitro* and *in vivo* anti-fertility tests of these leaves show competitive and reversible inhibition of the hyaluronidase enzyme in spermatozoa [4]. Therefore, this plant has the potential to be developed into a non-hormonal male contraceptive drug [5,6]. Capsule extracts from the Gendarussa as an anti-fertility have been tested in phase III clinical trials [7].

Gendarussa leaves contain potassium, flavonoids, justicin, steroids and alkaloids [4]. In addition, these leaves also contain gendarusin A (6,8-di-C- α -L-arabinopyranosyl-4',5,7-trihydroxyflavone) as a major component, gendarusin B (6-C- α -L-arabinopyranosyl-4',5,7-trihydroxy-8-C- β -D-cylopyranosyl-flavone), gendarusin C, gendarusin D, and gendarusin E as minor components [8]. Gendarusin is a compound of the more acid-resistant C-glycosides flavonoid group [9]. Analysis to determine the compounds contained in the plant is necessary to help identify the side effects of these active compounds so as to improve the quality of the drug [10,11].

Quality control is a process involved in maintaining the quality and validity of the resulting products [12]. This process needs to be done in a production activity to produce phytopharmaceutical with better quality [13]. Efforts to ensure the quality and safety of the drug must be carried out from the beginning of the production process when the selection of crude drug until the

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

end of production and then the product is circulated in the community [14]. The economic value added in a crude drug that meets the standard is much greater than the unstandardized crude drug [15]. Standardization of crude drugs is carried out to determine the quality, safety, and efficacy requirements of the Gendarussa crude drug leaves. Standardization parameters consist of specific and non-specific parameters.

Research on the quality control test of Gendarussa crude drug has been conducted in several areas, for example, Kediri, Mojokerto, and Ponorogo [16,17]. The quality of some of these crude drugs varies. This is because drugs produced from natural materials have varying compound content. These variations occur due to several factors, such as genetic, environmental, agronomy engineering, and harvest [18]. Therefore, it is necessary to standardize in order to obtain crude drugs with good quality as a first step in the process of developing drugs from natural materials.

ASUS 28/1/22 11.40

Comment [2]: -Tujuan penelitian review ini apa?
-Kalimat terakhir ini menyatakan pentingnya standarisasi, bukan pentingnya penelitian review ini.

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

2. Methods

1.1. Article Selection Criteria and Sources

Article searches were conducted on electronic databases namely PubMed, SCOPUS, ScienceDirect, Google Scholar, and Universitas Airlangga Repository. The searches were limited until September 2020. The keywords used in the search were Gendarussa (*Justicia gendarussa* Burm. f.), quality control, and crude drug in research articles/dissertation/thesis. From the searching results, the topic that focused on the research on quality control of Gendarussa crude drug was selected in this review. Gendarussa plant review articles, Gendarussa plant research articles that did not contain crude drug quality control, and crude drug quality control articles without the use of Gendarussa plants were further excluded.

1.2. Output Research & Data Extraction

The research output data used were specific and non-specific parameters that included macroscopic and microscopic tests, total ash content, extract content, water content, and compound levels. The data of subsequent research articles extracted in this case included the author, year of publication, research location, research design, and research results.

3. Results and discussion

Crude drug quality control is done by analyzing several specific and non-specific parameters. The specific parameters include macroscopic, microscopic, powder identification, water-soluble content, ethanol-soluble content, essential oil

ASUS 28/1/22 11.46

Comment [3]: Kriteria inklusi dan eksklusi tolong di state lebih jelas dan rinci

ASUS 28/1/22 13.22

Comment [4]: Penyampaian hasil dan pembahasan kurang sistematis. Sebaiknya dikelompokkan ke dalam parameter spesifik dan non

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

content, and gendarusin A levels. Gendarussa leaves harvested on average 9 months old will obtain high gendarusin A levels [19].


Macroscopic tests are conducted by observing the morphology of the leaves covering the shape, colour, and smell of the crude drug. They were single leaves with stalks of about 0.5-2 cm in size, long lancet-shaped, smooth hairless surface, and the edges of the leaves slightly edged. Then the ends of the leaves and the base of the leaves were spiky and dark green. The length of the leaves was about 5-20 cm, the width of the leaves was about 1-3.5 cm, and the leaves' bone was squinting and purple [20]. The morphology of Gendarussa plant is shown in Figure 1.



Figure 1. *J. gendarussa* Burm. f. Plants [17]

Microscopic tests are carried out using a microscope to determine the anatomical structure of the leaves. Gendarussa leaves are sorted to separate the leaves from other unwanted parts of the plant. After that, the leaves are washed to remove dust and dried at room temperature and it is called a crude drug. This is done to avoid crude drug damage due to enzymatic reactions and microbial growth [21]. Microscopic tests were conducted by anatomically observing the cross-section of the Gendarussa leaves (Figure 2).

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id



ASUS 28/1/22 13.20

Comment [6]: Resolusi gambar juga kurang baik

Figure 2. Transverse slices of Gendarussa leaf [17] with ea = upper epidermis, xi = xylem, fl = phloem, kl = kolenkim, k = cuticle, pl = palisade, rk = glandular hair, bk = coral flower, eb = lower epidermis [21]

The crude drug is smoothed until it becomes powder. Microscopic identification of powder is carried out to determine the tissues characteristics of the crude drugs. Crude drug powder was yellowish-green, had characteristic aromatic smells, and tasted bitter [21]. The identification of Gendarussa powder showed the absence of stomata, stair thickening xylem, sponge tissue, systolic, and glandular hair (Figure 3).

ASUS 28/1/22 13.26

Comment [7]: Absence? Jadi apa yang menjadi penciri mikroskopik daun ini?

Crude drug powder is soaked by citric acid at pH 3. This acidification aims to eliminate alkaloids inside the crude drug because it causes toxic effects [4]. After that, the crude drug is washed with running water to remove the remaining acid. The crude drug that does not contain alkaloids is referred to as an alkaloid-free crude drug. The content of metabolites in the crude drugs before and after acidification is different. Crude drug before acidification contains amino benzyl alkaloid derivatives that are not detected in alkaloid-free samples [22-26].

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id




Figure 3. Gendarussa leaf powder fragments [21] with se = epidermis cells, rk = glandular hair, st = stomata, sis = systolic, xy = xylem, c.o = calcium oxalate Crystals, tr = trachea, js = sponge tissue [21]


Gendarussa contains several other compounds such as essential oils, aromatic amines, stigmasterol, tannins, saponins, amino acids, lupeol, and flavonoids [27-31]. The flavonoid compound used as a marker in this plant is gendarusin A (6,8-di-C- α -L-arabinopyranosyl-4',5,7-trihydroxyflavone, or 6,8-di-C- α -L-arabinocylapigenin). Therefore, gendarusin A levels (as a major compound) in crude drugs need to be measured in quality control tests. There are other compounds such as 6-C- α -L-arabinopyranosyl-4',5,7-trihydroxy-8-C- β -

ASUS 28/1/22 13.26

Comment [8]: Resolusi gambar juga kurang bagus

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

D-cylopyranosyl-flavone or 6-C- α -L-arabinocyl-8-C- β -D-cylocilapigenin (gendarusin B), gendarusin C, gendarusin D, and gendarusin E (Figure 4) [8]. But, That compounds are minor compounds in Gendarussa leaves.



ASUS 28/1/22 13.28

Comment [9]: Struktur kimia senyawa hendaknya Digambar sendiri dengan software/aplikasi agar jelas dan seragam

Figure 4. The structure of gendarusin compounds in *Justicia gendarussa* Burm. f.

There is another compound in Gendarussa, which is an alkaloid. There are several alkaloid compounds that have been isolated from the Gendarussa leaves, namely Justidrusamida A, Justidrusamida B, Justidrusamida C, and Justidrusamida D (Figure 5) [22].

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id




Figure 5. Alkaloid structure of *Justicia gendarussa* Burm. f.

Each plant has a variety of ingredients that have different efficacy and safety [32]. The variability of chemical content in herbal medicine is influenced by various factors, such as genetic (internal) and environmental (external) that cause differences in the chemical compositions of plants, both qualitatively and quantitatively, causing differences in activities. Several studies have shown that this plant has antibacterial, antifungal, anti-HIV, anticancer, antioxidant, hepatoprotective, antiangiogenic, and anthelmintic activities [33-44] that are related to gendarusin A.

Analysis of chemical content in gendarussa leaves from several different places (Purwodadi, Pacet, and Makassar) using LC-MS instruments results, in different chemical content and different metabolite profiles [24]. In addition, there are differences in metabolites concentrations of Justidrusamid A, Justidrusamid B, and Gendarusin A from various regions in Indonesia [26]. Thirty-six metabolite profiles have been found on Gendarussa leaves using LC-HR-MS/MS [45].

Therefore, it is important to determine and know the phytochemical content contained in herbal medicine to ensure reliability, repeatability in clinical and pharmacological studies, as well as to know the bioactivity and side effects that

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

may occur. The appropriate methods for quality control of herbal products namely by metabolite profiling [46]. It can be applied as a quality control tool for phytopharmaceutical, especially if the specifications of marker compounds have not been determined [47]. Chromatography and spectroscopy techniques can be used for the complete analysis of phytopharmaceutical products by producing metabolic fingerprinting that contributes to the determination of equations and differences of various samples [48].

There were several other parameters measured in quality control, in addition to macroscopic, microscopic, and levels of major compounds in crude drugs which will be used as phytopharmaceutical drugs. That drugs must be manufactured according to the standards required to produce good quality products. The production process must comply with GMP standards. One of the most important GMP requirements is raw material [49].

Table 1 showed the quality of the Gendarussa crude drug varied by region. Based on the results of quality control from five locations (Kediri, Mojokerto-1, Mojokerto-2, Mojokerto-3, and Ponorogo), the best crude drug according to the standard test requirements was a crude drug derived from Mojokerto-3. The value of water-soluble extract of Gendarussa crude drug in Kediri and Mojokerto-1 and the value of acid-insoluble ash of Gendarussa crude drug in Mojokerto-2, Mojokerto-3, and Ponorogo have not met the requirements of *Materia Medika* Indonesia. Then, the values of ash content and water content in all five locations do not match the standards. Ash content does not meet the standard because the manufacturing process of crude drugs, especially washing, is less than perfect where there is still a lot of dirt sticking to the leaves. The high water content is caused by incomplete drying of the crude drug and the humidity level

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

of the environment during storage. Drying time is also influenced by the method of drying. Sequentially, the effective methods which are used are oven, sun & blower, blower, sun and win. But Leaf drying as a crude drug should not be directly exposed to sunlight because it will change chlorophyll compounds in the leaf [50]. In addition, the length of storage before inspection can also affect the water content [17]. The temperature in Kediri is 27.2 °C, Mojokerto is 24.8 °C, and Ponorogo is 27.8 °C. In addition, the average humidity at all locations is between 70-85%.

Contamination contained in crude drugs such as heavy metal contamination, microbes, and pesticide residues can also be measured to determine the quality of a crude drug (Table 2). The rate of contamination should not exceed the standards set by the WHO. Based on research results, microbial contamination parameters and pesticide residues at all locations are suitable with WHO requirements. The total plate count of the molds and yeast is no more than 10³ colonies/mL. The examination results of several bacteria such as *Salmonella* sp., *E. coli*, *P. aeruginosa*, and *S. aureus* and pesticide residues against the carbamate, organophosphate, and organochlorine groups showed negative results. However, heavy metal contamination (Hg and Cd) in Ponorogo does not meet the standards. The maximum heavy metal content for lead is 10 ppm, mercury is 1 ppm, cadmium is 0.02 ppm and arsenic is 10 ppm [51]. These nonspecific parameters can be applied to simple standardization of herbal medicines as QA markers if their chemical profile has not been determined [52].

4. Conclusions

The quality of *Justicia gendarussa* Burm. f. leaves crude drug is different in

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

ASUS 28/1/22 13.34

Comment [11]: Parameter mana yang dimaksud dapat menggantikan QA markers

ASUS 28/1/22 13.40

Comment [12]: Kesimpulan belum menjawab masalah penelitian

each location. This is due to differences in environmental conditions in which plants grow. Mojokerto-3 has better crude drug quality than other locations, based on physical parameters and contamination.

Keterbatasan penelitian ini apa?

Acknowledgement

The author would like to deliver our gratitude to the Head of Master Pharmaceutical Sciences Program, Faculty of Pharmacy, Universitas Airlangga for all support and to the researchers who have researched the Gendarussa plant.

References

- [1] Ayob Z, Bohari SPM, Samad AA, Jamil S. Cytotoxic activities against breast cancer cells of local *Justicia gendarussa* crude extracts. *Evidence-Based Complementary and Alternative Medicine*. 2014;1-12.
- [2] Raghu MG, Agrawal P. The isolation and structural determination of flavonoids from *Justicia gendarussa*. *Journal of Pharmacy and Biological Sciences*. 2016;11(6):73-79.
- [3] Correa GM, Alcantara AFC. Chemical constituents and biological activities of species of *Justicia* - A Review. *Brazilian Journal of Pharmacognosy*. 2012;22(1):220-238.
- [4] Prajogo BEW. Aktivitas antifertilitas flavonoid daun *Gendarussa vulgaris* Ness. (Penelitian eksperimental pencegahan penetrasi spermatozoa menci dalam proses fertilisasi *in vitro*). *Dissertation*. Surabaya: Program Pascasarjana Universitas Airlangga; 2002.

ASUS 28/1/22 13.39

Comment [13]: Darimana bisa menyimpulkan pengaruh factor lingkungan? Apakah umur tanaman, dll sudah dipastikan sama? Apakah dari 3 penelitian sudah cukup kuat untuk menyimpulkan?

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

- [5] Prajogo BEW, Juliaan F, Hinting A, Pramesti MPBD, Anggraeni M, Radjaram A, Musta'ina S. Laporan pelaksanaan uji klinik fase I. Surabaya: Universitas Airlangga dan Badan Koordinasi Keluarga Berencana Nasional; 2008.
- [6] Prajogo BEW, Juliaan F, Hinting A, Pramesti MPBD, Anggraeni M, Radjaram A, Musta'ina S. Laporan pelaksanaan uji klinik fase II. Surabaya: Universitas Airlangga dan Badan Koordinasi Keluarga Berencana Nasional; 2009.
- [7] Prajogo BEW, Juliaan F, Hinting A, Pramesti MPBD, Anggraeni M, Radjaram A, Musta'ina S. Laporan pelaksanaan uji klinik fase III. Surabaya: Universitas Airlangga dan Badan Koordinasi Keluarga Berencana Nasional; 2011.
- [8] Saifuddin A, Siswandono, Prajogo BEW. Studi *in silico* gendarusin A, B, C, D, dan E untuk prediksi absorpsi dan aktivitas terhadap hialuronidase (Ec 3.2.1.35). *Jurnal Farmasi dan Ilmu Kefarmasian Indonesia*. 2014;1(2):42-47.
- [9] Markham KR. Cara mengidentifikasi flavonoid. Bandung: ITB; 1988.
- [10] Schmidt BM, Ribnicky DM, Libsky PE, Raskin I. Revisiting the ancient concept of botanical therapeutics. *Nature Chemical Biology*. 2007;3:360-366.
- [11] Williamson EM. Synergy and other interactions in phytomedicines. *Phytomedicine*. 2001;8:401-409.
- [12] Aqil AF, Owais M. Modern phytomedicine. Turning medicinal plants into drugs. 2006.
- [13] Saller R, Reichling J. Phytotherapie. In: Melchart D, Brenke R, Dobos G,

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

- Gaisbauer M, Saller R. (Eds.). *Naturheilverfahren*. Schattauer GmbH, Stuttgart. 2002;180–293.
- [14]Kunle, Folashade O, Omoregie H, Ochogu P. Standardization of herbal medicines. *Internatonal Journal of Biodiversity and Conservation*. 2012;4(3):101–112.
- [15]Pranowo D, Noor E, Haditjaroko L, Maddu A. Karakterisasi simplisia dan ekstrak daun gedi (*Abelmoschus manihot* l.) sebagai bahan sediaan obat. *Prosiding Seminar Agroindustri dan Lokakarya Nasional FKPT-TPI*. 2015;175-184.
- [16]Kurniasari RL. Standarisasi simplisia daun *Gendarussa vulgaris* Nees. *Skripsi*. Surabaya: Fakultas Farmasi Universitas Airlangga; 2001.
- [17]Rizqa OD. Standardisasi simplisia daun *Justicia gendarussa* Burm. f. dari berbagai tempat tumbuh (Daerah Mojokerto lahan 1, Mojokerto lahan 2, dan Ponorogo). *Skripsi*. Surabaya: Fakultas Farmasi Universitas Airlangga; 2010.
- [18]Departemen Kesehatan RI. Parameter standar umum ekstrak tumbuhan obat. Jakarta: Departemen Kesehatan Republik Indonesia; 2000.
- [19]Prajogo BEW, Dudi S, Mulya HS. Analisis gendarusin A pada tanaman budidaya *Justicia gendarussa* Burm. f. *Jurnal Farmasi Indonesia*. 2007;3:176-180.
- [20]Prajogo BEW. Autentik tanaman *Justicia gendarussa* Burm. f. sebagai bahan baku obat kontrasepsi pria. Surabaya: Airlangga University Press; 2014.
- [21]Riwanti P. Perbandingan komposisi laktosa dan pati jagung pada formulasi

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

- granul kapsul ekstrak etanol 70% daun gendarusa (*Justicia gendarussa*) sebagai kontrasepsi pria. Tesis. Surabaya: Program Pascasarjana Universitas Airlangga; 2016.
- [22] Kiren Y, Deguchi J, Hirasawa Y, Morita H, Prajogo BEW. Justidrusamides A-D, new 2-aminobenzyl alcohol derivatives from *Justicia gendarussa*. *Journal of Natural Medicines*. 2014;68:754-758.
- [23] Souza LGS, Macia CS, Almeida, Lemos TLG, Ribeiro PRV, Canuto KM, Filho RB, Cistia CND, Sant'Anna CMR, Barretog FS, Moraes MO. Brazoides A-D, new alkaloids from *Justicia gendarussa* Burm. f. species. *J. Braz. Chem. Soc.* 2017;28(7):1281-1287.
- [24] Ningsih IY, Purwanti DI, Wongso S, Prajogo BEW, Indrayanto G. Metabolite profiling of *Justicia gendarussa* Burm. f. leaves using UPLC-UHR-QTOF-MS. *Sci. Pharm.* 2015;83:489–500.
- [25] Indrayoni P, Purwanti DI, Wongso S, Prajogo BEW, Indrayanto G. Metabolite profiles in various plant organs of *Justicia gendarussa* Burm. f. and its *in vitro* cultures. *Sci. Pharm.* 2016;84:555–566.
- [26] Mnatsakanyan MM, Queiroz EF, Marcourt L, Prajogo BEW, Wolfender JL. Quantitative evaluation of various preparations and extracts of the male contraceptive *Justicia gendarussa* and identification of a new aminobenzyl derivative. *Planta Med Int Open*. 2018;4:1–9.
- [27] Uddin MR, Sinha S, Hossain MA, Kaiser MA, Hossain MK, Rashid MA. Chemical and biological investigations of *Justicia gendarussa* Burm. f. *Dhaka University Journal of Pharmaceutical Sciences*. 2011;10(1):53–57.
- [28] Mustafa RA, Hamid AA, Mohamed S, Bakar FA. Total phenolic compounds,

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

- flavonoids, and radical scavenging activity of 21 selected tropical plants. *Journal of Food Science*. 2010;75(1):28–35.
- [29]Prajogo BEW, Guliet D, Ferreira QE, Wolfender JL, Zaini NC, Aucky H, Hostettmann K. Isolation of male antifertility compound in n-butanol fraction of *Justicia gendarussa* Burm. f. leaves. *Folia Medica Indonesiana*. 2009;45(1):28–31.
- [30]Chakravarty AK, Dastidar PPG, Pakrashi SC. Simple aromatic amines from *Justicia gendarussa* 13C NMR spectra of the bases and their analogues. *Tetrahedron*. 1982;38 (12):1797–1802.
- [31]Ratnasooriya WD, Deraniyagala SA, Dehigaspitiya DC. Antinociceptive activity and toxicological study of aqueous leaf extract of *Justicia gendarussa* Burm. f. in rats. *Pharmacognosy Magazine*. 2007;3(11):145–155.
- [32]Taylor DA. Botanical supplements: weeding out the health risks. *Environmental Health Perspectives*. 2004;112(13):A750-A753.
- [33]Periyanayagam K, Umamaheswari B, Suseela L, Padmini M, Ismail M. Evaluation of antiangiogenic effect of the leaves of *Justicia gendarussa* Burm. f. (Acanthaceae) by chorio allontoic membrane method. *American Journal of Infectious Diseases*. 2009;5(3):187–189.
- [34]Krishna KL, Mehta TA, Patel JA. *In-vitro* hepatoprotective activity of *Justicia gendarussa* stem on isolated rat hepatocytes. *Pharmacologyonline*. 2010;2:9–13.
- [35]Sharma KK, Saikia R, Kotoky J, Kalita JC, Devi R. Antifungal activity of *Solanum melongena* L., *Lawsonia inermis* L. and *Justicia gendarussa* B.

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

- against dermatophytes. *International Journal of PharmTech Research*. 2011;3(3):1635–1640.
- [36]Fazaludeen MF, Manickam C, Ashankyty IMA, Ahmed MQ, Beg QZ. Synthesis and characterizations of gold nanoparticles by *Justicia gendarussa* Burm. f. leaf extract. *Journal of Microbiology and Biotechnology Research*. 2012;2(1):23–34.
- [37]Saha MR, Debnath PC, Rahman MA, Islam MAU. Evaluation of *in vitro* snthelmintic activities of leaf and stem extracts of *Justicia gendarussa*. *Bangladesh Journal of Pharmacology*. 2012;7(1):50–53.
- [38]Subramanian N, Jothimanivannan C, Moorthy K. Antimicrobial activity and preliminary phytochemical screening of *Justicia gendarussa* Burm. f. against human pathogens. *Asian Journal of Pharmaceutical and Clinical Research*. 2012;5(3):229–233.
- [39]Sudhanandh VS, Arjun JK, Faisal AK, Ani MV, Renjini VS, Babu KN. *In-vitro* antibacterial screening of selected folklore Indian medicinal plants with few clinical pathogens. *Indian Journal of Pharmaceutical Education and Research*. 2012;46(2):174–178.
- [40]Kowsalya D, Sankaranarayanan S. Efficacies of bactericidal *Justicia gendarussa* extract inhibiting protein synthesis against methicilin resistant *Staphylococcus aureus*. *IOSR Journal of Pharmacy and Biological Sciences*. 2012;4(2):32–41.
- [41]Sinansari R, Bambang PE, Widiyanti P. *In silico* screening and biological evaluation of the compounds of *Justicia gendarussa* leaves extract as interferon gamma inducer: A study of anti Human Immunodeficiency Virus

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

- (HIV) development. *African Journal of Infectious Diseases*. 2018;12(1).
- [42] Hikmawanti NPE, Widiyanti P, Bambang PEW. *In vitro* anti-HIV activity of ethanol extract from gandarusa (*Justicia gendarussa* Burm. f.) leaves. *Infect Dis Rep*. 2020;doi: 10.4081/idr.2020.8730.
- [43] Prajogo BEW, Widiyanti P, Nasronudin N, Aksono B. The effect of gendarussin A isolates of *Justicia gendarussa* Burm. f. leaf in reverse transcriptase inhibition of HIV type I *in vitro*. *Indonesian Journal of Tropical and Infectious Disease*. 2015;5(5);doi: 10.20473/ijtid.v5i5.307.
- [44] Prajogo BEW, Widiyanti P, Riza H. Effect of free alkaloid and non-free alkaloid ethanol 70% extract of *Justicia gendarussa* Burm. f. leaves against reverse transcriptase HIV enzyme *in vitro* and chemical compound analysis. *Indonesian Journal of Tropical and Infectious Disease*. 2016;6(1).
- [45] Ratih GAM, Imawati MF, Nugroho RR, Purwanti DI, Wongso S, Prajogo BEW, Indrayanto G. Phytochemicals of gandarusa (*Justicia gendarussa*) and its preparations. *Natural Product Communications*. 2019;1-10.
- [46] Liang ZY, Xie P, Chan K. Quality control of herbal medicines. China: Research Center of Modernization of Chinese Medicines, Institute of Chemometrics and Intelligent Analytical Instruments; 2004.
- [47] Ratih GAM, Imawati MF, Purwanti DI, Nugroho RR, Wongso S, Prajogo B, Indrayanto G. Metabolite profiling of *Justicia gendarussa* herbal drug preparations. *Natural Product Communications*. 2019;1-5.
- [48] Xie PS. A feasible strategy for applying chromatography fingerprint to assess quality of Chinese Herbal Medicine. *Traditional Chinese Drug Research & Clinical Pharmacology*. 2001;12:141–169.

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

- [49]Prajogo BEW, Hendradi E, Riwanti P. Composition ratio of lactose and corn starch in granule capsule formulation of 70% ethanol extract *Justicia gendarussa* leaves as male contraceptive. *BROMO Conference*. 2018;183-189.
- [50]Sulasmı ES, Indriwati SE, Suarsini E. Preparation of various type of medicinal plants simplicia as material of jamu herbal. *International Conference on Education*. 2016;1014-1024.
- [51]WHO. Quality control methods for herbal materials. Switzerland: World Health Organization; 1998.
- [52]Indrayanto G. Recent development of quality control methods for herbal derived drug preparations. *Natural Product Communications*. 2018;13(12):1599-1606.

Table 1. Quality control results of crude drug of Gendarussa leaves

Author, year	Research design	Research results	Location
Kurniasari, 2001	Leaves were cleaned from impurities, dried by the wind, and made into powder	- Ash content 12.27±0.02%	Kediri
		- Acid-insoluble ash 1.79±0.02%	
- Water-soluble extract 23.29±0.17%			
- Ethanol soluble extract 7.15±0.08%			
- Water content 11.00%			
- Essential oil content 0.06%			
- Tannin levels 1.55±0.06%			
- Flavonoid levels 0.16±5.77x10 ⁻³ %			
- Gendarusin A levels (not measured)			
- Ash content 12.65±0.05%	Mojokerto-1		
- Acid-insoluble ash 1.93±0.07%			
- Water-soluble extract 18.30±0.13%			
- Ethanol soluble extract 6.90±0.07%			
- Water content 12.00%			
- Essential oil content 0.06%			
- Tannin levels 1.30%			
- Flavonoid levels 0.15±5.77x10 ⁻³ %			
- Gendarusin A levels (not			

ASUS 28/1/22 13.36

Comment [14]: Persyaratan dari standard (FHI misalnya) mana?

ASUS 28/1/22 13.35

Comment [15]: -Lokasi yang dimaksud lokasi penelitian atau lokasi tanaman?
-Waktu pengambilan tanaman

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

		measured)	
Rizqa, 2010	Leaves were cleaned from impurities, dried by wind and made into powder	- Ash content 12.02±0.10%	Mojokerto-2
		- Acid-insoluble ash 0.91±0.05%	
		- Water-soluble extract 40.92±0.17%	
		- Ethanol soluble extract 4.92±0.07%	
		- Water content 10.22±0.39%	
		- Essential oil content 0.04%	
		- Tannin levels (not measured)	
		- Flavonoid levels (not measured)	
		- Gendarusin A levels 0.15±0.01%	
		- Ash content 13.99±0.86%	Mojokerto-3
		- Acid-insoluble ash 0.67±0.06%	
		- Water-soluble extract 48.28±0.26%	
		- Ethanol soluble extract 7.40±0.45%	
		- Water content 12.89±0.20%	
		- Essential oil content 0.04%	
		- Tannin levels (not measured)	
		- Flavonoid levels (not measured)	
		- Gendarusin A levels 0.21%	
		- Ash content 13.53±1.08%	Ponorogo
		- Acid-insoluble ash 0.71±0.04%	
		- Water-soluble extract 42.76±1.29%	
		- Ethanol soluble extract 5.61±0.39%	
		- Water content 10.11±0.19%	
		- Essential oil content 0.04%	
		- Tannin levels (not measured)	
		- Flavonoid levels (not measured)	
		- Gendarusin A levels 0.27±0.01%	

Table 2. Contamination of heavy metal, microbes, and pesticides contained in crude drug of *Justicia gendarussa* Burm.f.

Author, year	Research results			Location
	Contamination of heavy metal	Contamination of microbes	Contamination of pesticides	
Rizqa, 2010	- Lead (Pb) 0.382 ppm - Mercury (Hg) 0.000 ppm - Cadmium (Cd) 0.107 ppm - Arsenic (As) 0.000 ppm	- Total Plate Count (TPC) 36700 col/ml - TPC of molds 700 col/ml - TPC of yeast 0 col/ml - <i>Salmonella sp.</i> (negative) - <i>E. coli</i> (negative) - <i>P. aeruginosa</i> (negative) - <i>S. aureus</i> (negative)	- Carbamate (negative) - Organophosphate (negative) - Organochlorine (negative)	Mojokerto-2
	- Lead (Pb) 0.427 ppm	- Total Plate Count (TPC) 26700	- Carbamate (negative) - Organophosphate	Mojokerto-3

ASUS 28/1/22 13.37

Comment [16]: Persyaratan?

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

	<ul style="list-style-type: none"> - Mercury (Hg) 0.000 ppm - Cadmium (Cd) 0.098 ppm - Arsenic (As) 0.000 ppm 	<ul style="list-style-type: none"> col/ml - TPC of molds 780 col/ml - TPC of yeast 0 col/ml - <i>Salmonella</i> sp. (negative) - <i>E. coli</i> (negative) - <i>P. aeruginosa</i> (negative) - <i>S. aureus</i> (negative) 	<ul style="list-style-type: none"> (negative) - Organochlorine (negative) 	
	<ul style="list-style-type: none"> - Lead (Pb) 0.000 ppm - Mercury (Hg) 1.829 ppm - Cadmium (Cd) 0.438 ppm - Arsenic (As) 0.000 ppm 	<ul style="list-style-type: none"> - Total Plate Count (TPC) 27700 col/ml - TPC of molds 800 col/ml - TPC of yeast 0 col/ml - <i>Salmonella</i> sp. (negative) - <i>E. coli</i> (negative) - <i>P. aeruginosa</i> (negative) - <i>S. aureus</i> (negative) 	<ul style="list-style-type: none"> - Carbamate (negative) - Organophosphate (negative) - Organochlorine (negative) 	Ponorogo
Riwanti, 2016	<ul style="list-style-type: none"> - Lead (Pb) 0.097 ppm - Mercury (Hg) 0.000 ppm - Cadmium (Cd) 0.012 ppm - Arsenic (As) <0.001 ppm 	<ul style="list-style-type: none"> - Total Plate Count (TPC) 7200 col/ml - TPC of molds 30 col/ml - TPC of yeast 4,200 col/ml - <i>Salmonella</i> sp. (negative) - <i>E. coli</i> (negative) - <i>P. aeruginosa</i> (negative) - <i>S. aureus</i> (negative) 	<ul style="list-style-type: none"> - Carbamate (negative) - Organophosphate (negative) - Organochlorine (negative) 	Mojokerto-4

Figure captions

Fig. 1. *J. gendarussa* Burm. f. Plants.

Fig. 2. Transverse slices of Gendarussa leaf.

Fig. 3. Gendarussa leaf powder fragments.

Fig. 4. The structure of gendarusin compounds in *Justicia gendarussa* Burm. f.

Fig. 5. Alkaloid structure of *Justicia gendarussa* Burm. f.

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

Mini Review: Quality Control Study of Crude Drug of *Justicia gendarussa* Burm. f. Leaves as Male Contraceptive

Rokhmatul Ummah^a, Bambang Prajogo EW^b, Retno Widyowati^{b,1*}

^aMaster of Pharmaceutical Science Program, Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia

^bDepartment of Pharmaceutical Sciences, Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia

Abstract

Justicia gendarussa Burm. f. (Gendarussa) leaves contain gendarusin A, which is a potential anti-fertility agent. Preclinical researches (*in vitro*, *in vivo*, & toxicity) and clinical trials have been carried out proving that Gendarussa leaves are safe to be used as a male contraceptive. Quality control of medicine must be carried out from the beginning when selecting crude drugs until the production process. Quality control of the crude drug is needed to maintain the quality, safety, and efficacy of its crude drug. That standardization has been done in Kediri, Mojokerto, and Ponorogo. The research output data used were specific and non-specific parameters (macroscopic and microscopic assay, determination of ash content, compound levels, etc.). The quality of crude drugs is different in each region. Crude drug from Mojokerto is better than crude drug from Kediri and Ponorogo. This is influenced by environmental difference factors on how plants grow. This review describes the differences in the quality control results of crude drug of Gendarussa from several areas. Hence, in the

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

future, the quality of this plant can be guaranteed to be used as a contraceptive drug.

Key Words: *Justicia gendarussa* Burm. f.; quality control; crude drug

1. Introduction

Justicia gendarussa Burm. f. (Gendarussa) is a plant from the Acanthaceae family spread in Indonesia and several Asian countries such as Sri Lanka, India, and Malaysia [1]. This plant is often used as a folk remedy to cure rheumatic, bronchitis, eczema, fever, etc. [2]. Apigenin and vitexin glycosides in Gendarussa can be used as anti-inflammatory and anti-tumor [3]. However, *in vitro* and *in vivo* anti-fertility tests of these leaves show competitive and reversible inhibition of the hyaluronidase enzyme in spermatozoa [4]. Therefore, this plant has the potential to be developed into a non-hormonal male contraceptive drug [5,6]. Capsule extracts from the Gendarussa as an anti-fertility have been tested in phase III clinical trials [7].

Gendarussa leaves contain potassium, flavonoids, justicin, steroids and alkaloids [4]. In addition, these leaves also contain gendarusin A (6,8-di-C- α -L-arabinopyranosyl-4',5,7-trihydroxyflavone) as a major component, gendarusin B (6-C- α -L-arabinopyranosyl-4',5,7-trihydroxy-8-C- β -D-cylopyranosyl-flavone), gendarusin C, gendarusin D, and gendarusin E as minor components [8]. Gendarusin is a compound of the more acid-resistant C-glycosides flavonoid group [9]. Analysis to determine the compounds contained in the plant is necessary to help identify the side effects of these active compounds so as to

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

improve the quality of the drug [10,11].

Quality control is a process involved in maintaining the quality and validity of the resulting products [12]. This process needs to be done in a production activity to produce phytopharmaceutical with better quality [13]. Efforts to ensure the quality and safety of the drug must be carried out from the beginning of the production process from the selection of crude drug until the end of production so that the product is then ready to be distributed in the community [14]. The economic value added in a crude drug that meets the standard is much greater than the unstandardized crude drug [15]. Standardization of crude drugs is carried out to determine the quality, safety, and efficacy requirements of the Gendarussa leaves crude drug. Standardization parameters consist of specific and non-specific parameters.

Research on the quality control test of Gendarussa crude drug has been conducted in several areas, for example, Kediri, Mojokerto, and Ponorogo [16,17]. The quality of some of these crude drugs varies. This is because drugs produced from natural materials have varying compound content. These variations occur due to several factors, such as genetic, environmental, agronomy engineering, and harvest [18]. Therefore, this review will discuss the information related to the quality control of Gendarussa crude drugs as a reference in the process of developing drugs from natural materials.

dapodik 19/3/22 09.22
Deleted: when

dapodik 19/3/22 09.24
Deleted: and then

dapodik 19/3/22 09.24
Deleted: circulated

dapodik 19/3/22 09.25
Deleted: crude drug

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

2. Method

1.1. Article Selection Criteria and Sources

The search of scientific articles were conducted on electronic databases namely PubMed, SCOPUS, ScienceDirect, Google Scholar, and Universitas Airlangga Repository. The search was limited until September 2020. The keywords used in the search were Gendarussa (*Justicia gendarussa* Burm. f.), quality control, and crude drug which were included in research articles/dissertation/thesis. From the searching results, the topic that focused on the research on quality control of Gendarussa crude drug was selected and used in this review. The inclusion criteria used as a reference in this review are scientific articles on quality control of Gendarussa with the research year no less than 2001. The exclusion criteria in this study were articles describing the quality control of Gendarussa extracts or granules.

1.2. Output Research & Data Extraction

The research output data used were specific and non-specific parameters that included macroscopic and microscopic tests, total ash content, extract content, water content, and compound levels. The data of subsequent research articles extracted in this case included the author, year of publication, research location, research design, and research results.

3. Results and discussion

Crude drug quality control is done by analyzing several specific and non-specific parameters. The specific parameters include macroscopic, microscopic,

dapodik 19/3/22 09.39

Deleted: s

dapodik 19/3/22 09.42

Deleted: Article searches

dapodik 19/3/22 09.44

Deleted: es

dapodik 19/3/22 09.44

Deleted: ere

dapodik 19/3/22 09.50

Deleted: t

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

powder identification, water-soluble content, ethanol-soluble content, essential oil content, and gendarusin A levels. Gendarussa leaves harvested on average 9 months old will obtain high gendarusin A levels [19].

Specific Parameters of Quality Control

Macroscopic tests are conducted by observing the morphology of the leaves covering the shape, color, and smell of the crude drug. They were single leaves with stalks of about 0.5-2 cm in size, long lancet-shaped, smooth hairless surface, and the edges of the leaves slightly edged. Then the ends of the leaves and the base of the leaves were spiky and dark green. The length of the leaves was about 5-20 cm, the width of the leaves was about 1-3.5 cm, and the leaves' bone was squinting and purple [20]. The morphology of Gendarussa plant is shown in Figure 1.



Figure 1. *J. gendarussa* Burm. f. plants [17]

Microscopic tests are carried out using a microscope to determine the anatomical structure of the leaves. Gendarussa leaves are sorted to separate the leaves from other unwanted parts of the plant. After that, the leaves are washed to remove dust and dried at room temperature which then is called a crude drug (Figure 2). This is done to avoid crude drug damage due to enzymatic reactions

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

dapodik 19/3/22 09.53

Deleted: colour

dapodik 19/3/22 09.55

Deleted: and it

and microbial growth [21]. Microscopic tests were conducted by anatomically observing the cross-section of the Gendarussa leaves (Figure 3).



Figure 2. Crude drugs (left) and leaf powder (right) of *J. gendarussa* Burm. f. [21]




Figure 3. Transverse slices of Gendarussa leaf [17] with ue = upper epidermis, xy = xylem, ph = phloem, k = kolenkim, c = cuticle, pl = palisade, gh = glandular hair, cf = coral flower, le = lower epidermis [21]

The crude drug is mashed until it becomes powder. Microscopic identification of powder is carried out to determine the tissues characteristics of the crude drugs. Crude drug powder was yellowish-green, had typical aromatic smells, and tasted bitter [21]. The identification of Gendarussa powder showed the presence of stomata fragments, stair thickening xylem, sponge tissue, systolic, and glandular hair (Figure 4).

dapodik 19/3/22 10.00

~~Deleted: smoothed~~

dapodik 19/3/22 10.01

~~Deleted: characteristic~~

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

Crude drug powder is soaked by citric acid at pH 3. This acidification aims to eliminate alkaloids inside the crude drug because it causes toxic effects [4]. After that, the crude drug is washed with running water to remove the remaining acid. The crude drug that does not contain alkaloids is referred to as an alkaloid-free crude drug. The content of metabolites in the crude drugs before and after acidification is different. Crude drug before acidification contains amino benzyl alkaloid derivatives that are not detected in alkaloid-free samples [22-26].





Figure 4. Gendarussa leaf powder fragments [21] with ec = epidermis cells, gh = glandular hair, s = stomata, sy = systolic, xy = xylem, co = calcium oxalate crystals, tr = trachea, st = sponge tissue [21]

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id



Gendarussa contains several other compounds such as essential oils, aromatic amines, stigmasterol, tannins, saponins, amino acids, lupeol, and flavonoids [27-31]. The flavonoid compound used as a marker in this plant is gendarusin A (6,8-di-C- α -L-arabinopyranosyl-4',5,7-trihydroxyflavone, or 6,8-di-C- α -L-arabinocylapigenin). Therefore, gendarusin A levels (as a major compound) in crude drugs need to be measured in quality control tests as marker. There are other compounds such as 6-C- α -L-arabinopyranosyl-4',5,7-trihydroxy-8-C- β -D-cylopyranosyl-flavone or 6-C- α -L-arabinocyl-8-C- β -D-cylocilapigenin (gendarusin B), gendarusin C, gendarusin D, and gendarusin E (Figure 5) [8]. **However, those** compounds are minor compounds in Gendarussa

leaves.




Gendarusin A

Gendarusin B



Gendarusin C

Gendarusin D



Gendarusin E

dapodik 19/3/22 10.12

Deleted: But

dapodik 19/3/22 10.12

Deleted: at

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

Figure 5. The structure of gendarusin compounds in *Justicia gendarussa* Burm. f.

There is another compound in Gendarussa, which is alkaloid. There are several alkaloid compounds that have been isolated from the Gendarussa leaves, namely Justidrusamida A, Justidrusamida B, Justidrusamida C, and Justidrusamida D (Figure 6) [22].




Figure 6. Alkaloid structure of *Justicia gendarussa* Burm. f.

Each plant has a variety of ingredients that have different efficacy and safety [32]. The variability of chemical content in herbal medicine is influenced by various factors, such as genetic (internal) and environmental (external) that cause differences in the chemical compositions of plants, both qualitatively and quantitatively, causing differences in activities. Several studies have shown that this plant has antibacterial, antifungal, anti-HIV, anticancer, antioxidant, hepatoprotective, antiangiogenic, and anthelmintic activities [33-44] that are related to gendarusin A.

Analysis of chemical content in gendarussa leaves from several different places (Purwodadi, Pacet, and Makassar) using LC-MS instruments results in different chemical content and different metabolite profiles [24]. In addition, there are differences in metabolites concentrations of Justridusamid A, Justridusamid B, and Gendarusin A from various regions in Indonesia [26]. Thirty-six metabolite

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

profiles have been found on Gendarussa leaves using LC-HR-MS/MS [45].

Therefore, it is important to determine and know the phytochemical content contained in herbal medicine to ensure reliability, repeatability in clinical and pharmacological studies, as well as to know the bioactivity and side effects that may occur. The appropriate method for quality control of herbal products namely by metabolite profiling [46]. It can be applied as a quality control tool for phytopharmaceutical, especially if the specifications of marker compounds have not been determined [47]. Chromatography and spectroscopy techniques can be used for the complete analysis of phytopharmaceutical products by producing metabolic fingerprinting that contributes to the determination of equations and differences of various samples [48].

There were several other parameters measured in quality control, in addition to macroscopic, microscopic, and levels of major compounds in crude drugs which will be used as phytopharmaceutical drugs. That drugs must be manufactured according to the standards required to produce good quality products. The production process must comply with GMP standards. One of the most important GMP requirements is raw material [49].

Non-specific Parameters of Quality Control

The non-specific parameters include acid-insoluble ash, ash content, water content, contamination of heavy metal, microbes, and pesticides. Table 1 showed the quality of the Gendarussa crude drug varied by region. Based on the results of quality control from five locations (Kediri, Mojokerto-1, Mojokerto-2, Mojokerto-3, and Ponorogo), the best crude drug according to the standard test requirements was a crude drug derived from Mojokerto-3. The value of water-

dapodik 19/3/22 10.17

Deleted: s

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

soluble extract of Gendarussa crude drug in Kediri and Mojokerto-1 and the value of acid-insoluble ash of Gendarussa crude drug in Mojokerto-2, Mojokerto-3, and Ponorogo have not met the requirements of Materia Medika Indonesia. Then, the values of ash content and water content in all five locations do not match the standards. Ash content does not meet the standard because the manufacturing process of crude drugs, especially washing, is less than perfect where there is still a lot of dirt sticking to the leaves. The high water content is caused by incomplete drying of the crude drug and the humidity level of the environment during storage. Drying time is also influenced by the method of drying. Sequentially, the effective methods which are used are oven, sun & blower, blower, sun and wind. But Leaf drying as a crude drug should not be directly exposed to sunlight because it will change chlorophyll compounds in the leaf [50]. In addition, the length of storage before inspection can also affect the water content [17]. The temperature in Kediri is 27.2 °C, Mojokerto is 24.8 °C, and Ponorogo is 27.8 °C. Moreover, the average humidity at all locations is between 70-85%.

Contamination contained in crude drugs such as heavy metal contamination, microbes, and pesticide residues can also be measured to determine the quality of a crude drug (Table 2). The rate of contamination should not exceed the standards set by WHO. Based on the research results, microbial contamination parameters and pesticide residues at all locations are suitable with WHO requirements. The total plate count of the molds and yeast is no more than 10³ colonies/mL. The examination results of several bacteria such as *Salmonella* sp., *E. coli*, *P. aeruginosa*, and *S. aureus* and pesticide residues against the carbamate, organophosphate, and organochlorine groups showed negative

dapodik 19/3/22 10.28

Deleted: In addition

dapodik 19/3/22 10.29

Deleted: the

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

results. However, heavy metal contamination (Hg and Cd) in Ponorogo does not meet the standards. The maximum heavy metal content for lead is 10 ppm, mercury is 1 ppm, cadmium is 0.02 ppm and arsenic is 10 ppm [51]. These nonspecific parameters can be applied to simple standardization of herbal medicines if QA marker (chemical profiles) have not been determined [52].

4. Conclusions

The quality of *Justicia gendarussa* Burm. f. leaves crude drug is different in each location. This is due to differences in environmental conditions in which plants grow, but still need other research such as the effect of plant's age for the quality of Gendarussa crude drugs. The limitation of this review is the limited number of articles that discuss about QC of Gendarussa.

Acknowledgement

The author would like to deliver gratitude to the Head of Master Pharmaceutical Sciences Program, Faculty of Pharmacy, Universitas Airlangga for all support and to the researchers who have researched the Gendarussa plant.

References

- [1] Ayob Z, Bohari SPM, Samad AA, Jamil S. Cytotoxic activities against breast cancer cells of local *Justicia gendarussa* crude extracts. *Evidence-Based Complementary and Alternative Medicine*. 2014;1-12.
- [2] Raghu MG, Agrawal P. The isolation and structural determination of flavonoids from *Justicia gendarussa*. *Journal of Pharmacy and Biological*

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

Sciences. 2016;11(6):73-79.

- [3] Correa GM, Alcantara AFC. Chemical constituents and biological activities of species of *Justicia* - A Review. *Brazilian Journal of Pharmacognosy*. 2012;22(1):220-238.
- [4] Prajogo BEW. Aktivitas antifertilitas flavonoid daun *Gendarussa vulgaris* Ness. (Penelitian eksperimental pencegahan penetrasi spermatozoa mencit dalam proses fertilisasi *in vitro*). *Dissertation*. Surabaya: Program Pascasarjana Universitas Airlangga; 2002.
- [5] Prajogo BEW, Juliaan F, Hinting A, Pramesti MPBD, Anggraeni M, Radjaram A, Musta'ina S. Laporan pelaksanaan uji klinik fase I. Surabaya: Universitas Airlangga dan Badan Koordinasi Keluarga Berencana Nasional; 2008.
- [6] Prajogo BEW, Juliaan F, Hinting A, Pramesti MPBD, Anggraeni M, Radjaram A, Musta'ina S. Laporan pelaksanaan uji klinik fase II. Surabaya: Universitas Airlangga dan Badan Koordinasi Keluarga Berencana Nasional; 2009.
- [7] Prajogo BEW, Juliaan F, Hinting A, Pramesti MPBD, Anggraeni M, Radjaram A, Musta'ina S. Laporan pelaksanaan uji klinik fase III. Surabaya: Universitas Airlangga dan Badan Koordinasi Keluarga Berencana Nasional; 2011.
- [8] Saifuddin A, Siswandono, Prajogo BEW. Studi *in silico* gendarusin A, B, C, D, dan E untuk prediksi absorpsi dan aktivitas terhadap hialuronidase (Ec 3.2.1.35). *Jurnal Farmasi dan Ilmu Kefarmasian Indonesia*. 2014;1(2):42-47.
- [9] Markham KR. Cara mengidentifikasi flavonoid. Bandung: ITB; 1988.

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

- [10]Schmidt BM, Ribnicky DM, Libsky PE, Raskin I. Revisiting the ancient concept of botanical therapeutics. *Nature Chemical Biology*. 2007;3:360-366.
- [11]Williamson EM. Synergy and other interactions in phytomedicines. *Phytomedicine*. 2001;8:401-409.
- [12]Aqil AF, Owais M. Modern phytomedicine. Turning medicinal plants into drugs. 2006.
- [13]Saller R, Reichling J. Phytotherapie. In: Melchart D, Brenke R, Dobos G, Gaisbauer M, Saller R. (Eds.). *Naturheilverfahren*. Schattauer GmbH, Stuttgart. 2002;180–293.
- [14]Kunle, Folashade O, Omoregie H, Ochogu P. Standardization of herbal medicines. *Internatonal Journal of Biodiversity and Conservation*. 2012;4(3):101–112.
- [15]Pranowo D, Noor E, Haditjaroko L, Maddu A. Karakterisasi simplisia dan ekstrak daun gedi (*Abelmoschus manihot* l.) sebagai bahan sediaan obat. *Prosiding Seminar Agroindustri dan Lokakarya Nasional FKPT-TPI*. 2015;175-184.
- [16]Kurniasari RL. Standarisasi simplisia daun *Gendarussa vulgaris* Nees. *Skripsi*. Surabaya: Fakultas Farmasi Universitas Airlangga; 2001.
- [17]Rizqa OD. Standardisasi simplisia daun *Justicia gendarussa* Burm. f. dari berbagai tempat tumbuh (Daerah Mojokerto lahan 1, Mojokerto lahan 2, dan Ponorogo). *Skripsi*. Surabaya: Fakultas Farmasi Universitas Airlangga; 2010.
- [18]Departemen Kesehatan RI. Parameter standar umum ekstrak tumbuhan

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

obat. Jakarta: Departemen Kesehatan Republik Indonesia; 2000.

- [19]Prajogo BEW, Dudi S, Mulya HS. Analisis gendarusin A pada tanaman budidaya *Justicia gendarussa* Burm. f. *Jurnal Farmasi Indonesia*. 2007;3:176-180.
- [20]Prajogo BEW. Autentik tanaman *Justicia gendarussa* Burm. f. sebagai bahan baku obat kontrasepsi pria. Surabaya: Airlangga University Press; 2014.
- [21]Riwanti P. Perbandingan komposisi laktosa dan pati jagung pada formulasi granul kapsul ekstrak etanol 70% daun gendarusa (*Justicia gendarussa*) sebagai kontrasepsi pria. *Tesis*. Surabaya: Program Pascasarjana Universitas Airlangga; 2016.
- [22]Kiren Y, Deguchi J, Hirasawa Y, Morita H, Prajogo BEW. Justidrusamides A-D, new 2-aminobenzyl alcohol derivatives from *Justicia gendarussa*. *Journal of Natural Medicines*. 2014;68:754-758.
- [23]Souza LGS, Macia CS, Almeida, Lemos TLG, Ribeiro PRV, Canuto KM, Filho RB, Cistia CND, Sant'Anna CMR, Barretog FS, Moraes MO. Brazoides A-D, new alkaloids from *Justicia gendarussa* Burm. f. species. *J. Braz. Chem. Soc.* 2017;28(7):1281-1287.
- [24]Ningsih IY, Purwanti DI, Wongso S, Prajogo BEW, Indrayanto G. Metabolite profiling of *Justicia gendarussa* Burm. f. leaves using UPLC-UHR-QTOF-MS. *Sci. Pharm.* 2015;83:489–500.
- [25]Indrayoni P, Purwanti DI, Wongso S, Prajogo BEW, Indrayanto G. Metabolite profiles in various plant organs of *Justicia gendarussa* Burm. f. and its *in vitro* cultures. *Sci. Pharm.* 2016;84:555–566.

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

- [26] Mnatsakanyan MM, Queiroz EF, Marcourt L, Prajogo BEW, Wolfender JL. Quantitative evaluation of various preparations and extracts of the male contraceptive *Justicia gendarussa* and identification of a new aminobenzyl derivative. *Planta Med Int Open*. 2018;4:1–9.
- [27] Uddin MR, Sinha S, Hossain MA, Kaiser MA, Hossain MK, Rashid MA. Chemical and biological investigations of *Justicia gendarussa* Burm. f. *Dhaka University Journal of Pharmaceutical Sciences*. 2011;10(1):53–57.
- [28] Mustafa RA, Hamid AA, Mohamed S, Bakar FA. Total phenolic compounds, flavonoids, and radical scavenging activity of 21 selected tropical plants. *Journal of Food Science*. 2010;75(1):28–35.
- [29] Prajogo BEW, Guliet D, Ferreira QE, Wolfender JL, Zaini NC, Aucky H, Hostettmann K. Isolation of male antifertility compound in n-butanol fraction of *Justicia gendarussa* Burm. f. leaves. *Folia Medica Indonesiana*. 2009;45(1):28–31.
- [30] Chakravarty AK, Dastidar PPG, Pakrashi SC. Simple aromatic amines from *Justicia gendarussa* ¹³C NMR spectra of the bases and their analogues. *Tetrahedron*. 1982;38 (12):1797–1802.
- [31] Ratnasooriya WD, Deraniyagala SA, Dehigaspitiya DC. Antinociceptive activity and toxicological study of aqueous leaf extract of *Justicia gendarussa* Burm. f. in rats. *Pharmacognosy Magazine*. 2007;3(11):145–155.
- [32] Taylor DA. Botanical supplements: weeding out the health risks. *Environmental Health Perspectives*. 2004;112(13):A750-A753.
- [33] Periyanyagam K, Umamaheswari B, Suseela L, Padmini M, Ismail M.

^{1*} Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

- Evaluation of antiangiogenic effect of the leaves of *Justicia gendarussa* Burm. f. (Acanthaceae) by chorio allontoic membrane method. *American Journal of Infectious Diseases*. 2009;5(3):187–189.
- [34] Krishna KL, Mehta TA, Patel JA. *In-vitro* hepatoprotective activity of *Justicia gendarussa* stem on isolated rat hepatocytes. *Pharmacologyonline*. 2010;2:9–13.
- [35] Sharma KK, Saikia R, Kotoky J, Kalita JC, Devi R. Antifungal activity of *Solanum melongena* L., *Lawsonia inermis* L. and *Justicia gendarussa* B. against dermatophytes. *International Journal of PharmTech Research*. 2011;3(3):1635–1640.
- [36] Fazaludeen MF, Manickam C, Ashankyty IMA, Ahmed MQ, Beg QZ. Synthesis and characterizations of gold nanoparticles by *Justicia gendarussa* Burm. f. leaf extract. *Journal of Microbiology and Biotechnology Research*. 2012;2(1):23–34.
- [37] Saha MR, Debnath PC, Rahman MA, Islam MAU. Evaluation of *in vitro* snthelmintic activities of leaf and stem extracts of *Justicia gendarussa*. *Bangladesh Journal of Pharmacology*. 2012;7(1):50–53.
- [38] Subramanian N, Jothimanivannan C, Moorthy K. Antimicrobial activity and preliminary phytochemical screening of *Justicia gendarussa* Burm. f. against human pathogens. *Asian Journal of Pharmaceutical and Clinical Research*. 2012;5(3):229–233.
- [39] Sudhanandh VS, Arjun JK, Faisal AK, Ani MV, Renjini VS, Babu KN. *In-vitro* antibacterial screening of selected folklore Indian medicinal plants with few clinical pathogens. *Indian Journal of Pharmaceutical Education and*

^{1*} Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

Research. 2012;46(2):174–178.

- [40]Kowsalya D, Sankaranarayanan S. Efficacies of bactericidal *Justicia gendarussa* extract inhibiting protein synthesis against methicilin resistant *Staphylococcus aureus*. *IOSR Journal of Pharmacy and Biological Sciences*. 2012;4(2):32–41.
- [41]Sinansari R, Bambang PE, Widiyanti P. *In silico* screening and biological evaluation of the compounds of *Justicia gendarussa* leaves extract as interferon gamma inducer: A study of anti Human Immunodeficiency Virus (HIV) development. *African Journal of Infectious Diseases*. 2018;12(1).
- [42]Hikmawanti NPE, Widiyanti P, Bambang PEW. *In vitro* anti-HIV activity of ethanol extract from gandarusa (*Justicia gendarussa* Burm. f.) leaves. *Infect Dis Rep*. 2020;doi: 10.4081/idr.2020.8730.
- [43]Prajogo BEW, Widiyanti P, Nasronudin N, Aksono B. The effect of gendarussin A isolates of *Justicia gendarussa* Burm. f. leaf in reverse transcriptase inhibition of HIV type I *in vitro*. *Indonesian Journal of Tropical and Infectious Disease*. 2015;5(5);doi: 10.20473/ijtid.v5i5.307.
- [44]Prajogo BEW, Widiyanti P, Riza H. Effect of free alkaloid and non-free alkaloid ethanol 70% extract of *Justicia gendarussa* Burm. f. leaves against reverse transcriptase HIV enzyme *in vitro* and chemical compound analysis. *Indonesian Journal of Tropical and Infectious Disease*. 2016;6(1).
- [45]Ratih GAM, Imawati MF, Nugroho RR, Purwanti DI, Wongso S, Prajogo BEW, Indrayanto G. Phytochemicals of gandarusa (*Justicia gendarussa*) and its preparations. *Natural Product Communications*. 2019;1-10.
- [46]Liang ZY, Xie P, Chan K. Quality control of herbal medicines. China:

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

Research Center of Modernization of Chinese Medicines, Institute of Chemometrics and Intelligent Analytical Instruments; 2004.

- [47]Ratih GAM, Imawati MF, Purwanti DI, Nugroho RR, Wongso S, Prajogo B, Indrayanto G. Metabolite profiling of *Justicia gendarussa* herbal drug preparations. *Natural Product Communications*. 2019;1-5.
- [48]Xie PS. A feasible strategy for applying chromatography fingerprint to assess quality of Chinese Herbal Medicine. *Traditional Chinese Drug Research & Clinical Pharmacology*. 2001;12:141–169.
- [49]Prajogo BEW, Hendradi E, Riwanti P. Composition ratio of lactose and corn starch in granule capsule formulation of 70% ethanol extract *Justicia gendarussa* leaves as male contraceptive. *BROMO Conference*. 2018;183-189.
- [50]Sulasmis ES, Indriwati SE, Suarsini E. Preparation of various type of medicinal plants simplicia as material of jamu herbal. *International Conference on Education*. 2016;1014-1024.
- [51]WHO. Quality control methods for herbal materials. Switzerland: World Health Organization; 1998.
- [52]Indrayanto G. Recent development of quality control methods for herbal derived drug preparations. *Natural Product Communications*. 2018;13(12):1599-1606.

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

Table 1. Quality control results of crude drug of Gendarussa leaves

Author (year)	Research design	Sampling time	Sampling location	Research results	Requirements*
Kurniasari (2001)	Leaves were cleaned from impurities, dried by the wind, and made into powder	2009	Kediri	<ul style="list-style-type: none"> - Ash content 12.27±0.02% - Acid-insoluble ash 1.79±0.02% - Water-soluble extract 23.29±0.17% - Ethanol soluble extract 7.15±0.08% - Water content 11.00% - Essential oil content 0.06% - Tannin levels 1.55±0.06% - Flavonoid levels 0.16±5.77×10⁻³% - Gendarusin A levels (not measured) 	<ul style="list-style-type: none"> - Ash content less than 8% - Acid-insoluble ash more than 1% - Water-soluble extract more than 24% - Ethanol-soluble extract more than 6% - Water content less than 10%
			Mojokerto-1	<ul style="list-style-type: none"> - Ash content 12.65±0.05% - Acid-insoluble ash 1.93±0.07% - Water-soluble extract 18.30±0.13% - Ethanol soluble extract 6.90±0.07% - Water content 12.00% - Essential oil content 0.06% - Tannin levels 1.30% - Flavonoid levels 0.15±5.77×10⁻³% - Gendarusin A levels (not measured) 	
Rizqa (2010)	Leaves were cleaned from impurities, dried by wind and made into powder	February, 2009	Mojokerto-2	<ul style="list-style-type: none"> - Ash content 12.02±0.10% - Acid-insoluble ash 0.91±0.05% - Water-soluble extract 40.92±0.17% - Ethanol soluble extract 4.92±0.07% - Water content 10.22±0.39% - Essential oil content 0.04% - Tannin levels (not measured) - Flavonoid levels (not measured) - Gendarusin A levels 0.15±0.01% 	
			Mojokerto-3	<ul style="list-style-type: none"> - Ash content 13.99±0.86% - Acid-insoluble ash 0.67±0.06% - Water-soluble extract 48.28±0.26% - Ethanol soluble extract 7.40±0.45% 	

*Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

		<ul style="list-style-type: none"> - Water content 12.89±0.20% - Essential oil content 0.04% - Tannin levels (not measured) - Flavonoid levels (not measured) - Gendarusin A levels 0.21%
		<ul style="list-style-type: none"> - Ash content 13.53±1.08% - Acid-insoluble ash 0.71±0.04% - Water-soluble extract 42.76±1.29% - Ethanol soluble extract 5.61±0.39%
	Ponorogo	<ul style="list-style-type: none"> - Water content 10.11±0.19% - Essential oil content 0.04% - Tannin levels (not measured) - Flavonoid levels (not measured) - Gendarusin A levels 0.27±0.01%

* based on Materia Medika Indonesia

Table 2. Contamination of heavy metal, microbes, and pesticides contained in crude drug of *Justicia gendarussa* Burm.f.

Author (year)	Sampling time	Sampling location	Research results			Requirements*
			Contamination of heavy metal	Contamination of microbes	Contamination of pesticides	
Rizqa (2010)	February, 2009	Mojokerto-2	<ul style="list-style-type: none"> - Lead (Pb) 0.382 ppm - Mercury (Hg) 0.000 ppm - Cadmium (Cd) 0.107 ppm - Arsenic (As) 0.000 ppm 	<ul style="list-style-type: none"> - Total Plate Count (TPC) 36700 col/ml - TPC of molds 700 col/ml - TPC of yeast 0 col/ml - <i>Salmonella</i> sp. (negative) - <i>E. coli</i> (negative) - <i>P. aeruginosa</i> (negative) - <i>S. aureus</i> (negative) 	<ul style="list-style-type: none"> - Carbamate (negative) - Organophosphate (negative) - Organochlorine (negative) 	<ul style="list-style-type: none"> - Lead (Pb) less than 10 ppm - Mercury (Hg) less than 0.5 ppm - Cadmium (Cd) less than 0.3 ppm - Arsenic (As) less than 5 ppm - TPC of molds or yeast less than 10³/g
		Mojokerto-3	<ul style="list-style-type: none"> - Lead (Pb) 0.427 ppm - Mercury (Hg) 0.000 ppm - Cadmium (Cd) 0.098 ppm - Arsenic (As) 	<ul style="list-style-type: none"> - Total Plate Count (TPC) 26700 col/ml - TPC of molds 780 col/ml - TPC of yeast 0 col/ml - <i>Salmonella</i> sp. (negative) 	<ul style="list-style-type: none"> - Carbamate (negative) - Organophosphate (negative) - Organochlorine (negative) 	<ul style="list-style-type: none"> - <i>E. coli</i> less than 10/g - <i>P. aeruginosa</i> negative - <i>S. aureus</i> negative

^{1*} Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

			0.000 ppm	- <i>E. coli</i> (negative) - <i>P. aeruginosa</i> (negative) - <i>S. aureus</i> (negative)	
		Ponorogo	- Lead (Pb) 0.000 ppm - Mercury (Hg) 1.829 ppm - Cadmium (Cd) 0.438 ppm - Arsenic (As) 0.000 ppm	- Total Plate Count (TPC) 27700 col/ml - TPC of molds 800 col/ml - TPC of yeast 0 col/ml - <i>Salmonella</i> sp. (negative) - <i>E. coli</i> (negative) - <i>P. aeruginosa</i> (negative) - <i>S. aureus</i> (negative)	- Carbamate (negative) - Organophosp hate (negative) - Organochlorin e (negative)
Riwanti (2016)	May, 2015	Mojokerto-4	- Lead (Pb) 0.097 ppm - Mercury (Hg) 0.000 ppm - Cadmium (Cd) 0.012 ppm - Arsenic (As) <0.001 ppm	- Total Plate Count (TPC) 7200 col/ml - TPC of molds 30 col/ml - TPC of yeast 4,200 col/ml - <i>Salmonella</i> sp. (negative) - <i>E. coli</i> (negative) - <i>P. aeruginosa</i> (negative) - <i>S. aureus</i> (negative)	- Carbamate (negative) - Organophosp hate (negative) - Organochlorin e (negative)

* based on WHO

Figure captions

Fig. 1. *J. gendarussa* Burm. f. Plants.

Fig. 2. Crude drugs and leaf powder of *J. gendarussa* Burm. f.

Fig. 3. Transverse slices of Gendarussa leaf.

Fig. 4. Gendarussa leaf powder fragments.

Fig. 5. The structure of gendarusin compounds in *Justicia gendarussa* Burm. f.

Fig. 6. Alkaloid structure of *Justicia gendarussa* Burm. f.

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

Mini Review: Quality Control Study of Crude Drug of *Justicia gendarussa* Burm. f. Leaves as Male Contraceptive

Rokhmatul Ummah^a, Bambang Prajogo EW^b, Retno Widyowati^{b,1*}

^aMaster of Pharmaceutical Science Program, Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia

^bDepartment of Pharmaceutical Sciences, Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia

Abstract

Justicia gendarussa Burm. f. (Gendarussa) leaves contain gendarusin A, which is a potential anti-fertility agent. Preclinical researches (*in vitro*, *in vivo*, & toxicity) and clinical trials have been carried out proving that Gendarussa leaves are safe to be used as a male contraceptive. Quality control of medicine must be carried out from the beginning when selecting crude drugs until the production process. Quality control of the crude drug is needed to maintain the quality, safety, and efficacy of their crude drug. That standardization has been done in Kediri, Mojokerto, and Ponorogo. The research output data used were specific and non-specific parameters (macroscopic and microscopic assay, determination of ash content, compound levels, etc.). The quality of crude drugs is different in each region. Crude drug from Mojokerto is better than crude drug from Kediri and Ponorogo. This is influenced by environmental difference factors in how plants grow. This review describes the differences in the quality control results of crude drug of Gendarussa from several areas. So in the future,

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

the quality of this plant can be guaranteed quality to be used as a contraceptive drug.

Key Words: *Justicia gendarussa* Burm. f.; quality control; crude drug

1. Introduction

Justicia gendarussa Burm. f. (Gendarussa) is a plant from the Acanthaceae family spread in Indonesia and several Asian countries such as Sri Lanka, India, and Malaysia [1]. This plant is often used as a folk remedy to cure rheumatic, bronchitis, eczema, fever, etc. [2]. Apigenin and vitexin glycosides in Gendarussa can be used as anti-inflammatory and anti-tumor [3]. However, *in vitro* and *in vivo* anti-fertility tests of these leaves show competitive and reversible inhibition of the hyaluronidase enzyme in spermatozoa [4]. Therefore, this plant has the potential to be developed into a non-hormonal male contraceptive drug [5,6]. Capsule extracts from the Gendarussa as an anti-fertility have been tested in phase III clinical trials [7].

Gendarussa leaves contain potassium, flavonoids, justicin, steroids and alkaloids [4]. In addition, these leaves also contain gendarusin A (6,8-di-C- α -L-arabinopyranosyl-4',5,7-trihydroxyflavone) as a major component, gendarusin B (6-C- α -L-arabinopyranosyl-4',5,7-trihydroxy-8-C- β -D-cylopyranosyl-flavone), gendarusin C, gendarusin D, and gendarusin E as minor components [8]. Gendarusin is a compound of the more acid-resistant C-glycosides flavonoid group [9]. Analysis to determine the compounds contained in the plant is necessary to help identify the side effects of these active compounds so as to

improve the quality of the drug [10,11].

Quality control is a process involved in maintaining the quality and validity of the resulting products [12]. This process needs to be done in a production activity to produce phytopharmaceutical with better quality [13]. Efforts to ensure the quality and safety of the drug must be carried out from the beginning of the production process when the selection of crude drug until the end of production and then the product is circulated in the community [14]. The economic value added in a crude drug that meets the standard is much greater than the unstandardized crude drug [15]. Standardization of crude drugs is carried out to determine the quality, safety, and efficacy requirements of the Gendarussa crude drug leaves. Standardization parameters consist of specific and non-specific parameters.

Research on the quality control test of Gendarussa crude drug has been conducted in several areas, for example, Kediri, Mojokerto, and Ponorogo [16,17]. The quality of some of these crude drugs varies. This is because drugs produced from natural materials have varying compound content. These variations occur due to several factors, such as genetic, environmental, agronomy engineering, and harvest [18]. Therefore, it is necessary to standardize in order to obtain crude drugs with good quality as a first step in the process of developing drugs from natural materials. This review aims to provide information related to the quality control of Gendarussa crude drugs that have been carried out by researchers, so it can be used as a reference for developing male contraceptive with the raw material of Gendarussa leaves.

2. Methods

1.1. Article Selection Criteria and Sources

Article searches were conducted on electronic databases namely PubMed, SCOPUS, ScienceDirect, Google Scholar, and Universitas Airlangga Repository. The searches were limited until September 2020. The keywords used in the search were Gendarussa (*Justicia gendarussa* Burm. f.), quality control, and crude drug in research articles/dissertation/thesis. From the searching results, the topic that focused on the research on quality control of Gendarussa crude drug was selected in this review. The inclusion criteria used as a reference in this review are scientific articles on quality control of Gendarussa, there are no limitations on research locations and years of research. The exclusion criteria in this study were articles describing the quality control of Gendarussa extract or granules.

1.2. Output Research & Data Extraction

The research output data used were specific and non-specific parameters that included macroscopic and microscopic tests, total ash content, extract content, water content, and compound levels. The data of subsequent research articles extracted in this case included the author, year of publication, research location, research design, and research results.

3. Results and discussion

Crude drug quality control is done by analyzing several specific and non-specific parameters. The specific parameters include macroscopic, microscopic,

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

powder identification, water-soluble content, ethanol-soluble content, essential oil content, and gendarusin A levels. Gendarussa leaves harvested on average 9 months old will obtain high gendarusin A levels [19].

Specific Parameters of Quality Control

Macroscopic tests are conducted by observing the morphology of the leaves covering the shape, colour, and smell of the crude drug. They were single leaves with stalks of about 0.5-2 cm in size, long lancet-shaped, smooth hairless surface, and the edges of the leaves slightly edged. Then the ends of the leaves and the base of the leaves were spiky and dark green. The length of the leaves was about 5-20 cm, the width of the leaves was about 1-3.5 cm, and the leaves' bone was squinting and purple [20]. The morphology of Gendarussa plant is shown in Figure 1.



Figure 1. *J. gendarussa* Burm. f. plants [17]

Microscopic tests are carried out using a microscope to determine the anatomical structure of the leaves. Gendarussa leaves are sorted to separate the leaves from other unwanted parts of the plant. After that, the leaves are washed to remove dust and dried at room temperature and it is called a crude drug

(Figure 2). This is done to avoid crude drug damage due to enzymatic reactions and microbial growth [21]. Microscopic tests were conducted by anatomically observing the cross-section of the Gendarussa leaves (Figure 3).



Figure 2. Crude drugs (left) and leaf powder (right) of *J. gendarussa* Burm. f. [21]



Figure 3. Transverse slices of Gendarussa leaf [17] with ea = upper epidermis, xi = xylem, fl = phloem, kl = kolenkim, k = cuticle, pl = palisade, rk = glandular hair, bk = coral flower, eb = lower epidermis [21]

The crude drug is smoothed until it becomes powder. Microscopic identification of powder is carried out to determine the tissues characteristics of the crude drugs. Crude drug powder was yellowish-green, had characteristic aromatic smells, and tasted bitter [21]. The identification of Gendarussa powder showed the presence of stomata fragments, stair thickening xylem, sponge

tissue, systolic, and glandular hair (Figure 4).

Crude drug powder is soaked by citric acid at pH 3. This acidification aims to eliminate alkaloids inside the crude drug because it causes toxic effects [4]. After that, the crude drug is washed with running water to remove the remaining acid. The crude drug that does not contain alkaloids is referred to as an alkaloid-free crude drug. The content of metabolites in the crude drugs before and after




Figure 4. Gendarussa leaf powder fragments [21] with se = epidermis cells, rk = glandular hair, st = stomata, sis = systolic, xy = xylem, c.o = calcium oxalate Crystals, tr = trachea, js = sponge tissue [21]

Gendarussa contains several other compounds such as essential oils, aromatic amines, stigmasterol, tannins, saponins, amino acids, lupeol, and flavonoids [27-31]. The flavonoid compound used as a marker in this plant is gendarusin A (6,8-di-C- α -L-arabinopyranosyl-4',5,7-trihydroxyflavone, or 6,8-di-C- α -L-arabinocylapigenin). Therefore, gendarusin A levels (as a major compound) in crude drugs need to be measured in quality control tests as marker. There are other compounds such as 6-C- α -L-arabinopyranosyl-4',5,7-trihydroxy-8-C- β -D-cylopyranosyl-flavone or 6-C- α -L-arabinocyl-8-C- β -D-cylocilapigenin (gendarusin B), gendarusin C, gendarusin D, and gendarusin E (Figure 5) [8]. But, that compounds are minor compounds in Gendarussa leaves.




Figure 5. The structure of gendarusin compounds in *Justicia gendarussa* Burm. f.

There is another compound in Gendarussa, which is an alkaloid. There are several alkaloid compounds that have been isolated from the Gendarussa leaves, namely Justidrusamida A, Justidrusamida B, Justidrusamida C, and Justidrusamida D (Figure 6) [22].




Figure 6. Alkaloid structure of *Justicia gendarussa* Burm. f.

Each plant has a variety of ingredients that have different efficacy and safety [32]. The variability of chemical content in herbal medicine is influenced by various factors, such as genetic (internal) and environmental (external) that cause differences in the chemical compositions of plants, both qualitatively and quantitatively, causing differences in activities. Several studies have shown that this plant has antibacterial, antifungal, anti-HIV, anticancer, antioxidant, hepatoprotective, antiangiogenic, and anthelmintic activities [33-44] that are related to gendarusin A.

Analysis of chemical content in gendarussa leaves from several different places (Purwodadi, Pacet, and Makassar) using LC-MS instruments results, in different chemical content and different metabolite profiles [24]. In addition, there are differences in metabolites concentrations of Justridusamid A, Justridusamid B, and Gendarusin A from various regions in Indonesia [26]. Thirty-six metabolite profiles have been found on Gendarussa leaves using LC-HR-MS/MS [45].

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

Therefore, it is important to determine and know the phytochemical content contained in herbal medicine to ensure reliability, repeatability in clinical and pharmacological studies, as well as to know the bioactivity and side effects that may occur. The appropriate methods for quality control of herbal products namely by metabolite profiling [46]. It can be applied as a quality control tool for phytopharmaceutical, especially if the specifications of marker compounds have not been determined [47]. Chromatography and spectroscopy techniques can be used for the complete analysis of phytopharmaceutical products by producing metabolic fingerprinting that contributes to the determination of equations and differences of various samples [48].

There were several other parameters measured in quality control, in addition to macroscopic, microscopic, and levels of major compounds in crude drugs which will be used as phytopharmaceutical drugs. That drugs must be manufactured according to the standards required to produce good quality products. The production process must comply with GMP standards. One of the most important GMP requirements is raw material [49].

Non-specific Parameters of Quality Control

The non-specific parameters include acid-insoluble ash, ash content, water content, contamination of heavy metal, microbes, and pesticides. Table 1 showed the quality of the Gendarussa crude drug varied by region. Based on the results of quality control from five locations (Kediri, Mojokerto-1, Mojokerto-2, Mojokerto-3, and Ponorogo), the best crude drug according to the standard test requirements was a crude drug derived from Mojokerto-3. The value of water-soluble extract of Gendarussa crude drug in Kediri and Mojokerto-1 and the

value of acid-insoluble ash of Gendarussa crude drug in Mojokerto-2, Mojokerto-3, and Ponorogo have not met the requirements of Materia Medika Indonesia. Then, the values of ash content and water content in all five locations do not match the standards. Ash content does not meet the standard because the manufacturing process of crude drugs, especially washing, is less than perfect where there is still a lot of dirt sticking to the leaves. The high water content is caused by incomplete drying of the crude drug and the humidity level of the environment during storage. Drying time is also influenced by the method of drying. Sequentially, the effective methods which are used are oven, sun & blower, blower, sun and win. But Leaf drying as a crude drug should not be directly exposed to sunlight because it will change chlorophyll compounds in the leaf [50]. In addition, the length of storage before inspection can also affect the water content [17]. The temperature in Kediri is 27.2 °C, Mojokerto is 24.8 °C, and Ponorogo is 27.8 °C. In addition, the average humidity at all locations is between 70-85%.

Contamination contained in crude drugs such as heavy metal contamination, microbes, and pesticide residues can also be measured to determine the quality of a crude drug (Table 2). The rate of contamination should not exceed the standards set by the WHO. Based on research results, microbial contamination parameters and pesticide residues at all locations are suitable with WHO requirements. The total plate count of the molds and yeast is no more than 10^3 colonies/mL. The examination results of several bacteria such as *Salmonella* sp., *E. coli*, *P. aeruginosa*, and *S. aureus* and pesticide residues against the carbamate, organophosphate, and organochlorine groups showed negative results. However, heavy metal contamination (Hg and Cd) in Ponorogo does not

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

meet the standards. The maximum heavy metal content for lead is 10 ppm, mercury is 1 ppm, cadmium is 0.02 ppm and arsenic is 10 ppm [51]. These nonspecific parameters can be applied to simple standardization of herbal medicines if QA marker (chemical profiles) have not been determined [52].

4. Conclusions

The quality of *Justicia gendarussa* Burm. f. leaves crude drug is different in each location. This is due to differences in environmental conditions in which plants grow, but still need other research such as the effect of plant age for the quality of Gendarussa crude drugs. The limitation of this review is the limited number of articles that discuss about QC of Gendarussa.

Acknowledgement

The author would like to deliver our gratitude to the Head of Master Pharmaceutical Sciences Program, Faculty of Pharmacy, Universitas Airlangga for all support and to the researchers who have researched the Gendarussa plant.

References

- [1] Ayob Z, Bohari SPM, Samad AA, Jamil S. Cytotoxic activities against breast cancer cells of local *Justicia gendarussa* crude extracts. *Evidence-Based Complementary and Alternative Medicine*. 2014;1-12.
- [2] Raghu MG, Agrawal P. The isolation and structural determination of flavonoids from *Justicia gendarussa*. *Journal of Pharmacy and Biological Sciences*. 2016;11(6):73-79.

- [3] Correa GM, Alcantara AFC. Chemical constituents and biological activities of species of *Justicia* - A Review. *Brazilian Journal of Pharmacognosy*. 2012;22(1):220-238.
- [4] Prajogo BEW. Aktivitas antifertilitas flavonoid daun *Gendarussa vulgaris* Ness. (Penelitian eksperimental pencegahan penetrasi spermatozoa mencit dalam proses fertilisasi *in vitro*). *Dissertation*. Surabaya: Program Pascasarjana Universitas Airlangga; 2002.
- [5] Prajogo BEW, Juliaan F, Hinting A, Pramesti MPBD, Anggraeni M, Radjaram A, Musta'ina S. Laporan pelaksanaan uji klinik fase I. Surabaya: Universitas Airlangga dan Badan Koordinasi Keluarga Berencana Nasional; 2008.
- [6] Prajogo BEW, Juliaan F, Hinting A, Pramesti MPBD, Anggraeni M, Radjaram A, Musta'ina S. Laporan pelaksanaan uji klinik fase II. Surabaya: Universitas Airlangga dan Badan Koordinasi Keluarga Berencana Nasional; 2009.
- [7] Prajogo BEW, Juliaan F, Hinting A, Pramesti MPBD, Anggraeni M, Radjaram A, Musta'ina S. Laporan pelaksanaan uji klinik fase III. Surabaya: Universitas Airlangga dan Badan Koordinasi Keluarga Berencana Nasional; 2011.
- [8] Saifuddin A, Siswandono, Prajogo BEW. Studi *in silico* gendarusin A, B, C, D, dan E untuk prediksi absorpsi dan aktivitas terhadap hialuronidase (Ec 3.2.1.35). *Jurnal Farmasi dan Ilmu Kefarmasian Indonesia*. 2014;1(2):42-47.
- [9] Markham KR. Cara mengidentifikasi flavonoid. Bandung: ITB; 1988.
- [10] Schmidt BM, Ribnicky DM, Libsky PE, Raskin I. Revisiting the ancient

- concept of botanical therapeutics. *Nature Chemical Biology*. 2007;3:360-366.
- [11]Williamson EM. Synergy and other interactions in phytomedicines. *Phytomedicine*. 2001;8:401-409.
- [12]Aqil AF, Owais M. Modern phytomedicine. Turning medicinal plants into drugs. 2006.
- [13]Saller R, Reichling J. Phytotherapie. In: Melchart D, Brenke R, Dobos G, Gaisbauer M, Saller R. (Eds.). *Naturheilverfahren*. Schattauer GmbH, Stuttgart. 2002;180–293.
- [14]Kunle, Folashade O, Omoregie H, Ochogu P. Standardization of herbal medicines. *Internatonal Journal of Biodiversity and Conservation*. 2012;4(3):101–112.
- [15]Pranowo D, Noor E, Haditjaroko L, Maddu A. Karakterisasi simplisia dan ekstrak daun gedi (*Abelmoschus manihot* L.) sebagai bahan sediaan obat. *Prosiding Seminar Agroindustri dan Lokakarya Nasional FKPT-TPI*. 2015;175-184.
- [16]Kurniasari RL. Standarisasi simplisia daun *Gendarussa vulgaris* Nees. *Skripsi*. Surabaya: Fakultas Farmasi Universitas Airlangga; 2001.
- [17]Rizqa OD. Standardisasi simplisia daun *Justicia gendarussa* Burm. f. dari berbagai tempat tumbuh (Daerah Mojokerto lahan 1, Mojokerto lahan 2, dan Ponorogo). *Skripsi*. Surabaya: Fakultas Farmasi Universitas Airlangga; 2010.
- [18]Departemen Kesehatan RI. Parameter standar umum ekstrak tumbuhan obat. Jakarta: Departemen Kesehatan Republik Indonesia; 2000.

^{1*}Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

- [19]Prajogo BEW, Dudi S, Mulya HS. Analisis gendarusin A pada tanaman budidaya *Justicia gendarussa* Burm. f. *Jurnal Farmasi Indonesia*. 2007;3:176-180.
- [20]Prajogo BEW. Autentik tanaman *Justicia gendarussa* Burm. f. sebagai bahan baku obat kontrasepsi pria. Surabaya: Airlangga University Press; 2014.
- [21]Riwanti P. Perbandingan komposisi laktosa dan pati jagung pada formulasi granul kapsul ekstrak etanol 70% daun gendarusa (*Justicia gendarussa*) sebagai kontrasepsi pria. *Tesis*. Surabaya: Program Pascasarjana Universitas Airlangga; 2016.
- [22]Kiren Y, Deguchi J, Hirasawa Y, Morita H, Prajogo BEW. Justidrusamides A-D, new 2-aminobenzyl alcohol derivatives from *Justicia gendarussa*. *Journal of Natural Medicines*. 2014;68:754-758.
- [23]Souza LGS, Macia CS, Almeida, Lemos TLG, Ribeiro PRV, Canuto KM, Filho RB, Cistia CND, Sant'Anna CMR, Barretog FS, Moraes MO. Brazoides A-D, new alkaloids from *Justicia gendarussa* Burm. f. species. *J. Braz. Chem. Soc*. 2017;28(7):1281-1287.
- [24]Ningsih IY, Purwanti DI, Wongso S, Prajogo BEW, Indrayanto G. Metabolite profiling of *Justicia gendarussa* Burm. f. leaves using UPLC-UHR-QTOF-MS. *Sci. Pharm*. 2015;83:489–500.
- [25]Indrayoni P, Purwanti DI, Wongso S, Prajogo BEW, Indrayanto G. Metabolite profiles in various plant organs of *Justicia gendarussa* Burm. f. and its *in vitro* cultures. *Sci. Pharm*. 2016;84:555–566.
- [26]Mnatsakanyan MM, Queiroz EF, Marcourt L, Prajogo BEW, Wolfender JL.

Quantitative evaluation of various preparations and extracts of the male contraceptive *Justicia gendarussa* and identification of a new aminobenzyl derivative. *Planta Med Int Open*. 2018;4:1–9.

[27]Uddin MR, Sinha S, Hossain MA, Kaiser MA, Hossain MK, Rashid MA. Chemical and biological investigations of *Justicia gendarussa* Burm. f. *Dhaka University Journal of Pharmaceutical Sciences*. 2011;10(1):53–57.

[28]Mustafa RA, Hamid AA, Mohamed S, Bakar FA. Total phenolic compounds, flavonoids, and radical scavenging activity of 21 selected tropical plants. *Journal of Food Science*. 2010;75(1):28–35.

[29]Prajogo BEW, Guliet D, Ferreira QE, Wolfender JL, Zaini NC, Aucky H, Hostettmann K. Isolation of male antifertility compound in n-butanol fraction of *Justicia gendarussa* Burm. f. leaves. *Folia Medica Indonesiana*. 2009;45(1):28–31.

[30]Chakravarty AK, Dastidar PPG, Pakrashi SC. Simple aromatic amines from *Justicia gendarussa* 13C NMR spectra of the bases and their analogues. *Tetrahedron*. 1982;38 (12):1797–1802.

[31]Ratnasooriya WD, Deraniyagala SA, Dehigaspitiya DC. Antinociceptive activity and toxicological study of aqueous leaf extract of *Justicia gendarussa* Burm. f. in rats. *Pharmacognosy Magazine*. 2007;3(11):145–155.

[32]Taylor DA. Botanical supplements: weeding out the health risks. *Environmental Health Perspectives*. 2004;112(13):A750-A753.

[33]Periyanayagam K, Umamaheswari B, Suseela L, Padmini M, Ismail M. Evaluation of antiangiogenic effect of the leaves of *Justicia gendarussa*

- Burm. f. (Acanthaceae) by chrio allontoic membrane method. *American Journal of Infectious Diseases*. 2009;5(3):187–189.
- [34] Krishna KL, Mehta TA, Patel JA. *In-vitro* hepatoprotective activity of *Justicia gendarussa* stem on isolated rat hepatocytes. *Pharmacologyonline*. 2010;2:9–13.
- [35] Sharma KK, Saikia R, Kotoky J, Kalita JC, Devi R. Antifungal activity of *Solanum melongena* L., *Lawsonia inermis* L. and *Justicia gendarussa* B. against dermatophytes. *International Journal of PharmTech Research*. 2011;3(3):1635–1640.
- [36] Fazaludeen MF, Manickam C, Ashankyty IMA, Ahmed MQ, Beg QZ. Synthesis and characterizations of gold nanoparticles by *Justicia gendarussa* Burm. f. leaf extract. *Journal of Microbiology and Biotechnology Research*. 2012;2(1):23–34.
- [37] Saha MR, Debnath PC, Rahman MA, Islam MAU. Evaluation of *in vitro* snthelmintic activities of leaf and stem extracts of *Justicia gendarussa*. *Bangladesh Journal of Pharmacology*. 2012;7(1):50–53.
- [38] Subramanian N, Jothimanivannan C, Moorthy K. Antimicrobial activity and preliminary phytochemical screening of *Justicia gendarussa* Burm. f. against human pathogens. *Asian Journal of Pharmaceutical and Clinical Research*. 2012;5(3):229–233.
- [39] Sudhanandh VS, Arjun JK, Faisal AK, Ani MV, Renjini VS, Babu KN. *In-vitro* antibacterial screening of selected folklore Indian medicinal plants with few clinical pathogens. *Indian Journal of Pharmaceutical Education and Research*. 2012;46(2):174–178.

- [40]Kowsalya D, Sankaranarayanan S. Efficacies of bactericidal *Justicia gendarussa* extract inhibiting protein synthesis against methicilin resistant *Staphylococcus aureus*. *IOSR Journal of Pharmacy and Biological Sciences*. 2012;4(2):32–41.
- [41]Sinansari R, Bambang PE, Widiyanti P. *In silico* screening and biological evaluation of the compounds of *Justicia gendarussa* leaves extract as interferon gamma inducer: A study of anti Human Immunodeficiency Virus (HIV) development. *African Journal of Infectious Diseases*. 2018;12(1).
- [42]Hikmawanti NPE, Widiyanti P, Bambang PEW. *In vitro* anti-HIV activity of ethanol extract from gandarusa (*Justicia gendarussa* Burm. f.) leaves. *Infect Dis Rep*. 2020;doi: 10.4081/idr.2020.8730.
- [43]Prajogo BEW, Widiyanti P, Nasronudin N, Aksono B. The effect of gendarussin A isolates of *Justicia gendarussa* Burm. f. leaf in reverse transcriptase inhibition of HIV type I *in vitro*. *Indonesian Journal of Tropical and Infectious Disease*. 2015;5(5);doi: 10.20473/ijtid.v5i5.307.
- [44]Prajogo BEW, Widiyanti P, Riza H. Effect of free alkaloid and non-free alkaloid ethanol 70% extract of *Justicia gendarussa* Burm. f. leaves against reverse transcriptase HIV enzyme *in vitro* and chemical compound analysis. *Indonesian Journal of Tropical and Infectious Disease*. 2016;6(1).
- [45]Ratih GAM, Imawati MF, Nugroho RR, Purwanti DI, Wongso S, Prajogo BEW, Indrayanto G. Phytochemicals of gandarusa (*Justicia gendarussa*) and its preparations. *Natural Product Communications*. 2019;1-10.
- [46]Liang ZY, Xie P, Chan K. Quality control of herbal medicines. China: Research Center of Modernization of Chinese Medicines, Institute of

Chemometrics and Intelligent Analytical Instruments; 2004.

- [47]Ratih GAM, Imawati MF, Purwanti DI, Nugroho RR, Wongso S, Prajogo B, Indrayanto G. Metabolite profiling of *Justicia gendarussa* herbal drug preparations. *Natural Product Communications*. 2019;1-5.
- [48]Xie PS. A feasible strategy for applying chromatography fingerprint to assess quality of Chinese Herbal Medicine. *Traditional Chinese Drug Research & Clinical Pharmacology*. 2001;12:141–169.
- [49]Prajogo BEW, Hendradi E, Riwanti P. Composition ratio of lactose and corn starch in granule capsule formulation of 70% ethanol extract *Justicia gendarussa* leaves as male contraceptive. *BROMO Conference*. 2018;183-189.
- [50]Sulasmi ES, Indriwati SE, Suarsini E. Preparation of various type of medicinal plants simplicia as material of jamu herbal. *International Conference on Education*. 2016;1014-1024.
- [51]WHO. Quality control methods for herbal materials. Switzerland: World Health Organization; 1998.
- [52]Indrayanto G. Recent development of quality control methods for herbal derived drug preparations. *Natural Product Communications*. 2018;13(12):1599-1606.

Table 1. Quality control results of crude drug of Gendarussa leaves

Author (year)	Research design	Sampling time	Sampling location	Research results	Requirements*
Kurniasari (2001)	Leaves were cleaned from impurities, dried by the wind, and made into powder	2009	Kediri	<ul style="list-style-type: none"> - Ash content 12.27±0.02% - Acid-insoluble ash 1.79±0.02% - Water-soluble extract 23.29±0.17% - Ethanol soluble extract 7.15±0.08% - Water content 11.00% - Essential oil content 0.06% - Tannin levels 1.55±0.06% - Flavonoid levels 0.16±5.77x10⁻³% - Gendarusin A levels (not measured) 	
			Mojokerto-1	<ul style="list-style-type: none"> - Ash content 12.65±0.05% - Acid-insoluble ash 1.93±0.07% - Water-soluble extract 18.30±0.13% - Ethanol soluble extract 6.90±0.07% - Water content 12.00% - Essential oil content 0.06% - Tannin levels 1.30% - Flavonoid levels 0.15±5.77x10⁻³% - Gendarusin A levels (not measured) 	<ul style="list-style-type: none"> - Ash content less than 8% - Acid-insoluble ash more than 1% - Water-soluble extract more than 24%
		Mojokerto-2	<ul style="list-style-type: none"> - Ash content 12.02±0.10% - Acid-insoluble ash 0.91±0.05% - Water-soluble extract 40.92±0.17% - Ethanol soluble extract 4.92±0.07% - Water content 10.22±0.39% - Essential oil content 0.04% - Tannin levels (not measured) - Flavonoid levels (not measured) - Gendarusin A levels 0.15±0.01% 	<ul style="list-style-type: none"> - Ethanol-soluble extract more than 6% - Water content less than 10% 	
Rizqa (2010)	Leaves were cleaned from impurities, dried by wind and made into powder	February, 2009	Mojokerto-3	<ul style="list-style-type: none"> - Ash content 13.99±0.86% - Acid-insoluble ash 0.67±0.06% - Water-soluble extract 48.28±0.26% - Ethanol soluble extract 7.40±0.45% - Water content 12.89±0.20% - Essential oil content 0.04% 	

*Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

				<ul style="list-style-type: none"> - Tannin levels (not measured) - Flavonoid levels (not measured) - Gendarusin A levels 0.21%
				<ul style="list-style-type: none"> - Ash content 13.53±1.08% - Acid-insoluble ash 0.71±0.04% - Water-soluble extract 42.76±1.29% - Ethanol soluble extract 5.61±0.39%
		Ponorogo		<ul style="list-style-type: none"> - Water content 10.11±0.19% - Essential oil content 0.04% - Tannin levels (not measured) - Flavonoid levels (not measured) - Gendarusin A levels 0.27±0.01%

* based on Materia Medika Indonesia

Table 2. Contamination of heavy metal, microbes, and pesticides contained in crude drug of *Justicia gendarussa* Burm.f.

Author (year)	Sampling time	Sampling location	Research results			Requirements*
			Contamination of heavy metal	Contamination of microbes	Contamination of pesticides	
Rizqa (2010)	February, 2009	Mojokerto-2	<ul style="list-style-type: none"> - Lead (Pb) 0.382 ppm - Mercury (Hg) 0.000 ppm - Cadmium (Cd) 0.107 ppm - Arsenic (As) 0.000 ppm 	<ul style="list-style-type: none"> - Total Plate Count (TPC) 36700 col/ml - TPC of molds 700 col/ml - TPC of yeast 0 col/ml - <i>Salmonella</i> sp. (negative) - <i>E. coli</i> (negative) - <i>P. aeruginosa</i> (negative) - <i>S. aureus</i> (negative) 	<ul style="list-style-type: none"> - Carbamate (negative) - Organophosp hate (negative) - Organochlorine (negative) 	<ul style="list-style-type: none"> - Lead (Pb) less than 10 ppm - Mercury (Hg) less than 0.5 ppm - Cadmium (Cd) less than 0.3 ppm - Arsenin (As) less than 5 ppm
		Mojokerto-3	<ul style="list-style-type: none"> - Lead (Pb) 0.427 ppm - Mercury (Hg) 0.000 ppm - Cadmium (Cd) 0.098 ppm - Arsenic (As) 0.000 ppm 	<ul style="list-style-type: none"> - Total Plate Count (TPC) 26700 col/ml - TPC of molds 780 col/ml - TPC of yeast 0 col/ml - <i>Salmonella</i> sp. (negative) - <i>E. coli</i> (negative) - <i>P. aeruginosa</i> (negative) 	<ul style="list-style-type: none"> - Carbamate (negative) - Organophosp hate (negative) - Organochlorine (negative) 	<ul style="list-style-type: none"> - TPC of molds or yeast less than 10³/g - <i>E. coli</i> less than 10/g - <i>P. aeruginosa</i> negative - <i>S. aureus</i> negative

¹*Corresponding author at e-mail address: rr-retno-w@ff.unair.ac.id

				- <i>S. aureus</i> (negative)
		Ponorogo	- Lead (Pb) 0.000 ppm - Mercury (Hg) 1.829 ppm - Cadmium (Cd) 0.438 ppm - Arsenic (As) 0.000 ppm	- Total Plate Count (TPC) 27700 col/ml - TPC of molds 800 col/ml - TPC of yeast 0 col/ml - <i>Salmonella</i> sp. (negative) - <i>E. coli</i> (negative) - <i>P. aeruginosa</i> (negative) - <i>S. aureus</i> (negative) - Carbamate (negative) - Organophosp hate (negative) - Organochlorin e (negative)
Riwanti (2016)	May, 2015	Mojokerto-4	- Lead (Pb) 0.097 ppm - Mercury (Hg) 0.000 ppm - Cadmium (Cd) 0.012 ppm - Arsenic (As) <0.001 ppm	- Total Plate Count (TPC) 7200 col/ml - TPC of molds 30 col/ml - TPC of yeast 4,200 col/ml - <i>Salmonella</i> sp. (negative) - <i>E. coli</i> (negative) - <i>P. aeruginosa</i> (negative) - <i>S. aureus</i> (negative) - Carbamate (negative) - Organophosp hate (negative) - Organochlorin e (negative)

* based on WHO

Figure captions

Fig. 1. *J. gendarussa* Burm. f. Plants.

Fig. 2. Transverse slices of Gendarussa leaf.

Fig. 3. Gendarussa leaf powder fragments.

Fig. 4. The structure of gendarusin compounds in *Justicia gendarussa* Burm. f.

Fig. 5. Alkaloid structure of *Justicia gendarussa* Burm. f.



rr retno widyowati <rr-retno-w@ff.unair.ac.id>

[MPI] Editor Decision

2 messages

Kartini <kartini@staff.ubaya.ac.id> Fri, Mar 18, 2022 at 10:04 AM
To: Rokhmatul Ummah <rokhmatul.ummah-2018@ff.unair.ac.id>, Bambang Prajogo <prajogo_ew@yahoo.com>, Retno Widyowati <rr-retno-w@ff.unair.ac.id>

Rokhmatul Ummah, Bambang Prajogo, Retno Widyowati:

We have reached a decision regarding your submission to MPI (Media Pharmaceutica Indonesiana), "Review: Quality Control Study of Crude Drug of Justicia gendarussa Burm. f. Leaves as Male Contraceptive".

Our decision is to: Accept Submission

Kartini
kartini@staff.ubaya.ac.id

[MPI \(Media Pharmaceutica Indonesiana\)](#)

rr retno widyowati <rr-retno-w@ff.unair.ac.id> Tue, Mar 22, 2022 at 2:22 PM
To: Kartini <kartini@staff.ubaya.ac.id>


Selamat siang,


Terima kasih atas informasi dan kesempatan pada artikel kami yang akan dipublikasi pada MPI. Berikut ini saya kirimkan hasil proofread yang telah kami lakukan dan bukti proofread. Terima kasih.

Salam,

Rr. Retno Widyowati, PhD
[Quoted text hidden]

2 attachments

 **2 4838-Article Text-13054-1-15-20220226 (Proffread).doc**
965K

 **2 4838-Article Text-13054-1-15-20220226-final.doc**
962K



rr retno widyowati <rr-retno-w@ff.unair.ac.id>

[MPI] Editor Decision

1 message

Kartini <kartini@staff.ubaya.ac.id>

Wed, Mar 23, 2022 at 9:03 PM

To: Rokhmatul Ummah <rokhmatul.ummah-2018@ff.unair.ac.id>, Bambang Prajogo <prajogo_ew@yahoo.com>, Retno Widyowati <rr-retno-w@ff.unair.ac.id>

Rokhmatul Ummah, Bambang Prajogo, Retno Widyowati:

The editing of your submission, "Review: Quality Control Study of Crude Drug of *Justicia gendarussa* Burm. f. Leaves as Male Contraceptive," is complete. We are now sending it to production.

Submission URL: <https://journal.ubaya.ac.id/index.php/MPI/authorDashboard/submission/4838>

Kartini

kartini@staff.ubaya.ac.id

[MPI \(Media Pharmaceutica Indonesiana\)](#)



rr retno widyowati <rr-retno-w@ff.unair.ac.id>

[MPI] New notification from MPI (Media Pharmaceutica Indonesiana)

1 message

Kartini <kartini@staff.ubaya.ac.id>

Wed, Jun 1, 2022 at 12:40 PM

Reply-To: Kartini <mpi@unit.ubaya.ac.id>

To: Retno Widyowati <rr-retno-w@ff.unair.ac.id>

You have a new notification from MPI (Media Pharmaceutica Indonesiana):

You have been added to a discussion titled "galley proof check" regarding the submission "Review: Quality Control Study of Crude Drug of *Justicia gendarussa* Burm. f. Leaves as Male Contraceptive".

Link: <https://journal.ubaya.ac.id/index.php/MPI/authorDashboard/submission/4838>

Kartini

MPI (Media Pharmaceutica Indonesiana)