

[ISSUES AND ARTICLE](#) +[ABOUT THIS JOURN](#) +[FOR AUTHORS](#) +[SUBSCRIBE](#) +**OPEN ACCESS**

YOUR ACCOUNT

[Update your registration details](#)[Update your areas of interest](#)[Modify your password](#)

YOUR ORDERS

[Order to be completed](#)[Completed orders](#)

SHOPPING BASKET

Items: 0

Total amount: € 0,00

[Order details and checkout](#)

HOW TO ORDER

[Journals](#)[Books](#)

YOUR SUBSCRIPTIONS

[Activate](#)[View](#)[Contact subscription department](#)

YOUR ARTICLES

[View](#)

YOUR EBOOKS

[View](#)

COUPON

[Enable your coupon](#)

ACCESSIBILITY

[Standard viewing](#)[Larger font](#)[Text only](#)[High-contrast layout](#)**Italian Journal of Vascular and Endovascular Surgery 2023 March;30(1)**

ORIGINAL ARTICLE

Italian Journal of Vascular and Endovascular Surgery 2023 March;30(1):1-6

How to improve results in carotid surgery. Analysis of a 20-year case history of more than 7000 operations

Brigida BIANCOFIORE *, Serena DE BLASIS, Roberto FALSO, Lapo DOLFI, Marta PANELLA, Rossella DI DOMENICO, Walter DORIGO, Raffaele PULLI, Carlo PRATESI

[Abstract](#) [HTML](#) [PDF](#)

ORIGINAL ARTICLE

Italian Journal of Vascular and Endovascular Surgery 2023 March;30(1):7-10

Comparative analysis of different endovascular modalities to treat splenic artery aneurysms

Giulia BERTAGNA, Nicola TROISI *, Lorenzo TORRI, Francesco CANOVARO, Francesca TOMEI, Daniele ADAMI, Roberto CIONI, Orsola PERRONE, Raffaella BERCHIOLLI

[Abstract](#) [HTML](#) [PDF](#)

ORIGINAL ARTICLE


Italian Journal of Vascular and Endovascular Surgery 2023 March;30(1):11-4

Carotid bifurcation stenosis: functional importance in physical exercises

Marina FEDORINA *, Igor DAVYDKIN, Giuseppe GALATI, Giuseppe BIONDI-ZOCCAI, Olga GERMANOVA

[Abstract](#) [HTML](#) [PDF](#)

REVIEW



JOURNAL TOOLS

[Publishing options](#)[eTOC](#)[To subscribe](#)[Submit an article](#)[Recommend to your librarian](#)



Copyright © 2023 Edizioni Minerva Medica

[Corporate information](#) [Privacy policy](#) [Terms and conditions](#)



[ISSUES AND ARTICLE](#) +[ABOUT THIS JOURNA](#) +[FOR AUTHORS](#) +[SUBSCRIBE](#) +**OPEN ACCESS**

YOUR ACCOUNT

[Update your registration details](#)[Update your areas of interest](#)[Modify your password](#)

YOUR ORDERS

[Order to be completed](#)[Completed orders](#)[SHOPPING BASKET](#)

Items: 0

Total amount: € 0,00

[Order details and checkout](#)

HOW TO ORDER

[Journals](#)[Books](#)

YOUR SUBSCRIPTIONS

[Activate](#)[View](#)[Contact subscription department](#)

YOUR ARTICLES

[View](#)

YOUR EBOOKS

[View](#)

COUPON

[Enable your coupon](#)

ACCESSIBILITY

[Standard viewing](#)[Larger font](#)[Text only](#)[High-contrast layout](#)

ITALIAN JOURNAL OF VASCULAR AND ENDOVASCULAR SURGERY

Hybrid journal

Official Journal of the **Italian Society of Vascular and Endovascular Surgery**

Indexed/Abstracted in: EMBASE, Emerging Sources Citation Index, Scopus

Quarterly

pISSN 1824-4777

eISSN 1827-1847

EDITORIAL BOARD**Chief Editor**

Mauro Gargiulo

University of Bologna, Bologna, Italy

Associate Editors

Roberto Chiesa

"Vita-Salute" University, Scientific Institute San Raffaele, Milan, Italy

Carlo Pratesi


University of Florence, Florence, Italy

Flavio Peinetti

"U. Parini" Hospital, Aosta, Italy

Carlo Setacci

University of Florence, Florence, Italy



JOURNAL TOOLS

[Publishing options](#)[eTOC](#)[To subscribe](#)[Submit an article](#)[Recommend to your librarian](#)

Angelo Argenterì
University of Padua, Padua, Italy

Enrico Ascher
Vascular Institute of New York, Brooklyn, New York, USA

Dittmar Böckler
University of Heidelberg, Heidelberg, Germany

Stefano Camparini
"G. Brotzu" Hospital, Cagliari, Italy

Piergiorgio Cao
Azienda Ospedaliera San Camillo - Forlanini, Rome, Italy

Patrizio Castelli
University of Insubria School of Medicine, Varese, Italy

Gioachino Coppi
University of Modena and Reggio Emilia, Modena, Italy

Alberto Cremonesi
Maria Cecilia Hospital, Ravenna, Italy

Gianmarco De Donato
University of Siena, Siena, Italy

Jean-Paul De Vries
University Medical Centre Groningen, Groningen, the Netherlands

Robert S. Dieter
Loyola University Medical Center, Maywood, Illinois, USA

Walter Dorigo
University of Florence, Florence, Italy

Vittorio Dorrucchi
University of Verona, Verona, Italy

Gianluca Faggioli
University of Bologna, Bologna, Italy

Antonio Freyrie
University of Parma, Parma, Italy

Paolo Frigatti
University of Padua, Padua, Italy

Andrea Gaggiano
Ospedale Mauriziano Umberto I, Turin, Italy

Enrico Gallitto
University of Bologna, Bologna, Italy

Peter Gloviczki
Mayo Clinic, Rochester, Minnesota, USA

Bruno Gossetti
Sapienza University, Rome, Italy

Franco Grego
University of Padua, Padua, Italy

Glenn M. La Muraglia
Massachusetts General Hospital, Boston, Massachusetts, USA

George Hamilton
UCL Medical School, London, United Kingdom

Stéphan Haulon
Lille University, Lille, France

Hans-Henning Eckstein
TUM: The Entrepreneurial University, München, Germany

Armando Lobato
Vascular & Endovascular Surgery Institute, Sao Paulo, Brazil

Martin Malina
Skåne University, Malmö, Sweden

Nicola Mangialardi
San Filippo Neri Hospital, Rome, Italy

Armando Mansilha
Hospital de São João, Porto, Portugal

Enrico Marone
University of Pavia, Pavia, Italy

Germano Melissano
San Raffaele Hospital, Milan, Italy

Stefano Michelagnoli
Ospedale S. Giovanni di Dio, Florence, Italy

Giovanni Nano
IRCCS Policlinico San Donato, Milan, Italy

Franco Nessi
Ospedale Mauriziano Umberto I, Turin, Italy

Domenico Palombo
IRCCS Policlinico San Martino Hospital, Genoa, Italy

Paolo Perini
University of Parma, Parma, Italy

Rodolfo Pini
University of Bologna, Bologna, Italy

Raffaele Pulli
University of Bari, Bari, Italy

Claudio Rabbia
A.O.U. Città della Salute e della Scienza di Torino, Turin, Italy

T. Resh
Skåne Univerisyt, Malmö, Sweden

Jean-Baptiste Ricco
University of Poitiers, Poitiers, France

Pietro Rispoli
University of Turin, Turin, Italy

Carlo Ruotolo
AORN Antonio Cardarelli, Naples, Italy

Francesco Speziale
Sapienza University, Rome, Italy

Francesco Spinelli
Campus Bio-Medico University, Rome, Italy

Maurizio Taurino
Sapienza University, Rome, Italy

Giovanni Torsello
University of Muenster, Münster, Germany

Santi Trimarchi
IRCCS Ca' Granda Foundation Maggiore Policlinico Hospital, Milan, Italy

Carlos Vaquero Puerta
University of Valladolid, Valladolid, Spain

Frank Vermassen
Ghent University Hospital, Ghent, Belgium

Fabio Verzini
A.O.U. Città della Salute e della Scienza di Torino, Turin, Italy

Editorial Secretary

Aaron Fargion
University of Florence, Florence, Italy

Bianca Pane
University of Genoa, Genoa, Italy

Gabriele Piffaretti
University of Insubria School of Medicine, Varese, Italy

Rodolfo Pini
University of Bologna, Bologna, Italy

Past Chief Editor

Andrea Stella
University of Bologna, Bologna, Italy

Founder Editor

M. D'Addato
University of Bologna, Bologna, Italy

Managing Editor

Alberto Oliaro
University of Turin, Turin, Italy

[top of page](#)



Copyright © 2023 Edizioni Minerva Medica

[Corporate information](#) [Privacy policy](#) [Terms and conditions](#)





SJR

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)

Italian Journal of Vascular and Endovascular Surgery

COUNTRY

Italy

Universities and
research institutions in
Italy

Media Ranking in Italy

SUBJECT AREA AND CATEGORY

Medicine
Cardiology and
Cardiovascular
Medicine
Surgery

PUBLISHER

Edizioni Minerva Medica
S.p.A.

H-INDEX

7

PUBLICATION TYPE

Journals

ISSN

18244777, 18271847

COVERAGE

2004-2021

INFORMATION

[Homepage](#)[How to publish in this
journal](#)journals6.dept@minervamedica.it

SCOPE


The Italian Journal of Vascular and Endovascular Surgery publishes scientific papers on vascular surgery. Manuscripts may be submitted in the form of editorials, original articles, review articles, case reports, therapeutical notes, special articles and letters to the Editor.

[Join the conversation about this journal](#)


FIND SIMILAR JOURNALS ?

<p>1 Vascular</p> <p>GBR</p> <p>93% similarity</p>	<p>2 Vascular and Endovascular Surgery</p> <p>USA</p> <p>91% similarity</p>	<p>3 Annals of Vascular Surgery</p> <p>USA</p> <p>89% similarity</p>	<p>4 Angiologia</p> <p>ESP</p> <p>8 s</p>
--	---	--	---


SJR




Total Documents




Total Cites Self-Cites



Citations per document





External Cites per Doc Cites per Doc



% International Collaboration


Citable documents

Non-citable documents



Cited documents

Uncited documents



Italian Journal of Vascular and Endovascular Surgery

Q4 Cardiology and Cardiovascular Medicine
best quartile

SJR 2021
0.14

powered by scimagojr.com


Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```

SCImago Graphica

Explore, visually communicate and make sense of data with our **new data visualization tool.**



Metrics based on Scopus® data as of April 2022



Loading comments...

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2022. Data Source: Scopus®



[Cookie settings](#)

[Cookie policy](#)

REVIEW

Patency durability of non-tunneled hemodialysis' catheter in adult patients with chronic kidney disease stadium 5 with diabetes mellitus and non-diabetes mellitus

I Gusti PUTRA ¹, Heroe SOEBROTO ² *, Yan E. SEMBIRING ², Artaria TJEMPAKASARI ³

¹Department of Thoracic, Cardiac and Vascular Surgery, Dr. Soetomo Academic General Hospital, Airlangga University, Surabaya, Indonesia; ²Department of Thoracic, Cardiac and Vascular Surgery, Faculty of Medicine, Dr. Soetomo General Academic Hospital, Airlangga University, Surabaya, Indonesia; ³Department of Internal Medicine, Faculty of Medicine, Dr. Soetomo Academic General Hospital, Airlangga University, Surabaya, Indonesia

*Corresponding author: Heroe Soebroto, Department of Thoracic, Cardiac and Vascular Surgery, Faculty of Medicine, Dr. Soetomo General Academic Hospital, Airlangga University, Surabaya, Indonesia. E-mail: heroesurgeon@yahoo.com

ABSTRACT

INTRODUCTION: Chronic kidney disease (CKD) is a condition of kidney that causes the need for kidney replacement therapy in the form of hemodialysis or kidney transplantation. One of the prerequisites and things that are very essential in the care of patients with hemodialysis is the existence of adequate hemodialysis vascular access. Temporary and short-term use usually use a non-tunneled hemodialysis catheter (NTHC) often be used in emergency cases. In patient with diabetic, complication often occurs as the occurrence of thrombosis in patients with the use of vascular access in DM is twice as high as those without comorbid DM.

EVIDENCE ACQUISITION: Literatures reviewed in our study were gathered from PubMed and Google Scholar. Advanced search was conducted on both Google Scholar and PubMed Builder, with the keywords used were ("Patency" AND "catheter" AND "durability" AND "hemodialysis;" "non-tunneled hemodialysis catheter" AND "chronic kidney disease" AND "diabetic") also words "patency of tunneled catheters in hemodialysis patients;" "patency of tunneled catheters in diabetic chronic kidney disease underwent hemodialysis". We reviewed the articles cited within the literatures to broaden the search results.

EVIDENCE SYNTHESIS: Non-tunneled hemodialysis catheters (NTHCs) are typically used when vascular access is required for urgent renal replacement therapy (RRT). Metabolic conditions in DM patients damage blood vessels and can induce abnormalities such as thrombosis and can be prevented by administering antithrombotic prophylaxis Maintenance of patency in diabetic patients related to the durability of NTHC by administration of thrombolytic agents or anticoagulant.

CONCLUSIONS: A non-tunneled hemodialysis catheter may be used as a temporary catheter while waiting for a placement of an arteriovenous fistula. However, NTDCs have several disadvantages. Diabetes Melitus significantly increase the risk of recurrent catheter-related thrombosis and infection site of catheter, which lead to more thrombosis and infection cases of NTHC in diabetic patients than non-diabetic.

(Cite this article as: Putra IG, Soebroto H, Sembiring YE, Tjempakasari A. Patency durability of non-tunneled hemodialysis' catheter in adult patients with chronic kidney disease stadium 5 with diabetes mellitus and non-diabetes mellitus. Ital J Vasc Endovasc Surg 2023;30:15-9. DOI: 10.23736/S1824-4777.22.01568-6)

KEY WORDS: Patency; Durability; Hemodialysis; Catheter; Non-tunneled hemodialysis catheter; Diabetes.

Introduction

Chronic kidney disease (CKD) is a condition of kidney that causes the need for kidney replacement therapy in the form of hemodialysis or kidney transplantation.¹ One of the treatments for CKD is hemodialysis. As a form of renal replacement therapy, the need for hemodialysis continues to increase from year to year with the number of ac-

tive hemodialysis patients as many as 77,892 patients with the addition of new patients in 2017 as many as 30,831. Hemodialysis is a treatment used for kidney function as an excretory organ to remove excess water filtrate, toxins, and other solutes. Hemodialysis can also maintain the physiology of the kidneys.² Hemodialysis has access to blood vessels. This access is a transcutaneous channel that is used as access to the intravascular space for renal replacement

therapy and is in the form of a central venous catheter. This hemodialysis therapy has a broad spectrum such as hemodialysis, hemofiltration, hemodiafiltration or ultrafiltration. One of the prerequisites and things that are very essential in the care of patients with hemodialysis is the existence of adequate hemodialysis vascular access. The use of a central venous catheter is used in various cases that require extracorporeal procedures. In cases of emergency, temporary and short-term use usually use a non-tunneled hemodialysis catheter (NTHC) and in the medium and long term use a tunneled hemodialysis catheter (THC) before the use of arteriovenous fistula.³ Diabetes mellitus (DM) is a multipathological degenerative disease so that complications that occurred previously will continue and can only be controlled by the rate of progression. This vulnerability triggers conditions related to treatment methods related to DM itself. One of them is a temporary hemodialysis catheter which is a vascular access in patients with CKD.⁴ DM conditions can affect the duration of hemodialysis catheter placement due to the occurrence of immunopathology and the coagulation cascade. These comorbidities lead to infection and thrombosis so that catheter patency is compromised. Impaired catheter patency causes repositioning or reimplantation of the catheter before an arteriovenous fistula can be performed.⁵ The use of NTHC in DM patients is more likely to experience thrombotic conditions than non-DM patients. The occurrence of thrombosis in patients with the use of vascular access in DM is twice as high as those without comorbid DM. Thrombotic conditions lead to catheter reimplantation conditions. Based on research, patients with DM who use temporary vascular access are considered more susceptible to infection and sepsis.⁶ Installation of NTHC is an action that is often done in RSUD Dr. Soetomo Surabaya and there has been no study on the duration of the hemodialysis catheter patency installed in RSUD Dr. Soetomo, Surabaya. Data collection and publications related to the condition of existing patients are limited in RSUD Dr. Soetomo, Surabaya. The factors that influence the duration of this patent were not clear at the time of this study. The femoral vein is the preferred location for insertion to reduce the risk of infection for staff. Selection of the insertion site in the femoral vein is not recommended despite the fact that there is a hypercoagulable condition.

Evidence acquisition

Search strategy

This study reviewed the literatures gathered from two major databases, PubMed, and Google Scholar. Advanced

search was conducted on PubMed with the keywords used were (“Patency” AND “catheter” AND “durability” AND “hemodialysis;” “non-tunneled hemodialysis catheter” AND “chronic kidney disease” AND “diabetic”). Advanced search was also conducted on Google Scholar with all the words “patency of tunneled catheters in hemodialysis patients;” “patency of tunneled catheters in diabetic chronic kidney disease underwent hemodialysis”. The showed literatures were further selected based on the publication year, by which we set a 10-year-study, starting from 2012 until 2022. This attempt resulted in 59 and 312 literatures gathered from PubMed and Google Scholar, respectively. We also reviewed the articles cited on the references within the article to broaden the search results.

Data extraction

The information reviewed from the articles consisted of: author (with study location and publication year); study design; population age (in years, with the format as mean±SD or median (IQR), except stated otherwise within the table); insertion site(s); duration of catheterization (in days, same format as population age, except stated otherwise); evaluation/measurement of catheter tip recirculation; location of the tip; and brief discussion regarding the results. The results are showed as in Table I.⁷⁻¹³

Evidence synthesis

Chronic kidney disease

Chronic kidney disease (CKD) is defined as a state of kidney damage or an estimated glomerular filtration rate (eGFR) of less than 60 mL/minute/1.73 m² that persists for three months or more regardless the causes. CKD patients should be assessed for risk factors of kidney disease, including prior exposure to potential nephrotoxins (e.g., nonsteroidal anti-inflammatory drugs [NSAIDs], phosphate-based bowel preparations, herbal remedies such as those containing aristolochic acid, antibiotic therapies such as gentamicin, and chemotherapies), history of nephrolithiasis or recurrent urinary tract infections, presence of comorbidities (e.g., hypertension, diabetes, autoimmune disease, chronic infections), family history of kidney disease, and, if available, other known genetic risk factors.¹⁴ Management of patients with CKD should include decreasing the rate of progression of CKD with regular monitoring of renal function, management of complications in CKD such as anemia and electrolyte disturbances as well as cardiovascular complications, preparation for

TABLE I.—Characteristics of the literatures reviewed.

N.	Author	Study design	Population	Vascular access	Outcome
1	Biesen <i>et al.</i> ⁷	Systematic literature review	Diabetes (type I or II) of end-stage renal disease or a superimposed condition	Tunneled catheters, grafts, or autogenous fistulas placed in any position	AVF appears to be the access of choice for younger and healthier HD diabetic patient
2	Cholewa ⁸	Retrospective analysis	Patient with permanent venous catheter thrombosis treated with urokinase	Right and left internal jugular vein, right and left femoral vein and inferior vena cava	The occurrence of DM significantly increases the risk of recurrent catheter-related thrombosis and in consequence decrease one-year catheter survival.
3	Clark <i>et al.</i> ⁹	Literature review	Patient with non-tunneled temporary hemodialysis catheter (NTHC)	Femoral site	The complications of NTHC insertion are frequent and can be fatal. Infection-control “bundles” of specific evidence-based practices to reduce the risk of CLABSIs and exit-site infections should be implemented in all settings in which NTHCs are inserted and used. The femoral site may not be associated with a higher risk of infection and is possibly even preferable in patients with diabetes
4	Tahir <i>et al.</i> ¹⁰	Observational (retrospective)	Patients with diabetes or hypertension or hypothyroidism or IHD received tunnel dialysis catheter	Right internal jugular vein, left internal jugular vein, right femoral vein, right subclavian vein	TDCs are the best available alternative for short to intermediate term use for hemodialysis when all preventive measures are taken to avoid catheter related complications either infectious or noninfectious until permanent dialysis access is achieved or renal transplant is done.
5	Halbert <i>et al.</i> ¹¹	Systematic literature review and meta-analysis	Diabetic and non-diabetic hemodialysis patients who underwent arteriovenous graft placements	ePTFE arteriovenous graft placements	Although fistulas are currently the gold standard for dialysis access, in part because of their low infection rates, certain subgroups of dialysis patients are burdened with higher use of ePTFE grafts and catheters. The observed rate of 9% infection per patient-year is consistent with other data sources pointing to worsened infectious outcomes in patients, especially women, who utilize ePTFE grafts for dialysis.
7	Cambria <i>et al.</i> ¹²	Systematic literature review and meta-analysis	Patency, mortality, access infection, and maturation of vascular access in adults requiring long-term dialysis	Arteriovenous grafts (AVGs) and catheters (tunneled and non-tunneled)	Autogenous access is the best approach when feasible. This study provides incidence rates in various subgroups to inform shared decision making and facilitate the conversation with patients about access planning
8	Liemarto ¹³	Observational analytic research	Hemodialysis patient at Dr. Soetomo Surabaya	Arteriovenous fistula (AVF)	In routine HD patients with DM, there is an increased risk factor for AVF. patent failure

renal replacement therapy, and immunization, especially hepatitis B.¹⁴ The KDIGO guidelines recommend that patients with CKD be referred to a nephrologist when eGFR falls below 30 mL/min/1.73 m² (stage G4) and/or urine ACR increases above 300 mg per 24 hours (stage A3). Additional indications for referral include the following: presence of greater than 20 red blood cells per high-power field of unclear etiology, red blood cell casts on urine microscopy or other indication of glomerulonephritis, CKD with uncontrolled hypertension despite 4 or more antihypertensive medications, persistent hypokalemia or hyperkalemia, anemia requiring erythropoietin replacement, recurrent or extensive kidney stones, hereditary kidney disease, acute kidney injury, and rapid CKD progression

(a decrease in eGFR ≥ 25% from baseline or a sustained decline in eGFR > 5 mL/min/1.73 m²).¹⁴

Diabetes mellitus in CKD and hemodialysis

Diabetes mellitus is a disease of inadequate control of blood levels of glucose. It has many subclassifications, including type 1, type 2, maturity-onset diabetes of the young (MODY), gestational diabetes, neonatal diabetes, and steroid-induced diabetes. Type 1 and 2 DM are the main subtypes, each with different pathophysiology, presentation, and management, but both have a potential for hyperglycemia.¹⁵ Diabetes mellitus (DM) is the most common cause of chronic kidney disease and has been identi-

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The production of derivative works from the Article is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo, or other proprietary information of the Publisher.

fied as a contributing factor to vascular access dysfunction, so it is very important to know the effect of DM and AVF patency in the HD patient population. DM conditions in CKD patients can be caused by diabetic kidney disease (DKD) conditions.¹³ DM could lead to a kidney complications and complications due to the installation of temporary access to a central venous catheter. It is associated with macrovascular damage, microvascular and immune deficiency in patients with DM. Metabolic conditions in DM patients damage blood vessels and can induce abnormalities such as thrombosis and this can be prevented by administering antithrombotic prophylaxis. Uncontrolled diabetes mellitus is also a risk factor for infection. Infection occurs in DM patients due to decreased immune cell function. This occurs due to dysregulation of the innate immune system, namely defects in the function of neutrophils and macrophages.¹⁶

Non-tunneled hemodialysis catheter

Non-tunneled hemodialysis catheters (NTHCs) are typically used when vascular access is required for urgent renal replacement therapy (RRT). In the setting of acute and chronic kidney injury, when the duration of RRT is difficult to predict, NTHCs are the recommended initial vascular access. However, due to an increased risk of complications, NTHCs are the least preferred access for chronic hemodialysis patients.¹⁷ Types of NTHC can be distinguished by design, purpose, and duration of use. The catheter can be divided into short term/non-cuffed (temporary) and long term/tunneled cuffed. Designs are distinguished according to their curvature, namely straight or precurved and tip types; namely staggered, split, and symmetric.¹⁸ Patients who use temporary vascular access in patients with CKD have a schedule for hemodialysis. The number of cycles is dependent on the target, the hemodialysis capability of the machine, and the patient's associated condition. The number of cycles or the frequency of hemodialysis does not affect the durability of existing vascular access. Maintenance of patency is related to the durability of NTHC by administration of thrombolytic agents or anticoagulants.¹⁹

Complications of NTHCs

Non-tunneled catheters for short-term extracorporeal RRT are preferentially inserted under ultrasound guidance, as this allows subcutaneous structures to be visualized and reduces the complication rates. Mechanical complications related to NTHC insertion are common with vascular inju-

ry or hematoma occurring in up to 5% of insertions. Other mechanical complications such as pneumothorax, pneumopericardium, and air and guidewire embolism occur less often but can be fatal. Although the life-threatening complications of NTHC insertion are typically related to insertions at the internal jugular site, fatal complications related to femoral NTHC insertion have been reported. Severe bleeding, typically retroperitoneal, occurs in approximately 0.5% of femoral insertions.²⁰ Infection in the vascular access is an infection that occurs as a primary infection that occurs in the vascular access as evidenced by clinical signs of fever >38.0 °C, chills or, hypotension accompanied by evidence of infection in the blood that is not associated with infection elsewhere. Infection can occur due to various factors whose relationship has not been explained, but it is known that DM is also a factor that influences the occurrence of infection in temporary catheters.²⁰ Obstruction due to thrombosis is a problem that occurs in various types of vascular access, both permanent and temporary. Thrombosis occurs due to various factors, especially the activation of platelets around the lumen of the catheter. Clinical signs of obstruction include pain and swelling in the distal area but no sign of infection. Impaired patency can be overcome by giving heparin flush and anticoagulant therapy.²⁰

Conclusions

A non-tunneled hemodialysis catheter may be used as a temporary catheter while waiting for a placement of an arteriovenous fistula, a peritoneal dialysis catheter, or a tunneled catheter. Compared to TDCs, NTDCs have several disadvantages such as a 5-fold higher rate of infection than TDCs and a greater likelihood of inadequate blood flow rate. Diabetes mellitus significantly increase the risk of recurrent catheter-related thrombosis and infection site of catheter, which lead to more thrombosis and infection cases of NTHC in diabetic patients than non-diabetic.

References

1. Webster AC, Nagler EV, Morton RL, Masson P. Chronic Kidney Disease. *Lancet* 2017;389:1238–52.
2. Hn M, Anjum F. Hemodialysis. Treasure Island, FL: StatPearls Publishing LLC; 2022. p.1–11.
3. Gb B, Nr A. Dialysis Catheter Contraindications. Treasure Island, FL: StatPearls Publishing LLC; 2021. p.1–7.
4. Galicia-Garcia U, Benito-Vicente A, Jebari S, Larrea-Sebal A, Siddiqi H, Uribe KB, *et al.* Pathophysiology of Type 2 Diabetes Mellitus. *Int J Mol Sci* 2020;21:6275.

5. Rafik H, Bahadi A, Aatif T, Sobhi A, Kabbaj D El. Bacteremia and Thrombotic Complications of Temporary Hemodialysis Catheters : Experience of a Single Center in Morocco. *Ibnos J Med Biomed Sci* 2022;159–63.
6. Xiong Y, Yu Y, Yang Y, Wang L, Shi P, Deng Y, *et al.* Impacts of age, diabetes, gender, and access type on costs associated with vascular access among Chinese patients on hemodialysis. *Int J Artif Organs* 2021;44:302–9.
7. Coentrão L, Van Biesen W, Nistor I, Tordoir J, Gallieni M, Marti Monros A, *et al.* Preferred haemodialysis vascular access for diabetic chronic kidney disease patients: a systematic literature review. *J Vasc Access* 2015;16:259–64.
8. Wójtowicz D, Cholewa D, Faba AM, Domańska B, Kokoszka J, Kopacz K, *et al.* Diabetes decreases patency of tunneled catheters in hemodialysis patients after first effective thrombolysis with urokinase. *Ren Fail* 2018;40:384–9.
9. Clark EG, Barsuk JH. Temporary hemodialysis catheters: recent advances. *Kidney Int* 2014;86:888–95.
10. Tahir M, Ali M. Outcomes of Tunneled Dialysis Catheter In Hemodialysis Patients In A Tertiary Care Hospital; 2022 [Internet]. Available from: <https://assets.researchsquare.com/files/rs-1129951/v1/aa5b22d6-4f9c-42cf-9be5-882d029f5296.pdf?c=1648712671> [cited 2022, Dec 5].
11. Halbert RJ, Nicholson G, Nordyke RJ, Pilgrim A, Niklason L. Patency of ePTFE Arteriovenous Graft Placements in Hemodialysis Patients: Systematic Literature Review and Meta-Analysis. *Kidney360* 2020;1:1437–46.
12. Almasri J, Alsawas M, Mainou M, Mustafa RA, Wang Z, Woo K, *et al.* Outcomes of vascular access for hemodialysis: A systematic review and meta-analysis. *J Vasc Surg* 2016;64:236–43.
13. Liemarto AK. Hubungan Diabetes Mellitus Dengan Patensi Arteriovenous Fistula (AVF) Pada Pasien Hemodialisis Rutin. *Perpust Univ Airlangga*; 2018 [Internet]. Available from: <https://repository.unair.ac.id/82250/> [cited 2022, Dec 5].
14. Cross SJ, Linker, Kay E, Leslie FM. HHS Public Access. *Physiol Behav* 2016;176:100–6.
15. Sapra A, Bhandari P. Diabetes Mellitus. Treasure Island, FL. StatPearls Publ 2022;LLC:1–11.
16. Frydrych LM, Bian G, O’Lone DE, Ward PA, Delano MJ. Obesity and type 2 diabetes mellitus drive immune dysfunction, infection development, and sepsis mortality. *J Leukoc Biol* 2018;104:525–34.
17. Vats HS. Complications of catheters: tunneled and nontunneled. *Adv Chronic Kidney Dis* 2012;19:188–94.
18. Banerjee S, Engineer D, Hirpara J, Shah N, Dave R, Sil K, *et al.* Dialysis Vascular Access: Where do Tunneled Catheters Stand? - A Single-Center Experience. *Indian J Nephrol* 2021;31:235–9.
19. Mandel-Shorer N, Tzvi-Behr S, Harvey E, Revel-Vilk S. Central venous catheter-related venous thrombosis in children with end-stage renal disease undergoing hemodialysis. *Thromb Res* 2018;172:150–7.
20. Sirvent AE, Enríquez R, Millán I, García-Marco JM, Rodríguez-Czaplicki E, Redondo-Pachón MD, *et al.* Severe hemorrhage because of delayed iliac vein rupture after dialysis catheter placement: is it preventable? *Hemodial Int* 2012;16:315–9.

Conflicts of interest.—The authors certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

Authors' contributions.—All authors equally contributed to the manuscript, read and approved the final version of the manuscript.

History.—Manuscript accepted: November 29, 2022. - Manuscript received: November 7, 2022.

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The production of derivative works from the Article is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo, or other proprietary information of the Publisher.