



Table of Contents

[Next Issue](#)

[Previous Issue](#)



January-March 2021
Volume 5 | Issue 1
Page Nos. 1-59

Online since Monday, February 8, 2021

Accessed 40,623 times.

PDF access policy

Journal allows immediate open access to content in HTML + PDF

[View issue as eBook](#)

[Issue citations](#)

[Issue statistics](#)

[RSS](#)

[POPULAR ARTICLES](#)


[JOIN AS REVIEWER](#)

[GET EMAIL ALERTS](#)

[RECOMMEND](#)

[Show all abstracts](#) [Show selected abstracts](#) [Export selected to](#) [Add to my list](#)

ORIGINAL ARTICLES



Effect of supplemental caudal analgesia on intraoperative blood glucose levels in pediatric patients undergoing urological surgery under general anesthesia: An observational study p. 1

Apurva Sadhoo, Sunil Rajan, Sowmya Jain, Kaushik Barua, Lakshmi Kumar
DOI:10.4103/BJOA.BJOA_143_20


[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[Sword Plugin for Repository\]](#)^{Beta}



Acute kidney injury incidence following cardiac surgery: A risk factor analysis p. 6

Andi Adil, Philia Setiawan, Yan Efrata Sembiring, Syafri Kamsul Arif, Hisbullah Amin
DOI:10.4103/BJOA.BJOA_202_20


[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[Sword Plugin for Repository\]](#)^{Beta}



Comparative study between betamethasone gel and lignocaine jelly applied over the tracheal tube to reduce postoperative airway complications p. 11

Guriqbal Singh, Priti Jadeja, R Yashwanth Patnaik, Sugadev Ravinbothayan, Vijaylaxmi Singh, Raghav Dhawan
DOI:10.4103/BJOA.BJOA_159_20


[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Sword Plugin for Repository\]](#)^{Beta}



To determine the efficacy of intrathecal dexmedetomidine and fentanyl as adjuvants to spinal anesthesia for lower abdominal surgeries below the level of umbilicus - A prospective randomized controlled double blinded study p. 15

Jadi Laxmikanth, Sivakumar Segaran, Sagiev Koshy George, Ashwin John, Jemie Rachel Johns, Nikita Mani
DOI:10.4103/BJOA.BJOA_160_20

[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Sword Plugin for Repository\]](#)^{Beta}



A comparative randomized clinical trial to assess the efficacy between dexmedetomidine and midazolam infusions for procedural sedation during septoplasty p. 21

Siyam Sundar, Sameera M Jahagirdar, VR Hemanth Kumar, N Krishnaveni
DOI:10.4103/BJOA.BJOA_169_20

[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Sword Plugin for Repository\]](#)^{Beta}

REVIEW ARTICLE

Fluid management in kidney disease patients for nontransplant and transplantation surgeries p. 26

Amal Francis Sam, Sandeep Sahu, Karthik T Ponnappan
DOI:10.4103/BJOA.BJOA_203_20

[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[Sword Plugin for Repository\]](#)^{Beta}

CASE REPORTS

- Dexmedetomidine versus clonidine as adjuvants in epidural analgesia in gynecological laparotomy: Case series** p. 33

Dewa Ayu Nyoman Isma Pratiwi, I Made Agus Kresna Sucandra, Tjokorda Gde Agung Senapathi
DOI:10.4103/BJOA.BJOA_129_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}

- Ropivacaine 0.75% for peribulbar block in vitrectomy** p. 38

Tjokorda Gde Agung Senapathi, I Made Gede Widnyana, Christopher Ryalino, Ida Bagus Gita Dharma Wibawa
DOI:10.4103/BJOA.BJOA_146_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}

- Perioperative management in obstetric patients with suspected COVID-19 at Bali, Indonesia: Case series** p. 40



Madyline Victorya Katipana, Made Wiryana, EM Tjahya Aryasa, Adinda Putra Pradhana
DOI:10.4103/BJOA.BJOA_166_20

[ABSTRACT] [HTML Full text] [PDF] [Citations (1)] [Sword Plugin for Repository]^{Beta}

- Low-cost techniques for bilateral intermittent transversus abdominis plane block using intravenous cannula following cesarean section** p. 45



Syahruil Mubarak Danar Sumantri
DOI:10.4103/BJOA.BJOA_179_20


[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}

- Intubation quality and hemodynamic response in endotracheal intubation with ultrasound-guided bilateral superior laryngeal nerve block: Case series** p. 50

Agus Tri, I Made Gede Widnyana, Made Agus Kresna Sucandra, Tjokorda Gde Agung Senapathi
DOI:10.4103/BJOA.BJOA_191_20

[ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}

- Anesthesia management of congenital diaphragmatic hernia in neonates** POPULAR p. 53



Harry Pranata, Putu Kurniyanta
DOI:10.4103/BJOA.BJOA_192_20

[ABSTRACT] [HTML Full text] [PDF] [Sword Plugin for Repository]^{Beta}

LETTERS TO EDITOR

- Inhospital counseling sustaining the well-being of patients and family members during the COVID-19 pandemic** p. 57

Poonam Kumari, Amarjeet Kumar, Chandni Sinha, Ajeet Kumar
DOI:10.4103/BJOA.BJOA_121_20

[HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}

- Anesthesia for fetal intervention for congenital aortic stenosis** p. 58



Rekha Varghese, Nitu Puthenveetil, Sunil Rajan, Lakshmi Kumar
DOI:10.4103/BJOA.BJOA_170_20

[HTML Full text] [PDF] [Mobile Full text] [EPub] [Citations (1)] [Sword Plugin for Repository]^{Beta}



Editorial Board

[About the journal](#)

Editor-in-Chief

Prof. Made Wiryana, Dr. dr. SpAn, KIC, KAO.

Scopus ID: [36099337800](#)

ORCID ID: [0000-0002-6552-8194](#)

Professor, Department of Anesthesiology and Intensive Care
Faculty of Medicine, Udayana University
Sanglah General Hospital
Bali, Indonesia

Editorial Board

Nishith Govil, MD, MBBS.

Scopus ID: [57202997171](#)

ORCID ID: [0000-0003-3749-6217](#)

Professor and Head of Department of Anaesthesiology,
Shri Guru Ram Rai Institute of Medical and Health Sciences
Dehradun, India

See Jee Jian, Dr, MBBS, MMed (Anaes)

Scopus ID: [8666451000](#)

Associate Professor, Head of Anaesthesia & Perioperative Medicine Department
Tan Tock Seng Hospital
Singapore, Singapore

Prof. Kazuya Morikawa

Scopus ID: [7202244107](#)

ORCID ID: [0000-0002-1503-7151](#)

Faculty of Medicine, University of Tsukuba
Tsukuba, Ibaraki, Japan

I Ketut Sinardja, dr., SpAn, KIC.

Scopus ID: [57191345194](#)

ORCID ID: [0000-0001-8609-8798](#)

Department of Anesthesiology and Intensive Care
Faculty of Medicine, Udayana University
Sanglah General Hospital
Bali, Indonesia

Prof. Adrian Antonius Jozef van Zundert, MD, PhD, FRCA, EDRA, FANZCA.

Scopus ID: [7003716417](#)

ORCID ID: [0000-0002-1836-6831](#)

Professor, Department of Anaesthesia & Perioperative Medicine
The University of Queensland
Royal Brisbane & Women's Hospital

Prof. Stephen P. Gatt, FRCA, FJFICM, FANZCA, FFA, FFARACS, LRCP, MRCS.

Scopus ID: [7004816461](#)

Associate Professor, Department of Anaesthesia
Prince of Wales Hospitals
Sydney Children's Hospital
Sydney, Australia

[POPULAR ARTICLES](#)

[JOIN AS REVIEWER](#)

[GET EMAIL ALERTS](#)

[RECOMMEND](#)

Susilo Chandra, dr. SpAn, FRCA.

Scopus ID: [55202737900](#)

Department of Anesthesiology and Intensive Care
Faculty of Medicine, University of Indonesia
Cipto Mangunkusumo General Hospital
Jakarta, Indonesia

Ratna Farida, Dr. dr. SpAn, KAKV.

Department of Anesthesiology and Intensive Care
Faculty of Medicine, University of Indonesia
Cipto Mangunkusumo General Hospital
Jakarta, Indonesia



Source details

Bali Journal of Anesthesiology

Open Access ⓘ

Scopus coverage years: from 2017 to Present

Publisher: Wolters Kluwer Health

E-ISSN: 2549-2276

Subject area:

Source type: Journal

CiteScore 2021

0.4 ⓘ

SNIP 2021

0.279 ⓘ

[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology ✕

CiteScore 2023 counts the citations received in 2020-2023 to articles, reviews, conference papers, book chapters and data papers published in 2020-2023, and divides this by the number of publications published in 2020-2023. [Learn more >](#)



CiteScore rank ⓘ

Category Rank Percentile

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site ↗](#)

About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

ELSEVIER

[Terms and conditions](#) ↗ [Privacy policy](#) ↗

Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗.



Acute Kidney Injury Incidence Following Cardiac Surgery: A Risk Factor Analysis

Andi Adil, Philia Setiawan, Yan Efrata Sembiring¹, Syafri Kamsul Arif, Hisbullah Amin

Department of Anesthesiology, Intensive Therapy, and Pain Management, Faculty of Medicine, Hasanuddin University, Makassar, ¹Department of Cardiothoracic and Vascular Surgery, Faculty of Medicine, Airlangga University, Surabaya, Indonesia

Abstract

Background: Acute decline of renal function following cardiac surgery is a common and complex problem with an incidence of up to 30% in Europe and leading to increased mortality rate in immediate and long term. The aim of this study is to analyze risk factors regarding acute kidney injury (AKI) incidence following cardiac surgery. **Patients and Methods:** This was a descriptive cross-sectional study conducted at Dr. Wahidin Sudirohusodo Central General Hospital, Makassar, Indonesia. Serum creatinine level was measured to diagnose AKI. Hemoglobin level, blood transfusion, vasopressor, and inotropic needs before surgery were recorded. Hemodynamic parameters that included mean arterial pressure and heart rate were also recorded. **Results:** Of all 33 subjects who underwent adult cardiac surgery that included coronary artery bypass graft, heart valve, and congenital disease surgery, 21 (63.6%) patients had AKI and 12 (36.4%) patients did not. Several perioperative factor analyses between AKI and non-AKI group revealed that only age showed a significant difference between the two groups ($P = 0.047$). **Conclusion:** Age is the most significant factor in AKI incidence following cardiac surgery. Further studies are required to analyze other perioperative factors that might induce AKI.

Keywords: Acute kidney injury, cardiac surgery, cardiopulmonary bypass

INTRODUCTION

Acute decline of renal function following cardiac surgery is a common and complex problem with an incidence of up to 30% in Europe and leading to increased mortality rate in immediate and long term.^[1] Data on acute kidney injury (AKI) incidence following cardiac surgery in Indonesia from previous study showed an estimated incidence of 36.4% in pediatric cardiac surgery.^[2]

Lewington *et al.* stated that the causes of AKI following cardiac surgery are very complex and multifactorial, including endogenous and exogenous toxins, metabolic factors, ischemic reperfusion, neurohormonal activation, inflammation, and oxidative stress.^[3] Distinctive features that occur during cardiac surgery procedure such as cardiopulmonary bypass (CPB) machine and aortic cross-clamp usage, increased blood transfusion volume, and high dose of vasopressor administration will increase AKI incidence following cardiac surgery compared to noncardiac surgery.^[1]

The aim of this study is to analyze risk factors regarding AKI incidence following cardiac surgery; therefore, this study might

give insight to improve AKI management following cardiac surgery in the future.

PATIENTS AND METHODS

This descriptive cross-sectional was conducted on October 2019–February 2020 at Dr. Wahidin Sudirohusodo Central General Hospital, Makassar, Indonesia. Population in this study was patients who underwent adult on-pump cardiac surgery. Samples that met inclusion criteria gave their written informed consent. Exclusion criteria in this study were patients with presurgery AKI who underwent hemodialysis, patients with elevated presurgery creatinine level, and patients with a history of kidney transplantation.

Address for correspondence: Dr. Andi Adil, Medical Faculty of Hasanuddin University, Perintis Kemerdekaan No. Km. 10, Tamalanrea Indah, Tamalanrea, Makassar 90245, South Sulawesi, Indonesia. E-mail: adil_zanetti@yahoo.co.id

Submitted: 30-Aug-2020

Revised: 05-Oct-2020

Accepted: 30-Oct-2020

Published: 08-Feb-2021

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Adil A, Setiawan P, Sembiring YE, Arif SK, Amin H. Acute kidney injury incidence following cardiac surgery: A risk factor analysis. *Bali J Anaesthesiol* 2021;5:6-10.

Access this article online

Quick Response Code:



Website:
www.bjoaonline.com

DOI:
10.4103/BJOA.BJOA_202_20

Before this study was conducted, we requested ethical clearance from the Ethical Committee of Biomedical Study on Humans of Medical Faculty of Hasanuddin University and research permit from Dr. Wahidin Sudirohusodo Central General Hospital, where both of them were approved on July 25, 2019, with registration number: 538/UN4.6.4.5.31/PP36/2019. This study is in accordance with Helsinki Declaration 2000. Patients who agreed to join in this study signed their informed consent.

Serum creatinine level was measured 1 day before surgery and once per day for a week postsurgery. Additional measurements were performed based on clinician's advice. Creatinine level postsurgery that was measured for a week was used to diagnose AKI and to determine its grade. AKI based on Kidney Disease Improving Global Outcome (KDIGO) criteria is defined as an elevated absolute creatinine level of 0.3 mg/dl at 48 h or 1.5–1.9 times from baseline level for 7 days. Hemoglobin level, blood transfusion, vasopressor, and inotropic needs before surgery were recorded. Hemodynamic parameters that included mean arterial pressure (MAP) and heart rate were also recorded.

Data of patients' characteristics were presented as frequencies and percentages. Comparison of AKI and non-AKI patients' characteristics was analyzed using Chi-square test. Statistical analysis was performed using SPSS package version 25 for Windows (IBM Corporation, Armonk, NY, USA).

RESULTS

Of all 33 subjects who underwent adult cardiac surgery that included coronary artery bypass graft (CABG), heart valve, and congenital disease surgery, 21 patients (63.6%) had AKI and 12 patients (36.4%) did not.

Table 1 shows the characteristics of cardiac surgery patients. The age of patients varied from 11 to 70 years, with 11–20 years as the group with the lowest percentage and the highest percentage being from group 41–50 years. From 33 samples, 19 were male and 21 were female.

There were several types of cardiac surgery that were conducted: CABG, atrial septal defect closure, mitral valve replacement (MVR), MVR in combination with aortic valve replacement, and ventricular septal defect closure in combination with pulmonary stenosis dilation. Twenty people had heart failure, two people had ejection fraction (EF) <35%, and one person had a previous history of cardiac surgery. Before surgery, one person had intra-aortic balloon pump installation, ten people received chronic obstructive pulmonary disease (COPD) treatment, and five people received insulin.

Table 2 shows on-pump surgery parameter characteristics. CPB was conducted on average for 94.33 ± 30.68 min and cross-clamp was conducted on average for 78.39 ± 30.27 min. The mean blood transfusion volume that was given was 305.74 ± 110.11 ml and the mean hemoglobin level during on-pump procedure was 6.93 ± 1.35 g/dL.

In Table 3, several presurgery factors were analyzed between AKI and non-AKI group, where it was found that age being

Table 1: Demographics of the study subjects

Variables	Category	Frequency (n=33), n (%)
Age (years)	11-20	1 (3.0)
	21-30	4 (12.1)
	31-40	2 (6.1)
	41-50	14 (42.4)
	51-60	8 (24.2)
	61-70	4 (12.1)
Sex	Men	19 (57.6)
	Women	14 (42.4)
Surgery type	CABG	22 (66.7)
	ASD closure	6 (18.2)
	MVR	3 (9.1)
	MVR + AVR	1 (3.0)
	VSD closure + PS dilation	1 (3.0)
Heart failure	Yes	20 (60.6)
	No	13 (39.4)
Ejection fraction <35%	Yes	2 (6.1)
	No	31 (93.9)
Cardiac surgery history	Yes	1 (3.0)
	No	32 (97.0)
Presurgery IABP	Yes	1 (3.0)
	No	32 (97.0)
Presurgery COPD therapy	Yes	10 (30.3)
	No	23 (69.7)
Presurgery insulin therapy	Yes	5 (15.2)
	No	28 (84.8)
History of cardiac surgery	CABG	22 (66.7)
	ASD closure	6 (18.2)
	MVR	3 (9.1)

CABG: Coronary artery bypass graft, ASD: Atrial septal defect, MVR: Mitral valve replacement, AVR: Aortic valve replacement, VSD: Ventricular septal defect, PS: Pulmonary stenosis, IABP: Intra-aortic balloon pump, COPD: Chronic obstructive pulmonary disease

Table 2: On-pump parameter characteristics in cardiac surgery patients

On-pump	Mean \pm SD	Minimum	Maximum
CPB (min)	94.33 \pm 30.68	29.00	165.00
X-clamp (min)	78.39 \pm 30.27	16.00	147.00
Transfusion (ml)	305.74 \pm 110.11	197.00	559.00
Hb (g %)	6.93 \pm 1.35	5.10	10.90

CPB: Cardiopulmonary bypass, Hb: Hemoglobin, SD: Standard deviation

the only characteristic that had significant difference between AKI incidence (50.57 ± 11.97 vs. 40.50 ± 13.28 , $P = 0.047$). Comparison between sex revealed that there were 12 (57.1%) males and 9 (42.9%) females in the AKI group and 7 (58.3%) males and 5 (41.7%) females in the non-AKI group. AKI incidence in CABG procedure was in 16 (76.2%) patients compared to 6 (50%) patients who did not have AKI, and in heart valve surgery, there were 2 (9%) patients who had AKI and 2 (16%) patients who did not. In the AKI group, there were 15 (71.4%) patients with heart failure and 1 (4.8%) patient with EF <35%, whereas in the non-AKI group, there were 5 (41.7%) patients with heart failure and 1 (8.3%) patient with EF <35%.

Table 4 shows that there were no significant differences between two groups in terms of on-pump characteristics. The mean CPB and cross-clamp aorta duration in the AKI group were 101.19 ± 23.91 min and 84.67 ± 25.57 min, respectively, whereas in the non-AKI group, the mean durations were 82.33 ± 38.11 min and 67.42 ± 35.63 min, respectively. The mean blood transfusion volume and hemoglobin level in the AKI group were 310.76 ± 105.22 ml and 6.76 ± 1.25 g%, respectively, and in the non-AKI group were 291.50 ± 132.61 ml and 7.25 ± 1.53 g%, respectively.

DISCUSSION

In our study, 21 out of 33 (63%) patients experienced AKI following cardiac surgery. AKI incidence in our study was higher than several studies.^[1,4-6] Differences in diagnostic techniques in determining AKI could be one of the reasons for

different AKI incidences from each study. We used KDIGO criteria in determining AKI diagnosis in our study, using both creatinine and urine production criteria. The use of risk/injury/failure/loss/end-stage (RIFLE) and AKI network (AKIN) criteria following cardiac surgery without correcting changes in serum creatinine due to fluid balance will make AKI diagnosis less detectable. KDIGO criteria is associated with higher sensitivity in diagnosing AKI and predicting mortality compared to RIFLE and AKIN. KDIGO criteria, which is a combination of AKIN and RIFLE classifications, is commonly used for AKI diagnosis following cardiac surgery.^[7-12]

Several perioperative factor analyses between AKI and non-AKI group revealed that only age showed a significant difference between the two groups ($P = 0.047$). Factors behind elderly to more likely develop AKI might be due to decreased kidney function, patient's comorbidities, or related medical treatment received by elderly patients. Decreased renal mass due to reduced weight and decreased of glomeruli volume and number are morphological and anatomical factor due to renal age.^[13-17]

This study is in accordance with a study conducted by Guan *et al.*, which stated that age is one of the factors associated with AKI incidence following cardiac surgery.^[18] Age factor varied widely in several other studies as a risk for AKI following cardiac surgery.^[18-21] Jiang *et al.* showed that although the numbers differed, they did not show a statistically significant difference.^[19]

The use of angiotensin-converting enzyme (ACE) inhibitor and hypertension history are risk factors for AKI incidence following cardiac surgery. In this study, the use of ACE inhibitor did not show significant differences between AKI (13 patients = 61.9%) and non-AKI group (three patients = 25%). The use of ACE inhibitor as a risk factor has been studied by several experts. Arora *et al.* carried out a cohort study of 1358 patients and found an increased risk of AKI by 27.6%.^[22] This is in accordance with other studies conducted by Kristovic *et al.*, which showed an association between ACE inhibitor use and AKI incidence following cardiac surgery and Whiting *et al.* who conducted a meta-analysis study of perioperative ACE inhibitor use.^[23,24]

In our study, there was no significant difference for comorbid hypertension between AKI (20 patients = 95.2%) and non-AKI (ten patients = 83.3%) group. Hypertension is one of the AKI predictors following cardiac surgery. This is largely due to renal vascular damage due to chronic hypertension and a shift in autoregulation exacerbated by CPB use with low-pressure MAP condition.^[25]

There was no significant difference for diabetes mellitus (DM) comorbid between AKI (five patients = 23.8%) versus non-AKI (one patient = 8.3%) group. This is in accordance with the studies reported by Ferreiro and Lombardi and Kanji *et al.*^[20,25] Several studies have shown that DM is a risk factor for AKI following cardiac surgery. This is due to diabetic nephropathy

Table 3: Comparison of characteristics between acute kidney injury and nonacute kidney injury group

Variables	AKI incidence		P*
	Yes (n=21)	No (n=12)	
Age (years)	50.57±11.97	40.50±13.28	0.047
Sex			
Male	12 (57.1)	7 (58.3)	1.000
Female	9 (42.9)	5 (41.7)	
CABG surgery	16 (76.2)	6 (50.0)	0.149
Heart failure	15 (71.4)	5 (41.7)	0.142
Ejection fraction <35%	1 (4.8)	1 (8.3)	1.000
History of previous cardiac surgery	1 (4.8)	0	1.000
History of DM	5 (23.8)	1 (8.3)	0.379
History of hypertension	20 (95.2)	10 (83.3)	0.538
IABP presurgery	1 (4.8)	0	1.000
COPD treatment presurgery	5 (23.8)	5 (41.7)	0.433
Insulin presurgery	4 (19.0)	1 (8.3)	0.630
ACE-inhibitor presurgery	13 (61.9)	3 (25)	0.071
Valve surgery	2 (9%)	2 (16%)	0.610
Dobutamine intra-surgery	3.1±4.9	1.4±2	0.449
Norepinephrine intra-surgery	0.08±0.08	0.06±0.07	0.699

*Chi-square test. AKI: Acute kidney injury, CABG: Coronary artery bypass graft, DM: Diabetes mellitus, IABP: Intra-aortic balloon pump, COPD: Chronic obstructive pulmonary disease, ACE: Angiotensin converting enzyme

Table 4: Procedure duration and laboratory results during on-pump procedure in cardiac surgery patients

Variable	AKI incidence		P*
	Yes (n=21)	No (n=12)	
CPB (min)	101.19±23.91	82.33±38.11	0.100
X-clamp (min)	84.67±25.57	67.42±35.63	0.123
Transfusion volume (ml)	310.76±105.22	291.50±132.61	0.708
Hb (g %)	6.76±1.25	7.25±1.53	0.326

*Chi-square test. AKI: Acute kidney injury, CPB: Cardiopulmonary bypass, Hb: Hemoglobin

which will cause damage to tubules and glomerulus.^[26] Insulin-dependent DM is a strong risk factor in AKI incidence following cardiac surgery, which was not in accordance with this study. Arun *et al.* also reported that 80% of DM patients suffered from AKI compared to 20% of non-DM patients who underwent CABG surgery.^[27]

Patients with low EF (<35%) in our study did not differ significantly between AKI and non-AKI group. Low EF is a risk factor because it reduces perfusion to kidneys. Pieri *et al.* stated that low EF increases AKI risk following cardiac surgery. This was more prominent in patients with EF <35%.^[28] Omar *et al.* in their study found that patients with lower EF suffered from AKI more compared to non-AKI patients.^[29]

Our study also did not show a significant difference for COPD comorbid between AKI (five patients = 23.8%) versus non-AKI (five patients = 41.7%). The mechanism of COPD effect on AKI incidence of AKI is still unclear. This is probably because of a decrease in arterial oxygen content which affects oxygen delivery to kidney tissue. Several studies have shown the effect of COPD risk factors on AKI incidence following cardiac surgery, although the results are not significant.^[19,30]

In this study, the mean CPB and cross-clamp durations in AKI cases were 101.19 ± 23.91 min and 84.67 ± 25.57 min, respectively, while in the non-AKI group, the durations were 82.33 ± 38.11 min and 67.42 ± 35.63 min, respectively. CPB and aortic cross-clamp duration were associated with AKI incidence following cardiac surgery. This is due to elevated inflammatory factors and ischemic reperfusion injury.^[31] In their study, Sirvinskas *et al.* demonstrated that AKI incidence following cardiac surgery was significant if CPB duration was around 134 min and aortic cross-clamp duration was 75 min.^[32] Karim *et al.* showed that the risk of AKI increased with CPB duration >70 min and aortic cross-clamp duration >60 min.^[33]

Of all AKI patients, 16 (76.2%) underwent CABG surgical procedures compared to 6 (50%) patients in the non-AKI group. Two (9%) AKI patients underwent heart valve surgery compared to 2 (16%) patients in the non-AKI group. There was no significant differences between the two groups in terms of CABG or valve surgery. Valve surgery is a complex procedure, with prolonged surgery that might cause hemodynamic instability in renal perfusion. Ramos and Dias reported that valve replacement surgery was a strong predictor of AKI incidence following cardiac surgery (odds ratio [OR] = 4.7, $P = 0.002$; 95% confidence interval [CI] = 1.76–12.62).^[34]

In this study, PRC transfusion volume and Hb levels were 310.76 ± 105.22 ml and 6.76 ± 1.25 g%, respectively, in the AKI group, and 291.50 ± 132.61 ml and 7.25 ± 1.53 g%, respectively, in the non-AKI group. Although it is not statistically significant, transfusion volume that was given was larger in the AKI group and Hb level in AKI patients was lower compared to non-AKI patients. Intraoperative blood transfusion might increase oxygen delivery, but also increase AKI risk following cardiac surgery. This is because red blood

cell transfusion will cause an increase of free heme due to hemolysis process in stored blood, therefore damaging the kidneys. Crosina *et al.* reported that intraoperative hematocrit level was a factor that increased AKI incidence's area under curve that was combined with Cleveland clinical score, with hematopoietic cell transplantation = 0.25 ± 0.03 versus 0.28 ± 0.04 ($P = 0.001$) and transfusion volume = 1.51 ± 2.13 versus 0.6 ± 1.44 ($P = 0.001$).^[35] A research conducted by Baranauskas *et al.* showed that anemia incidence was around 7.59%, and there was a significant positive correlation between perioperative anemia and AKI ($r = 0.50$, $P < 0.05$).^[36] Ng *et al.* found that low hematocrit levels and preoperative anemia were potential modifiable risk factors associated with AKI following cardiac surgery in the Asian population, whereas administration of blood transfusions did not affect AKI incidence.^[37]

Inotropes and vasopressors use (dobutamine and norepinephrine) did not show significant differences between AKI (dobutamine = 3.1 ± 4.9 mcg/kg/min), (norepinephrine = 0.08 ± 0.08 mcg/kg/min) and non-AKI (dobutamine = 1.4 ± 2 mcg/kg/min), (norepinephrine = 0.06 ± 0.07 mcg/kg/min). The use of inotropes and vasopressors is associated with postsurgery AKI incidence due to renal ischemia and reperfusion due to hemodynamic instability and decreased cardiac function. Ramos and Dias found that norepinephrine (OR = 3.32, $P = 0.013$; 95% CI = 1.29–8.58) and dobutamine (OR = 5.3, $P = 0.019$, 95% CI = 1,32–21,64) were predictors of AKI incidence following cardiac surgery.^[34]

This study had limited number of samples (33 patients). Furthermore, other perioperative factors that induce AKI following cardiac surgery should be analyzed in future studies.

CONCLUSION

Age is the most significant factor in AKI incidence following cardiac surgery. Further studies are required to analyze other perioperative factors that might induce AKI.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Lagny MG, Jouret F, Koch JN, Blaffart F, Donneau AF, Albert A, *et al.* Incidence and outcomes of acute kidney injury after cardiac surgery using either criteria of the RIFLE classification. *BMC Nephrol* 2015;16:76.
2. Heriwardito A, Alatas A, Perdana A, Sutedja AD. Incidence and risk factor of acute kidney injury post open heart surgery in pediatric patients. *Bali J Anesthesiol* 2018;2:48-50.
3. Lewington AJ, Cerdá J, Mehta RL. Raising awareness of acute kidney injury: A global perspective of a silent killer. *Kidney Int* 2013;84:457-67.
4. Bastin AJ, Ostermann M, Slack AJ, Diller GP, Finney SJ, Evans TW. Acute kidney injury after cardiac surgery according to risk/injury/failure/loss/end-stage, acute kidney injury network, and kidney disease: Improving global outcomes classifications. *J Crit Care* 2013;28:389-96.
5. Machado MN, Nakazone MA, Maia LN. Prognostic value of acute

- kidney injury after cardiac surgery according to kidney disease: Improving global outcomes definition and staging (KDIGO) criteria. *PLoS One* 2014;9:e98028.
6. Jorge-Monjas P, Bustamante-Munguira J, Lorenzo M, Heredia-Rodríguez M, Fierro I, Gómez-Sánchez E, *et al.* Predicting cardiac surgery-associated acute kidney injury: The CRATE score. *J Crit Care* 2016;31:130-8.
 7. Vives M, Hernandez A, Parramon F, Estanyol N, Pardina B, Muñoz A, *et al.* Acute kidney injury after cardiac surgery: Prevalence, impact and management challenges. *Int J Nephrol Renovasc Dis* 2019;12:153-66.
 8. Englberger L, Suri RM, Li Z, Casey ET, Daly RC, Dearani JA, *et al.* Clinical accuracy of RIFLE and acute kidney injury network (AKIN) criteria for acute kidney injury in patients undergoing cardiac surgery. *Crit Care* 2011;15:R16.
 9. Khwaja A. KDIGO clinical practice guidelines for acute kidney injury. *Nephron Clin Pract* 2012;120:c179-84.
 10. Luo X, Jiang L, Du B, Wen Y, Wang M, Xi X, *et al.* A comparison of different diagnostic criteria of acute kidney injury in critically ill patients. *Crit Care* 2014;18:R144.
 11. National Kidney Foundation. K/DOQI clinical practice guidelines for chronic kidney disease: Evaluation, classification, and stratification. *Am J Kidney Dis* 2002;39:S1-266.
 12. Koeze J, Keus F, Dieperink W, van der Horst IC, Zijlstra JG, van Meurs M. Incidence, timing and outcome of AKI in critically ill patients varies with the definition used and the addition of urine output criteria. *BMC Nephrol* 2017;18:70.
 13. Pedersen AB, Gammelager H, Kahlert J, Sørensen HT, Christiansen CF. Impact of body mass index on risk of acute kidney injury and mortality in elderly patients undergoing hip fracture surgery. *Osteoporos Int* 2017;28:1087-97.
 14. Chronopoulos A, Cruz DN, Ronco C. Hospital-acquired acute kidney injury in the elderly. *Nat Rev Nephrol* 2010;6:141-9.
 15. Chronopoulos A, Rosner MH, Cruz DN, Ronco C. Acute kidney injury in elderly intensive care patients: A review. *Intensive Care Med* 2010;36:1454-64.
 16. Hsu RK, Siew ED. The growth of AKI: Half empty or half full, it's the size of the glass that matters. *Kidney Int* 2017;92:550-3.
 17. Yokota LG, Sampaio BM, Rocha EP, Balbi AL, Sousa Prado IR, Ponce D. Acute kidney injury in elderly patients: Narrative review on incidence, risk factors, and mortality. *Int J Nephrol Renovasc Dis* 2018;11:217-24.
 18. Guan C, Li C, Xu L, Zhen L, Zhang Y, Zhao L, *et al.* Risk factors of cardiac surgery-associated acute kidney injury: Development and validation of a perioperative predictive nomogram. *J Nephrol* 2019;32:937-45.
 19. Jiang W, Xu J, Shen B, Wang C, Teng J, Ding X. Validation of four prediction scores for cardiac surgery-associated acute kidney injury in chinese patients. *Braz J Cardiovasc Surg* 2017;32:481-6.
 20. Ferreira A, Lombardi R. Acute kidney injury after cardiac surgery is associated with mid-term but not long-term mortality: A cohort-based study. *PLoS One* 2017;12:e0181158.
 21. Lee CC, Chang CH, Cheng YL, Kuo G, Chen SW, Li YJ, *et al.* Diagnostic performance of cyclophilin A in cardiac surgery-associated acute kidney injury. *J Clin Med* 2019;9:108.
 22. Arora P, Rajagopalam S, Ranjan R, Kolli H, Singh M, Venuto R, *et al.* Preoperative use of angiotensin-converting enzyme inhibitors/angiotensin receptor blockers is associated with increased risk for acute kidney injury after cardiovascular surgery. *Clin J Am Soc Nephrol* 2008;3:1266-73.
 23. Kristovic D, Horvatic I, Husedzinovic I, Sutlic Z, Rudez I, Baric D, *et al.* Cardiac surgery-associated acute kidney injury: Risk factors analysis and comparison of prediction models. *Interact Cardiovasc Thorac Surg* 2015;21:366-73.
 24. Whiting P, Morden A, Tomlinson LA, Caskey F, Blakeman T, Tomson C, *et al.* What are the risks and benefits of temporarily discontinuing medications to prevent acute kidney injury? A systematic review and meta-analysis. *BMJ Open* 2017;7:e012674.
 25. Kanji HD, Schulze CJ, Hervas-Malo M, Wang P, Ross DB, Zibdawi M, *et al.* Difference between pre-surgery and cardiopulmonary bypass mean arterial pressure is independently associated with early cardiac surgery-associated acute kidney injury. *J Cardiothorac Surg* 2010;5:71.
 26. Patschan D, Müller GA. Acute kidney injury in diabetes mellitus. *Int J Nephrol* 2016;2016:6232909.
 27. Arun O, Celik G, Oc B, Unlu A, Celik JB, Oc M, *et al.* Renal effects of coronary artery bypass graft surgery in diabetic and non-diabetic patients: A study with urinary neutrophil gelatinase-associated lipocalin and serum cystatin C. *Kidney Blood Press Res* 2015;40:141-52.
 28. Pieri M, Belletti A, Monaco F, Pisano A, Musu M, Dalessandro V, *et al.* Outcome of cardiac surgery in patients with low preoperative ejection fraction. *BMC Anesthesiol* 2016;16:97.
 29. Omar AS, Mahmoud K, Hanoura S, Osman H, Sivadasan P, Sudarsanan S, *et al.* Acute kidney injury induces high-sensitivity troponin measurement changes after cardiac surgery. *BMC Anesthesiol* 2017;17:15.
 30. Thakar CV, Arrigain S, Worley S, Yared JP, Paganini EP. A clinical score to predict acute renal failure after cardiac surgery. *J Am Soc Nephrol* 2005;16:162-8.
 31. Paarmann H, Charitos EI, Beilharz A, Heinze H, Schön J, Berggreen A, *et al.* Duration of cardiopulmonary bypass is an important confounder when using biomarkers for early diagnosis of acute kidney injury in cardiac surgical patients. *Appl Cardiopulm Pathophysiol* 2013;17:284-97.
 32. Sirvinskas E, Andrejaitiene J, Raliene L, Nasvytis L, Karbonskiene A, Pilvinis V, *et al.* Cardiopulmonary bypass management and acute renal failure: Risk factors and prognosis. *Perfusion* 2008;23:323-7.
 33. Karim HM, Yunus M, Saikia MK, Kalita JP, Mandal M. Incidence and progression of cardiac surgery-associated acute kidney injury and its relationship with bypass and cross clamp time. *Ann Card Anaesth* 2017;20:22-7.
 34. Ramos KA, Dias CB. Acute kidney injury after cardiac surgery in patients without chronic kidney disease. *Braz J Cardiovasc Surg* 2018;33:454-61.
 35. Crosina J, Lerner J, Ho J, Tangri N, Komenda P, Hiebert B, *et al.* Improving the prediction of cardiac surgery-associated acute kidney injury. *Kidney Int Rep* 2017;2:172-9.
 36. Baranauskas T, Kaunienė A, Švagždienė M, Širvinskas E, Lenkutis T. The correlation of post-operative acute kidney injury and perioperative anaemia in patients undergoing cardiac surgery with cardiopulmonary bypass. *Acta Med Litu* 2019;26:79-86.
 37. Ng RR, Chew ST, Liu W, Shen L, Ti LK. Identification of modifiable risk factors for acute kidney injury after coronary artery bypass graft surgery in an Asian population. *J Thorac Cardiovasc Surg* 2014;147:1356-61.