

Left Anterior Descendent Coronary Artery Fistula

by Lyndon Darwin

Submission date: 10-Apr-2023 09:18PM (UTC+0800)

Submission ID: 2060533213

File name: Left_Anterior_Descendent_Coronary_Artery_Fistula.pdf (1.3M)

Word count: 3077

Character count: 17796

¹⁷ Left Anterior Descendent Coronary Artery Fistula to Main Pulmonary Artery with Triple Vessel Disease: A Report of Two Cases

Lyndon Darwin¹, Yan Efrata Sembiring², Oky Revianto Sediono Pribadi²

³⁸ ⁷ ¹⁵
¹Resident, ²Senior Attendant, Department of Thoracic, Cardiac and Vascular Surgery, Airlangga University School of Medicine-Dr. Soetomo General Hospital, Surabaya, Indonesia

Abstract

¹⁵
Coronary artery fistula is a rare anomaly of coronary artery. It represents connection between one or more of coronary arteries and cardiac chamber or great vessel. It can also cause significant hemodynamic changes. To report cases of left anterior descending coronary artery (LAD) fistula to main pulmonary artery (MPA) with concomitant triple vessel disease. Case 1 was 59-year old male presented with intermittent chest discomfort for a year. Echocardiogram showed severe MR with cleft at A1 and A2 and coronary angiogram result showed LAD fistulation to MPA. Case 2 was 57-year old male presented with chest discomfort and at the left shoulder. Echocardiogram showed trivial MR and coronary angiogram showed triple vessel disease along with tortuous fistulation of LAD to MPA. Both patients underwent teflonpledget-reinforced direct suturing of fistula origin and 3-grafts coronary artery bypass graft. Patient no 1 also underwent mitral annuloplasty and valvuloplasty. Post-operative period was uneventful and both patients were discharged after 17 days and 5 days respectively. Short term follow-up showed improvement of symptoms and no residual fistulation. In short term follow-up teflonpledget-reinforced direct suturing of coronary fistula origin result satisfactory. Larger study and further follow up is necessary.

Keywords: Coronary artery fistula, direct pledget suturing, coronary artery disease, triple vessel disease, mitral regurgitation.

Introduction

Coronary artery fistulae are rare anomaly of coronary artery and usually occurs in isolation. It is defined as a connection between one or more of coronary arteries and cardiac chamber or great vessel¹. The prevalence of coronary artery fistula detected with computed tomography (CT) angiogram was found to be 0.19%-0.9% while conventional angiographic findings was 0.05-0.25%^{2,3}. Most are asymptomatic but some may cause hemodynamically significance problem. It may lead to serious complications including myocardial ischemia, heart failure, arrhythmias, thromboembolism, superior vena cava syndrome and even death⁴.

Case Description

Case #1

A 59-year old male was referred from a secondary

referral hospital. He presented with intermittent chest discomfort for about a year. Chest discomfort was aggravated with exertion and relieved at rest. He had controlled hypertension but no history or diabetes mellitus or stroke. Patient also had no history of smoking or alcohol consumption.

Physical exam showed grade IV/VI systolic murmur at apex radiated to axilla. Echocardiogram findings were severe MR (Carpentier Type I), dilatation of left atrium, normal systolic function of left ventricle (LV), concentric LV hypertrophy and ejection fraction of 67%. Coronary angiogram showed tortuous and dilated vessel arising superiorly from left anterior descending (LAD) coronary artery (Figure 1a and 1b). Computed tomography angiography (CTA) was also performed. CTA finding showed tortuous, dilated connection between LAD and main pulmonary artery, 2.3 cm in diameter.




Figure 1. Coronary angiogram showing dilated and tortuous vessel arising from left anterior descending coronary artery (yellow arrow). 1a)Cranial right anterior oblique projection; 1b)Caudal left anterior oblique projection.

Surgical repair was then carried out for this patient. Operative finding was tortuous and dilated fistula originated from left anterior descending (LAD) coronary artery (Figure 2a) draining to main pulmonary artery (MPA) along with calcification at proximal to mid right coronary artery (RCA), chronic total occlusion (CTO) in distal RCA, calcification at mid left circumflex (LCx) artery, calcification at anterior mid-LAD and annular dilatation of mitral valve and chordal rupture of P2 and P3, cleft at A1 and A2 of mitral valve.

Three procedures were performed for this patient. Ligation of coronary artery fistula was first carried out, followed by mitral valve repair and then coronary artery bypass graft. Mammary graft was performed later after mitral valve repair.

Coronary artery fistula was identified and then ligated at proximal and distal end of the fistula (Figure 2b). Origin of fistula was ligated using Teflon pledget reinforced direct suture as well as distal end of the fistula

just before MPA. Coronary bypass graft (CABG) using left internal mammary artery and two saphenous vein conduits and mitral valvuloplasty and annuloplasty were also performed. Two saphenous vein conduits and left internal mammary artery were grafted, one vein conduit to Right Posterior Descending Artery, (distal), one vein conduit to distal Right Coronary Artery, and left internal mammary artery to distal left anterior descending (LAD) coronary artery. Valvuloplasty was performed by triangular resection of P2 and closure of cleft between A1 and A2 using 6-0 Prolene. Physio ring was implanted for annuloplasty. Physio ring was stitched using 12 non-pledget suture with 2-0 Premicron® suture. Duration of cardiopulmonary bypass was 256 minutes and duration of aortic cross clamp was 185 minutes. Post-operative blood pressure was 118/68 mmHg, heart rate 100 bpm, central venous pressure 9 cmH₂O and 99% of peripheral oxygen saturation. Post-operative hemodynamic support was 1 mg/hour of Nitroglycerin and 5 gammas of Dobutamine.




Figure 2. 2a) (left) Operative findings yellow arrow shows dilated and tortuous segment arising from LAD; 2b) (right) Yellow arrow shows ligated proximal end of coronary artery fistula.

After surgery patient was admitted to ICU for four days and then discharged on post-operative day 17 with no cardiovascular related complication.

After one-year follow-up patient underwent echocardiography and cardiac MSCT Scan. Echocardiogram showed normal cardiac chamber and kinetic, as well as normal left and right ventricular function and ejection fraction was 82%. No obvious fistulation into pulmonal trunk was shown in cardiac MSCT scan (Figure 3b). Multiple patent graft was also shown. (Figure 4).




Figure 3. 3a) (left) CT angiogram demonstrated connection of fistula to main pulmonary artery (MPA) (red arrow); 3b) (right) Axial slice of cardiac MSCT angiogram one-year post operatively showed no obvious fistulation to main pulmonary artery (MPA).




Figure 4. Three-dimensional computed tomography image of after one year follow up showed multiple patent grafts (SVG to RPDA, SVG to distal RCA, LIMA to LAD).

Case #2

Fifty-seven-year-old male presented with symptoms of discomfort around left area of the neck. Discomfort was felt for almost a year. Patient then had himself check to neurologist and then referred to cardiologist before then referred to our hospital.

No abnormalities were found during physical examination. Echocardiography showed normal cardiac chambers and valves, with 64% ejection fraction and normal systolic and diastolic function. Fistulation from

proximal LAD was shown and connect to MPA on cardiac catheterization (Figure 5a and 5b). Chronic total occlusion at proximal of LAD and proximal of RCA, 99% stenosis at distal of LCx were also shown.

During operation, tortuous fistulation from LAD to MPA was shown (Figure 6). The aforementioned fistula was then ligated by Teflon-pledged-reinforced direct suture at the LAD side. Coronary artery bypass graft was also performed which are left internal mammary artery to LAD, saphenous vein graft to distal of RCA, and saphenous vein graft to distal part of LCx.




Figure 5a and 5b. Fistulation from LAD to MPA was shown on cranial right anterior oblique projection and right anterior oblique projection of coronary catheterization (yellow arrow) along with chronic total occlusion of proximal LAD and 99% stenosis at distal of LCx.

²⁹ Aortic cross clamp time of operation was 79 minutes and 127 minutes of cardiopulmonary bypass time. No problem was encountered during operation and patient was hemodynamically supported by 0.5 nano of nitroglycerine and 50 nano of norepinephrine.

Post-operative echocardiogram was performed and trivial mitral regurgitation and trivial tricuspid regurgitation was shown. Left ventricular systolic function was 65% by Teich and 67% by biplane.

Patient was then admitted to intensive care unit and later admitted to elective ward after 3 days. On the seventh day of care patient was then discharge without any significant complications. Short-term follow-up showed improvements of symptoms without any limitations of daily activities.

Discussion

Coronary artery fistula can occur congenitally or acquired. Most coronary artery fistula are congenital. Acquired coronary artery fistula are extremely rare and usually iatrogenic, posttraumatic, or caused by Takayasu arteritis, Kawasaki disease or chest irradiation^{5,6,7}. The feeding artery of the fistula may drain from a main coronary artery or one of its branches and is usually

a dilated and tortuous artery terminating in one of the cardiac chambers of vessel¹. It may also present with aneurysm of the vessel itself⁸. The fistula may originate from right coronary artery (52%), left anterior descending coronary artery (30%) and circumflex artery (18%)⁸. Fistulas originated from both coronary arteries have also been reported^{9,10,11}. The types of coronary artery fistula are coronary to pulmonary artery (76.8%), coronary to bronchial artery fistula (8.9%), coronary artery to cardiac chamber fistula (8.9%), combined coronary to pulmonary and coronary to bronchial artery (3.6%), and coronary artery to superior vena cava fistula (1.8%)³. Another congenital anomaly like single coronary artery has been reported concomitant with coronary artery fistula¹².

¹⁰ Regarding findings in these cases, coronary artery fistula may present with concomitant coronary artery disease. No direct relationship was reported in literature about coronary artery fistula and coronary artery disease, but in most cases coronary artery fistula was found during coronary angiogram for diagnosis of coronary artery disease. Inoue et al reported coronary-pulmonary artery fistula with coronary artery disease related to Kawasaki disease⁶.

Mitral regurgitation found in case #1 was probably related to ischemic heart disease caused by coronary artery disease. In patients with coronary artery disease especially coronary disease involving left anterior descending coronary artery or left circumflex artery or right posterior descending artery (depending on dominance) for the supply papillary muscles^{13,14}. No literature was found regarding relationship of mitral regurgitation and coronary artery fistula, yet coronary artery fistula may complicate into mitral valve endocarditis¹⁵.

Coronary artery fistula may produce hemodynamic changes. When the fistula drains to the right side of the heart, the volume load is increased to the right heart as well as to the pulmonary vascular bed, the left atrium and the left ventricle. When the fistula drains into the left atrium or the left ventricle, there is volume overloading of these chambers but no increase in the pulmonary blood flow. The size of the shunt is determined by the size of the coronary chamber into which the fistula drains¹.

Coronary arterial fistula is usually asymptomatic in the first two decades, especially when they are hemodynamically small. Even though most patients are asymptomatic, it may also cause serious consequences i.e. myocardial ischemia, thromboembolism, cardiac failure, arrhythmias, rupture, and endocarditis^{16,17}. Spontaneous rupture of the aneurysmal fistula causing hemopericardium has also been reported¹. It is also reported to be associated with bronchiectasis and hypoplasia of pulmonary artery.

Diagnosis of coronary artery fistula can be made by two-dimensional echocardiography and color doppler while coronary angiography and cardiac catheterization are necessary investigations to decide the exact location and thus the mode of intervention to close the fistula¹⁸. Coronary Computed Tomography Angiography (CTA) is also useful for making diagnosis of coronary artery fistula³. Gundogdu et al recommend multi detector computed tomography to be used to precise delineation of coronary fistulas¹⁹.

Without concomitant heart condition, the indications for treatment of coronary arterial fistulas include the present of a large or increasing left-to-

right shunt, left ventricular overload, myocardial ischemia, left ventricular dysfunction, congestive heart failure and for prevention of endocarditis/ endarteritis. Volume overload of any heart chamber producing secondary valve disease, heart failure, significant left-to-right shunt, disturbed cardiac rhythms and evidence of ischemia necessitate early closure. Indications for interventions in asymptomatic patients are controversial but presence of increased left atrial diameter and Qp/Qs ratio more than 1.3 need early interventions to prevent further complication. Surgery for asymptomatic patients remains controversy but some surgeons advocate operative treatment of coronary artery fistula due to the possibility of sudden death without any warrant symptoms⁹.

Coronary artery fistula may be closed surgically or using coil embolization via trans-catheter approach²⁰. Saboo *et al.* point out indications for transcatheter closure and indications for surgical management. Transcatheter closure is indicated when a CAF is proximally located with single drainage, no complicated communications and single drainage site. Transcatheter closure can be performed using coil embolization and may followed by stent implantation if patient has concomitant coronary artery disease²¹. Surgical management is indicated when there are distal fistulas; possible accidental occlusion of important vessels by embolization, multiple complex communications; tortuous arteries, prominent aneurysm, wide drainage site and drainage to coronary sinus⁵. Approach for surgical management may be median sternotomy, left anterior mini-thoracotomy or left parasternal approach^{22,23,24}. Some reports also stated that surgical closure is the gold standard for treatment with excellent prognosis^{4,18}. In these cases we performed telfonpledget reinforced direct suture to close fistulation of coronary artery. Telfonpledget was used for reinforcement so that the suture will not cut or severe the vessel or the fistula.

Even though outcome from most surgical closure and transcatheter interventions are satisfactory, recurrence of coronary artery fistula has been reported. The exact mechanism of recurrence is still unknown but Kostis *et al.*²⁵ pointed out that the pulsatile aortic pressure transmitted to the probable small residual

lumen between the in-folded walls of the vessel at the level of the ligature resulted in reestablishment of the fistula, recommending that division and over-sewing should be done whenever feasible to prevent coronary artery fistula from recurrence. Despite ligation and division recurrence may still be occurred therefore long-term follow-up of patient is essential.

Conclusion

³ Coronary artery fistula is a rare anomaly of coronary artery that usually are congenital. Most case are asymptomatic but it may cause hemodynamic changes that may result in harmful complications. Diagnostic tools for coronary artery fistula are echocardiography, coronary angiography and CT angiography. Management of coronary artery fistula can be transcatheter closure or surgical repair. Long term follow-up is essential since recurrence may occur. In our cases surgical management with by Teflon-pledget-reinforced direct suture of coronary artery fistula of left descending coronary artery to pulmonary provide satisfactory outcome with uneventful result and remarkable recovery of patients.

¹³ **Conflict of Interest :** The author declare that they have no conflict of interest.

Source of Funding : None.

Acknowledgements: Special thanks toEJA Team, Indonesia for editing the manuscript.

³⁶ **Ethical Approval:** This study approved by Dr. Soetomo General Hospital, Surabaya, Indonesia.

References

1. Qureshi SA. Coronary Artery Fistulas. *Oprhanet Journal of Rare Disease*. 2006; 1(151).
2. Zhou K, Kong L, Wang Y, *et al*. Coronary Artery fistula in Adults Evaluation with Dual-Source CT Coronary Angiography. *BR Journal Radiology*. 2-15; 88.
3. Lim JJ, Jung IJ, Lee BY, *et al*. Prevalence and Types of Coronary Artery Fistula Detected With Coronary Angiography. *AJR*. 2014; 203.
4. Chen PF, Tang L, Liu Zz, *et al*. Superior Vena Cava Syndrome Secondary to Recurrent Coronary Artery Fistula. *Medicine*. 2017; 96.

5. Saboo SS, Juan YH, Khandelwal A, *et al*. MDCT of Congenital Coronary Artery Fistula. *AJR*. 2014; 203.
6. Inoue T, Miyake T, Mushiake S. A Case of Coronary-Pulmonary Artery Fistula with Coronary Artery Aneurysm Detected for Kawasaki disease Remote Phase. *J Clin Ultrasound*. 2019.
7. Gentry J, Summers M, Harb S, *et al*. Iatrogenic Left Circumflex Coronary Artery Fistula After Mitral Valve Replacement. *CASE: Cardiovascular Imaging Case Report*. 2018; 2(5).
8. McNamara J, Gross R. Congenital Coronary Artery Fistula. *Surgery*. 1969; 65.
9. Carrel T, Tkebuchava T, Jenni R, *et al*. Congenital Coronary Fistulas in Children and Adults: Diagnosis, Surgical Technique and Results. *Cardiology*. 1996; 87.
10. da Silva FP, Donato P, Donato H, *et al*. Symptomatic Bilateral Coronary Artery Fistula to Pulmonary Artery in Elderly Patient. *European Journal of Radiology Open*. 2016; 3.
11. Kim CY, Choi JY, Kim KS. A Coronary Artery Fistula Having Connection Between 2 Coronary Arteries And The Left Ventricle: A Case Report. *Medicine*. 2017; 96(45).
12. Xie LJ, Zhang L, Xiao TTSJ. Coronary Artery Fistula Between Single Right Coronary Artery and Right Pulmonary Artery: A Case Report And Literature Review. *BMC Cardiovascular Disorders*. 2015; 15.
13. Chiewe J, Cooke D, Weiss A. *Oxford Specialist Handbooks in Surgery: Cardiothoracic Surgery*. 2nd ed. Oxford: Oxford University Press; 2013.
14. Berdajs D, Turina MI. *Operative Anatomy of The Heart*. 1st ed. Laussane: Springer; 2011.
15. Shi M, Wang Y, Pang L, *et al*. Coronary Artery Fistula Associated with Mitral Valve Endocarditis. *J Card Surg*. 2012; 27.
16. Said SM, Burkhart HM, Schaff HV, *et al*. Late Outcome of Repair of Congenital Coronary Artery Fistulas-A Word of Caution. *The Journal of Thoracic and Cardiovascular Surgery*. 2012; 145(2).
17. Santos EL, Gouveia MM, Silva RF, *et al*. Coronary Artery Fistula Causing Acute Myocardial Infarction and Right Ventricle Thrombus. *Cureus*. 2018; 10(3).
18. Waqar T, Rizvi HMFAAMZA. Surgical Repair of Congenital Coronary Artery Fistula: A Case Report.

- J Pak Med Assoc. 2018; 28.
19. Gundogdu F, Arslan S, Buyukkaya E. Coronary artery Fistula in A Patient with Coronary Artery Disease: Evaluation by Coronary Angiography and Multidetector Computed Tomography. *Int J Cardiovasc Imaging*. 2007; 23.
 20. Al-Douri A, Cedars A, Tran D. Coronary Artery Fistula Between The Left Anterior Descending Artery and Pulmonary Artery. *Baylor University Medical Center Precoeding*. 2018; 31(1).
 21. Lee WC, Fang HY, Wu CJ. How to Treat The Combination of Coronary Artery Fistula and Occluded Coronary Artery. *Medicine*. 2018; 97(14).
 22. Mitropoulos FA, Kanakis MA, Chatzis A, Contrafouris C, Sodianidou IA, Luoulias AG. Minimal Invasive Coronary Artery Fistula Ligation. *Korean J Thorac Cardiovasc Surg*. 2014; 47.
 23. Zhang Q, Duan Y, Hongxin L, *et al*. An Innovative Technique of Periventricular Device Closure of A Coronary Artery Fistula Through A Left Parasternal Approach. *European Heart Journal*. 2017.
 24. Kondo N, Totsugawa T, Yoshitaka H, *et al*. A Case of Coronary Artery Fistula Ligation in Minimally Invasive Mitral Valve Plasty via Right Anterolateral Minithoracotomy. *Sage*. 2019; 14(2).
 25. Kostis JB, Burns JJ, Moreyra AE, Pichard AD. Recurrent Coronary Artery Fistula. *Clinical Cardiology*. 1984; 7:307-313.

Left Anterior Descendent Coronary Artery Fistula

ORIGINALITY REPORT

17%

SIMILARITY INDEX

9%

INTERNET SOURCES

16%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Nosakhare O. Erhunmwunse, Timeyin J. Pajiah, Ogbodo Victory, Endurance Ewere. "Does short-term exposure to a Neonicotinoid insecticide trigger Biochemical and Physiological responses in Juvenile catfish?", Research Square Platform LLC, 2022
Publication 1%
- 2** Xinyu Chang, Uchechukwu Edna Obianwuna, Jing Wang, Haijun Zhang, Guanghai Qi, Kai Qiu, Shugeng Wu. "Glycosylated proteins with abnormal glycosylation changes are potential biomarkers for early diagnosis of breast cancer", International Journal of Biological Macromolecules, 2023
Publication 1%
- 3** Xie, Mingxing, Ling Li, Tsung O. Cheng, Zhenxing Sun, Xinfang Wang, Qing Lv, Hua Peng, Li Yuan, Jing Wang, and Li Zhang. "Coronary artery fistula: Comparison of diagnostic accuracy by echocardiography versus coronary arteriography and surgery in 63 patients studied between 2002 and 2012 in

a single medical center in China",
International Journal of Cardiology, 2014.

Publication

4

www.researchsquare.com

Internet Source

1 %

5

Gabin Yun, Tae Hyun Nam, Eun Ju Chun.
"Coronary Artery Fistulas: Pathophysiology,
Imaging Findings, and Management",
RadioGraphics, 2018

Publication

1 %

6

V.H. Contreras-Gutiérrez, R.A. Téllez-Ramírez.
"Coronary arterio-venous fistula associated
acute coronary syndrome: A case-report and
review of literature", Revista Médica del
Hospital General de México, 2017

Publication

1 %

7

fk.unair.ac.id

Internet Source

1 %

8

www.mp.pl

Internet Source

1 %

9

Luca Favero. "Cardiac and extracardiac
complications during CTO interventions:
prevention and management", Interventional
Cardiology, 06/2010

Publication

1 %

10

Niyazi Cebi. "Congenital coronary artery
fistulas in adults: concomitant pathologies

1 %

and treatment", The International Journal of
Cardiovascular Imaging, 04/2008

Publication

11

www.accessdata.fda.gov

Internet Source

<1 %

12

Negin Yavari, Mina Ghorbanpour Landy,
Yasaman Motevali, Elham Tavousi Tabatabaei
et al. "Iatrogenic left circumflex coronary
occlusion following mitral valve replacement
surgery: A case report", Clinical Case Reports,
2021

Publication

<1 %

13

Shingo Kunioka, Hiroto Kitahara, Seima Ohira,
Yuki Tada, Nobuyuki Akasaka, Hiroyuki
Kamiya. "Surgical treatment of multiple
coronary-to-pulmonary artery fistulas with
severe coronary artery stenosis", General
Thoracic and Cardiovascular Surgery, 2019

Publication

<1 %

14

Velvis, H.. "Diagnosis of coronary artery fistula
by two-dimensional echocardiography, pulsed
doppler ultrasound and color flow imaging",
Journal of the American College of Cardiology,
198910

Publication

<1 %

15

www.wjgnet.com

Internet Source

<1 %

16

Anne Cecilie K. Larstorp, Camilla Lund Søråas, Theis Tønnessen, Carl Müller, Sverre E. Kjeldsen, Arild Mangschau. "Scintigraphic demonstration of myocardial perfusion and ischaemia associated with coronary artery bypass grafting", *Scandinavian Cardiovascular Journal*, 2009

Publication

<1 %

17

Manolis Vavuranakis, Charles A. Bush, Harisios Boudoulas. "Coronary artery fistulas in adults: Incidence, angiographic characteristics, natural history", *Catheterization and Cardiovascular Diagnosis*, 1995

Publication

<1 %

18

Yiginer, O.. "Demonstration of coronary-to-pulmonary fistula with MDCT and conventional angiography", *International Journal of Cardiology*, 20090529

Publication

<1 %

19

www.umm.edu

Internet Source

<1 %

20

T. Carrel, T. Tkebuchava, R. Jenni, U. Arbenz, M. Turina. "Congenital Coronary Fistulas in Children and Adults: Diagnosis, Surgical Technique and Results", *Cardiology*, 1996

Publication

<1 %

- | | | |
|----|--|------|
| 21 | cyberleninka.org
Internet Source | <1 % |
| 22 | eurointervention.pcronline.com
Internet Source | <1 % |
| 23 | www.cct.gr.jp
Internet Source | <1 % |
| 24 | www.karger.com
Internet Source | <1 % |
| 25 | Hitoshi Hirose, Masatake Takagi, Na.
"Coronary Atherosclerosis with Dual Coronary
Artery Fistulas", Scandinavian Cardiovascular
Journal, 2009
Publication | <1 % |
| 26 | J. G. Brockmann, A. Vaidya, S. Reddy, P. J.
Friend. "Retrieval of abdominal organs for
transplantation", British Journal of Surgery,
2006
Publication | <1 % |
| 27 | Mohamed Sayyoub, Elizabeth Lee, Nicole
Bhave, Karen Kim, Prachi P. Agarwal. "Imaging
and Management of Coronary Artery
Anomalies", Current Treatment Options in
Cardiovascular Medicine, 2020
Publication | <1 % |
| 28 | Novick, R.J.. "Analysis of the learning curve in
telerobotic, beating heart coronary artery | <1 % |

bypass grafting: a 90 patient experience", The Annals of Thoracic Surgery, 200309

Publication

29

Ryoji Iida, Tomonori Yamamoto, Noriko Kondo, Takahiro Suzuki, Shigeru Saeki, Setsuro Ogawa. "Identification of the Site of Drainage of Left Main Coronary Artery to Right Atrium Fistula With Intraoperative Transesophageal Echocardiography", Journal of Cardiothoracic and Vascular Anesthesia, 2005

Publication

<1 %

30

academic.oup.com

Internet Source

<1 %

31

archive.org

Internet Source

<1 %

32

bmccardiovascdisord.biomedcentral.com

Internet Source

<1 %

33

cardiothoracicsurgery.biomedcentral.com

Internet Source

<1 %

34

www.invasivecardiology.com

Internet Source

<1 %

35

Kevin Kalisz, Alison E. Sanders, Ryan Avery, Bradley D. Allen. "Coronary Artery Fistulas", Journal of Thoracic Imaging, 2021

Publication

<1 %

36

Priskila Christy, Hanna Evelina Sidjabat, Anggia Augustasia Lumban Toruan, Emmanuel Jairaj Moses et al. "Comparison of Laboratory Diagnosis of Urinary Tract Infections Based on Leukocyte and Bacterial Parameters Using Standardized Microscopic and Flow Cytometry Methods", International Journal of Nephrology, 2022

Publication

<1 %

37

Wei-Chieh Lee, Hsiu-Yu Fang, Chiung-Jen Wu. "How to treat the combination of coronary artery fistula and occluded coronary artery", Medicine, 2018

Publication

<1 %

38

"Thoracic Surgery", Springer Science and Business Media LLC, 2020

Publication

<1 %

39

Chiu, S.N.. "Acquired coronary artery fistula after open heart surgery for congenital heart disease", International Journal of Cardiology, 20050818

Publication

<1 %

40

Masafumi Takahashi. "Multicystic aneurysmal dilatation of bilateral coronary artery fistula", Catheterization and Cardiovascular Diagnosis, 04/1994

Publication

<1 %

41

ŞİŞMAN, Seviye Bora, DOĞAN, Demet, ŞAHİN, Sinan, ŞAHAN, Hasan and EMRE, Ayşe.

"Koronar arter fistüllerinin tanısında çok kesitli bilgisayarlı tomografinin yeri", Türk Göğüs Kalp Damar Cerrahisi Derneği, 2018.

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Left Anterior Descendent Coronary Artery Fistula

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8
