



Vol. 33 No. 1 (2023): MAJALAH BIOMORFOLOGI

Current Issue

E-ISSN: 2716-0920

P-ISSN: 0215-8833

MAJALAH BIOMORFOLOGI

Biomorphology Journal

Vol. 33 No. 1 January 2023


Published by: Universitas Airlangga
Jl. Dr. Ir. H. Soekarno, Mulyorejo, Kec. Mulyorejo,
Surabaya, Indonesia 60115
in collaboration with
(Ikatan Dokter Indonesia - IDI Surabaya)
(PAAI- Perhimpunan Ahli Anatomi Indonesia)




Vol. 33 No. 1 (2023): MAJALAH BIOMORFOLOGI


Published: 2023-01-10

RESEARCH



THORACIC CT-SCAN WITH EGFR MUTATIONS IN PULMONARY ADENOCARCINOMA PATIENTS WITH PULMONARY METASTASES AT DR. MOEWARDI HOSPITAL, SURAKARTA, INDONESIA

 DOI : 10.20473/mbiom.v33i1.2023.1-6

 Viska Wahyu Arianti , Widiastuti Soewondo , Sulistiyani Kusumaningrum , Muhammad Lukman Hermansyah , Yusup Subagio Sutanto , Hari Wujoso

 1-6

 Abstract : 198

 PDF : 103



CLINICAL PROFILE OF TYPE-1 DIABETES PEDIATRIC PATIENTS IN DR. SOETOMO GENERAL ACADEMIC HOSPITAL SURABAYA: CORRELATION OF GROWTH STATUS AND METABOLIC CONTROL

 DOI : 10.20473/mbiom.v33i1.2023.7-13


 Grawira Drana Putra Respati , Muhammad Faizi , Agung Pranoto , Mia Ratwita Andarsini

 7-13

 Abstract : 135

 PDF : 82





BODY MASS INDEX WITH SYSTOLIC AND DIASTOLIC BLOOD PRESSURE AT A PUBLIC HEALTH CENTER IN SURABAYA, INDONESIA

 DOI : 10.20473/mbiom.v33i1.2023.14-19

 Bella Patricia Simanjong , Florentina Sustini , Sundari Indah Wiyasihati

 14-19

 Abstract : 126


 PDF : 46



MOLECULAR DOCKING OF CROCUS SATIVUS PHYTOCHEMICALS AGAINST INDUCIBLE NITRIC OXIDE SYNTHASE AND PHOSPHODIESTERASE-9 IN HEART FAILURE PRESERVED EJECTION FRACTION

 DOI : 10.20473/mbiom.v33i1.2023.20-29

 Syafira Yasmine , Neissya Nastiti Firmanto , Annisa Maya Sabrina , Siti Khaerunnisa

 20-29


 Abstract : 144


 PDF : 91






THE HDL LEVEL IN DIABETIC FOOT ULCER PATIENTS WITH AND WITHOUT AMPUTATION AT DR. SEOTOMO GENERAL ACADEMIC HOSPITAL SURABAYA

 DOI : 10.20473/mbiom.v33i1.2023.30-37

 Ahmed Eliaan Shaker Abuajwa , Hermina Novida , Heroe Soebroto , Yan Efrata Sembiring

 30-37

 Abstract : 153


 PDF : 91


 PDF



INSULIN THERAPY IN T2DM PATIENTS IN DIABETES OUTPATIENT CLINIC, DR. SOETOMO GENERAL ACADEMIC HOSPITAL, SURABAYA, INDONESIA

 DOI : 10.20473/mbiom.v33i1.2023.38-43

 Adinda Ayu Dyah Rahadini , Sri Murtini , Sidarti Soehita

 38-43

 Abstract : 146

 PDF : 75


 PDF

REVIEW



DIFFERENTIATION OF SPINAL TUBERCULOSIS AND METASTATIC SPINAL TUMOR USING MRI FEATURE: A SYSTEMATIC REVIEW

 DOI : 10.20473/mbiom.v33i1.2023.44-51

 Justicia Izza Tsuroya , Muhammad Faris , Paulus Rahardjo

 44-51

 Abstract : 133

 PDF : 167


 PDF



HYPERBARIC OXYGEN THERAPY AS AN ADJUVANT TREATMENT IN HYDROCHLORIC ACID POISONING: A LITERATURE REVIEW

 DOI : 10.20473/mbiom.v33i1.2023.52-58

 Desi Rahmah Sari , Ismi Dian Meiliana , Dewangga Satria Sakti Kinasih , Hamidah Kurniasari , Purwo Sri Rejeki

 52-58

 Abstract : 137

 PDF : 102

 PDF

Login

Username *

Password *

[Forgot your password?](#)

Keep me logged in

Login

Register

National Accreditation



[View certificate here](#)

Accreditation Number: 0010/E5/KI.02.04/2022

Instruction for Author

[Guidelines for Authors](#)

[Online submissions](#)

[Submission Template](#)

[Authors' Declaration](#)

[Patient Consent for Publication](#)

Journal Policy

[Focus and Scope](#)

[Publication Ethics](#)

[Article Processing Charge](#)

[Editorial Team](#)

Peer Review Process	Peer Reviewers
Open Access Policy	Archiving
Plagiarism Statement	Copyright
Repository Policy	ORCID ID Policy
License Term	Contact
Old Website	

Meet Our Editorial Team



Viskasari P. Kalanjati
 Editor in Chief
 Universitas Airlangga, Surabaya, Indonesia
 54388384000



Abdurrachman Abdurrachman
 Advisory Editor
 Universitas Airlangga, Surabaya, Indonesia
 57205111921



Tania Ardiani
 Associate Editor
 Indonesian Association of Anatomist (IAA), Indonesia
 -

[Read More](#)

Tools



Journal Standing

Scopus[®]

In Collaboration



Links



Visitors



Indexed In





Information

For Readers

For Authors

For Librarians

Keywords



Address

Majalah Biomorfologi (Biomorphology Journal)
UKJF, Faculty of Medicine, Universitas Airlangga
Jl. Mayjen. Prof. Dr. Moestopo 47, Surabaya 60131, Indonesia

Contact Info:

Email: mbiomj@fk.unair.ac.id



Lembaga Inovasi, Pengembangan Jurnal,
Penerbitan dan Hak Kekayaan Intelektual

LIPJPHKI

Gedung AUP, Kampus C, Universitas Airlangga, Kota Surabaya, Jawa Timur, 60115



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).



Editorial Team



Viskasari P. Kalanjati

Editor in Chief

Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

 0000-0002-7005-0025

Google Scholar

54388384000

 5975075



Abdurachman

Advisory Editor

Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

 0000-0002-8166-3898

Google Scholar

57193135762

 5992333



Tania Ardiani

Associate Editor

Indonesian Association of Anatomist (IAA), Indonesia

 -

Google Scholar

-

 6074585



Irmawan Farindra

Editorial Team

Faculty of Medicine, Universitas Nahdatul Ulama Surabaya,

Surabaya, Indonesia

 -

Google Scholar

-

 6644853



Tjahja Nurrobi

Editorial Team

Indonesian Association of Anatomist (IAA), Indonesia

 0000-0002-1433-9932

Google Scholar

-



-



Elies Fitriani

Editorial Team

Indonesian Association of Anatomist (IAA), Indonesia

 0000-0001-7129-0226

Google Scholar

-



6748930



Ulfah Dian Indrayani

Editorial Team

Indonesian Association of Anatomist (IAA), Indonesia

 -

Google Scholar

-



6587260



Azlin Muhammad

Editorial Team

Department of Medical Parasitology and Entomology, Faculty of Medicine, Universiti Kebangsaan Malaysia, Malaysia

 0000-0001-9308-4175

-

8625546000



-



Stephanie Miller

Editorial Team

Centre for Clinical Research, The University of Queensland, Australia

 0000-0002-5565-2001

-

35082353900



-



Dzanuar Rahmawan

Editorial Team

Institut Ilmu Kesehatan Bhakti Wiyata, Kediri, Indonesia



Google Scholar

-



-

Ira Anggar Kusuma

Editorial Team

Faculty of Dental Medicine, Universitas Diponegoro, Semarang, Indonesia



Google Scholar

-



6794040

Etha Rambung

Editorial Team

Faculty of Medicine, University of Ciputra, Indonesia



Google Scholar

-



6664132

Risdiansyah Risdiansyah

Editorial Team

Indonesian Medical Association (IDI), Surabaya, Indonesia



Google Scholar

-



-

Peppy Nawangsasi

Editorial Team

Faculty of Medicine, Universitas Hangtuh Surabaya, Indonesia



Google Scholar

-



-

Rudi Irawan

Editorial Team

Institut Ilmu Kesehatan Bhakti Wiyata, Kediri, Indonesia




Google Scholar

-



-






Mujaddidah Mujaddidah

Editorial Team

Faculty of Medicine, Universitas Muhammadiyah, Surabaya, Indonesia

 0000-0002-6753-7190
Google Scholar


 6163373



Dion Krismashogi Dharmawan

Editorial Team

Faculty of Medicine, Universitas Jember, Indonesia

 0000-0003-0218-3712
Google Scholar


 6067402



Dwi Martha

Editorial Team

Faculty of Medicine, Universitas Surabaya, Surabaya, Indonesia

 0000-0002-3550-0289
Google Scholar


 -




Jesslyn Kristanti

Editorial Team

Faculty of Medicine, Universitas Hang Tuah, Surabaya, Indonesia

 0000-0002-3472-3085
Google Scholar


 -



Astri Dewayani

Editorial Team

Oita University, Japan

 0000-0001-9577-8373
Google Scholar

 6751213



Anggie Kusumawardhani

Assistant Editor

Unit of Journal Consortium and Folia Medica Indonesiana,
Universitas Airlangga, Surabaya

-

-

-



Endro Sugiharto

Assistant Editor

Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

-

-

-



Moch. Zuhdy

Assistant Editor

Unit of Journal Consortium and Folia Medica Indonesiana,
Universitas Airlangga, Surabaya, Indonesia

-

-

-



Moch. Rudy

Assistant Editor

Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

-

-

-



Dimas B. Bagus Pamungkas

Assistant Editor

Faculty of Medicine, Universitas Muhammadiyah, Surabaya,
Indonesia

-

-

-



Login

Username *

Password *

[Forgot your password?](#)

Keep me logged in

Login

Register

National Accreditation



[View certificate here](#)

Accreditation Number: 0010/E5/KI.02.04/2022

Instruction for Author

[Guidelines for Authors](#)

[Online submissions](#)

[Submission Template](#)

[Authors' Declaration](#)

[Patient Consent for Publication](#)

Journal Policy

[Focus and Scope](#)

[Publication Ethics](#)

[Article Processing Charge](#)

[Editorial Team](#)

Peer Review Process	Peer Reviewers
Open Access Policy	Archiving
Plagiarism Statement	Copyright
Repository Policy	ORCID ID Policy
License Term	Contact
Old Website	

Meet Our Editorial Team



Viskasari P. Kalanjati
 Editor in Chief
 Universitas Airlangga, Surabaya, Indonesia
 54388384000



Abdurrachman Abdurrachman
 Advisory Editor
 Universitas Airlangga, Surabaya, Indonesia
 57205111921



Tania Ardiani
 Associate Editor
 Indonesian Association of Anatomist (IAA), Indonesia
 -

[Read More](#)

Tools



Journal Standing

Scopus[®]

In Collaboration



Links



Visitors



Indexed In






Information

For Readers

For Authors

For Librarians

Keywords



Address

Majalah Biomorfologi (Biomorphology Journal)
UKJF, Faculty of Medicine, Universitas Airlangga
Jl. Mayjen. Prof. Dr. Moestopo 47, Surabaya 60131, Indonesia

Contact Info:

Email: mbiomj@fk.unair.ac.id



Lembaga Inovasi, Pengembangan Jurnal,
Penerbitan dan Hak Kekayaan Intelektual

LIPJPHKI

Gedung AUP, Kampus C, Universitas Airlangga, Kota Surabaya, Jawa Timur, 60115



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).



MAJALAH BIOMORFOLOGI

UNIVERSITAS AIRLANGGA

P-ISSN : 02158833 < > E-ISSN : 27160920

0.16666
7
Impact Factor40
Google CitationsSinta 4
Current
Accreditation
[Google Scholar](#)
[Garuda](#)
[Website](#)
[Editor URL](#)

History Accreditation

2020 2021 2022 2023 2024

[Garuda](#) [Google Scholar](#)

INSULIN THERAPY IN T2DM PATIENTS IN DIABETES OUTPATIENT CLINIC, DR. SOETOMO GENERAL ACADEMIC HOSPITAL, SURABAYA, INDONESIA

Universitas Airlangga, Surabaya, Indonesia [Majalah Biomorfologi Vol. 33 No. 1 \(2023\): MAJALAH BIOMORFOLOGI 38-43](#)

2023 DOI: [10.20473/mbiom.v33i1.2023.38-43](https://doi.org/10.20473/mbiom.v33i1.2023.38-43) Accred : Sinta 4

MOLECULAR DOCKING OF CROCUS SATIVUS PHYTOCHEMICALS AGAINST INDUCIBLE NITRIC OXIDE SYNTHASE AND PHOSPHODIESTERASE-9 IN HEART FAILURE PRESERVED EJECTION FRACTION

Universitas Airlangga, Surabaya, Indonesia [Majalah Biomorfologi Vol. 33 No. 1 \(2023\): MAJALAH BIOMORFOLOGI 20-29](#)

2023 DOI: [10.20473/mbiom.v33i1.2023.20-29](https://doi.org/10.20473/mbiom.v33i1.2023.20-29) Accred : Sinta 4

HYPERBARIC OXYGEN THERAPY AS AN ADJUVANT TREATMENT IN HYDROCHLORIC ACID POISONING: A LITERATURE REVIEW

Universitas Airlangga, Surabaya, Indonesia [Majalah Biomorfologi Vol. 33 No. 1 \(2023\): MAJALAH BIOMORFOLOGI 52-58](#)


2023 DOI: [10.20473/mbiom.v33i1.2023.52-58](https://doi.org/10.20473/mbiom.v33i1.2023.52-58) Accred : Sinta 4

CLINICAL PROFILE OF TYPE-1 DIABETES PEDIATRIC PATIENTS IN DR. SOETOMO GENERAL ACADEMIC HOSPITAL SURABAYA: CORRELATION OF GROWTH STATUS AND METABOLIC CONTROL

Universitas Airlangga, Surabaya, Indonesia [Majalah Biomorfologi Vol. 33 No. 1 \(2023\): MAJALAH BIOMORFOLOGI 7-13](#)

2023 DOI: [10.20473/mbiom.v33i1.2023.7-13](https://doi.org/10.20473/mbiom.v33i1.2023.7-13) Accred : Sinta 4

Citation Per Year By Google Scholar



Journal By Google Scholar

	All	Since 2018
Citation	40	36
h-index	3	3
i10-index	0	0

[THORACIC CT-SCAN WITH EGFR MUTATIONS IN PULMONARY ADENOCARCINOMA PATIENTS WITH PULMONARY METASTASES AT DR. MOEWARDI HOSPITAL, SURAKARTA, INDONESIA](#)

Universitas Airlangga, Surabaya, Indonesia [Majalah Biomorfologi Vol. 33 No. 1 \(2023\): MAJALAH BIOMORFOLOGI 1-6](#)

2023 [DOI: 10.20473/mbiom.v33i1.2023.1-6](#) [Accred : Sinta 4](#)

[THE HDL LEVEL IN DIABETIC FOOT ULCER PATIENTS WITH AND WITHOUT AMPUTATION AT DR. SEOTOMO GENERAL ACADEMIC HOSPITAL SURABAYA](#)

Universitas Airlangga, Surabaya, Indonesia [Majalah Biomorfologi Vol. 33 No. 1 \(2023\): MAJALAH BIOMORFOLOGI 30-37](#)

2023 [DOI: 10.20473/mbiom.v33i1.2023.30-37](#) [Accred : Sinta 4](#)

[DIFFERENTIATION OF SPINAL TUBERCULOSIS AND METASTATIC SPINAL TUMOR USING MRI FEATURE: A SYSTEMATIC REVIEW](#)

Universitas Airlangga, Surabaya, Indonesia [Majalah Biomorfologi Vol. 33 No. 1 \(2023\): MAJALAH BIOMORFOLOGI 44-51](#)

2023 [DOI: 10.20473/mbiom.v33i1.2023.44-51](#) [Accred : Sinta 4](#)

[BODY MASS INDEX WITH SYSTOLIC AND DIASTOLIC BLOOD PRESSURE AT A PUBLIC HEALTH CENTER IN SURABAYA, INDONESIA](#)

Universitas Airlangga, Surabaya, Indonesia [Majalah Biomorfologi Vol. 33 No. 1 \(2023\): MAJALAH BIOMORFOLOGI 14-19](#)

2023 [DOI: 10.20473/mbiom.v33i1.2023.14-19](#) [Accred : Sinta 4](#)

[SERUM AFP \(ALPHA FETO PROTEIN\) LEVELS PROFILE OF HEPATOCELLULAR CARCINOMA PATIENTS IN DR. SOETOMO GENERAL ACADEMIC HOSPITAL, SURABAYA, INDONESIA](#)

Universitas Airlangga, Surabaya, Indonesia [Majalah Biomorfologi Vol. 32 No. 1 \(2022\): MAJALAH BIOMORFOLOGI 6-12](#)

2022 [DOI: 10.20473/mbiom.v32i1.2022.6-12](#) [Accred : Sinta 4](#)

[BACTERIAL AND SENSITIVITY PATTERN OF PATHOGENS CAUSING VENTILATOR-ASSOCIATED PNEUMONIA IN INTENSIVE CARE UNIT](#)

Universitas Airlangga, Surabaya, Indonesia [Majalah Biomorfologi Vol. 32 No. 1 \(2022\): MAJALAH BIOMORFOLOGI 22-28](#)

2022 [DOI: 10.20473/mbiom.v32i1.2022.22-28](#) [Accred : Sinta 4](#)

[View more ...](#)

Get More with
SINTA Insight

[Go to Insight](#)

Citation Per Year By Google Scholar

Journal By Google Scholar

	All	Since 2018
Citation	40	36
h-index	3	3
i10-index	0	0

ORIGINAL RESEARCH

The HDL Level in Diabetic Foot Ulcer Patients with and without Amputation at Dr. Soetomo General Academic Hospital Surabaya

Ahmed Eliaan Shaker Abuajwa¹, Hermina Novida², Heroe Soebroto³,
Yan Efrata Sembiring³

¹Medical Program, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia.

²Department of Internal Medicine, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia; Dr. Soetomo General Academic Hospital, Surabaya, Indonesia.

³Department of Thoracic and Cardiovascular Surgery, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia; Dr. Soetomo General Academic Hospital Surabaya, Indonesia.

Article Info

Article history:

Received Sep 1, 2022
Revised Oct 18, 2022
Accepted Nov 9, 2022
Published Jan 10, 2023

Keywords:

Amputation
Diabetes
Lipid profile
HDL
Ulcer

*Corresponding author:

Yan Efrata Sembiring
yan-e-s@fk.unair.ac.id

ABSTRACT

Background: Diabetic foot ulcers (DFU) are microvascular and macrovascular consequences of diabetes and are associated with neuropathy, vascular disease, and immunosuppression. Increased blood glucose levels may result in the glycosylation of proteins in the nerve cells of the foot, leading to ischemia and cellular disturbances that can damage muscular, sensory, and autonomic neurons. **Objective:** To compare the lipid profiles of amputated and non-amputated DFU patients. **Material and Method:** This was a cross-sectional, analytic, retrospective research using total sample approach to analyze the medical records of all diabetes patients admitted to Dr. Soetomo General Academic Hospital between February 2018 and December 2020, with or without amputation. **Result:** The average age of DFU patients at Dr. Soetomo General Academic Hospital was 55.6 9.4 years, and the majority of patients were female (57%). Diabetic foot ulcers were most prevalent in the age group ≥ 50 years old (74%), amputated DFU patients had larger mean total cholesterol, TG, LDL, and HDL than non-amputated DFU patients, although the difference was not statistically significant. While the mean LDL/HDL ratio was greater in non-amputated DFU patients, the difference was statistically insignificant. The HDL level was low in 84% of amputated DFU patients, whereas in non-amputated DFU patients it was 81%. **Conclusion:** This research showed no significant variation in total cholesterol, TG, LDL, HDL, or the LDL/HDL ratio between amputated and non-amputated DFU patients. Most DFU patients with and without amputations had low HDL values.

How to cite:

Abuajwa, A.E.S., Novida, H., Soebroto, H., Sembiring, Y.E. 2023. The HDL Level in Diabetic Foot Ulcer Patients with and without Amputation at Dr. Soetomo General Academic Hospital Surabaya. *Majalah Biomorfologi*, 33(1): 30-37

Majalah Biomorfologi (Biomorphology Journal) p.ISSN:0215-8833, e.ISSN: 2716-0920

doi: [10.20473/mbiom.v33i1.2023.30-37](https://doi.org/10.20473/mbiom.v33i1.2023.30-37)



Copyright: © 2023 by the authors. Open access publication under the terms and condition of the Creative Commons Attribution 4.0 International license (CC.BY 4.0).

Highlights

1. Diabetes mellitus is a category of metabolic disease characterized by uncontrolled hyperglycemia.
2. This study revealed no statistically significant differences in total cholesterol, TG, LDL, HDL, or LDL/HDL ratio between amputated and non-amputated DFU patients.

BACKGROUND

Diabetes mellitus is a group of metabolic illnesses defined by the presence of untreated hyperglycemia. The diverse etiopathology includes abnormalities in insulin production, insulin action, or both, and problems in carbohydrate, lipid, and protein metabolism (World Health Organization, 2019). Diabetes mellitus (DM) is a global issue whose incidence is rising globally. Morbidity and mortality of diabetes induced by microvascular and metabolic factors. Moreover, macrovascular problems are more prevalent (Romadhon., et al. 2019). Diabetes is a metabolic condition characterized by high levels of blood glucose. Uncontrolled hyperglycemia might result in complications such as diabetic foot (Dewi., et al. 2020). Diabetic foot ulcers are the most widespread and serious diabetes complications. These ulcers are linked to an increased risk of mortality, morbidity, and foot amputation (Jarl, et al., 2019).

DFU is one of the most frequent consequences of diabetes. Amputation is 10 to 20 times more prevalent among diabetics compared to non-diabetics, and it is estimated that a lower leg is amputated partially or entirely every 30 seconds worldwide (International Diabetes Federation, 2021).

Diabetes mellitus pathophysiology is characterized by neuropathy, vasculopathy, and immunosuppression. Increased blood glucose levels may result in the glycosylation of proteins in the nerve cells of the foot, leading to ischemia and cellular disturbances that can damage muscular, sensory, and autonomic neurons. As the sensory and motor function of the foot deteriorates, the risk of infection, skin ulcers, and gangrene increases. Immune changes in diabetic foot ulcers (DFU) will impede the healing process (Aumiller & Dollahite, 2015). Infections may increase the likelihood of amputations, disability, extended hospitalization, and mortality in diabetic patients (Bekele & Chelkeba, 2020).

Due to the high expense of treatment and care, DFU patients suffer stigma, job loss, social isolation, unhappiness with their body image, and a loss of productivity (Atosona & Larbie, 2019). To avoid these health and social issues, we must identify the risk factors for DFU as well as preventative measures against its sequelae. Age, male gender, lipid abnormalities, hypertension, neuropathy, nephropathy, poor glycemic control, white blood cell count, and ulcer size are risk factors for amputation in diabetic patients (Namgoong, et al., 2016).

Cardiovascular disease (CVD) accompanied by dyslipidemia and atherogenic disorders is common in individuals with type 2 diabetes. Dyslipidemia is one of the most common risk factors for DFU (Shankarprasad, et al., 2015). High total cholesterol (T Chol), high triglycerides (TG), low high-density lipoprotein cholesterol (HDL-C), and elevated levels of tiny dense LDL particles are characteristic of diabetes individuals with lipid abnormalities, often known as "diabetic dyslipidemia." Additionally, low-density lipoprotein cholesterol (LDL-C) levels may rise or remain constant (Bhowmik, et al., 2018).

According to Pei et al. (2014), low HDL cholesterol levels are associated with diabetic foot, thus, efforts should be made to increase HDL cholesterol levels to avoid diabetic foot. According to Ikura et al. (2015), in patients with DFU, decreased HDL cholesterol levels may be an independent predictor of lower extremity amputation (LEA) and wound-related mortality. A high amount of TG is the most prevalent blood lipid anomaly in diabetes populations, according to another research. Increased TG levels are caused by hyperinsulinemia, not hyperglycemia, which is offset by insulin resistance (Hirano, 2018). While, according to Robbins (2017), dyslipidemia, mainly high LDL and HDL, is considered to have a role in atherogenesis, which may lead to ischemia and ultimately necrosis in DFU.

Numerous studies have related lipid profiles to the risk of diabetes mellitus. However, there is less evidence tying the lipid profile of a diabetic foot ulcer to the risk of amputation in the most current literature. Furthermore, not all of those studies are comparable in terms of lipid profiles in diabetics. Knowing the difference between the lipid profiles of amputated and non-amputated diabetic foot ulcer patients is of clinical and public health significance, and such information could form the basis for prevention programs and prognoses for diabetic foot ulcer patients and patients at high risk for amputation in Indonesia.

OBJECTIVE

This study was conducted to compare the difference in the lipid profiles of amputated and non-amputated diabetic foot ulcer patients at the Dr. Soetomo General Academic Hospital in Surabaya.

MATERIAL AND METHOD

This was a cross-sectional, observational, analytic research using the total sampling approach. This study included T2DM patients admitted to inpatient wards at Dr. Soetomo General Academic Hospital in Surabaya, Indonesia, for patients with diabetic foot ulcers with or without amputation hospitalized in the Internal Medicine and Thoracic and Cardiovascular Departments from January 2019 to December 2020. Consideration was given to maintaining the confidentiality of patient-identifiable information.

All DFU patients hospitalized between January 2019 and December 2020 who met the inclusion criteria had full medical records, comprehensive lipid profiles, and patient administration. The exclusion criteria were inadequate or ambiguous information. From the medical record, the age, gender, and lipid profile (total cholesterol, TG, LDL, HDL, and the LDL/HDL ratio) were extracted. The lipid profile utilized in this study was the most recent lipid profile found in the medical records of patients admitted to Dr. Soetomo General Hospital Surabaya. Data collected were compiled and presented as mean \pm standard deviation analyzed using Mann Whitney test with $p < 0.05$ considered statistically significant using SPSS.

RESULT

The criteria were met by 202 of the 465 DFU patients admitted between January 2019 and December 2020. Sixty-nine patients (34.2%) of them were amputated, and the other was non-amputated (Table 1). The patients generally came in with a mean age of 55.61 ± 9.48 years old. The youngest patient was 32 years old and the oldest patient was 83 years old, and there were slightly more female patients (57%) than male DFU patients. Patients aged 50 and up were the most common demographic (Table 1).

The mean values of total cholesterol, triglyceride, low-density lipoprotein, and high-density lipoprotein were higher in amputated DFU patients than in the non-amputated group, although they were not statistically significant. However, there was no difference in HDL levels between the two groups. Sixty-six percent of males and 76% of females with amputations had low HDL levels, compared to 85% of males and 78% of females with DFU who did not have amputations (Figure 3).




Figure 1. Mean values of lipid profile in non-amputated and amputated DFU patients

The total cholesterol levels of all non-amputated DFU patients were within normal limits while in amputated DFU patients 90% were in normal range (Figure 2). The mean LDL/HDL ratio in non-amputated DFU patients was 3.6, while in patients with amputation it was 3.52 (Figure 2).




Figure 2. Mean value of LDL/HDL ratio in non-amputated and amputated DFU patients

Table 1. Demographic variables among non-amputated and amputated DFU patients

Variables	Group	Non-amputated frequency	%	Amputated frequency	%
Gender	Male	60	45%	27	39.4 %
	Female	73	55%	42	60.2 %
Age	< 50 years old	36	27%	16	23 %
	≥ 50 years old	97	73%	53	77 %
Total		133		69	

Triglyceride levels differed between DFU patients with amputated limbs and those without. Twelve percent of non-amputated DFU patients had elevated TG levels, compared to 18% of amputated patients (Figure 3). Figure 3 shows that 58% of DFU patients with and without amputation have an ideal LDL/HDL ratio, whereas 42% of same patients have an elevated LDL/HDL ratio.

Table 2. The range of lipid profile in amputated and non-amputated DFU

Parameters	DFU status	
	133 Non-amputated	69 Amputated
Total cholesterol (mg/dl)		
Normal range <200	133 (100%)	63 (90%)
Borderline 200-239	0	3(5%)
High ≥ 240	0	3 (5%)
TG (mg/dl)		
Normal range <150	94 (70%)	45 (64%)
Borderline 150-199	25 (18%)	12 (18%)
High ≥ 200	14 (12%)	12 (18%)
LDL (mg/dl)		
Optimal <100	89 (67%)	49 (71%)
Borderline 100-160	44 (33%)	17 (24%)
High ≥ 160	0	3 (5%)
HDL (mg/dl)		
Low <40	108 (81%)	58 (84%)
Borderline 40-59	23 (17%)	5 (7%)
High ≥60	2 (2%)	6 (9%)
LDL/HDL ratio		
Optimal <3.5	78 (58.6%)	40 (58%)
High ≥3.5	55 (41.4%)	29 (42%)

Table 3. The characteristics of lipid profile in amputated and non-amputated DFU

Parameters	Group	N	Mean \pm SD	Range	P
Total cholesterol (mg/dl)	Non-amputated	133	131.52 \pm 36.23	43-193	0.922
	Amputated	69	138.5 \pm 52.4	69-391	
TG (mg/dl)	Non-amputated	133	125.2 \pm 64.32	14-543	0.461
	Amputated	69	134 \pm 71.1	30-333	
LDL (mg/dl)	Non-amputated	133	81.14 \pm 28	14-148	0.899
	Amputated	69	85.2 \pm 43.8	14-297	
HDL (mg/dl)	Non-amputated	133	28.6 \pm 15	3-106	0.621
	Amputated	69	31 \pm 19	6-100	
LDL/HDL ratio (mg/dl)	Non-amputated	133	3.60 \pm 2.46	0.54-21.33	0.837
	Amputated	69	3.52 \pm 2.27	0.62-11.29	

The mean LDL for DFU patients with amputation was 85.3 mg/dl, which was higher than the mean LDL for DFU patients without amputation (81.1 mg/dl) but statistically insignificant (Table 3). Most of the DFU patients with amputated legs had below-the-knee amputations, 36% above-the-knee amputations, and 5% had toe amputations.

DISCUSSION

According to the findings, there were more female than male DFU patients. This differed from prior research, such as an Egyptian study in which 67.3% of DFU patients at high risk for amputation were male, whereas 50% of DFU patients at low risk were male (Soliman, et al., 2022).

In our study, the mean age of amputated DFU patients was 56.89 \pm 9.1 years old, and the mean age of non-amputated DFU patients was 54.9 \pm 9.6 years old. Similar to a research conducted in the United Arab Emirates, diabetics with foot ulcers are more prevalent among those aged 50 to 59 (Manda, et al., 2012), which is similar to our study. In contrast, neither age nor gender were significant predictors of amputation in a Japanese research (Ikura, et al., 2015).

In our study in Indonesia, the total cholesterol, triglyceride, LDL, and HDL differences between the amputated and non-amputated DFU groups were not statistically significant. These findings were consistent with those from other studies. A study in Saudi Arabia detected no relation between the risk for amputation and LDL, triglycerides, or total cholesterol (Musa, et al., 2018). According to a Japanese study, neither triglyceride levels nor LDL cholesterol levels were indicators of the risk of amputation in the future (Ikura, et al., 2015). Moreover, there was no difference in the levels of triglycerides, LDL, or total cholesterol between diabetics with and without ulcers, but only the DFU patients had low HDL levels (Manda, et al., 2012). According to a research conducted in Turkey, TG, total cholesterol, LDL cholesterol, and ox-LDL did not vary significantly between amputated and non-amputated DFU patients (Muhtaroglu, et al., 2016). There was no significant correlation between diabetic foot and LDL-cholesterol, TC, or TG levels, according to a meta-analysis of case-control studies. This outcome did not match the American Diabetes Association's recommendations (American Diabetes Association, 2012). However, decreased HDL cholesterol was associated with increased risk of developing DFU (Pei, et al., 2014).

According to a study, determining insulin resistance can be done with high accuracy using the low-density lipoprotein/high-density lipoprotein cholesterol ratio (LDL/HDL ratio) (Zhang, et al., 2015). Another study found that the risk of prediabetes is positively connected with an elevated LDL/HDL ratio (Kuang, et al., 2022). All of these results suggested that the LDL:HDL ratio might be a reliable indicator of blood glucose metabolism. However, it is not yet apparent how the LDL/HDL ratio relates to diabetes. Other studies are required to determine the level of the LDL/HDL ratio in amputated and non-amputated DFU patients to determine whether this ratio can be used to predict ulcers in diabetes patients.

Although in our study there was no difference in HDL levels between amputated and non-amputated DFU patients, our study revealed that the HDL level was low in both groups. Low HDL levels were

seen in 96% of males and 76% of females who were amputated patients, while among the non-amputated patients, 85% of males and 78% of females had a low HDL level. Other studies have indicated a substantial association between low HDL cholesterol levels and the prevalence of minor and major extremities amputations and ulcer-related mortality (Ikura, et al., 2015). Nevertheless, a growing body of evidence shows that a lowered blood HDL level is a significant risk factor for the onset of DFU. In particular, other studies in Egypt report that a low HDL level is a predictor for limb amputation in DFU (Soliman, et al., 2022).

The dysfunction of HDL is ultimately caused by a substantial reduction in structural integrity, which has been commonly reported in both malignant and cardiometabolic illnesses (Vekic, et al., 2022). Consequently, HDL functioning and composition should be monitored and evaluated in a wider sense as indications of metabolic alterations associated with a range of clinical illnesses, particularly in DFU patients. HDL stimulates cholesterol outflow from foam cells, hence reducing cholesterol accumulation in atherosclerotic lesions (Zeljko, et al., 2021). Glycation and oxidation of apoA-I are increased by prolonged hyperglycemia, and the accumulation of acute phase proteins in HDL is stimulated by a pro-inflammatory state that is often present in diabetic patients (McEneny, et al., 2015). The absence of a significant difference in the lipid profile in amputated and non-amputated DFU patients may be due to earlier treatment for DFUs and good compliance. Most diabetic patients with foot ulcers have one or more of the following risk factors: neuropathy, vascular disease, foot deformity, or peripheral neuropathy. Microvascular disease and inadequate glycemic control are the main factors. Socio-demographic factors appear to be critical in the emergence of diabetic foot ulcers (Khan, et al., 2020). Treatment also plays a big role in reducing the risk of the lipid profile, especially after the patient's diagnosis with DFU. These drugs may include statins, bile acid sequestrants, or fibrates (International Diabetes Federation, 2020). Dyslipidemia is not the only risk factor for amputation. Many factors must be considered in foot amputation, including hypertension, diabetes, chronic renal disease, and smoking (Firnhaber & Powell, 2019), and the main cause of amputation in diabetic individuals is a delay in wound healing. Hyperglycemia, ongoing inflammation, micro- and macrovascular dysfunction, and neuropathy all contribute to the delayed healing of DFU (Baltzis, et al., 2014; Husakova, et al., 2022).

Foot ulcers and amputation are common complications of diabetes mellitus. Patients with improper glucose control with PAD presence and hypertension are a risk factor for amputation (Pemayun, et al., 2015). While we did not investigate the association between diabetes duration and foot ulcers in our study, a longer duration of diabetes was likely associated with more diabetic complications, such as micro- and macrovascular complications, which were likely to play crucial roles in the skin breach and persistence of the ulcer. The risk of neuropathy may be reduced by avoiding risk factors such as inadequate blood glucose management. Neuropathy sufferers must improve their glycemic control, practice proper foot care, and educate the general public in order to reduce their risk of foot ulceration and amputation. To provide complete treatment for people with diabetes, a multidisciplinary healthcare team is required.

Strength and limitations

The physicians will be able to make a prognosis about the health status of DFU patients according to their lipid profile, the population of DFU patients with amputation, and the clinical significance of HDL levels in DFU. Our study also had some limitations, such as the fact that the confounding factors and the full description of the foot ulcer were not given priority in this study. During the follow-up period, we did not notice any time-dependent changes in serum lipid profiles, and we did not evaluate apolipoproteins and proinflammatory cytokines such as interleukin-6 and tumor necrosis factor (TNF).

CONCLUSION

There was no difference in total cholesterol, triglycerides, LDL, HDL, or the LDL/HDL ratio between amputated and non-amputated DFU patients. However, this study found that the majority of DFU patients had low HDL cholesterol levels.

Acknowledgment

None

Conflict of Interest

All authors have no conflict of interest.

Ethic Consideration

The research protocol was approved by the Committee on Medical Research Ethics in the Faculty of Medicine, Universitas Airlangga (0753/LOE/301.4.2/I/2022).

Funding Disclosure

This research was self-funded.

Author Contribution

All authors have contributed to all processes in this research, including preparation, data gathering, analysis, drafting, and approval for publication of this manuscript.

REFERENCES

- American Diabetes Association. 2012. Medical advice for people with diabetes in emergency situations, ADA.
- Atosona, A., Larbie, C. 2019. Prevalence and determinants of diabetic foot ulcers and lower extremity amputations in three selected tertiary hospitals in Ghana. *Journal of Diabetes Research*, 2019, pp. 1–9. doi: 10.1155/2019/7132861.
- Aumiller, W. D., Dollahite, H. A. 2015. Pathogenesis and management of diabetic foot ulcers. *Journal of the American Academy of Physician Assistants*, 28(5), pp. 28–34. doi: 10.1097/01.JAA.0000464276.44117.b1.
- Baltzis, D., Eleftheriadou, I., Veves, A. 2014. Pathogenesis and treatment of impaired wound healing in diabetes mellitus: New insights. *Advances in Therapy*, 31(8): 817–836. doi: 10.1007/s12325-014-0140-x.
- Bekele, F., Chelkeba, L. 2020. Amputation rate of diabetic foot ulcer and associated factors in diabetes mellitus patients admitted to Nekemte referral hospital, western Ethiopia: prospective observational study. *Journal of Foot and Ankle Research*, 13(1): 65. doi: 10.1186/s13047-020-00433-9.
- Bhowmik, B., Siddiquee, T., Mujumder, A., Afsana, F., Ahmed, T., et al. 2018. Serum lipid profile and its association with diabetes and prediabetes in a rural Bangladeshi population. *International Journal of Environmental Research and Public Health*, 15(9): 1944. doi: 10.3390/ijerph15091944.
- Dewi, G. A. A. I. K., Wibisono, S., Pawana, I. P. A. 2020. Risk factors for lower extremity amputation in diabetic foot ulcer patients: A case-control study. *JUXTA: Jurnal Ilmiah Mahasiswa Kedokteran Universitas Airlangga*, 11(2): 83–85. doi: 10.20473/juxta.V11I22020.83-85.
- Firnhaber, J. M., Powell, C. S. 2019. Lower extremity peripheral artery disease: Diagnosis and treatment. *American family physician*, 99(6): 362–369.
- Hirano, T. 2018. Pathophysiology of diabetic dyslipidemia. *Journal of Atherosclerosis and Thrombosis*, 25(9): 771–782. doi: 10.5551/jat.RV17023.
- Husakova, J., Bem, R., Fejfarova, V., Jirkovska, A., Woskova, V., et al. 2022. Factors influencing the risk of major amputation in patients with diabetic foot ulcers treated by autologous cell therapy. *Journal of Diabetes Research*. Edited by A. C. Panayi, 2022: 1–8. doi: 10.1155/2022/3954740.
- Ikura, K., Hanai, K., Shinjyo, T., Uchigata, Y. 2015. HDL cholesterol as a predictor for the incidence of lower extremity amputation and wound-related death in patients with diabetic foot ulcers. *Atherosclerosis*, 239(2): 465–469. doi: 10.1016/j.atherosclerosis.2015.02.006.
- International Diabetes Federation. 2020. Type 2 diabetes prevention. International Diabetes Federation.
- International Diabetes Federation. 2021. IDF diabetes atlas. International Diabetes Federation.
- Jarl, G., Alnemo, J., Tranberg, R., Lundqvist, L. 2019. Gender differences in attitudes and attributes of people using therapeutic shoes for diabetic foot complications. *Journal of Foot and Ankle Research*, 12(1): 21. doi: 10.1186/s13047-019-0327-0.
- Khan, T., Plotkin, A., Magee, G., Shin, L., Woelfel, S., et al. 2020. Functional ambulatory status as a potential adjunctive decision-making tool following wound, level of ischemia, and severity of foot

- infection assessment. *Journal of Vascular Surgery*, 72(2): 738–746. doi: 10.1016/j.jvs.2019.11.064.
- Kuang, M., Peng, N., Qiu, J., Zhong, Y., Zou, Y., et al. 2022. Association of LDL:HDL ratio with prediabetes risk: a longitudinal observational study based on Chinese adults. *Lipids in Health and Disease*, 21(1): 44. doi: 10.1186/s12944-022-01655-5.
- Manda, V., Sreedharan, J., Muttappallymyalil, J., Das, R., Hisamatsu, E. 2012. Foot ulcers and risk factors among diabetic patients visiting Surgery Department in a University Teaching Hospital in Ajman, UAE. *International Journal of Medicine and Public Health*, 2(3): 34–38. doi: 10.5530/ijmedph.2.3.8.
- McEneny, J., Daniels, J., McGowan, A., Gunness, A., Moore, K., et al. 2015. A cross-sectional study demonstrating increased serum amyloid a related inflammation in high-density lipoproteins from subjects with type 1 diabetes mellitus and how this association was augmented by poor glycaemic control. *Journal of diabetes research*, 2015: 351601. doi: 10.1155/2015/351601.
- Muhtaroglu, S., Keti, D. B., Ünlühizarci, K. 2016. Investigation of ischemia-modified albumin levels and some atherosclerosis-related serum parameters in patients with diabetic foot. *Turkish Journal of Medical Sciences*, 46: 126–132. doi: 10.3906/sag-1406-38.
- Musa, I. R. Ahmed, M., Sabir, E., Alsheneber, I., Ibrahim, E., et al. 2018. Factors associated with amputation among patients with diabetic foot ulcers in a Saudi population. *BMC Research Notes*, 11(1): 260. doi: 10.1186/s13104-018-3372-z.
- Namgoong, S., Jung, S., Han, S., Jeong, S., Dhong, E., et al. 2016. Risk factors for major amputation in hospitalised diabetic foot patients. *International Wound Journal*, 13: 13–19. doi: 10.1111/iwj.12526.
- Pei, E., Li, J., Lu, C., Xu, J., Tang, T., et al. 2014. Effects of lipids and lipoproteins on diabetic foot in people with type 2 diabetes mellitus: A meta-analysis. *Journal of Diabetes and its Complications*, 28(4): 559–564. doi: 10.1016/j.jdiacomp.2014.04.002.
- Pemayun, T. G. D., Naibaho, R., Novitasari, D., Amin, N., Minuljo, T. 2015. Risk factors for lower extremity amputation in patients with diabetic foot ulcers: a hospital-based case–control study. *Diabetic Foot & Ankle*, 6(1): 29629. doi: 10.3402/dfa.v6.29629.
- Robbins, B. 2017. *The beneficiary*. Durham: Duke University Press.
- Romadhon, P.Z, Sutjahjo, A., Novida, H., Soelistijo, S.A., Wibisono, S., Prajitno, J.H., Tjokroprawiro, A. 2019. HBA1C and plasma transforming growth factor-beta 1 in type-2 diabetes mellitus patients. *New Armenian Medical Journal*, 13(1): 69-73.
- Shankarprasad, D. , Gundali, S., Kashinakunti, S., et al. 2015. Lipid profile in diabetes mellitus. *Indian Journal of Pathology and Oncology*, 2(4): 290. doi: 10.5958/2394-6792.2015.00030.7.
- Soliman, M. M., Gaber, M. A.-E., Bassaly, B. F. S. 2022. High-density lipoprotein level may be a risk factor for major cardiac events and lower extremity amputation in patients with a diabetic foot ulcer. *Journal of Current Medical Research and Practice*, 7(2): 116–120. doi: 10.4103/jcmrp.jcmrp_123_20.
- Vekic, J., Zeljkovic, A., Al Rasadi, K., Cesur, M., Silva-Nunes, J., et al. 2022. A new look at novel cardiovascular risk biomarkers: The role of atherogenic lipoproteins and innovative antidiabetic therapies. *Metabolites*, 12(2): 108. doi: 10.3390/metabo12020108.
- World Health Organization. 2019. *Classification of diabetes mellitus*. World Health Organization.
- Zeljkovic, A., Vekic, J., Mihajlovic, M., Gojkovic, T., Vladimirov, S., et al. 2021. Revealing the role of high-density lipoprotein in colorectal cancer. *International Journal of Molecular Sciences*, 22(7): 3352. doi: 10.3390/ijms22073352.
- Zhang, L., Chen, S., Deng, A., Liu, X., Liang, Y., et al. 2015. Association between lipid ratios and insulin resistance in a Chinese population. *PLOS ONE*. Edited by M. Thearle, 10(1): e0116110. doi: 10.1371/journal.pone.0116110.