

# Evaluation of biofloc technology the risk of giving

*by* Endang Dm

---

**Submission date:** 17-Apr-2023 11:24AM (UTC+0800)


**Submission ID:** 2066678522

**File name:** Evaluation\_of\_biofloc\_technology\_the\_risk\_of\_giving.pdf (257.7K)

**Word count:** 9375

**Character count:** 49688

**AACL Bioflux, Volume 12(2)**  
**April 30, 2019**



**Contents**

22

Ransangan J., Soon T. K., Duisan L., 2019 Population dynamics of marsh clam, *Polymesoda* spp. (Bivalvia: Corbiculidae) in Marudu Bay, Malaysia. AAFL Bioflux 12(2):395-403.

7

Kurdomanov A., Sirakov I., Stoyanova S., Velichkova K., Nedeva I., Staykov Y., 2019 The effect of diet supplemented with ProvioticR on growth, blood biochemical parameters and meat quality in rainbow trout (*Oncorhynchus mykiss*) cultivated in recirculation system. AAFL Bioflux 12(2):404-412.

5

Macale A. M. B., Nieves P. M., 2019 Stakeholders' perception on the status of blue swimming crabs *Portunus pelagicus* (Linnaeus, 1758) and performance of lying-in hatchery concept in San Miguel Bay, Philippines. AAFL Bioflux 12(2):413-416.

8

Supono, Wardiyanto, Harpeni E., Khotimah A. H., Ningtyas A., 2019 Identification of *Vibrio* sp. as cause of white feces diseases in white shrimp *Penaeus vannamei* and handling with herbal ingredients in East Lampung Regency, Indonesia. AAFL Bioflux 12(2):417-425.

Buban I. C. R., Soliman V. S., 2019 Review of scallop grow-out methods in tropical and temperate marine waters. AAFL Bioflux 12(2):426-436.

Ardiansyah M., Suharno, Susilowati I., 2019 Estimating the conservation value of mangrove forests in Marine Protected Areas: special reference to Karimunjawa waters, Indonesia. AAFL Bioflux 12(2):437-447.

12

Noor N. M., Nursyam H., Widodo M. S., Risjani Y., 2019 Biological aspects of green mussels *Perna viridis* cultivated on raft culture in Pasaran coastal waters, Indonesia. AAFL Bioflux 12(2):448-456.

9

Musel J., Anuar A., Mohamad S., Mustapa N., Hassan M. H., Rajali H., Anuar A., 2019 Morphometric relationship and the spawning season of *Acetes intermedius* from the coast of Miri Sarawak, Northwestern Borneo. AAFL Bioflux 12(2):457-471.

24

Adipu Y., Lumenta C., Mangindaan R. E. P., Manoppo H., 2019 Growth performance of *Litopenaeus vannamei* grown in biofloc system produced from different carbohydrate sources. AAFL Bioflux 12(2):472-479.

Hastuti S., Subandiyono, Windarto S., 2019 Blood performance of jaundice catfish *Clarias gariepinus*. AAFL Bioflux 12(2):480-489.

Fuad, Baskoro M. S., Riyanto M., Mawardi W., 2019 Catch characteristics on stationary lift net using light emitting diode (LED) and kerosene lights in Pasuruan waters. *AACL Bioflux* 12(2):490-501.

13 Paulangan Y. P., Fahrudin A., Sutrisno D., Bengen D. G., 2019 Distribution and condition of coral reef ecosystem in Tanah Merah Bay, Jayapura, Papua, Indonesia. *AACL Bioflux* 12(2):502-512.

10 Mohamad-Zulkifli N. F. N., Yong A. S. K., Kawamura G., Lim L.-S., Senoo S., Devic E., Mustafa S., Shapawi R., 2019 Apparent digestibility coefficient of black soldier fly (*Hermetia illucens*) larvae in formulated diets for hybrid grouper (*Epinephelus fuscoguttatus x Epinephelus lanceolatus*). *AACL Bioflux* 12(2):513-522.

Guntur G., Asadi M. A., Jullanda M. S. H., Luthfi O. M., Bintoro G., 2019 Ecology of bivalves in the intertidal area of Ngemboh, Gresik, East Java, Indonesia. *AACL Bioflux* 12(2):523-534.

Serdiati N., Arfiati D., Widodo M. S., Lelono T. D., Toha A. H. A., 2019 Genetic characteristics of ricefish from Lake Poso, Central Sulawesi, Indonesia. *AACL Bioflux* 12(2):535-552.

87 Goda A. M. A. S., Srour T. M., Mansour A. T., Baromh M. Z., Sallam G. R., Baromh A. Z., 2019 Assessment of stressful ambient water salinity on growth, feed utilization and hematological indices of European sea bass, *Dicentrarchus labrax*, juveniles. *AACL Bioflux* 12(2):553-563.

Cahyadinata I., Fahrudin A., Sulistiono, Kurnia R., 2019 Household welfare of mud crab fishermen in small outermost islands. Case study: Enggano Island, Bengkulu Province, Indonesia. *AACL Bioflux* 12(2):564-574.

14 Chilmawati D., Hutabarat J., Anggoro S., Suminto S., 2019 Biomolecular identification and optimization of growth performance and egg production in *Oithona sp.* under different salinity culture conditions. *AACL Bioflux* 12(2):575-585.

15 Rumampuk N. D. C., Schaduw J. N. V., Lintang R. A. J., Rompas R. M., 2019 Imposex phenomenon in gastropods from Bitung waters, North Sulawesi, Indonesia. *AACL Bioflux* 12(2):586-592.

Dewi D. A. N. N., Iskandar D. D., 2019 Cooperative or individually? Impact of fishing expedition decision on income. *AACL Bioflux* 12(2):593-600.

Rasyid A., Handayani T., 2019 Evaluation of the biochemical composition of tropical red seaweeds *Galaxaura rugosa* and *Gelidiella acerosa* from Ujung Genteng waters, Indonesia. *AACL Bioflux* 12(2):601-609.

Krisnafi Y., Yusrizal, Halim S., Santoso H., Suharto, Waluyo A. S., Kusdinar A., Danapraja S., Pickassa F. I., Alamsah S., Fadly Z. R., 2019 CPUE analysis of crab resources in Karangantu, Serang Banten, Indonesia. *AACL Bioflux* 12(2):610-617.

Ben Yahkoub Y., Fekhaoui M., El Qoraychy I., Yahyaoui A., 2019 Current state of knowledge on Louisiana crawfish (*Procambarus clarkii* Girard, 1852) in Morocco. *AACL Bioflux* 12(2):618-628.

Natan Y., Pattikawa J. A., Tomia B., 2019 Biological aspects of jumping halfbeak (*Hemiramphus archipelagicus*) in the waters of Kelang Island, Western Seram, Indonesia. *AACL Bioflux* 12(2):629-635.

Khan A., Rizal A., Dewanti L. P., Apriliani I. M., Junianto, Supriyadi D., Ghiffary W., Nasution A. M., Gray T. S., Mill A. C., Polunin N. V. C., 2019 Skipjack (*Katsuwonus pelamis*) tuna pole-and-line marketing supply chains in Indonesia: case study in Pulau Bacan. *AACL Bioflux* 12(2):636-641.

Achmad D. S., Ali S. A., Sudirman, Indar Y. N., 2019 The gonad maturity development and spawning season of orange-spotted grouper (*Epinephelus coioides*) at Kwandang Bay, Gorontalo Province, Indonesia. *AACL Bioflux* 12(2):642-649.

Asriyana, Irawati N., 2019 Assessment of the trophic status in Kendari Bay, Indonesia: a case study. *AACL Bioflux* 12(2):650-663.

Ompi M., Lumoindong F., Tamanampow Y., Undap N., Papu A., Wagele H., 2019 Monitoring marine *Heterobranchia* in Lembeh Strait, North Sulawesi (Indonesia), in a changing environment. *AACL Bioflux* 12(2):664-677.

Kreckhoff R. L., Ngangi E. L. A., Undap S. L., Kusen D. J., 2019 Crude extracts of *Kappaphycus alvarezii* algae cultivated in several seaweed production centers in North Sulawesi, Indonesia a immunostimulant. *AACL Bioflux* 12(2):678-686.

Subair N., Haris R., 2019 Factors that motivate Mappakasunggu women of seaweed farmers to develop a family economic survival strategy. *AACL Bioflux* 12(2):687-695.

Yulius, Kepel T. L., Hadiwijaya S. L., Heriati A., Ramdhan M., 2019 Preliminary study of lead and mercury concentrations in seven commercial seafood at Lombok Island, Indonesia. *AACL Bioflux* 12(2):696-705.

Nugroho L. A., Masithah E. D., Satyanrini W. H., 2019 Evaluation of biofloc technology: the risk of giving different commercial probiotics to C:N and N:P ratio and quality of seawater. *AACL Bioflux* 12(2):706-715.

Pamula O. Y. T., Prayogo, Sudarno., 2019 Optimization on survival and growth rate of African catfish (*Clarias* sp.) using water spinach (*Ipomoea aquatica*)-based aquaponics system. *AACL Bioflux* 12(2):716-723.

Hutajulu J., Kusumo T., Saputra A., Mualim R., Handri M., Sugriwa E., Nainggolan C., Syamsuddin S., 2019 Financial analysis in the exploitation of blue swimming crab *Portunus pelagicus* in Banten Bay, West Java, Indonesia. *AACL Bioflux* 12(2):724-734.

36  
MSc in Sustainable Development and Environmental Management. A 2-year full time programme taught in English. p.735.

**International Standard Serial Number  
Online ISSN 1844–9166; Print ISSN 1844-8143**

**Published by Bioflux – six issues/year (bimonthly); in cooperation with The Natural Sciences Museum Complex (Constanta, Romania)**

The journal includes original papers, short communications, and reviews on Aquaculture (Biology, Technology, Economics, Marketing), Fish Genetics and Improvement, Aquarium Sciences, Fisheries, Ichthyology, Aquatic Ecology, Conservation of Aquatic Resources and Legislation (in connection with aquatic issues) from wide world.

**Editor-in-Chief**

Petrescu-Mag I. Valentin: USAMV Cluj, Cluj-Napoca, University of Oradea (Romania)  
Gavriloae Ionel-Claudiu: SC Bioflux SRL, Cluj-Napoca (Romania).

**Editors**

Adascalitei Oana: Maritime University of Constanta, Constanta (Romania)  
Amira Aicha Beya: Badji Mokhtar Annaba University, Annaba (Algeria)  
Arockiaraj A. Jesu: SRM University, Chennai (India)  
Appelbaum Samuel: Ben-Gurion University of the Negev (Israel)  
Baharuddin Nursalwa: Universiti Malaysia Terengganu, Terengganu (Malaysia)  
Boaru Anca: USAMV Cluj, Cluj-Napoca (Romania)  
Botha Miklos: Bioflux SRL, Cluj-Napoca (Romania)  
Breden Felix: Simon Fraser University (Canada)  
Burny Philippe: Universite de Liege, Gembloux (Belgium)  
Caipang Cristopher M.A.: Temasek Polytechnic (Singapore)  
Chapman Frank: University of Florida, Gainesville (USA)  
Cocan Daniel: USAMV Cluj, Cluj-Napoca (Romania)  
Coroian Cristian: USAMV Cluj, Cluj-Napoca (Romania)  
Creanga Steofil: USAMV Iasi, Iasi (Romania)  
Cristea Victor: Dunarea de Jos University of Galati, Galati (Romania)  
Das Simon Kumar: Universiti Kebangsaan Malaysia, Bangi, Selangor (Malaysia)  
Dimaggio Matthew A.: University of Florida (USA)  
Firica Cristian Manuel: Spiru Haret University Bucharest, Craiova (Romania)  
Georgescu Bogdan: USAMV Cluj, Cluj-Napoca (Romania)  
Karayucel Ismihan: University of Sinop, Sinop (Turkey)  
Khamesipour Faham: Shiraz University, Shiraz (Iran)  
Kosco Jan: Presov University, Presov (Slovakia)  
Mehrad Bahar: Gorgan University of Agricultural Sciences and Nat. Res. (Iran)  
Miclaus Viorel: USAMV Cluj, Cluj-Napoca (Romania)  
Mihociu Tamara: R&D National Institute for Food Bioresources (Romania)  
Molnar Kalman: Hungarian Academy of Sciences, Budapest (Hungary)  
Muchlisin Zainal Abidin: Universiti Sains (Malaysia), Syiah Kuala University (Indonesia)  
Nowak Michal: University of Agriculture in Krakow (Poland)  
Nyanti Lee: Universiti Malaysia Sarawak, Sarawak (Malaysia)  
Olivotto Ike: Universita Politecnica delle Marche, Ancona (Italy)  
Oroian Firuta Camelia: USAMV Cluj, Cluj-Napoca (Romania)  
Papadopol Nicolae: Natural Sciences Museum Complex, Constanta (Romania)  
Papuc Tudor: USAMV Cluj, Cluj-Napoca (Romania)  
Parvulescu Lucian: West University of Timisoara (Romania)  
Pasarin Benone: USAMV Iasi, Iasi (Romania)  
Pattikawa Jesaja Ajub: Pattimura University, Ambon (Indonesia)  
Petrescu Dacinia Crina: Babes-Bolyai University, Cluj-Napoca (Romania)  
Petrescu-Mag Ruxandra Malina: Babes-Bolyai University, Cluj-Napoca (Romania)  
Petrovici Milca: West University of Timisoara (Romania)  
Pratasik Silvester Benny: Sam Ratulangi University, Manado (Indonesia)  
Proorocu Marian: USAMV Cluj, Cluj-Napoca (Romania)  
Ray Sunuram: Khulna University (Bangladesh)  
Rhyne Andrew: Roger Williams University; New England Aquarium, Boston (USA)  
Safirescu Calin: USAMV Cluj, Cluj-Napoca (Romania)  
Serrano Jr. Augusto E.: University of the Philippines Visayas (Philippines)  
Sima Nicusor Flaviu: USAMV Cluj, Cluj-Napoca (Romania)  
Tlusty Michael E.: New England Aquarium, Boston (USA)  
Vesa Stefan Cristian: Iuliu Hatieganu UMF, Cluj-Napoca (Romania)  
Vintila Iuliana: Dunarea de Jos University of Galati, Galati (Romania)  
Wariaghli Fatima: University Mohammed V in Rabat, Rabat (Morocco)  
Yusli Wardiatno: Bogor Agricultural University, Bogor (Indonesia).

### Contact Publisher

SC Bioflux SRL, 54 Ceahlău Street, Cluj-Napoca, 400488, Romania, European Union.

Ioan Valentin Petrescu-Mag, e-mail: zoobiomag2004@yahoo.com

Note that there is also an electronic version of the journal. You may download the fulltext version of AACL Bioflux – Volume 12/2019 from <http://www.bioflux.com.ro/aacfl/> -open access.

All articles included in AACL Bioflux are peer reviewed (double blind peer-review is used). Each published article was seen before by two reviewers; the two peer-reviews are made independently.

**AACL Bioflux has a publishing agreement, or is indexed, abstracted or full text reproduced by/in the following scholar/scientific databases, search engines, libraries, publishers:**

ISI Web of Science – via CABI and Zoological Record

Zoological Record (direct submission/coverage)

Scopus - Elsevier; Sciverse

Scimago - Journal Rank

CAB International - CAB Abstracts

ProQuest

China Educational Publications Import & Export Corporation - SOCOLAR

Ulrich's Periodicals Directory

EBSCO – EBSCOhost Online Research Databases

CAB Direct (as part of CABI)

Wolters Kluwer - Ovid LinkSolver

CNCSIS Romania (Rank: B+)

Polish Ministry of Science and Higher Education – 2.00 pts/article

The National Science Digital Library - NSDL

The University of Hong Kong Libraries – HKUL Database

Deutsche Nationalbibliothek - ZDB Database

State Library of Ohio - OhioLINK Database

Smithsonian Institution Libraries

Biblioteka Główna Uniwersytetu Medycznego w Poznaniu

Google Scholar, Academic

Google (search engine)

Athabasca University - ICAAP Database

University of Southampton - ROAR

Georgetown University Library - NewJour

Università degli Studi di Modena e Reggio Emilia – SBA, Risorse Elettroniche

Simon Fraser University - PKP

University of Tsukuba - Tulips Database

Teikyo Scientific University- NTU Database

LIS Links: Link Library of Open Access English Language Journals

Fayetteville State University - Charles W. Chesnutt Library

Vrije Universiteit Brussel - VUB e-journal list

Russian CJDB

Universität Giessen - Digitale Bibliothek

Pace University Library

Tel Aviv University. Gitter-Smolarz Library of Life Sciences and Medicine

MALMAD – Israel Center for Digital Information Services

California State University – Monterey Bay Library

Réseau des Bibliothèques de l'Université Joseph Fourier et de l'Institut National Polytechnique de Grenoble

University of Saskatchewan – University Library (Electronic Journals)

YanXue Resource – YXRES

Feng Chia University – Electronic Library

Main Library of the University of Agriculture in Krakow

SUMMON Database

uOttawa Library

Università di Roma Tor Vergata - Digital Library

UTC Lupton Library

University of Notre Dame - Hesburgh Libraries

Guilford College - Hege Library

Eastern Michigan University Library - Periodicals locator

Universiteitsbibliotheek Gent - Tijdschriften

Chung Yuan Christian University Library

ArgosBiotech

Ecole Polytechnique Federale de Lausanne - Library

Rowan University – Library Services

James Cook University Library – Australia

Ministerstwo Nauki i Szkolnictwa Wyższego Warszawa

Kun Shan University Library

CCG-IBT Biblioteca

Reference Zone – Nigeria

Dayang Journal System

Riley-Hickingbotham Library



THOMSON REUTERS

Index to Organism Names (ION)



Politechnika Wroclawska  
Universite Paris Diderot – Revues Electroniques  
ERSA – Chinese Library.

## Our collaborators



33  
SciVerse Scopus is the world's largest abstract and citation database of peer-reviewed literature and quality web sources. Contains 41 million records, 70% with abstracts; Nearly 18,000 titles from 5,000 publishers worldwide; 70% of content is pulled from international sources; Includes over 3 million conference papers; Provides 100% Medline coverage; Offers sophisticated tools to track, analyze and visualize research.

## Socolar

The free scholastic search engine is established by China Educational Publications Import and Export Corporation (CEPIEC). Under the leadership of China Ministry of Education, CEPIEC is the only company dealing with publication ex-im business in the education field. CEPIEC has been serving colleges and universities, research institutions and public libraries in China for more than 20 years, providing them with excellent academic resources from all over the world. Contact: No 41 Zhong Guan Cun Street, Haidian District, Beijing, P.R.China 100080. Tel: +86 (0)10 6251 4068; e-mail: li\_yanxia@cepiec.com.cn Fax: +86 (0)10 6251 4068 [www.cepiec.com.cn](http://www.cepiec.com.cn)



CABI is a not-for-profit international organization that improves people's lives by providing information and applying scientific expertise to solve problems in agriculture and the environment. Their mission and direction is influenced by their member countries who help guide the activities they undertake. These include scientific publishing, development projects and research, and microbial services. 2010 marked 100 years of CABI. Since its beginnings as an entomological committee in 1910, it then developed into a Commonwealth organization before becoming a truly international service in agricultural information, pest identification and biological control in the mid 1980s.

EBSCOhost® databases are the most-used, premium online information resources for tens of thousands of institutions worldwide, representing millions of end-users.  
Address: 10 Estes Street  
Ipswich, MA 01938 USA  
Phone: (978) 356-6500  
Toll Free: (800) 653-2726 (USA & Canada)  
Fax: (978) 356-6565  
E-mail: [information@ebSCOhost.com](mailto:information@ebSCOhost.com)



## Zoological Record (part of Biosis)

ION contains the organism names related data gathered from the scientific literature for Thomson Reuters' *Zoological Record*® database. Viruses, bacteria and plant names will be added from other Thomson Reuters databases such as *BIOSIS Previews*® and *Biological Abstracts*®.




## Index to Organism Names (ION)



## ELSEVIER PRODUCTS

Elsevier B.V., Bibliographic Databases, Radarweg 29, 1043 NX Amsterdam, The Netherlands.





## Evaluation of biofloc technology: the risk of giving different commercial probiotics to C:N and N:P ratio and quality of seawater

Lutfi A. Nugroho, Endang D. Masithah, Woro H. Satyanrini

Department of Fisheries and Marine, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya, East Java, Indonesia. Corresponding author: E. D. Masithah, endangdewimashitah@yahoo.com

**Abstract.** Biofloc technology is the solution to the problem of decreasing water quality related to aquaculture activities. The composition of the microflora in this technology is influenced by carbon, nitrogen and phosphorus through C:N and N:P ratios. This study examined the risks of different commercial probiotics for the ratio of N:P and C:N to biofloc media and the quality of seawater produced. This experiment was carried out with a completely randomized design (CRD) for 8 weeks with 3 treatments, namely probiotics Petrofish (A), Probiotics Pro I (B), and without probiotics (C). After 6 replications, the data were analyzed descriptively and statistically. After giving treatment, ratio of C:N ratio was known in treatment A (2.46), B (1.83) and C (2.21) and N:P ratio in treatment A (22.01), B (22.08) and C (25.96). The treatment of different commercial probiotics in treatment A and treatment B of the ratio C:N and N:P was not significant ( $p > 0.05$ ). Seawater quality is based on pH parameters (7.94-8.11), dissolved oxygen ( $2.05-4.24 \text{ mg L}^{-1}$ ) water temperature ( $26.5-27.1^\circ\text{C}$ ), salinity (22.33-25.17 ppt) and ammonia level ( $1-1.6 \text{ mg L}^{-1}$ ). From the result, we emphasize that giving different commercial probiotics does not have an effect on the values of the C:N and N:P ratio and the seawater quality based on parameter is suitable to the standard.

**Key Words:** water management, seawater research, effectiveness of biofloc, environmental condition, aquaculture activities.

**Introduction.** Indonesia, one of the developing countries and an archipelago surrounded by the sea, enables the management of large natural and non-marine resources (FAO 2018). Development of aquaculture in Indonesia has enormous opportunities (Asmanah et al 2012). Developing countries become a source of 80 billion dollars in fishery exports, providing net trade income that is higher than the combined meat, tobacco, rice and sugar in 2016 (FAO 2016).

Global fish consumption per capita has increased to above 20 kilograms per year or doubled compared to the 1960s as a result of fisheries development (FAO 2018). Half of the seafood consumed globally comes from aquaculture. Aquaculture plays an important role in global food systems, the environment, and human health along with declining capture fisheries (Fry et al 2016). Aquaculture is one of the fastest growing food-producing sectors, supplying around 40% of fish food world (Cole et al 2012).

Apart from being beneficial to the community, this industry also has environmental degradation. Aquaculture practice arrangements, including site selection, pollution control, water quality, feed supply, and food security need to be done (Cole et al 2012; Martinez-Porchas & Martinez-Cordova 2012). Responsible fisheries governance, as well as national and international policies and laws will ensure sustainable harvests, maintain biodiversity and ecosystem functions, and adapt to climate change (Ekasari 2009; Garcia & Rosenberg 2015). So that, the result of deep sea water resources function as usual and can help overcome health problems (Nani et al 2016).

Water treatment in a bad cultivation system is one of the main obstacles that can lead to a decrease in water quality (Boyd & Green 2002). This reduction in water quality occurs due to waste from the rest of the cultivation activities. The remaining feed that is not eaten and feces will settle on the bottom of the water and is broken down by bacteria



into nitrogen and ammonia compounds. The amount of uncontrolled ammonia compounds in the waters can endanger cultured organisms (Avnimelech 2007; Ekasari & 2009).

The biofloc technology (BFT) application offers benefits in increasing aquaculture production that can contribute to the achievement of sustainable development. This technology can produce higher productivity with a smaller impact on the environment that can be developed and carried out in integration with other food production. This system is productively integrated to produce more food and feed from the same area of land with fewer inputs (Bossier & Ekasari 2017). In addition, BFT is one of the most beneficial alternative solutions to waste cultivation problems because it can reduce inorganic nitrogen waste (Crab et al 2007).

Probiotic parts cannot be separated from biofloc technology. The main principle applied in biofloc technology is management of water quality based on the ability of heterotrophic bacteria to utilize organic and inorganic N contained in the waters. (Purwanta & Firdayati 2002; Ekasari 2009; Gunarto & Suwoyo 2011). The principle of BFT is to grow microorganisms, especially heterotrophic bacteria in water conserving media. This growth is intended to absorb pollutant components, such as ammonia in the waters therefore this efforts can improve quality and benefit in environment. Heterotrophic bacteria will assimilate ammonia and nitrogen if the ratio of C:N to the media is well balanced (Schneider et al 2005). The application of BFT allows at least zero water to exchange practices during the cultivation period, and thus can increase sustainability, biosafety and production. Manipulating C/N ratio inputs has a significant influence on the development and characteristics of biofloc in zero-exchange systems by influencing inorganic nitrogen and phosphorus dynamics, and ultimately affecting performance (Xu et al 2016).

In addition to the C:N ratio, high nutrient concentrations in the waters will affect water productivity. The composition between nutrient components, namely the N to P ratio is often referred to as redfield ratio (Makmur et al 2012). Nitrogen and phosphorus are the most considered because they are related to CO<sub>2</sub>. Nitrogen and phosphorus, which are macro nutrients, both have benefits as limiting nutrients for phytoplankton growth (Souther & Rissik 2009).

A research to improve the effectiveness of biofloc technology and the quality of water produced is important, considering that it can be used as a support for future aquaculture technology. This study aims to study the risk of different commercial probiotics on the ratio of C:N and N:P to biofloc technology and the quality of seawater produced.

**Material and Method.** This experimental study was carried out for 8 weeks during June to July in 2015 and located in the Laboratory of the Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya, while the examination of N, P and C elements was carried out in the Environmental Quality Laboratory of the Faculty of Civil Engineering and Planning, Institut Teknologi 10 November, Surabaya.

Complete Randomized Design (CRD) was carried out with three treatments and six replications. The experimental design consisted of three treatments for each probiotic Petrofish and Pro-1 and without probiotics, namely treatments A, B, and C respectively, which were repeated six times so that there were 18 experimental units, namely: A1, A2, A3, A4, A5, A6, B1, B2, B3, B4, B5, B6, C1, C2, C3, C4, C5, C6. The research was carried out by the formation of biofloc with probiotic Petrofish, the formation of biofloc with probiotics Pro-1 and the formation of biofloc without probiotics as controls.

The stages of the research are largely carried out through the sterilization of tools and materials with liquid soap and chlorine, the making and breeding of small-scale biofloc in-door and the addition of probiotics. The biofloc characteristics used were aquariums measuring 40 x 20 x 35 cm<sup>3</sup> filled with 4.5 liters of sea water with 22 ppt salinity and 3 point aeration. In the aquarium, every 7 days there are several ingredients included, namely 230 grams of fine bran, 90 grams of fish flour, 4.5 grams of yeast, 10 grams of urea fertilizer, 5 grams of SP-36 fertilizer, 12 mL molasses, 4.5 grams of dolomite and probiotic lime 2 mg L<sup>-1</sup>.

The main parameter in this study is the C:N and N:P ratios. The Kjeldahl method was applied in nitrogen testing (Saez-Plaza et al 2013). The nitrogen content used in the ratio calculation is anorganic nitrogen in the form of nitrite, nitrate and ammonia. The method used for testing phosphate content is the spectrophotometric method by calculating the inorganic phosphate content of orthophosphate. The method applied in carbon testing uses the gravimetry analysis (Miyazawa et al 2000). All parameters were observed on a daily basis for 21 days. The degree of acidity (pH), dissolved oxygen (DO) and temperature were measured twice a day, at 6:00 and 17:00. Ammonia measurements were measured once a week together with salinity measurements. The pH was measured twice a day at 6:00 and 17:00. After all data have been collected, they were analyzed statistically and descriptively. Statistical analysis was done using ANOVA with SPSS 16 for Windows. The results can be concluded as significant if p-value < 0.05. This shows a significant difference in the use of probiotics for C:N ratio and N:P ratio on biofloc technology.

**Results.** The results of this study consisted of test data for the content of C, N, P elements as well as calculating the ratios of C:N and N:P to the seawater experiment. Water quality parameters were observed in the form of acidity (pH), DO, temperature, ammonia, and salinity during the study. From the experimental results, the average value of N content at the beginning of the study showed that the highest value in treatment B (4,362.69 mg L<sup>-1</sup>) and the lowest value in treatment A (2,892.79 mg L<sup>-1</sup>). However, the average value of N content at the end of the study showed that the highest value in treatment C (3,698.00 mg L<sup>-1</sup>) and the lowest value in treatment A (3,207.92 mg L<sup>-1</sup>). Calculation of the floc volume formed in treatment A (44.33 mL L<sup>-1</sup>) and B (33.33 mL L<sup>-1</sup>) showed higher than in treatment C (16.66 mL L<sup>-1</sup>).

Testing the value of P content at the beginning of the study showed that treatment A had the highest value of 291.43 mg L<sup>-1</sup> and the lowest treatment B (137.36 mg L<sup>-1</sup>). The average value of P content at the end of the study showed that treatment B had the highest value (155.48 mg L<sup>-1</sup>) and the lowest treatment C was 146.59 mg L<sup>-1</sup>.

C content testing at the beginning of the study showed the highest value was found in treatment B (6,900 mg L<sup>-1</sup>) and the lowest was in treatment C (5,948 mg L<sup>-1</sup>). At the end of the study the average value of C content showed that treatment C had the highest value (8,206.66 mg L<sup>-1</sup>) and the lowest in treatment B (5,442.00 mg L<sup>-1</sup>).

The C:N ratio was obtained from data of C and N content in the media, then compared to get the value of the ratio C:N. The results of the measurement of C:N ratio at the beginning of the study showed that the value of the treatment ratio A gave the highest result (2.3) and the lowest at treatment B (1.58). The final value of the C: N ratio in all treatments has increased. The highest value of the C:N ratio occurred in treatment A (2.46±1.73858) and the lowest in treatment B (1.83±0.78).

The N:P ratio value is obtained from the data of N and P content in the media, then compared to get the value of the ratio N:P. The results of the measurement of the N:P ratio at the beginning of the study showed the value of the ratio in treatment B gave the highest results of 31.76 and the lowest in treatment A (9.92). Value of N:P ratio on the last day of the study on treatments A and C showed an increase to 22.01±12.24 and 25.96±3.81 respectively, while in treatment B it decreased to 22.08±8.14.

Giving different probiotics namely petrofisch probiotics, pro 1 probiotics and control without probiotics to the ratio C:N (p: 0.662) and N:P (p: 0.678) showed a non-significant result (p > 0.05). Table 1 shows the average data for N, P and C content and C:N and N:P ratios.

The content of N, P and C and the value of the ratio N:P and C:N

Table 1

T	N (ppm)		P (ppm)		C (ppm)		N:P ratio		C:N ratio	
	Start	End	Start	End	Start	End	Start	End	Start	End
A	2,892.79	3,207.92	291.43	153.8	6.670	5,881.66	9.92	22.01 <sup>a</sup>	2.30	2.46 <sup>a</sup>
B	4,362.69	3,231.62	137.36	155.48	6.900	5,442	31.76	22.08 <sup>a</sup>	1.58	1.83 <sup>a</sup>
C	3,478.50	3,698.28	138.68	146.59	5.948	8,206.66	28.08	25.96 <sup>a</sup>	1.7	2.21 <sup>a</sup>

Note: the values with the same superscript in the same column show results that are not significantly different.

**Quality of seawater.** In the beginning of the study, the pH value was classified as acid, which ranged from 5.72 to 5.86 but at week 2 the pH value in all treatments experienced increases ranging from 7.40 to 7.47. Until the end of the study at week 3 the pH value in all treatments was relatively stable ranging from 7.94 to 8.11.

In the first week, the highest DO content was found in treatment C (2.85 mg L<sup>-1</sup>) and the lowest was treatment B (2.56 mg L<sup>-1</sup>). In the second week the DO content increased in all treatments, namely treatment A (3.36 mg L<sup>-1</sup>), treatment B (2.89 mg L<sup>-1</sup>) and treatment C (2.91 mg L<sup>-1</sup>). At the end of the study, which was the third week, all treatments still experienced an increase in DO with the highest in treatment A (3.92 mg L<sup>-1</sup>) and the lowest was treatment C (3.68 mg L<sup>-1</sup>). During this study DO was in the range of 2.05 to 4.24 mg L<sup>-1</sup> so that it can be said to be normal based on existing references (Becerril-Cortés et al 2018).

The water temperature in the study is known in the range of 26.5-27.1°C. A higher range is obtained from the results of the salinity assessment with an average during the study ranging from 22.33 to 25.17 ppt. Ammonia measurements tend to fluctuate even though at the beginning until the study ended the value of ammonia content ranged from 1 to 1.6 mg L<sup>-1</sup>.

Dissolved oxygen and pH at week 3 on all treatments were within acceptable limits even though at the beginning both parameters as in lower value. Oxygen, ammonia levels and salinity are relatively stable according to the limits determined from the first week to the end of the study. The quality of seawater compared with acceptable standard presented in Table 2.

Table 2  
Quality of seawater based on degree of acidity, dissolved oxygen, temperature, ammonia level and salinity

Treatment	Mean			Acceptable range
	Week 1	Week 2	Week 3	
<i>Degree of acidity (pH)</i>				
Treatment A	5.86±0.40	7.46±2.29	8.11±0.04	7-9.5 (Bhatnagar & Devi 2013);
Treatment B	5.72±0.20	7.40±0.33	7.94±0.15	6.8-8 (Becerril-Cortés et al 2018)
Treatment C	5.79±0.38	7.47±0.42	7.97±0.17	
<i>Dissolved oxygen (DO) (mg L<sup>-1</sup>)</i>				
Treatment A	2.70±0.45	3.36±0.88	3.92±0.32	3-5 (Bhatnagar & Devi 2013);
Treatment B	2.56±0.51	2.89±0.88	3.71±0.53	Above 4.0 (ideal) and at least 60%
Treatment C	2.85±0.40	2.91±0.77	3.68±0.29	saturation (Becerril-Cortés et al 2018)
<i>Temperature (°C)</i>				
Treatment A	27±0.43	26.8±0.34	26.8±0.26	15-35 (Bhatnagar & Devi 2013);
Treatment B	26.7±0.16	27.1±0.16	26.7±0.32	28-30 (Becerril-Cortés et al 2018)
Treatment C	26.8±0.18	26.5±0.37	26.8±0.31	
<i>Ammonia level (mg L<sup>-1</sup>)</i>				
Treatment A	1±0.27	1.5±0.54	1.25±0.54	1-1.6 (FAO 2002)
Treatment B	1±0.25	1.6±0.51	1±0.25	
Treatment C	1±0.27	1.6±0.51	1.3±0.54	
<i>Salinity (ppt)</i>				
Treatment A	22.50±0.54	24.00±0.89	24.67±0.81	35 (Alkhudhiri & Hilal 2018);
Treatment B	22.67±0.81	23.67±0.81	25.17±0.75	0-50 (Becerril-Cortés et al 2018)
Treatment C	22.33±0.51	23.50±0.54	24.67±0.51	

**Discussion.** Giving different probiotics does not have a significant effect on the C:N ratio and the N:P ratio in biofloc media. The effect of adding probiotics in biofloc is unclear, because the results depend on the concentration and type used, but most importantly, the impact on the microbiota community is developed in the biofloc system (Cienfuegos et al 2017). Although the bacterial community plays the most important role in recycling organic materials and metabolizing toxic nitrogen compounds, addition of probiotics does not improve water quality or productive responses from cultivation (Arias-Moscoso et al 2018).

Microbial control in biofloc technology leads to efficient material degradation of waste, efficient nitrification and through manipulation of the C:N ratio. This effort aims to facilitate control and recycling of nitrogen and multiply protein utilization (Luo et al 2013). A balanced C:N ratio will contribute naturally to the formation and stabilization of heterotrophic microbial communities through the availability of nutrients. These microorganisms play three main roles in maintaining water quality by absorbing nitrogen compounds that produce microbial proteins in situ, nutrition increases cultural feasibility by reducing feed conversion ratios (FCR) and decreasing feed costs and competition with pathogens (Becerril-Cortés et al 2018).

The highest C:N ratio value of this study only reached 2.46 in treatment A, this value is still much lower than previous studies conducted in fresh water which is 7.8. (Yurika 2015). This is allegedly due to the use of culture media which different, namely sea water and fresh water. Nitrogen content in marine waters is higher than in freshwaters. Nitrogen is a biological inert gas and the problem of excess N<sub>2</sub> in water is supersaturation. This gas is often saturated in waters which pumped from groundwater. If air saturation increases by more than 110%, this saturation can cause problems in some fish, by causing "gas embolism" or gas bubble disease. This problem can increase the depth of dissolved gas content, therefore, it has implications in using pumped water from the depths for aquaculture purposes (FAO 2018)

As it is known, seawater has a more abundant variety of solutes, which are between 35,000 and 42,000 ppm (Biyantoro & Basuki 2007). Source of nutrient abundance in marine waters comes from various organic compounds carried from the land entering through river flow. Organic compounds that enter in waters will settle and undergo decomposition into inorganic compounds (Rousseau et al 2002). Inorganic compounds contain nutrients in the form of phosphates, nitrates and silicates which are important components for marine biota (Marojahan 2009).

The highest value of C:N ratio in this study is 2.46, this value is much lower compared to previous studies which ranged from 15 to 25 in making biofloc and attempted up to 20 (Amin & Abdul 2010; Gunarto & Suwoyo 2011; Husain et al 2014). Source C in this study comes from molasses and fine bran while for source N comes from fish flour and urea fertilizer. The use of these ingredients still cannot increase the value of the C:N ratio. The low value of the C:N ratio in this study is thought to be due to urea fertilizer as a source of N. Use of ZA fertilizer as a source of N can be used (Gunarto & Suwoyo 2011). The use of urea fertilizer in making biofloc is not recommended because it will be consumed directly by blue-green algae, and can therefore have an effect on blooming (Aiyushirota 2009). When bacteria are fed with organic substrates that contain mainly carbon and little or no nitrogen (sugar, starch, molasses, etc.), consequently they must take nitrogen from the water to produce proteins that needed for cell growth and multiplication. The low C:N ratio is related to the carbon state which functions as a limiting nutrient for heterotrophic bacteria populations in aquaculture ponds (Luo et al 2017).

Comparison of C and N is an important element that must be known in the formation of biofloc elements. The content of C at the end of the study revealed that treatment A and B experienced a decrease, indicating that the presence of probiotic bacteria in treatment A and B made use of element C. Element C is the main element that plays a role in the preparation of bacterial cells (Shewfelt et al 2005). Whereas in treatment C, the content of C experienced an increase because probiotic bacteria were not added to treatment media so that the C content in the media was not widely used by bacteria growing on the media. The contribution of bacteria to the use of carbon substrates relies on changes in bacterial biomass processing CO<sub>2</sub> into C in the pros glucose treatment process (Rinnan & Baath 2009). Increased ratio 10-20 of C:N, increases the volume of biofloc but the ratio 15 C:N, turns out to increase biofloc production better. The C/N ratio causes the best balance of better water quality, nutritional value C and resistance (Dauda et al 2018). If C:N ratio is optimal, BFT can be used to optimize the composition of the bacterial community for optimal water quality and optimal shrimp health. This study shows the possibility of obtaining better

performance than *Litopenaeus vannamei* culture by adjusting the C:N ratio correctly in biofloc based systems (Panigrahi et al 2018).

The N content in treatments A and C has increased while in treatment B has decreased. The increase in N content in treatments A and C shows that the process of reforming organic matter (fish and bran flour) becomes N. However, the increase in N elements cannot be utilized by probiotic bacteria so that the N content in treatments A and C does not show a decrease. While in treatment B, the N content in the media has decreased. It is suspected that in treatment B probiotic bacteria use N more quickly. Besides the use of probiotics B (Pro-1) which have a combination of *Bacillus subtilis* bacteria, *Bacillus licheniformis* and *Lactobacillus plantarum* are able to utilize N better. The high N content in treatment A is suspected because of the combination of the bacteria *Nitrosomonas* sp. and *Bacillus* sp. found in petrofish probiotics can degrade organic nitrogen compounds better so as to increase the content of inorganic N in the waters. *Nitrosomonas* sp. is a nitrifying bacteria which acts to oxidize ammonium to nitrate (Titiresmi & Sopiiah 2006). *Bacillus* sp. is also able to degrade the organic nitrogen content into inorganic nitrogen and produce higher nitrate content (Irianto 2003). In productive systems with major biofloc technology elements for the growth of microorganisms derived from waste compounds from cultures of high density of species of interest for human consumption, which produce complex trophic networks in which nitrogen compounds become important molecules for the development of beneficial bacteria and food for culture (Jimenez-Ojeda et al 2018).

Sequentially the N content in treatment C had the highest content of 3698.28 mg L<sup>-1</sup> compared to other treatments. This is because the treatment of C is not given the addition of probiotic bacteria so that the growth of bacteria in the media is lower than that of treatments A and B. So that the N content in the media is not utilized optimally by bacteria, resulting in high N content in the C media, indicated by calculating the volume of floc formed. The N:P ratio at the end of the study ranged from 22.01 to 25.96. The value of the N:P ratio in this study is much higher than the study on fresh water which only reached 5.68 (Yurika 2015). This is presumably because the use of SP-36 fertilizer as a source of P and urea fertilizer as a source N can be utilized by blue-green algae and other microorganisms so that the value of the N:P ratio in the water becomes higher.

The highest value of the N:P ratio at the end of the study was seen in treatment C (25.96). The high value of the N:P ratio in treatment C shows that the N content in the media is relatively higher and the P content is lower. The increase in N content in treatment C was suspected because it was not done by adding probiotic bacteria so that N in the media was not utilized optimally compared to treatments A and B which were given the addition of probiotic bacteria. P element is one of the limiting factors for aquatic fertility that is closely related to phytoplankton composition (Reynolds et al 2001). The ratio of N:P will affect the autotroph community that will occur in the system (eg chlorophytes versus cyanophytes) (Becerril-Cortés et al 2018).

The P value in treatment C (146.59 ppm) was lower than in the other treatments. The low P value in this study is thought to be due to the phosphate content derived from sp-36 fertilizer. In addition, biofloc is a rich source of natural lipid-protein from food which available 24 hours per day because of the complex interactions between organic matter, physical substrate, and various types of microorganisms. This natural productivity plays an important role in recycling nutrients and maintaining water quality Becerril-Cortés et al 2018).

Water temperature and plankton are water quality parameters that can affect the growth of phytoplankton. At the beginning of the study the pH measurements for each treatment were acidic, ranging from 5.72 to 5.86. This is thought to be due to the process of decomposition of organic matter by decomposing bacteria. In general seawater has a pH value greater than 7 and tends to be alkaline, but under certain conditions the value can be lower than 7 and acidic (Marojahan 2009). But at the end of the study the pH of all treatments experienced an increase of around 7.97-8.11. The pH value at the end of this study still meets the quality standard threshold for the cultivation of marine biota, which ranges from 7 to 8.5. The increase in pH value in this study is

suspected because the use of dolomite lime in the media can increase the pH content in all treatments.

The optimum temperature for the growth of microorganisms is around 28-32°C (Becerril-Cortés et al 2018). This is in accordance with the temperature measurement data in this study experiencing fluctuations which ranged from 26.7 to 26.8°C. The temperature will increase the chemical reaction in photosynthesis by 2-3 times. The increase in the rate of photosynthesis will continue to occur to a certain extent, because the temperature that exceeds the maximum limit will cause a denaturation of proteins and enzymes so that the metabolism in the cell stops (Makmur et al 2012). Temperature has good basis and consequences for metabolism and growth of organisms and the strength of interactions with the environment. The effects of temperature can be limited by changing the maintenance of zooplankton and the availability of nutrients combined with an increase in temperature. The temperature effects also differ for colonies from different compositions, different species of algae and different groups in their optimal temperature. Abiotic conditions affect the biomass of community phytoplankton exposed to different temperature conditions (Striebel et al 2016).

Organic materials in the media will be utilized and decomposed by decomposing bacteria to produce new microbial cells and compounds of CO<sub>2</sub>, NH<sub>3</sub>, H<sub>2</sub>S, and CH<sub>4</sub> as well as other compounds such as phosphorus and sulfide acid components. In this study biofloc began to appear formed in the second week. According to Yurika (2015), bioflocs begin to appear in the second week. Characteristics of the emergence of biofloc, namely the emergence of foam on the surface of the water, the color of the water is brown and gives off an unpleasant aroma. The characteristics of biofloc that have been formed are that there is abundant foam on the surface of the water (Gunarto & Suwoyo 2011). Media that have formed biofloc will be brown and pH value in the range of 7 (Aiyushirota 2009). This is thought to be due to the degradation of organic matter by decomposing bacteria that produce H<sub>2</sub>S compounds. As a result of the degradation process of organic matter in the media resulting in a decrease in DO content. This is in accordance with the results of measurements of DO at the beginning of the study which is relatively low, which is around 2.56-2.85 mg L<sup>-1</sup>. But at the end of the study the measurement of DO increased to 3.69-3.92 mg L<sup>-1</sup>. This shows that the effort to give aeration at 3 points in the media succeeded in increasing the value of DO even though the value was still relatively small.

Ammonia content values range from 1 to 1.6 mg L<sup>-1</sup>. Ammonia levels are very toxic at levels of less than 0.1 mg L<sup>-1</sup> even, they cause toxic effects on some fish. Tropical species can withstand higher toxicity and tropical fish ponds develop ammonia. Levels above 3-4 mg L<sup>-1</sup> are supported by carp and tilapia. Ammonia toxicity is mainly caused by ionized ammonia (UIA) (NH<sub>3</sub>). NH<sub>3</sub> (UIA) is 300-400 times more toxic than NH<sub>4</sub>. The toxicity effect of ammonia is high at high pH, the proportion of ionized ammonia is higher at higher pH. This aspect has indeed been well studied recently mainly because of its importance in intensive cultural systems (FAO 2002).

Biofloc technology (BFT), an aquaculture technique that is environmentally friendly, is based on microorganism production in situ. Continuous water movement in the water column is fully needed to induce macroaggregate formation (biofloc). BFT is considered a new "blue revolution" because nutrition can continue to be recycled and reused in culture media, benefiting from minimum exchange or zero water. Biofloc consumption by shrimp or fish has shown many benefits such as increasing growth rates, decreasing FCR, and associated costs in bait (Becerril-Cortés et al 2018). Results of measurements of seawater quality indicate that they are still within the threshold accepted by the environment. Maintenance and monitoring of water quality in aquaculture is an important practice that aims at the success of the growing cycle. Temperature, DO, pH, salinity, alkalinity, and orthophosphate are some examples of parameters that must be continuously monitored, especially in biofloc technology. Water quality parameters and their interactions in biofloc technology are very important for the development and maintenance of the correct production cycle. The safe range of these parameters will improve the quality of health growth and reduce mortality due to short-term and long-term eating patterns (Becerril-Cortés et al 2018). Biofloc technology has

many benefits for freshwater aquaculture. This technology provides a sustainable tool that is able to solve together the problems related to the environment, social aspects and the economy (Choo & Caipang 2015).

**Conclusions.** The use of different commercial probiotics had no effect on C:N ratio and N:P ratio. The quality of water produced is still in accordance with the standards for biota safety. Efforts are needed to increase the ratio of C:N by knowing the optimum dose of addition of molasses as a source of C as well as knowing that the amount of addition of SP-36 fertilizer is efficient to increase the value of the ratio N:P.

## References

- Alkhudhiri A., Hilal N., 2018 Membrane distillation - principles, applications, configurations, design and implementation. In: Emerging technologies for sustainable desalination handbook. Gude V. G. (ed), Mathew Deans, India, pp. 55-106.
- Amin M., Abdul M., 2010 [The plankton growth in probiotic applications of windu shrimp maintenance (*Panaeus monodon* Fabricius) in controlled tubes]. Aquaculture Prosiding Forum Inovasi Teknologi, pp. 261-268. [in Indonesian]
- Arias-Moscoso J. L., Espinoza-Barron L. G., Miranda-Baeza A., Rivas-Vega M. E., Nieves-Soto M., 2018 Effect of commercial probiotics addition in a biofloc shrimp farm during the nursery phase in zero water exchange. Aquaculture Reports 11:47-52.
- Asmanah D., 2012 [Factors affecting aquaculture production in Central Java]. Thesis in BSc program, Universitas Padjadjaran, Bandung, pp. 1-6. [in Indonesian]
- Avnimelech Y., 2007 Feeding with microbial flocs by tilapia in minimal discharge bio-flocs technology ponds. Aquaculture 264:140-147.
- Becerril-Cortés D., Monroy-Dosta M. D., Emerenciano M. G. C., Castro-Mejía G., Schettino Bermúdez B. S., Correa G. V., 2018 Effect on nutritional composition of produced bioflocs with different carbon sources (molasses, coffee waste and rice bran) in biofloc system. International Journal of Fisheries and Aquatic Studies 6(2):541-547.
- Bhatnagar A., Devi P., 2013 Water quality guidelines for the management of pond fish culture. International Journal of Environmental Sciences 3(6):1980-2009.
- Biyantoro D., Basuki K. T., 2007 [Measurement and elemental analysis of Muria Sea water for primary water of PWR]. Prosiding PPI – PDIPTN. PWR, Yogyakarta, pp. 68-73. [in Indonesian]
- Bossier P., Ekasari J., 2017 Biofloc technology application in aquaculture to support sustainable development goals. Microbial Biotechnology 10(5):1012-1016.
- Boyd C. E., Green B. W., 2002 Coastal water quality monitoring in shrimp farming areas: an example from Honduras. A report prepared for the World Bank, network of Aquaculture Centres in Asia-Pacific, World Wildlife Fund and Agriculture Organization of the United Nations Consortium Program on Shrimp Farming and the Environment, 29 pp.
- Cienfuegos M. K., Monroy D. M., Hamdan P. A., Castro M. J., Becerril C. D., 2017 Probiotics used in biofloc system for fish and crustacean culture: a review. International Journal of Fisheries and Aquatic Science 5(5):120-125.
- Choo H. X., Caipang C. M. A., 2015 Biofloc technology (BFT) and its application towards improved production in freshwater tilapia culture. AACL Bioflux 8(3):362-366.
- Crab R., Avnimelech Y., Defoirdt T., Bossier P., Verstraete W., 2007 Nitrogen removal techniques in aquaculture for sustainable production. Aquaculture 270(1-4):1-14.
- Cole D. W., Cole R., Gaydos S. J., Gray J., Hyland G., Jacques M. L., Powell-Dunford N., Sawhney C., Au W. W., 2009 Aquaculture: environmental, toxicological, and health issues. International Journal of Hygiene and Environmental Health 212:369-377.
- Dauda A. B., Romano N., Ebrahimi M., Teh J. C., Ajadi A., Chong C. M., Karim M., Natrah I., Kamarudin M. S., 2018 Influence of carbon/nitrogen ratios on biofloc production and biochemical composition and subsequent effects on the growth, physiological status and disease resistance of African catfish (*Clarias gariepinus*) cultured in glycerol-based biofloc systems. Aquaculture 483:120-130.

- Ekasari J., 2009 [Biofloc technology: theory and practical in intensive aquaculture. system]. *Jurnal Akuakultur Indonesia* 8(2):117-126. [in Indonesian]
- FAO, 2002 Ammonia solution. pp. 2-3. Available at: [www.fao.org/fileadmin/user\\_upload/jecfa\\_additives/docs/.../Additive-019.pdf](http://www.fao.org/fileadmin/user_upload/jecfa_additives/docs/.../Additive-019.pdf). Accessed: March, 2019.
- FAO, 2018 The state of fisheries and aquaculture: meeting the sustainable development goals. FAO, Rome, 210 pp.
- Fry J. P., Love D. C., MacDonald G. K., West P. C., Engstrom P. M., Nachman K. E., Lawrence R. S., 2016 Environmental health impacts of feeding crops to farmed fish. *Environment International* 91:201-214.
- Garcia S. M., Rosenberg A. A., 2010 Food security and marine capture fisheries: characteristics, trends, drivers and future perspectives. *Philosophical Transactions of the Royal Society B* 365:2869-2880.
- Gunarto, Suwoyo H. S., 2011 [Biofloc production and its nutritional value on the laboratory scale]. *Prosiding Forum Inovasi Teknologi Akuakultur 2011*, pp. 1009-1017. [in Indonesian]
- Husain N., Putri B., Supono, 2014 [Carbon and nitrogen comparison in biofloc systems against red tilapia growth (*Oreochromis niloticus*)]. *Jurnal Rekayasa dan Teknologi Budidaya Perairan* 3(1):343-350. [in Indonesian]
- Irianto A., 2003 [Probiotic aquaculture]. Gadjah Mada University Press, Yogyakarta, 125 pp. [in Indonesian]
- Jimenez-Ojeda Y. K., Collazos-Lasso L. F., Arias-Castellanos J. A., 2018 Dynamics and use of nitrogen in Biofloc Technology - BFT. *AAFL Bioflux* 11(4):1107-1129.
- Luo G. Z., Avnimelech Y., Pan Y. F., Tan H. X., 2013 Effects of temperature on inorganic nitrogen dynamics in sequencing batch reactors using biofloc technology to treat aquaculture sludge. *Aquacultural Engineering* 52:73-79.
- Luo G., Li W., Tan H., Chen X., 2017 Comparing salinities of 0, 10 and 20 in biofloc genetically improved farmed tilapia (*Oreochromis niloticus*) production systems. *Aquaculture and Fisheries* 2(5):220-226.
- Makmur M., Kusnoputranto H., Moersidik S. S., Wisnubroto D. S., 2012 [Influence of the organic waste and its N/P ratio to phytoplankton diversity in Cilincing mussel farming]. *Jurnal Teknologi Pengelolaan Limbah* 15(2):59-64. [in Indonesian]
- Marojahan, 2009 [The relationship of physical and chemical environmental factors to the distribution of plankton in the waters of the eastern Pacific Islands of Bangka Belitung]. *Jurnal Perikanan Universitas Gajah Mada* 11(1):31-45. [in Indonesian]
- Martinez-Porchas M., Martinez-Cordova L. R., 2012 World aquaculture: environmental impacts and troubleshooting alternatives. *Scientific World Journal* 2012:389623.
- Miyazawa M., Pavan M. A., De Oliveira E. L., Ionashiro M., Silva A. K., 2000 Gravimetric determination of soil organic matter. *Brazilian Archives of Biology and Technology* 43(5):475-478.
- Nani S. Z. M., Majid F. A. A., Jaafar A. B., Mahdzir A., Musa M. N., 2016 Potential health benefits of deep sea water: a review. *Evidence-Based Complementary and Alternative Medicine* 2016:6520475, 18 pp.
- Panigrahi A., Saranya C., Sundaram M., Vinoth Kannan S. R., Das R. R., Satish Kumar R., Rajesh P., Otta S. K., 2018 Carbon:Nitrogen (C:N) ratio level variation influences microbial community of the system and growth as well as immunity of shrimp (*Litopenaeus vannamei*) in biofloc based culture system. *Fish and Shellfish Immunology* 81:329-337.
- Purwanta W., Firdayati M., 2002 [The effect of probiotic microbial applications on the chemical quality of shrimp pond waters]. *Jurnal Teknologi Lingkungan* 3(1):62-65. [in Indonesian]
- Reynolds C. S., Irish A. E., Elliott J. A., 2001 The ecological basis for simulating phytoplankton response to environmental change (PROTECH). *Ecological Modelling* 140:271-291.
- Rinnan R., Baath E., 2009 Differential utilization of carbon substrates by bacteria and fungi in tundra soil. *Applied and Environmental Microbiology* 75(11):3611-3620.



- Rousseau V., Leynaert A., Daoud N., Lancelot C., 2002 Diatom succession, silicification and silicic acid availability in Belgian coastal waters (southern North Sea). *Marine Ecology Progress Series* 236:61-73.
- Saez-Plaza P., Navas M. J., Wybraniec S., Michalowski T., Asuero A. G., 2013 An overview of the Kjeldahl method of nitrogen determination. Part II: sample preparation, working scale, instrumental finish, and quality control. *Critical Reviews in Analytical Chemistry* 43(4):224-272.
- Schneider O., Sereti V., Eding E. H., Verret J. A. C., 2005 Protein production by heterotrophic bacteria using carbon supplemented fish waste. Paper Presented in Bali World Aquaculture International Peace and Development through Aquaculture - Proceeding 2005, pp 562.
- Shewfelt K., Lee H., Zytner R. G., 2005 Optimization of nitrogen for bioventing of gasoline contaminated soil. *Journal of Environmental Engineering and Science* 4:29-42.
- Souther I. M., Rissik D., 2009 *Plankton: a guide to their ecology and monitoring for water quality*. CSIRO Publishing, Collingwood, 115 pp.
- Striebel M., Schabhutti S., Hodapp D., Hingsamer P., Hillebrand H., 2016 Phytoplankton responses to temperature increases are constrained by abiotic conditions and community composition. *Oecologia* 182(3):815-827.
- Titiresmi, Sopiah, 2006 [Biofilter technology for ammonia waste treatment]. *Jurnal Teknologi Lingkungan* 7(2):173-179. [in Indonesian]
- Xu W. J., Morris T. C., Samocha T. M., 2016 Effects of C/N ratio on biofloc development, water quality, and performance of *Litopenaeus vannamei* juveniles in a biofloc-based, high-density, zero-exchange, outdoor tank system. *Aquaculture* 453:169-175.
- Yurika D., 2015 [The effect of commercial probiotics differences to C:N ratio and N:P ratio in biofloc culture media at experimental tube]. BSc Thesis, Universitas Airlangga, Surabaya, 62 pp. [in Indonesian]

Received: 15 February 2019. Accepted: 30 March 2019. Published online: 30 April 2019.

Authors:

Luthfi Adi Nugroho, Department of Fisheries and Marine, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya, East Java, Indonesia, e-mail: lutfiadinugroho@yahoo.com

Endang Dewi Masithah, Department of Fisheries and Marine, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya, East Java, Indonesia, e-mail: endangdewimashitah@yahoo.com

Woro Hastuti Satyanrini, Department of Fisheries and Marine, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya, East Java, Indonesia, e-mail: wsatyanrini@yahoo.com

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Nugroho L. A., Masithah E. D., Satyanrini W. H., 2019 Evaluation of biofloc technology: the risk of giving different commercial probiotics to C:N and N:P ratio and quality of seawater. *AAFL Bioflux* 12(2):706-715.

# Evaluation of biofloc technology the risk of giving

---

## ORIGINALITY REPORT

---

20%

SIMILARITY INDEX

10%

INTERNET SOURCES

16%

PUBLICATIONS

0%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

- |   |   |    |
|---|---|----|
| 1 | <a href="http://www.pr.bioflux.com.ro">www.pr.bioflux.com.ro</a><br>Internet Source   | 1% |
| 2 | W H Satyantini, M Salsabila, D D Nindarwi, A M Sahidu, A T Mukti. " Provision of bacteria from shrimp pond sediment towards N/P ratio, plankton abundance, and total bacteria in the culture media of white shrimp ( ) ", IOP Conference Series: Earth and Environmental Science, 2020<br>Publication | 1% |
| 3 | Ahmed M. Aboseif et al.. "Feed restriction and compensatory growth of the Nile tilapia, <i>Oreochromis niloticus</i> under Biofloc system condition", Egyptian Journal of Aquatic Biology and Fisheries, 2022<br>Publication  | 1% |
| 4 | Mervat A. M. Ali et al.. "The effect of different stocking densities of marine shrimp larvae <i>Litopenaeus vannamei</i> on growth performance and survival rate using biofloc  | 1% |

# technology", Egyptian Journal of Aquatic Biology and Fisheries, 2021

Publication

---

5

[msp.fpik.ipb.ac.id](http://msp.fpik.ipb.ac.id)

Internet Source

1 %

---

6

Iis Diatin, Dadang Shafruddin, Nurul Hude, Mar'atus Sholihah, Ilmi Mutsmir. "Production performance and financial feasibility analysis of farming catfish (*Clarias gariepinus*) utilizing water exchange system, aquaponic, and biofloc technology", Journal of the Saudi Society of Agricultural Sciences, 2021

Publication

1 %

---

7

Abdelhamid Mohamed Abdelhamid, Mohamed Moaaz Refaey, Hazem Ga'far Mohammed El-Sayed, Gehad Ahmed Mo'aty. "Comparison Between Blood Haematology of Egyptian Freshwater Food Fish and Ornamental Fish", Journal of Applied Science, Engineering, Technology, and Education, 2020

Publication

1 %

---

8

Hailiang Wang, Xiaoyuan Wan, Guosi Xie, Xuan Dong, Xiuhua Wang, Jie Huang. "Insights into the histopathology and microbiome of Pacific white shrimp, *Penaeus vannamei*, suffering from white feces syndrome", Aquaculture, 2020

Publication

1 %

---

9

A Mashar, A P N Firdaussyia, M Krisanti, A A Hakim. "Biodiversity of macroinvertebrate in artificial substrate from several habitats at Ponelo Island, Gorontalo", IOP Conference Series: Earth and Environmental Science, 2021

Publication

---

<1 %

10

Bocheng Huang, Shuang Zhang, Xiaohui Dong, Shuyan Chi, Qihui Yang, Hongyu Liu, Beiping Tan, Shiwei Xie. "Effects of fishmeal replacement by black soldier fly on growth performance, digestive enzyme activity, intestine morphology, intestinal flora and immune response of pearl gentian grouper (*Epinephelus fuscoguttatus* ♀ × *Epinephelus lanceolatus* ♂)", *Fish & Shellfish Immunology*, 2022

Publication

---

<1 %

11

Birdie Scott Padam, Fook Yee Chye. "Seaweed components, properties, and applications", Elsevier BV, 2020

Publication

---

<1 %

12

Sri Rejeki, Adolphe O. Debrot, Anneke M. den Brink, Restiana W. Ariyati, Lestari Lakshmi Widowati. " Increased production of green mussels ( ) using longline culture and an economic comparison with stake culture on the north coast of Java, Indonesia ", *Aquaculture Research*, 2020

<1 %

13

Y P Paulangan, K Rumbiak, B Barapadang.  
"Fishing season and participatory mapping of  
the fishing ground of target fish in Depapre  
Bay, Jayapura Regency, Papua Indonesia", IOP  
Conference Series: Earth and Environmental  
Science, 2020

Publication

---

14

[ejournal.undip.ac.id](http://ejournal.undip.ac.id)

Internet Source

---

15

Russell G. Uc-Peraza, Ítalo B. Castro, Gilberto  
Fillmann. "An absurd scenario in 2021:  
Banned TBT-based antifouling products still  
available on the market", Science of The Total  
Environment, 2021

Publication

---

16

[eprints.ugd.edu.mk](http://eprints.ugd.edu.mk)

Internet Source

---

17

[udsspace.uds.edu.gh](http://udsspace.uds.edu.gh)

Internet Source

---

18

[event.ners.unair.ac.id](http://event.ners.unair.ac.id)

Internet Source

---

19

[ojs.omniakuatika.net](http://ojs.omniakuatika.net)

Internet Source

---

20

[www.jove.com](http://www.jove.com)

Internet Source

---

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

21 Brittany M. Cummings, Joseph A. Needoba, Tawnya D. Peterson. "Effect of metformin exposure on growth and photosynthetic performance in the unicellular freshwater chlorophyte, *Chlorella vulgaris*", PLOS ONE, 2018  
Publication <1 %

---

22 [eprints.ums.edu.my](http://eprints.ums.edu.my)  
Internet Source <1 %

---

23 [www.sciencegate.app](http://www.sciencegate.app)  
Internet Source <1 %

---

24 "Effect of Protein Levels on Growth Performance, Feed Utilization and Economic Evaluation of Fingerlings Nile Tilapia Fingerlings under Biofloc System", Journal of Animal, Poultry & Fish Production, 2020  
Publication <1 %

---

25 Ludson Guimarães Manduca, Marcos Antônio da Silva, Érika Ramos de Alvarenga, Gabriel Francisco de Oliveira Alves et al. "Effects of different stocking densities on Nile tilapia performance and profitability of a biofloc system with a minimum water exchange", *Aquaculture*, 2021  
Publication <1 %

---

26 [ndl.ethernet.edu.et](http://ndl.ethernet.edu.et)  
Internet Source <1 %

---

27

Wenwen Jiang, Wenjing Ren, Li Li, Shuanglin Dong, Xiangli Tian. "Light and carbon sources addition alter microbial community in biofloc-based *Litopenaeus vannamei* culture systems", *Aquaculture*, 2020

Publication

&lt;1 %

28

David O. Caldwell. "Sterile neutrinos and supernova nucleosynthesis", *Physical Review D*, 05/2000

Publication

&lt;1 %

29

[es.slideshare.net](http://es.slideshare.net)

Internet Source

&lt;1 %

30

[www.jurnal.unsyiah.ac.id](http://www.jurnal.unsyiah.ac.id)

Internet Source

&lt;1 %

31

B Pantjara, A Sarijanah, M Hermawan. "Improved pond productivity through integrated cultivation of red tilapia ( ), tiger shrimp ( ) and seaweed ( ) in Maros, South Sulawesi, Indonesia ", *IOP Conference Series: Earth and Environmental Science*, 2020

Publication

&lt;1 %

32

Maren Striebel, Stefanie Schabhüttl, Dorothee Hodapp, Peter Hingsamer, Helmut Hillebrand. "Phytoplankton responses to temperature increases are constrained by abiotic conditions and community composition", *Oecologia*, 2016

Publication

&lt;1 %

---

33 Mario Coccia, Diego Margon, Ugo Finardi. <math><1\%</math>  
"Chapter 14 CURRENT SCIENTIFIC  
TRAJECTORIES AND INTERACTIONS BETWEEN  
MAIN WORLD GEO-ECONOMIC PLAYERS IN  
NANOTECHNOLOGY RESEARCH", IntechOpen,  
2012  
Publication

---

34 [animalsciencejournal.usamv.ro](http://animalsciencejournal.usamv.ro) <math><1\%</math>  
Internet Source

---

35 PHYLLIS A. HICKS. "THE CARBON/NITROGEN  
RATIO IN THE WHEAT PLANT", New  
Phytologist, 4/1928 <math><1\%</math>  
Publication

---

36 [www.isumadecip.ro](http://www.isumadecip.ro) <math><1\%</math>  
Internet Source

---

37 Deswati Deswati, Elsa Yani, Safni Safni, Olly  
Norita Tetra, Hilfi Pardi. "Development  
methods in aquaponics systems using biofloc  
to improve water quality (ammonia, nitrite,  
nitrate) and growth of tilapia and samhong  
mustard", International Journal of  
Environmental Analytical Chemistry, 2020 <math><1\%</math>  
Publication

---

38 S. Souissi. "Spatial characterization of nutrient  
dynamics in the Bay of Tunis (south-western  
Mediterranean) using multivariate analyses:  
consequences for phyto- and zooplankton <math><1\%</math>



distribution", Journal of Plankton Research,  
11/01/2000

Publication

---

39

S. M. Garcia. "Food security and marine capture fisheries: characteristics, trends, drivers and future perspectives", Philosophical Transactions of The Royal Society B Biological Sciences, 09/27/2010

Publication

---

<1 %

40

Wellica G. Reis, Wilson Wasielesky, Paulo C. Abreu, Hellyjúnyor Brandão, Dariano Krummenauer. "Rearing of the Pacific white shrimp *Litopenaeus vannamei* (Boone, 1931) in BFT system with different photoperiods: Effects on the microbial community, water quality and zootechnical performance", Aquaculture, 2019

Publication

---

<1 %

41

[www.flowserve.com](http://www.flowserve.com)

Internet Source

---

<1 %

42

D Ningrum, M Arief, dan K T Pursetyo. "Comparative Test on Bacteria in the Digestive Tract of *Vannamei* Shrimp ( ) at Intensive and Extensive Ponds in Ujungpangkah, Gresik ", IOP Conference Series: Earth and Environmental Science, 2019

Publication

---

<1 %

43

Internet Source

&lt;1 %

44

[lumenpublishing.com](http://lumenpublishing.com)

Internet Source

&lt;1 %

45

Ravindra Sontakke, Virendra Kumar Tiwari, Paniprasad Kurcheti, Babitha Rani Asanaru Majeedkutty et al. " Yam - based biofloc system improves the growth, digestive enzyme activity, bacterial community structure and nutritional content in milkfish ( ) ", Aquaculture Research, 2021

Publication

&lt;1 %

46

Solmaz Aslanzadeh, Katherine Kho, Irnayuli Sitepu. "An Evaluation of the Effect of Takakura and Effective Microorganisms (EM) as Bio Activators on the Final Compost Quality", IOP Conference Series: Materials Science and Engineering, 2020

Publication

&lt;1 %

47

[journal.ipb.ac.id](http://journal.ipb.ac.id)

Internet Source

&lt;1 %

48

[ppgaquicultura.furg.br](http://ppgaquicultura.furg.br)

Internet Source

&lt;1 %

49

[repository.ju.edu.et](http://repository.ju.edu.et)

Internet Source

&lt;1 %

50

[www.bioflux.com.ro](http://www.bioflux.com.ro)

Internet Source

<1 %

51

Akeem Babatunde Dauda, Nicholas Romano, Mahdi Ebrahimi, Jun Chin Teh et al. "Influence of carbon/nitrogen ratios on biofloc production and biochemical composition and subsequent effects on the growth, physiological status and disease resistance of African catfish ( *Clarias gariepinus* ) cultured in glycerol-based biofloc systems", *Aquaculture*, 2018

<1 %

Publication

52

Asaduzzaman, M.. "C/N ratio control and substrate addition for periphyton development jointly enhance freshwater prawn *Macrobrachium rosenbergii* production in ponds", *Aquaculture*, 20080801

<1 %

Publication

53

Ashraf M. Soliman, Mohsen Abdel-Tawwab. "Effects of different carbon sources on water quality, biofloc quality, and the productivity of Nile tilapia reared in biofloc-based ponds", *Annals of Animal Science*, 2022

<1 %

Publication

54

Guozhi Luo, Wenyan Liang, Hongxin Tan, Chen Yao, Nan Zhang, Lu Lu. "Effects of calcium and magnesium addition on the start-up of sequencing batch reactor using biofloc

<1 %

technology treating solid aquaculture waste",  
Aquacultural Engineering, 2013

Publication

---

55

Md. Eilious Hosain, Sarker Mohd Nurul Amin, Mohd Salleh Kamarudin, Aziz Arshad, Nicholas Romano. " Effects of C - N ratio on growth, survival and proximate composition of post larvae reared under a corn starch based zero - exchange brackish water biofloc system ", Aquaculture Research, 2021

Publication

---

<1 %

56

Sweileh, W. M., S. H. Zyoud, S. Al-Khalil, S. W. Al-Jabi, and A. F. Sawalha. "Assessing the Scientific Research Productivity of the Palestinian Higher Education Institutions: A Case Study at An-Najah National University, Palestine", SAGE Open, 2014.

Publication

---

<1 %

57

Tran Nguyen Duy Khoa, Chau Tai Tao, Ly Van Khanh, Tran Ngoc Hai. "Super-intensive culture of white leg shrimp (*Litopenaeus vannamei*) in outdoor biofloc systems with different sunlight exposure levels: Emphasis on commercial applications", Aquaculture, 2020

Publication

---

<1 %

58

Van Hai, Ngo, Ravi K. Fotedar, and Nguyen Van Hao. "Penaeid Prawns", Recent Advances

<1 %

and New Species in Aquaculture  
Fotedar/Recent Advances and New Species in  
Aquaculture, 2011.

Publication

59

Y N Ihsan, B Nathanael, R Permana, Y Mulyani, T D K Pribadi. " Identification of lead (Pb(CH COO) )-reducing bacteria using 16S rRNA Gene ", IOP Conference Series: Earth and Environmental Science, 2020

Publication

<1 %

60

[iopscience.iop.org](http://iopscience.iop.org)

Internet Source

<1 %

61

[journals.plos.org](http://journals.plos.org)

Internet Source

<1 %

62

[link.springer.com](http://link.springer.com)

Internet Source

<1 %

63

"Alphabetical Index of Libraries", Walter de Gruyter GmbH, 2007

Publication

<1 %

64

Alfredo Gallardo-Collí, Carlos Iván Pérez-Rostro, Martha Patricia Hernández-Vergara, Ignacio Alejandro Pérez-Legaspi. "Microeukaryote community and the nutritional composition of the biofloc during Nile tilapia culture in water-reusing biofloc systems", Aquaculture International, 2019

Publication

<1 %

65

De Galan, S.. "Spatial and temporal trends in nutrient concentrations in the Belgian Continental area of the North Sea during the period 1993-2000", *Estuarine, Coastal and Shelf Science*, 200411

Publication

---

<1 %

66

Diana Chilmawati, Johannes Hutabarat, Sutrisno Anggoro, Suminto Suminto. " Effects of Aeration Flow Rate in the Culture Medium on the Growth Performance and Egg Production of Copepod Fed with Fermented Organic Diet ", *E3S Web of Conferences*, 2020

Publication

---

<1 %

67

Mazdak Aalimahmoudi, Hamid Mohammadiazarm. "Dietary protein level and carbon/nitrogen ratio manipulation in bioflocs rearing of *Cyprinus carpio* juvenile: Evaluation of growth performance, some blood biochemical and water parameters", *Aquaculture*, 2019

Publication

---

<1 %

68

Melody Allison. "Comparison of CINAHL® via EBSCOhost®, Ovid®, and ProQuest®", *Journal of Electronic Resources in Medical Libraries*, 04/21/2006

Publication

---

<1 %

69

Nayara Nunes Caldini, Hermano Hertz de Almeida Capistrano, Pedro Roberto Nogueira

<1 %

Rocha-Filho, Marcelo Vinícius do Carmo e Sá.  
"Partial replacement of artificial diets by  
wet bioflocs biomass in Nile tilapia culture  
tanks", *Acta Scientiarum. Animal Sciences*,  
2018

Publication

---

70

OA Artha, Sudarno, H Pramono, LA Sari.  
"Identification of extracellular enzyme-  
producing bacteria (proteolytic,  
cellulolytic, and amylolytic) in the  
sediment of extensive ponds in  
Tanggulrejo, Gresik", *IOP Conference  
Series: Earth and Environmental Science*,  
2019

Publication

---

71

Roselien Crab. "The effect of different  
carbon sources on the nutritional value of  
bioflocs, a feed for *Macrobrachium  
rosenbergii* postlarvae", *Aquaculture  
Research*, 03/2010

Publication

---

72

Sabiriba Alain Hema, Mamadou Traore,  
Kalifa Coulibaly, Bazoumana Koulibaly,  
Hassan Bismarck Nacro. "Agricultural Soil  
Fertilizing Potential of Dry Faecal Sludge  
from Treatment Plants in Burkina Faso",  
*Open Journal of Soil Science*, 2022

Publication

---

<1 %

<1 %

<1 %

73

Saranya Chakrapani, Akshaya Panigrahi, Jayashree Sundaresan, Mullaivanam R. Sivakumar, Rajesh Palanisamy, Vinodh Kumar. " Three different C: N ratios for Pacific white shrimp, under practical conditions: Evaluation of growth performance, immune and metabolic pathways ", Aquaculture Research, 2020

Publication

---

<1%

74

WANG, C.. "Variation of phytoplankton community before an induced cyanobacterial (Arthrospira platensis) bloom", Journal of Environmental Sciences, 2009

Publication

---

<1%

75

Wen-Chi Tseng, Chien-Hsiang Tang, Tsuei-Yun Fang. "The role of dextran conjugation in transfection mediated by dextran-grafted polyethylenimine", The Journal of Gene Medicine, 2004

Publication

---

<1%

76

Ying Liu, Des E. Walling, Mingyi Yang, Fengbao Zhang. "Sediment source fingerprinting and the temporal variability of source contributions", Journal of Environmental Management, 2023

Publication

---

<1%



77	Internet Source	<1 %
78	<a href="http://dumas.ccsd.cnrs.fr">dumas.ccsd.cnrs.fr</a> Internet Source	<1 %
79	<a href="http://dyuthi.cusat.ac.in">dyuthi.cusat.ac.in</a> Internet Source	<1 %
80	<a href="http://epdf.pub">epdf.pub</a> Internet Source	<1 %
81	<a href="http://irep.iium.edu.my">irep.iium.edu.my</a> Internet Source	<1 %
82	<a href="http://journal.uinjkt.ac.id">journal.uinjkt.ac.id</a> Internet Source	<1 %
83	<a href="http://science.buu.ac.th">science.buu.ac.th</a> Internet Source	<1 %
84	<a href="http://www.atlantis-press.com">www.atlantis-press.com</a> Internet Source	<1 %
85	<a href="http://www.frontiersin.org">www.frontiersin.org</a> Internet Source	<1 %
86	<a href="http://www.scielo.cl">www.scielo.cl</a> Internet Source	<1 %
87	El-Sayed Hemdan Eissa, Fatima S. Alaryani, Marwa S. Khattab, Amal Elfeky, Moaheda E.H. Eissa, Khaled Youssef AbouelFadl. "The effect of <i>Pediococcus acidilactici</i>	<1 %

probiotics supplementation in the feed of Nile Tilapia (*Oreochromis niloticus*) on water quality, growth performance, Hematological parameters, antioxidants, immunological responses, enzymatic Activity, and histology.", Research Square Platform LLC, 2022

Publication

---

88

Jinghao Li, Jiayao Li, Wei Li, Ying Sun, Xiaofei Liu, Mingming Liu, Yongxu Cheng. " Juvenile farmed using biofloc technology or commercial feed in zero - water exchange indoor tanks: A comparison of growth performance, enzyme activity and proximate composition ", Aquaculture Research, 2019

Publication

---

<1%

89

Shuiping Cheng, Feng Ren, Wolfgang Grosse, Zhenbin Wu. " Effects of Cadmium on Chlorophyll Content, Photochemical Efficiency, and Photosynthetic Intensity of Linn. ", International Journal of Phytoremediation, 2002

Publication

---

<1%

Exclude quotes  Off

Exclude matches  Off

Exclude bibliography  On

# Evaluation of biofloc technology the risk of giving

GRADEMARK REPORT

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14

PAGE 15

PAGE 16