

Table of contents

Volume 679

2021

◀ Previous issue Next issue ▶

The 1st International Conference on Biotechnology and Food Sciences 11 September 2020, Surabaya, Indonesia

Accepted papers received: 08 February 2021

Published online: 26 February 2021

Open all abstracts

Preface

OPEN ACCESS 011001

The 1st International Conference on Biotechnology and Food Sciences (INCOBIFS)
Surabaya Indonesia, 11 September 2020

+ Open abstract  View article  PDF

OPEN ACCESS 011002

Conference Photographs

+ Open abstract  View article  PDF

OPEN ACCESS 011003

Organizing Committee

+ Open abstract  View article  PDF

OPEN ACCESS 011004

Peer review declaration

+ Open abstract  View article  PDF

Papers

OPEN ACCESS 012001

Converting husbandry waste into liquid organic fertilizer using probiotic consortiums
(*Lactobacillus* sp., *Rhodopseudomonas* sp., *Actinomycetes* sp., *Streptomyces* sp.)

S Amrullah, M. Amin and M Ali

+ Open abstract  View article  PDF

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



OPEN ACCESS

OPEN ACCESS

A review: bioactive compounds of macroalgae and their application as functional beverages

012002

S G Widyaswari, Metusalach, Kasmiasi and N Amir

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012003

Improvement quality of sugar cane bagasse as fish feed ingredient

L H Suryaningrum and R Samsudin

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012004

The Growth and Yields of Shallot (*Allium Wakegi* Araki) CV. lembah palu Growing under Hydroponic Substrate Systems

R Yusuf, S A Lasmini, M Sandi, A Rahim and I Wahyudi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012005

Application of glycerol on bioplastic based carrageenan waste cellulose on biodegradability and mechanical properties bioplastic

S N Fauziyah, A S Mubarak and D Y Pujiastuti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012006

Gill and skin pathology of hybrid grouper (*E. fuscoguttatus* x *E. lanceolatus*) infested *Zeylanicobdella arugamensis* worms in different infestations degree

M Nisa, G Mahasri and L Sulmartiwi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012007

Growth performance of tambaqui (*Colossoma macropomum*) supplemented with honey prebiotic in stagnant peat ponds

H Silalahi, R Djauhari and S S Monalisa

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012008

The Concentration of polyethylen glycol (PeG) 400 on bioplastic cellulose based carrageenan waste on biodegradability and mechanical properties bioplastic

D S Maulana, A S Mubarak and D Y Pujiastuti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012009

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



Fish oil extraction as a by-product of Tilapia (*Oreochromis* sp.) fish processing with dry rendering method

S H Suseno, A K Rizkon, A M Jacob, Kamini and D Listiana

[+](#) Open abstract [View article](#) [PDF](#)

OPEN ACCESS

012010

Optimization of Extraction Time on The Characteristic of Gelatin from Scales of Red Snapper (*Lutjanus* sp.)

I Safi'i, W Tjahjaningsih and E D Masithah

[+](#) Open abstract [View article](#) [PDF](#)

OPEN ACCESS

012011

Screening acetylcholinesterase inhibitors from marine-derived actinomycetes by simple chromatography

M Kamaruddin, I Marzuki, A Burhan and R Ahmad

[+](#) Open abstract [View article](#) [PDF](#)

OPEN ACCESS

012012

Assessment of Heavy Metal Lead (Pb) Contents in Canned Crab Products by Atomic Absorption Spectrophotometry (AAS)

M Agustina, Mulyono and W Tjahjaningsih

[+](#) Open abstract [View article](#) [PDF](#)

OPEN ACCESS

012013

The concentration of sorbitol on bioplastic cellulose based carrageenan waste on biodegradability and mechanical properties bioplastic

M D Arief, A S Mubarak and D Y Pujiastuti

[+](#) Open abstract [View article](#) [PDF](#)

OPEN ACCESS

012014

Identification photoprotective activity of marine seaweed: *Eucheuma* sp

R D Kasitowati, A Wahyudi, R Asmara, D Aliviyanti, F Iranawati, M A P Panjaitan, D C Pratiwi and S Arsad

[+](#) Open abstract [View article](#) [PDF](#)

OPEN ACCESS

012015

Analysis of the density and intensity of metallothionein in *Crassostrea cucullata* oyster hulls in the coastal fishing port of Mayangan Probolinggo, East Java using immunohistochemical techniques

W Isoni, N Maulida and M I Mubaroqi

[+](#) Open abstract [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more,

[see our Privacy and Cookies policy.](#)

012016 

The effect of different bait on the catch of traps in the waters of the tip of Pangkah Gresik Regency, East Java

W Isoni, N Maulida and M Huda

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012017

The Identification and Distribution Components of Polycyclic Aromatic Hydrocarbon Contaminants at the Port of Paotere, Makassar, South Sulawesi

I Marzuki, I Pratama, H E Ismail, I. Paserangi, M Kamaruddin, M. Chaerul and R Ahmad

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012018

Koi (*Cyprinus carpio*) Hatchery techniques: its performance in BBI Boyolali

M G Laksono, Sugianta and M B Santanumurti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012019

Ectoparasite infestation and survival rate of pacific white shrimp (*Litopenaeus vannamei*) that immunized with crude protein *Zoothamnium penaei* in intensive ponds

G Mahasri, A T Mukti, M Nisa, G C Prakosa and W H Satyantini

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012020

Study of crude extract yield and gonad maturity level of sea cucumber *Phylloporus dobsoni* correlation

S Andriyono, E D Masithah and D Winarni

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012021

Techniques of additional *Kappaphycus alvarezii* on seaweed face mask production

N R Prima and S Andriyono

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012022

Ice cream properties affected by carrageenan from seaweed deference type drying methods

I Irawan and Fitriyana

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012023

The effect of washing on the making of surimi and kamaboko tilapia (*Oreochromis sp.*)

E Saputra, W Tjanjungsih and A A Abdillah

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012024

The effect of storage on the making of surimi and kamaboko tilapia (*Oreochromis* sp.)

E Saputra, W Tjahjaningsih and A A Abdillah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012025

Molecular identification and prevalence of endoparasite worms in Silver pompano (*Trachinotus blochii*) in floating net cages of Mari-culture Center, Lampung

L N F Haryanto, S Subekti, H B Ardiyanti, M K Amiin, R E K Akbar, I Achmadi and M A Yudarana

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012026

Microbiology safety of green mussel, *Perna viridis* after treated with boiling and *sous vide*

A S Samsudin and N U Karim

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012027

The utilization of chitosan from Comb-pen shell (*Atrina pectinata*) as an emulsion stabilizer in the production of hand body cream

Y Supratin, L Sulmartiwi and E Saputra

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012028

The potential of chitosan from comb-pen (*Atrina pectinata*) shell waste on the characteristics of hand body cream

F Fauzi, L Sulmartiwi and E Saputra

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012029

Spawning technique of yellowfin tuna (*Thunnus albacares*) infloating nets cage

B Bramantya, Gunawan and L A Sari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012030


Enlargement technique of humpback grouper (*Cromileptes altivelis*) with floating nets cage

G Y Pamungkas and L A Sari

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



-
- OPEN ACCESS** 012031
Microencapsulated fish oil powder by spray drying using combination of wall materials in Kasetsart University, Bangkok
I Aprilia, D Y Pujiastuti and W Klaypradit
[+](#) Open abstract [View article](#) [PDF](#)
-
- OPEN ACCESS** 012032
Effect of deacetylation conditions on physicochemical properties of chitosan derived from shrimp shell and squid pen
A R Basarah, D Y Pujiastuti and Yaowapha Waiprib
[+](#) Open abstract [View article](#) [PDF](#)
-
- OPEN ACCESS** 012033
Substitution of patin *Pangasius pangasius* flour in making sticks as an alternative of food high protein and source of calcium for autism patients
V Amelia, S Subekti and L Sulmartiwi
[+](#) Open abstract [View article](#) [PDF](#)
-
- OPEN ACCESS** 012034
Antioxidant properties from seaweeds *Kappaphycus alvarezii*, *Euchema spinosum* and *Sargasum* sp. using different solvent
A A Abdillah, M A Alamsjah and N E Nasution Sugijanto
[+](#) Open abstract [View article](#) [PDF](#)
-
- OPEN ACCESS** 012035
Effect of kappa-carrageenan on physicochemical properties of mantou (Chinese steamed bread)
M H C Putra and A A Abdillah
[+](#) Open abstract [View article](#) [PDF](#)
-
- OPEN ACCESS** 012036
The application of betacyanin microcapsules as natural food colorant on beverage model
S R Nurbaya, W D R Putri, E S Murtini and A Khamidah
[+](#) Open abstract [View article](#) [PDF](#)
-
- OPEN ACCESS** 012037
Microbial quality and diversity of *Caesio cunning* and *Scolopsis taenioptera* harvested by using trap and trawl fishing techniques
M Suhaimi and N U Karim
[+](#) Open abstract [View article](#) [PDF](#)
-
- OPEN ACCESS** 012038
See our Privacy Policy and Cookies policy. 

Characteristics physicochemical of melanin from squid ink (*loligo sp.*) extracted by ethanol

F Abidin, L Sulmartiwi and E Saputra

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012039

Analysis of fluctuating asymmetry of black strain tilapia *oreochromis niloticus* and red strain tilapia *Oreochromis niloticus* in Kabat Fish Hatchery Center Banyuwangi, East Java

D Kurniawan and A H Fasya

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012040

Utilization of fermented Seligi leaf flour *Phyllanthus buxifolius* toward the specific growth rate, daily growth rate and survival rate of siam catfish (*Pangasius pangasius*)

Y G Budiman, M Lamid, B S Rahardja and M Amin

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012041

Improving crude protein and crude fat content of Seligi leaf (*Phyllanthus buxifolius*) flour through probiotic fermentation

A K Nisa, M Lamid, W P Lokapirnasari and M Amin

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012042

Addition of turmeric in feed on growth and survival rate of Nilasa red tilapia (*Oreochromis sp.*)

R Cahyani, W H Satyantini, D D Nindarwi and Y Cahyoko

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012043

Characterization of semi-refined kappa-carrageenan from *Kappaphycus alvarezii* with different solvents in Tanjung Sumenep

H M Noor, M A Alamsjah and S Andriyono

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012044

The substitution effect of bone fish flour milkfish (*Chanos chanos*) physical and chemical characteristics of cookies

I Muzaki, H Suprpto and R Kusdarwati

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012045

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



Identification and the prevalence of fungal gouramy (*Osphronemus gouramy*) in modern market Surabaya

M S Andreas, R Kusdarwati and H Suprpto

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012046

Substitution of commercial feed with maggot meal (*Hermetia illucens*) to the growth rate, feed conversion ratio and feed efficiency catfish (*Pangasius pangasius*)

M Ulumiah, M Lamid and K T Pursetyo

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012047

Correlation between water quality with prevalence of black tiger shrimp (*Penaeus monodon*) infested by ectoparasite protozoa in traditional ponds of Wonorejo, Surabaya

M A Luthfi, S Subekti and G Mahasri

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012048

Study of nitrogen (N) and phosphorus (P) in the land of mangrove sediments in ecotourism area Wonorejo Surabaya and coastal area of Jenu Tuban

N Pradipta, M A Alamsjah and E D Masithah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012049

Corellation of water quality to the prevalence of ectoparasite in milkfish (*Chanos chanos*) in Sedati District, Sidoarjo

H Irawan, Kismiyati and G Mahasri

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012050

Effect of soaking time of blood cockle (*Anadara* sp.) shells powder with hydrochloric acid on the characteristics of nano calcium

D S Herlina, L Sulmartiwi and E D Masithah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012051

Addition of crude fish oil (CFO) in feed toward fat and energy retention of mud crab (*Scylla serrata*)

S Hadijah, Agustono and W P Lokapirnasari

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.

012052

Effectiveness of giving clove oil as an anaesthetic for survival rate and number of leucocytes in cantang grouper (*Epinephelus* sp.) in the closed transportation

S Nurkomaria, H Suprpto and Sudarno

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012053

Effect of the addition of crude fish oil (CFO) in feed to the content of EPA and DHA in mud crab (*Scylla serrata*)

T Wijaya, Agustono and M A A Arif

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012054

The effectiveness combination of maggot (*Hermetia illucens*) flour with commercial feed on growth rate, feed conversion ratio, and feed efficiency of tilapia (*Oreochromis niloticus*)

V Indriawati, B S Rahardja and Prayogo

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012055

Depuration of heavy metals Pb and Cd content in blood cockles (*Anadara antiquata*) with different filters

W Arifin, B S Rahardja and K T Pursetyo

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012056

Substitution of commercial feed with fermented banana peel flour (*Musaceaea* sp.) and fish meal to feed consumption level, specific growth rate, feed efficiency, fat retention, and energy retention in siam catfish (*Pangasius hypophthalmus*)

A Aisyah, A S Gustiningrum, Agustono and M A Al-Arif

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012057

The effect *Moringa oleifera* leaf extract and *Lactobacillus acidophilus* supplementation on crude protein and crude fat retention in Tambaqui, *Colossoma macropomum*

R A Pitri, E Suyanti, S Septayani, Agustono and W P Lokapirnasari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012058

Molecular identification and prevalence of ectoparasite worms in barramundi (*Lates calcarifer*) in Lampung Waters

I Achmadi, S Subekti, H B Ardiyanti, M K Amiin, L N F Haryanto, R E K Akbar and M A Yudarana

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



OPEN ACCESS

012059

Morphological profile of L2 *Anisakis typica* on Indian Mackerel (*Rastrelliger kanagurta*) from Sedati Fish Auction, Sidoarjo-East Java, Indonesia using Scanning Electron Microscope (SEM)

N Suryani, S Subekti, S Koesdarto and M K Amiin

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012060

Occurance of Anisakis of mackarel tuna (*Euthynnus affinis*) from Sendangbiru fishing auction place, East Java, Indonesia

R Bobsaid, P D W Sari and S Subekti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012061

The study of virus collation with the polymerase chain reaction (PCR) method in export fishery commodities

L Kurniawati and K T Pursetyo

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012062

Effect of Culture Combination on Growth and Carrageenan Content of *Kappaphycus alvarezii* Green and Red Variety

A A Abdillah and J Triastuti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012063

Nitrate and phosphate dynamics of phytoplankton abundance in Kanceng River, Sepuluh, Bangkalan, East Java, Indonesia

D D Nindarwi, S H Samara and M B Santanumurti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012064

Assessment of Seasonal Waters Quality Based on Abundance, Diversity, and Domination of Phytoplankton in Bajulmati Reservoir

E W Pertiwi, E D Masithah and Suciyo

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012065

Effect of sublethal dose organophosphate pesticides on embryogenesis and hatching rate of silver rasbora eggs (*Rasbora argyrotaenia*)

A B Prastika, L Sulmartiwi and L Lutfiyah

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.

[+ Open abstract](#) [View article](#) [PDF](#)



OPEN ACCESS

012066

The correlation between temperature and intensity of *Zeylanicobdella arugamensis* on cantang grouper (*E. fuscoguttatus* × *E. lanceolatus*) from traditional ponds in the Kampung Kerapu Lamongan East Java Indonesia

D D Afifah, G Mahasri and W H Satyantini

[+](#) Open abstract [View article](#) [PDF](#)

OPEN ACCESS

012067

Utilization of *Nitrosomonas* sp and *Nitrobacter* sp probiotic towards total suspended solid and ammonia level in Nile tilapia culturing using aquaponic system

K H Dwiardani, Prayogo and B S Rahardja

[+](#) Open abstract [View article](#) [PDF](#)

OPEN ACCESS

012068

The administration of *Caulerpa racemosa* extract on total bacteria and survival rates of white shrimp (*Litopenaeus vannamei*) after infected by *Vibrio parahaemolyticus*

A F Pratiwi, W H Satyantini, G Mahasri, L Sulmartiwi, A T Mukti and Sudarno

[+](#) Open abstract [View article](#) [PDF](#)

OPEN ACCESS

012069

Examination of Taura Syndrome Virus (TSV) in white shrimp (*Litopenaeus vannamei*) and tiger prawn (*Penaeus monodon*) with Polymerase Chain Reaction (PCR) method

A N Fadilah and A H Fasya

[+](#) Open abstract [View article](#) [PDF](#)

OPEN ACCESS

012070

Potential utilization of kombucha as a feed supplement in diets on growth performance and feed efficiency of catfish (*Clarias* sp.)

H U Ramadhan, Prayogo, H Kenconoajati, B S Rahardja, M H Azhar and D S Budi

[+](#) Open abstract [View article](#) [PDF](#)

OPEN ACCESS

012071

Application of edible film from chitosan as biodegradable packaging

E Saputra, W Tjahjaningsih and A A Abdillah

[+](#) Open abstract [View article](#) [PDF](#)

OPEN ACCESS

012072

Substitution of commercial feed with fermented banana peel flour (*Musaceaea* sp.) and fish meal to crude protein, energy, crude lipid and organic matter of meat in siamese catfish (*Pangasius hypophthalmus*)

R Z Darmawan, S M Ghaisani, Agustono and M A Al-Arif

[+](#) Open abstract [View article](#) [PDF](#)
This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



OPEN ACCESS

012073

The effect of garlic (*Allium sativum*) and turmeric (*Curcuma longa*) extract addition in commercial feed on feeding rate, feed efficiency and feed conversion ratio of gouramy fish (*Osphronemus gouramy*)

D Afifah, M Arief and M A Al-Arif

[+](#) [Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012074

The Variability in Population Structure of Gastropods in Sedati Waters, Sidoarjo Regency, East Java

S Fadliyah, L A Sari, K T Pursetyo, A Zein, M H Idris and Y Cahyoko

[+](#) [Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012075

The effect of pH and incubation time on crude protease enzymes activity of *Bacillus mycoides* from anchovy isolates (*Stolephorus* sp.)

D Nirmala, P Yudha and D Cahyanto

[+](#) [Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012076

The influence of garlic (*Allium sativum*) and turmeric (*Curcuma longa*) extract as attractant in commercial feed on feeding rate and fat retention in gouramy (*Osphronemus gouramy*)

G Y Pamungkas, M Arief and M A A Arif

[+](#) [Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012077

Beardless barb *Cyclocheilichthys apogon* (Valenciennes, 1842) (*Cypriniformes*, *Cyprinidae*): Distribution extension and first record from South Bali

V Hasan, A Wijayanti, M B Tamam, R A Islamy and M S Widodo

[+](#) [Open abstract](#) [View article](#) [PDF](#)

JOURNAL LINKS

[Journal home](#)

[Journal scope](#)

[Information for organizers](#)

[Information for authors](#)

[Contact us](#)

[Reprint services from Curran Associates](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



PAPER • OPEN ACCESS

Study of crude extract yield and gonad maturity level of sea cucumber *Phylloporus dobsoni* correlation

To cite this article: S Andriyono *et al* 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **679** 012020

View the [article online](#) for updates and enhancements.

You may also like

- [Gonad maturity level and somatic index of sea cucumber *Acaudina* sp. \(caudinidae, molpadida, holothuroidea\) In Delta Wulan Waters, Demak Regency, Indonesia](#)
W Widianingsih, R Hartati, M T Sibero et al.
- [Maturity Gonad Sea Cucumber *Holothuria scabra* Under The Month Cycle](#)
Santi Penina Tua Rahantoknam
- [Diversity and abundance of sea cucumber \(Holothuroidea\) in seagrass ecosystem at Suli Village, Maluku, Indonesia](#)
G D Manuputty, M M Pattinasarany, G V Limmon et al.



244th Electrochemical Society Meeting

October 8 – 12, 2023 • Gothenburg, Sweden

50 symposia in electrochemistry & solid state science

Abstract submission deadline:
April 7, 2023

Read the call for papers &
submit your abstract!

Study of crude extract yield and gonad maturity level of sea cucumber *Phylloporus dobsoni* correlation

S Andriyono^{1,3}, E D Masithah¹, and D Winarni²

¹Department of Marine, Faculty of Fisheries and Marine, Universitas Airlangga, Kampus C
Jalan Mulyorejo, Surabaya 60115 East Java, Indonesia

²Department of Biology, Faculty of Science and Technology, Universitas Airlangga, Kampus
C Jalan Mulyorejo, Surabaya 60115 East Java, Indonesia

³Corresponding author: sapto.andriyono@fpk.unair.ac.id

Abstract. The study of the yield of n-hexane extract on the body wall of *Phylloporus dobsoni* sp. Is the first research conducted in Indonesia. This study aims to measure the n-hexane extract of the body wall of sea cucumber *P. dobsoni* which is related to the maturity level of the gonads. The measurement results obtained that the average wet weight and dry weight of sea cucumber body wall was 23.58 grams of body wall wet weight, body wall dry weight 5.56, gonad weight 4.02, and GSI value 18.03. Judging from the histology of male sea cucumber gonads, it is known that they have grown and grown at an advanced stage with an average content of n-hexane extract at an average value of 0.202 grams and 0.266 grams, respectively. Whereas in the female group there are all stages of growth, continued growth, maturity, recovery and post-spawning. The content of n-hexane extract showed an increase in the range of 0.085; 0.2506; 0.278; 0.354 and 0.3045. The increase in extract content in male and female sea cucumbers is possible as a protective mechanism from predation and protection in preserving offspring.

1. Introduction

Sea cucumber has habitat widespread throughout the world, ranging from tidal zone to the deep sea. About 10% of the number of sea cucumbers species in the world inhabit in Indonesian waters [1]. There more than 50 species commonly traded [2], and about 27 species of sea cucumbers identified as commercially potential derived from Indonesian waters [3]. In the international trade of sea cucumbers, 80% goes to Hong Kong, Indonesia and the Philippines leading supplier. Hong Kong exports of these products to China, Singapore and Taiwan [4]. In specific markets, found in raw form, usually used as a Chinese cuisine which is believed to have specific properties. In Surabaya and the surrounding areas, sea cucumbers are found in the form of ordinary snack, which is marketed in the form of fried sea cucumber [5].

According to the FAO global statistics on sea cucumber, Indonesia is the largest sea cucumber exporter in the world [6]. Recorded foreign exports of 2.100 tons of reef invertebrates out of which 320 tons were sea cucumber in 1981. The essential collecting and export centre for marine product trade-in Indonesia are Ujung Pandang in Sulawesi [7]. Researchers showed that the rate of Indonesian exports of dried sea cucumbers was massive, was not offset by the many useful products made of sea cucumbers from Indonesia. Food supplement products made from sea cucumbers mentioned have the benefits for health and healing [8], are imported, and no one has even mentioned that the material was essentially a sea cucumber in Indonesian waters.

Although the region is now the largest producer of sea cucumbers are the beaches in eastern Indonesia [1]. Similarly, this phenomenon occurred in other places in Indonesia [9]. It is not possible over the next ten years that the situation in eastern Indonesia suffers the same fate as the east coast of Surabaya if the rate of exploitation is not balanced with the preservation and cultivation efforts [6].

Sea cucumber species collected in the study are not listed in the list of commercially valuable sea cucumbers in the world market. The distribution of those species are widespread, only one species with high abundance and distribution level, while others arranged from low to moderate. The preferred habitat is generally sandy. So far, *P. dobsoni* and *P. australis* are commonly consumed. Both are used for snacks or soup mix. The sea cucumbers are found, the type *P. dobsoni* has the highest abundance values comparing the other in the amount of 44.4%, and a high distribution rate is the distribution index of 1.9062 [13].

Based on a variety of the above information, it shows that the concern the development of species of *P. dobsoni* is still limited to the exploitation that is not sustainable. Based on morphological and molecular identification, this sea cucumber is *P. dobsoni* [10]. The attention that we paid in this study was an effort to develop cultivation level expected to be a developed laboratory and mass [5,11]. This study also constitutes active ingredients at the same owned *P. dobsoni* as a promising commodity in the future as pharmacological ingredients.

2. Material and methods

The sea cucumbers were purchased from local fisherman. They were collected from the east coastal water of Surabaya. The species of sea cucumber is *P. dobsoni*. The sea cucumbers are performed surgery has not performed the selection, and the entire sample was 50 sea cucumbers. Each measurement of sea cucumbers wet weight, dry weight and the amount of the resulting extract. Besides, gonad weight and histology of gonads on each sea cucumber were done separately. Wet weight measurements were done using analytical digital scales. The body wall of sea cucumbers was weighing and cutting with a small size in order to facilitate the freeze-drying. The freeze-drying process is done for 24 hours at the Tropical Disease Center, Universitas Airlangga.

2.1. Gonadal histology

Gonadal histology performed by means of fixing the gonads in a liquid fixative neutral buffered formalin solution for ≥ 24 hours [12]. Furthermore, gonad histology has taken five of each gonad tubules for further processing from the processing stage, the stage of infiltration and planting (embedding), stage of cutting (sectioning) and attachment (affixing), and the last stage of staining (staining). Subsequent gonadal histology results were observed under a microscope with 100x magnification and documented.

2.2. Samples extraction

Extraction of sea cucumbers *P. dobsoni* done by maceration method that starts with the preparation of sea cucumbers material *P. dobsoni* obtained from the waters Kenjeran Beach, Surabaya. Sea cucumbers *P. dobsoni* discarded entrails and washed with running water. Samples that have been cleaned, put in a blender and then blended. The next process is freeze-drying so that the water content in the sample decreases but does not damage the existing active ingredients. Freeze-drying carried out at -80°C . Maceration process is done using hexane solvent [13]. Soaking the sample and the solvent in the ratio 1:3 (w/v) or submerged entirely. Soaking was done twice every 72 hours. Results filtrate formed filtering [14], then evaporated using a rotary evaporator at a temperature of 38°C to form a viscous extract and have no smell of solvent [15]. The resulting extract yield calculation is done by calculating the percentage ratio of the resulting extract and extract raw materials, formulated in the following equation [14].

$$\text{Extract Yield (\%)} = (\text{weight extract} / \text{weight samples}) \times 100\%$$

2.3. Analysis of relationship gonads and hexane extract

Statistical analysis of the relationship and the yields of extract content based on gonad maturity level in each group of male and female sea cucumber do regression and graphs by Microsoft Excel [16]. From the result of the graph, the pattern extract content level can be obtained at each stage of gonad maturity both on sea cucumber: male and the female. The histology results are performed descriptively.

3. Result and discussion

The value of the yield of the crude extract (n-hexane) is about 0.242 ± 0.233 grams or 4.35% of the dry weight obtained can be recognized. The dry weight of sea cucumber body wall by an average of 5.56 ± 3.33 grams or about 23.57% of the average wet weight of 23.58 ± 6.49 grams of the body wall (Table 1).

Table 1. Data of average weight wet, dry weight and the gonads weight of sea cucumber *P. dobsoni* and weight of n-hexane extract

Parameters	Value \pm SD
Weight Wet (grams)	23.580 ± 6.490
Dry Weight (grams)	5.560 ± 3.330
Gonads Weight (grams)	4.020 ± 1.250
GSI	18.030 ± 7.870
n-Hexane Extract Weight (grams)	0.242 ± 0.233

Total results yield n-Hexane extract on sea cucumbers *P. dobsoni* This produced approximately 4.35% of the average weight of 5.56 grams of the dried body wall that produce an average weight of 0.242 ± 0.233 gramsof extract. The dry weight of sea cucumber body wall by an average of 5.56 grams or about 23.57% of the average wet weight of 23.58 grams of the body wall. These conditions do not reflect the value of the content of other secondary metabolites that have not been tested in detail. This phenomenon is in contrast with the results of the yield on the previous study, which get *Stichopus variegatus* sea cucumber extract with methanol solvent, which is only 2.78% [14]. This results confirmed that the sea cucumber extract produced is very low, so if this is done will increase the value of selling sea cucumbers that can be used as efforts to increase value-added products of sea cucumbers.

At gonads maturity level (GML), the analysis on 43 samples of collected sea cucumbers known those 27 males, 13 females, and 3 samples are hermaphrodite. Seven samples are broken and unable to histological analysis. In the group of male sea cucumbers, the largest proportion of GML was at 66.67% growth stage, the advanced growth was 29.63%, and 3.7% was in the recovery phase (Figure 1).




Figure 1. Histology of the gonads of sea cucumbers *P. dobsoni* males in the growth phase (A) and advanced growth (B). (dt: tubular wall; sz: spermatozoa; sg: spermatogonia) scale bar: $100\mu\text{m}$ with a magnification of 100 times.

In the group of female sea cucumbers, the highest proportion in the advanced stage of growth was 38.46%, 30.76% of it was in the mature stage, in the growth stage was 15.38% and the same percentage was in post-spawning stage and recovery of value which was 7.69% (Figure 4B). On histological analysis of the gonads, it was also found that there are three tails of sea cucumbers have two gonads (hermaphrodite), i.e. at the stage of growth, and advanced growth and recovery. The condition is known by the presence of spermatozoa (male marker) present in the gonads as well together as marker hermaphrodite oocytes (Figure 2).




Figure 2. Histology of the gonads of sea cucumbers *P. dobsoni* females in the gonad development stage of growth (A), advanced growth (B), mature (C), post-spawning (D), and recovery (E). (dr: tubular wall; l: lumen; OPV: previtellogenic oocytes; ov: vitellogenin oocytes; jl: jelly layer) scale bar: 100 μ m and a magnification of 100 times.




Figure 3. Gonadal histology *P. dobsoni* females who are hermaphroditic organisms on gonadal development stages of growth (A), advanced growth (B), and recovery (C) (dr: tubular wall; OPV: previtellogenesis oocytes; ov: vitellogenesis oocytes; sz: spermatozoa, l: lumen; and jl: jelly layer)) scale bar: 100 μ m and a magnification of 100 times.

In the male gonads, histology results showed only at an advanced stage of growth and growth with a mean of n-hexane extract produced by 0.202 ± 0.1324 grams and 0.266 ± 0.3505 grams (Figure 4A). Values of n-hexane extract number increased 0.064 grams at earlier stages of gonad maturity. The biological activity of sea cucumbers *P. dobsoni* in the metabolism may produce results which are also focused on the improvement of future gonads to be generated.




Figure 4. Graph content of n-Hexane extract the body wall *P. dobsoni* in the male group (A) and female (B) at different levels of gonadal maturity.

In the male group of sea cucumbers, found two stadia or the stage of maturity gonad growth and advanced growth. The second level of maturity gonad showed an increased amount of n-hexane extracts are produced. This condition proves that in the formation of the reproductive organs, sea cucumbers do protection mechanisms in the gonads produced in order to grow until the spawning stage. The secondary metabolites and organic acids in the body, internal organs and the gonads may play a role in self-defence. The active ingredients are also derived in the gonads and eggs, which allows preservation generation *Ascidians* group that studies [17]. This finding reinforces that sea cucumbers *P. dobsoni* is a benthic animal that enables very high-level predation because of its impaired mobility and has a soft body. This defence mechanism can be studied further in the content of secondary metabolites in the gonads that were not tested in the study this year.

Sea cucumber *P. dobsoni* females group are found throughout the stages of gonadal development of start growth, advanced growth, mature, post-spawning and recovery with the trend of increasing values of n-Hexane extract produced (Figure 4B). It also shows that the increase in future stages of gonad maturity of gonads will result in higher production of metabolites. This mechanism, both in males and females, occurs as a defence mechanism against other predators.




Figure 5. Regression chart between wet weight and dry weight of sea cucumbers

The graph above is the relationship between wet weight and dry weight of the body wall of sea cucumbers. The graph shows that the dry weight and wet weight has had a very close correlation between these parameters (Figure 5). This result shows that the weight of the sample in wet conditions, then dried samples obtained will also be more and more. The condition also occurs in relation to extracting obtained from the dry weight of the sample. The more dry samples are extracted, the more extracts that may be obtained. This finding can be seen in the graph below (Figure 6) with a correlation value of $R^2 = 0.909$.




Figure 6. Regression graph dry weight of sea cucumbers and n-hexane extracts yield.




Figure 7. Regression graph of gonads maturity stadia and yields extract in males (A) and female (B)

Regression graph above illustrates that the content of the extract is not related to the maturity level of gonads, either male or female in the sea cucumber (Figure 7). Although the extract obtained is useful metabolites in protection, but does not affect the maturity of gonads of sea cucumbers. Internal factors of sea cucumbers do not only influence the maturity of gonads of sea cucumbers, but external factors are also influential. Many environmental factors are thought to affect the reproductive aspects of sea cucumbers, including temperature, blooming phytoplankton, and the cycles of the moon [18].

4. Conclusion

The number of natural products produced from the body wall extract *P. dobsoni* with n-Hexane solvent is an average of 0.242 ± 0.233 grams or 4.35% of the dry weight obtained. The dry weight of sea cucumber body wall by an average of 5.56 grams or about 23.57% of the average wet weight of 23.58 grams of the body wall. Thus, the content of n-Hexane extract yield in the body wall *P. dobsoni* can be estimated by referring to this study. The correlation between wet weight and dry weight indicates a very close value. This result means that the weight of the body wall of sea cucumbers, the more will allow the amount of dry weight was obtained. The amount of the extract obtained also has a linear pattern on the dry weight. The regression equation concerning the dry weight of the body wall and extract the resulting number is $Y = 0,099X + 0.184$, with the value of x is the dry weight of the body wall of sea cucumbers and y as the number of n-hexane extracts are produced.

5. References

- [1] Samad M 2000 *JSTI* **3**(3), 52-53 [in Indonesian].
- [2] Purcell S W, Lovatelli A, Vasconcellos M, and Ye Y 2010 *Managing Sea Cucumber Fisheries with An Ecosystem Approach* (Rome: FAO Fisheries And Aquaculture Technical Paper No. 520).
- [3] Setyastuti A and Purwati P 2015 *Beche-De-Mer Information Bulletin* **35**, 19-25.
- [4] Conand C, Shea S and To A 2014 *PLoS ONE* **9**(4), e95075.

- [5] Andriyono S, Masithah E D, Rumiayati B, Triastuti J, and Winarni D 2016 *AJAS* **4**(4).
- [6] Tuwo A 2004 Advances in Sea Cucumber Aquaculture and Management: 49-56 *In* Lovatelli A (Ed) *Advances in Sea Cucumber Fisheries Aquaculture and Management* (Rome: Food and Agriculture Organization of The United Nations).
- [7] Choo P-S 2008 Population Status, Fisheries and Trade of Sea Cucumbers in Asia: 81-118. *In* Toral-Granda V, Lovatelli A and Vasconcellos M (Eds) *Sea cucumbers. A Global Review of Fisheries and Trade* (Rome: FAO Fisheries and Aquaculture Technical Paper. No. 516).
- [8] Farouk A E-A, Ghouse F A H, and Ridzwan B 2007 *Am J Biochem Biotechnol.* **3**(2), 60-65.
- [9] Darsono P 2003 *Oseana* **28**(2), 1-9.
- [10] Pratiwi A I 2018 *Studi Taksonomi Phyllophorus Dobsoni di Perairan Pantai Timur Surabaya Berdasarkan Karakter Morfologi dan Genetika Molekuler Skripsi* (Surabaya: Universitas Airlangga) [in Indonesian].
- [11] Andriyono S, Masruroh N, Masithah E D, Triastuti J, and Winarni D 2015 *J. Biosci.* **23**, 39-46.
- [12] Mumford S L 2004 *Histology of Finfish* (Olympia Washington: USFWS, Olympia Fish Health Center).
- [13] Winarni D, Affandi M, Masithah E D, and Kristanti A N 2009 *Eksplorasi Potensi Teripang Pantai Timur Surabaya Sebagai Modulator Imunitas Alami Terhadap Mycobacterium tuberculosis Laporan Penelitian* (Surabaya: Universitas Airlangga) [in Indonesian].
- [14] Rasyid A 2014 *OLDI* **40**(2), 49-55 [in Indonesian].
- [15] Nimah S, Ma'ruf W F, and Trianto A 2012 *JPBHP* **1**(1), 9-17 [in Indonesian].
- [16] Dretzke B 2008 *Statistics with Microsoft Excel* (New Jersey: Prentice Hall Press).
- [17] Pisut D P and Pawlik J R 2002 *J. Exp. Mar. Biol. Ecol.* **270**(2), 203-214.
- [18] Mackey A and Hentschel B 2001 *Factors that Influence The Reproduction of Sea Cucumbers* (San Diego: San Diego State University).

6. Acknowledgement

We want to deliver our gratitude to the Ministry of Education, through the Directorate General of Higher Education that provides Decentralization Grants of Higher Education Research DIPA BOPTN No.6386/UN3.6/KU/PPK/2014. We also thank Dr Dwi Winarni and students involved in the comprehensive studies.