

Table of contents

Volume 236

2019

◀ Previous issue Next issue ▶

The 1st International Conference on Fisheries and Marine Science 6 October 2018, East Java, Indonesia

Accepted papers received: 16 January 2019

Published online: 01 March 2019

Open all abstracts

Preface

OPEN ACCESS 011001

The 1st International Conference on Fisheries and Marine Science

+ Open abstract  View article  PDF

OPEN ACCESS 011002

Organizing Committee

+ Open abstract  View article  PDF

OPEN ACCESS 011003

Conference Photos

+ Open abstract  View article  PDF

OPEN ACCESS 011004

Peer review statement

+ Open abstract  View article  PDF

Papers

OPEN ACCESS 012001

The Increase in β -carotene Content in *Dunaliella salina* from the Application of Different Light Intensities

N Sugiati, E D Masithah, W Tjahjaningsih and A A Abdillah

+ Open abstract  View article  PDF

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more,

OPEN ACCESS and Cookies policy.

012002 

Ammonia-eliminating potential of *Gracilaria* sp. And zeolite: a preliminary study of the efficient ammonia eliminator in aquatic environment

M R Royan, M H Solim and M B Santanumurti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012003

Identification of extracellular enzyme-producing bacteria (proteolytic, cellulolytic, and amylolytic) in the sediment of extensive ponds in Tanggulangrejo, Gresik

OA Artha, Sudarno, H Pramono and LA Sari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012004

Addition of water from the treatment pond of pangasius fillet waste (*Pangasius* sp.) with different concentrations in the cultivation medium due to the population growth of *Daphnia* sp.

H P Alvian, E D Masithah and M H Azhar

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012005

The growth and survival rate in lettuce aquaponic systems (*Latuca sativa*) of eels in various stocking densities of eel (*Monopterus albus*)

N K Portalia, L Sulmartiwi and B S Rahardja

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012006

The prevalence of benedeniasis in humpback grouper (*Cromileptes altivelis*) in floating net cages in Situbondo Regency, East Java, Indonesia

S O Wijaya, S Subekti and Kismiyati

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012007

The prevalence of fungi on groupers (*Epinephelus* sp.) in cage mariculture systems of the northern coast of Surabaya, East Java

E Yuliasuti, R Kusdawarti and Sudarno

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012008

The spectrum of light and nutrients required to increase the production of phycocyanin *Spirulina platensis*

H A Wicaksono, W H Satyantini and E D Masithah

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more,

[see our Privacy and Cookies policy.](#)

012009

The growth and survival rate of the larvae of the sunu grouper (*Plectropomus leopardus*) in different temperatures

L Lutfiyah, D S Budi and M F Ulkhaq

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012010

Preservation of common carp (*Cyprinus carpio*) sperm using 0.9% NaCl and ringer's lactate solution

D S Budi, L A Adawiyah and L Lutfiyah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012011

Comparison of the efficiency (flash point, freezing point, and viscosity test) of biodiesels from *Sargassum* sp.

M B Santanumurti, M R Royan, S H Samara, S Sigit and M A Alamsjah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012012

Study of patterns in the relationship of ecdysis with the age of freshwater crayfish *Cherax quadricarinatus* aged 76 Days

A H Fasya

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012013

Maximum density in the *Moina macrocopa* culture able to produce parthenogenesis in female offspring

A S Mubarak, D Jusadi, M Z Junior and M A Suprayudi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012014

The oxygen content and dissolved oxygen consumption level of white shrimp *Litopenaeus vannamei* in the nanobubble cultivation system

D P Galang, A K Ashari, L Sulmatiwi, G Mahasri, Prayogo and LA Sari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012015

The effect of the epiphytes of *Chaetomorpha crassa* on the total chlorophyll-a and growth of *Gracilaria verrucosa*

A L L Handayani, R J Triastuti and L Sulmartiwi

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.

012016 

Growth monitoring of koi fish (*Cyprinus carpio*) in natural hatchery techniques in Umbulan, Pasuruan, East Java

F P Putri and N N Dewi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012017

Dynamic Ratio Correlation of N:P in relation to the Diatom Abundance in the Intensive System of the Vannamei (*Litopenaeus vannamei*) Shrimp Pond

E D Masithah, D D Nindarwi, T Rahma and dan R R Satrya P I

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012018

Dynamic ratio correlation of N:P on the abundance of Bluegreen algae in an intensive system in a white shrimp (*Litopenaeous vannamei*) pond

E D Masithah, D D Nindarwi, A L A Suyoso and D Husin

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012019

Dynamic ratio correlation of N:P toward phytoplankton explosions in intensive systems of white shrimp pond

E D Masithah, D D Nindarwi, D Husin and T Rahma

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012020

Development of water and nutrient management models to improve multitrophic seafarming productivity

J A Surbakti, I A L Dewi, M A Alamsjah and M Lamid

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012021

Pond soil characteristic in reclaimed tidal lowlands and its correlation with the water quality for aquaculture

M Fitriani, I Wudtisin, M Kaewnern and R H Susanto

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012022

The dynamics of total organic matter (tom) on sangkuriang catfish (*clarias gariepinus*) farming at upt ptpbp2kp and the effectiveness of freshwater bivalve (*anodonta woodiana*) in reducing the total organic matter with varying density

D Arfiati, C D G Putra, A H Tullah, S W A Permanasari and A W Puspitasari

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



OPEN ACCESS 012023

Growth and morphological changes in relation to the maturation of male Japanese eel, *Anguilla japonica* injected with human chorionic gonadotrophin (HCG) in the different interval in the tropical region

Y T Hee, F F Ching and S Senoo

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012024

Genetic diversity of the endangered species *Sphyrna lewini* (Griffith and Smith 1834) in Lombok based on mitochondrial DNA

S Hadi, N P Anggraini, E Muttaqin, B M Simeon, B Subhan and H Madduppa

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012025

Sex ratio and size at first maturity of razor clam *Solen* sp. in Pamekasan and Surabaya coastal area, East Java, Indonesia

N Trisyani, N I Wijaya and I Yuniar

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012026

Improving productivity and water quality of catfish, *Clarias* sp. cultured in an aquaponic ebb-tide system using different filtration

E Setiadi, I Taufik, Y R Widyastuti, I Ardi and D Puspaningsih

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012027

Different substrate of trickling filter on growth, survival rate, and water quality of common carp (*Cyprinus carpio*) cultivation by using an intensive recirculation system

E Setiadi, I Taufik, Y R Widyastuti, I Ardi and A Saputra

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012028

Water quality dynamic, production and profitability of catfish, *Clarias* sp. cultured at different design construction of aquaponic

Y R Widyastuti, E Setiadi, I Taufik and L Setijaningsih

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012029

Effect of C:N ratio on the spore production of *Bacillus* sp. indigenous shrimp pond

A Yuniarti, N B Arifin, M Fakhri and A M Hariati

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



OPEN ACCESS 012030

Comparative Test on Bacteria in the Digestive Tract of Vannamei Shrimp (*Litopenaeus vannamei*) at Intensive and Extensive Ponds in Ujungpangkah, Gresik

D Ningrum, M Arief and dan K T Pursetyo

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012031

Growth Performance of Laboratory-Scale *Chaetoceros calcitrans* in Different Containers

M Jannah, M F Ulkhaq, M H Azhar, Suciyo and dan W Soemarjati

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012032

Study of the Dynamic Density and Diversity of Plankton at Different Brackishwater Pond Managements in Gresik, East Java

O Tilahwatih, E D Masithah and dan B S Rahardja

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012033

The Effect of Demineralization Stage of Agar's Solid Waste on the Characterization of Activated Carbon

R Febrianto, Sudarno and R Kusdarwati

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012034

The Dynamic of Density and Diversity of Cyanophyta in Different Pond Bases in Educational Pond of Faculty of Fisheries and Marine Universitas Airlangga

S Z Cahyani, E D Masithah and Prayogo

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012035

Effect of Different Salinity Level within Water Against Growth Rate, Survival Rate (FCR) of Catfish (*Clarias* sp.)

D Prananingtyas, Prayogo and S Rahardja

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012036

Molecular identification and phylogenetic reconstruction of two fiddler crabs (*Uca forcipata* and *Uca triangularis*)

S Andriyono, H Pramono and H W Kim

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more,

[See our Privacy and Cookies policy.](#)

012037 

The molecular identification and phylogenetic reconstruction of Palaemonid and Penaeid shrimp from the southern part of Bangladesh

M J Alam, S Andriyono, A T M Eunos and H W Kim

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012038

Morphometric characteristics of Fur Cockles (*Anadara spp.*) in Wonokromo and Juanda Estuary, Surabaya

P B Pamungkas, K T Pursetyo, J Triastuti and N N Dewi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012039

Stock status of ark clams (*Anadara spp.*) based on dredge fishing of the east coast of Surabaya, Indonesia

N N Dewi, K T Pursetyo, O P Darmono, F R Fachri, F S Puspitasari and A Damora

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012040

The distribution patterns and biomass of bivalves in Segoro Tambak estuary, Sedati, Sidoarjo, East Java

S H Liyana, L A Sari, N N Dewi, E D Masithah, A M Sahidu and K T Pursetyo

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012041

Inventorization of reef fish on Tabuhan Island, Banyuwangi, East Java, Indonesia

Suciyono, M A Azhar, M F Ulkhaq and H Kenconoajati

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012042

Dynamic study on the effect of calcium hydroxide and sodium bicarbonate treatment on the N/P ratio and plankton abundance

M R N Tsany, E D Masithah, B S Rahardjo and D D Nindarwi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012043

Distribution patterns and the biomass of bivalves at Segoro Tambak estuary, Sedati, Sidoarjo, East Java

S H Liyana, L A Sari, N N Dewi, E D Masithah, A M Sahidu and K T Pursetyo

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012044

Optimization of diatom *Hyaloea ostrearia* cultivation in different mediums and nutrients

Seputri Pratiwi and Coomes pole



S Arsad, C Stavrakakis, V Turpin, P Rossa, Y Risjani, L A Sari, F S Prasetya and J-L Mouget

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012045

Coastal ecosystem model based on environmental suitability and carrying capacity of the fishpond in Banyuwangi Region, East Java, Indonesia

E W Setyaningrum, Maghdalena, A T K Dewi, M. Yuniartik and E D Masithah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012046

Development and succession of sessile macrofouling organisms on the artificial structure in the Shallow Coastal Waters of Sabah, Malaysia

M A M Affandy, J Madin, K P Jakobsen and M Auluck

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012047

Fish species difference around the light of metal halide lamps and LED lamps with mini purse seine operation

M A Sofijanto, D Arfiati, T D Lelono and A Muntaha

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012048

Status of coral diseases and compromised health syndromes on Pemuteran shallow reefs, North Bali island

W Karim

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012049

Management options for restoring artificial coral reefs in Indonesia: strengthening in institutional approach

Rudianto and Ahmad Zainul

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012050

Growth of salt-secretor and non-salt secretor mangrove seedlings with varying salinity and their relations to habitat zonation

M Basyuni, Ramayani, A Hayullah, Prayunita, M Hamka, L A Putri and S Baba

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012051

Distribution of *Ctenactis Echinata* and *Fungia Consinna* coral on Mamburit island,

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



Sawiya, D Arfiati, Guntur and U Zakiyah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012052

Diversity species and condition of seagrass ecosystem in Teluk Awur and Prawean Jepara

I Riniatsih, A Ambariyanto, E Yudiati, R Hartati, W Widianingsih and R T Mahendrajaya

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012053

Copper (Cu) and Cadmium (Cd) toxicity on growth, chlorophyll-a and carotenoid content of phytoplankton *Nitzschia* sp

D Hindarti and A.W. Larasati

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012054

Vegetation Characteristic and Micro Environment of Mangrove Rehabilitation Forest at Coastal Areas of East Sinjai, South Sulawesi

H Setiawan

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012055

The Use of Water Lettuce (*Pistia stratiotes*) as Phytoremediator for Concentration and Deposits of Heavy Metal Lead (Pb) Tilapia (*Oreochromis niloticus*) Gills

A A D Amalia, B S Rahardja and Rr J Triastuti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012056

The Effectiveness of Heavy Metals Pb, Cd and Zn Reduction in NPK Fertilizer Waste Combined with Biofilters of Seaweed (*Gracillaria* sp.), Blood Clam (*Anadara* sp.), and Zeolite

A R K Sari, R K Harryes, F A Anggraini, M A Alamsyah and dan A Ahadi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012057

The Effectiveness of Combining *Gracilaria* Sp. Seaweed Biofilter and *Anadara granosa* Shell with Zeolite in the Decrease in the Level of Mercury (Hg) Heavy Metal

J A Spesatri, B S Rahardja and A A Abdillah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012058

The Effectiveness of Combination of Seaweed (*Gracillaria* sp.), Blood Clam (*Anadara granosa*), and Zeolite as Biofilter in the Reduction of Heavy Metal Copper (Cu)

A A D Amalia, B S Rahardja and Rr J Triastuti

[+ Open abstract](#) [View article](#) [PDF](#)



[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012059

Bioaccumulation of Cadmium (Cd) Heavy Metal on Seaweed (*Gracilaria* sp.) in Traditional Fishpond of Jabon Subdistrict, Sidoarjo District

O Ardiyansyah, Sudarno and Rosmanida

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012060

Identification of Proteolytic Bacterial Isolates in Sediment Ecosystem of Gunung Anyar Mangrove Forest, Surabaya

P B Utomo, Sudarno and B S Rahardja

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012061

The Analysis of Cockle (*Anadara inaequalis*) Gonad Maturity Level in the Estuary of Banjar Kemuning River, Sedati, Sidoarjo

R F Saputra, E D Masithah and P D Wulansari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012062

Spatial and Temporal Variation of Biomass Blood Cockle (*Anadara* sp.) in Estuaries Dadapan, Sedati Sub-District, Sidoarjo, East Java

M R Ramadhan, K T Pursetyo, Prayogo and N N Dewi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012063

Distribution Patterns and Biomass of Bivalve in Juanda and Segoro Tambak Estuary in Sedati, Sidoarjo, East Java

S H Liyana, E D Masithah and A M Sahidu

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012064

Analysis of Cadmium (Cd) Heavy Metal on Sediment and Mangrove Leaves *Avicennia marina* at Mangrove Ecotourism Wonorejo, Surabaya

W C Dermawan, Prayogo and B S Rahardja

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012065

Analysis of Lead (Pb) Value Comparison on Seaweed (*Euचेuma cottonii*) in Bluto and Saronggi Sumenep Marine, Madura, East Java

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy](#) and [Cookies](#) policy.



[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012066

The potential addition of lemuru oil to commercial feed to increase the content of EPA and DHA in eels (*Monopterus albus*)

A Imanisa, M B Santanumurti, M Lamid and Agustono

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012067

Effect of lysine in addition to commercial feed on crude protein and the energy digestibility of gourami (*Osphronemus gouramy*)

D Setiyawan, S H Samara, Agustono and M A A Arif

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012068

Addition of the papain enzyme to commercial feed against protein retention and feed efficiency in eels (*Anguilla bicolor*)

DA Liono, M Arief, Prayogo and W Isoni

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012069

Combination of papain enzyme and phytase enzyme in commercial feed and the protein and energy retention of tilapia *Oreochromis niloticus*

A Saifulloh, M B Santanumurti, M Lamid and W P Lokapirnasari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012070

The effect of giving cake artificial feed on the survival rate, and growth of Common carp (*Cyprinus carpio*) larva in an Installation of Freshwater Culture (IBAT) in Punten, Batu.

I P Zainiyah, Rozi, W H Satyantini and A M Sahidu

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012071

The Utilization of Phytase Enzymes and SEM Analysis in order to increase the Quality of Rice Bran as a Layer and Fish Feed

M Lamid, Anam Al-Arif and S H Warsito

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012072

The dynamic relationship of phytoplankton abundance and diversity in relation to white shrimp (*Litopenaeus vannamei*) feed consumption in intensive ponds

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012073

Effect of partial replacement of fish meal with *Spirulina platensis* meal in practical diets and culture location on growth, survival, and color enhancement of percula clownfish *Amphiprion percula*

S Hudaidah, B Putri, S H Samara and Y T Adiputra

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012074

The Effect of Different level of Probiotic Addition on Commercial Feed against Digestibility and Efficiency of Nile Tilapia Feed (*Oreochromis Niloticus*)

D Taufik, M Arief and H Kenconoati

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012075

The Effect of Adding Synbiotics Into Commercial Feed Towards Protein Retention and Fat Retention of Dumbo Catfish (*Clarias sp.*)

H Syevidiana, M Arief and I S Hamid

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012076

The Effect of Adding Lysine in Commercial Feed on Growth Rate, Feed Efficiency, and Feed Conversion Ratio to Tambaqui (*Colossoma Macropomum*)

L V D Putra, U Agustono and S H Kenconoati

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012077

The Effect of Coconut Shell Liquid Smoke in Commercial Feed on Total Bacteria of *Pseudomonas Aeruginosa* in the Tilapia's Kidney (*Oreochromis niloticus*)

M Rahmawati, Sudarno and S Subekti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012078

The Effect of Coconut Shell Liquid Smoke in Commercial Feed Towards Total *Pseudomonasaeruginosa* Bacteria on Gastrointestinal Tract Tilapia (*Oreochromis Niloticus*)

S Rahmadini, Sudarno and S Subekti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012079

Antibacterial activity of honey in preserving high-pressure cooked milkfish stored at room temperature

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



D A Hakim, W Tjahjaningsih and Sudarno

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012080

Bacterial composition in the gastrointestinal tract of *Uca* spp crabs fed on *Avicennia marina* leaf litter

M A B Kareho, E D Masithah and W Tjahjaningsih

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012081

The correlation between ectoparasite infestation and the total plate count of *Vibrio* sp. in pacific white shrimp (*Litopenaeus vannamei*) in ponds

G Mahasri, Rozi, A T Mukti, W H Satyantini and N M Usuman

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012082

In vitro study of an ethanolic extract of coffea leaves to inhibit freshwater pathogenic bacteria

H Kenconoajati, MF Ulkhaq, DS Budi and MH Azhar

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012083

The effect of noni fruits (*Morinda citrifolia*) with different ripeness stages against the total erythrocytes and leukocytes of comet goldfish (*Carassius auratus*) infested by *Argulus*

E Setyaningsih, Kismiyati and S Subekti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012084

The protection capacity of the crude and whole protein spores of *Myxobolus koi* as an immunostimulant material development in goldfish (*Cyprinus carpio*) for preventing Myxobolus

G Mahasri, M Yusuf, R Woro and M B Santanumurti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012085

Identification of white spot syndrome virus (WSSV) in pacific white shrimps (*Litopenaeus vannamei*) from ponds postexposure to immunogenic membrane proteins (*Zoothamnium penaei*)

P A Wiradana, G Mahasri, R E R Sari, U C Marwiyah and R Prihadhana

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012086

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



Gills and swimming leg histopathologies in pacific white shrimp (*Litopenaeus vannamei*) from ponds exposed to the immunogenic membrane proteins of *Zoothamnium penaei*

R E R Sari, G Mahasri, P A Wiradana and U C Marwiyah

[+](#) [Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012087

Total plate count and identification of vibrio in pacific white shrimp (*Litopenaeus vannamei*) from ponds and in those exposed to immunogenic protein membrane *Zoothamnium penaei*

U C Marwiyah, G Mahasri, R E Ratnasari and P A Wiradana

[+](#) [Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012088

Effectiveness of *Nitrobacter* on the specific growth rate, survival rate and feed conversion ratio of dumbo catfish *Clarias* sp. with density differences in the aquaponic system

A S Taragusti, M B Santanumurti, B S Rahardja and Prayogo

[+](#) [Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012089

The erythrocyte and leucocyte profile of saline tilapia (*Oreochromis Niloticus*) in a cultivation system with nanobubbles

M Gunanti, P D Wulansari and K Kinzella

[+](#) [Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012090

The prevalence and intensity of ectoparasites infecting vanname shrimp (*Litopenaeus vannamei*) reared in different ponds

N Nurlatiffah, Kismiyati and MF Ulkhaq

[+](#) [Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012091

Protozoan parasites of Vannamei Shrimp (*Litopenaeus vannamei*) in farmed fish from Pasuruan, Indonesia

U Hafidloh and P D W Sari

[+](#) [Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012092

Growth of *Bacillus* sp. and *Flavobacterium* sp. in culture media with the addition of liquid whey tofu waste

W H Satyantini, R M Pratiwi, A M Sahidu and D D Nindarwi

[+](#) [Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



OPEN ACCESS

012093

The effect of noni *Morinda citrifolia* L. fruit extracts on the gill histopathological changes of Nile tilapia *Oreochromis niloticus*

A T Mukti, E Dewi, W H Satyantini, L Sulmartiwi, Sudarno and M Hassan

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012094

Prevalence and intensity of ectoparasites in Pacific white shrimp (*Litopenaeus vannamei*) seeds from a pond and hatchery

G Mahasri, T Hidayat and Sudarno

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012095

Prevalence and intensity of ectoparasites in gabus fish (*Channa striata*) at Cangkringan Fishery Cultivation Technology Development Center, Sleman, Yogyakarta

E N Fitriani, Rozi, M Arief and H Suprpto

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012096

Antibacterial activity of bitter melon (*Momordica charantia* L.) leaf extract against *Aeromonas hydrophila*

D A Masithoh, R Kusdarwati and D Handijatno

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012097

Determination of the *aerolysin* gene in *Aeromonas hydrophila* using the polymerase chain reaction (pcr) technique

G Christy, R Kusdawarti and D Handijatno

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012098

Bioaccumulation and histopathological effect on the gills and liver of silver barb (*Barbonymus gonionotus*) exposed to the heavy metal nickel

I Purwanti, W Arroisi, B S Rahardja and L Sulmartiwi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012099

Identification and prevalence of the ectoparasite *Octolasmis* in sand lobster (*Panulirus homarus*) and bamboo lobster (*Panulirus versicolor*) in Floating Net Cages in Sape, Bima Regency, West Nusa Tenggara Province, Indonesia

L. Yusgita, Kismiyati, S. Subekti, P D Wulansari and M K Amiin

[+ Open abstract](#) [View article](#) [PDF](#)
This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



OPEN ACCESS

012100

Treatment of *Chlorella sp.* extract on heat shock cluster (HSC) response from the tissue and bloodcells proliferation of *Epinephelus fuscoguttatus-lanceolatus* infected by *Viral Nervous Necrosis*

U Yanuhar, I Al-Hamidy and N R Caesar

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012101

Superoxide dismutase (SOD) and metallothionein (MT) *Tubifex tubifex* at the acute mercury exposure

I M Widiastuti, A M S Hertika, M Musa and D Arfiati

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012102

Antifungal activity of marine sponges (Class Demospongiae) collected from Biak, Indonesia

R D Kasitowati, K Wittriansyah, A Trianto, D C Pratiwi and M A P Panjaitan

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012103

Antibacterial activities of *Physalis angulata* herb extract on white feces diseases (WFD) in *Litopenaeus shrimp vannamei*

E Saraswati and AS Wijaya

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012104

The Effects of Mercury Chloride (Hgcl₂) on the Changes in Hematology and Blood Sugar Level in Carps (*Cyprinus carpio*)

I Setiyowati, H Suprpto and G Mahasri

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012105

Change in Two-Spot Catfish Histopathological Liver (*Mystus nigriceps*) Accumulated with Heavy Metal Cadmium (Cd) in Ketingan Estuary, Sidoarjo - East Java, Indonesia

L Anggitasari, H Suprpto and dan D D Nindarwi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012106

Effectiveness of Pepaya Leaf Extract (*Carica Papaya L.*) to Control Ectoparasite *Argulus* on Common Carp (*Cyprinus Carpio*)

L S Azizah, Kismiyati and A H Fasya

[+ Open abstract](#) [View article](#) [PDF](#)
This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.




-
- OPEN ACCESS** 012107
Bacterial Identification from Marine Ornamental Fish in Fish Quarantine, Quality Control and Fishery Products Safety Class I Denpasar, Bali
N R Rukmana, G Mahasri, S N Hidayah, M F Ulkhaq and H Kenconoajati
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012108
Prevalence and Intensity of Ectoparasites of Tilapia (*Oreochromis niloticus*) in Ponds with Low, Medium and High Stocking Density
M Indahsari, Kismiyati and M F Ulkhaq
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012109
Texture profile of the bread produced from composite flour *Bruguiera gymnorrhiza* flour (BGF) and wheat flour
M N G Amin, M N Hasan, Zakariya, S A Pralebda, H Pramono, E Saputra, S Subekti and M A Alamsjah
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012110
The potential of seaweed waste (*gracilaria* sp. and *eucheuma cottonii*) as a medium density fiberboard (mdf)-based pot material for better water use efficiency in tomato plants
H Kurnia, R R Rifadi, Agustono, M N G Amin, S A Sudjarwo and M A Alamsjah
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012111
Characterization of edible coating based on surimi fillet catfish as biodegradable packaging
E Saputra
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012112
Characteristics of the fish protein isolate recovered from Sardine by-products using the Isoelectric Solubilization- Precipitation method
H. Pramono, N T Irawan and M R A Firdaus
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012113
The potential of peptides derived from the chymotrypsin hydrolysate of soft shelled turtle yolk against the Angiotensin I Converting Enzyme
D Y Pujiastuti and J L Hsu
[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



-
- OPEN ACCESS** 012114
Kappa and iota carrageenan combination of *Kappaphycus alvarezii* and *Eucheuma spinosum* as a gelatin substitute in ice cream raw material product
I Suryani, D I Permata Sari, D M Astutik and A Abdillah
[+](#) Open abstract [View article](#) [PDF](#)
-
- OPEN ACCESS** 012115
Detection of antibiotic-resistant *Salmonella* sp. in the seafood products of Surabaya local market
H Pramono, A Kurniawan, N Andika, T F Putra, M A R Hazwin, S Utari, A P Kurniawan, E D Masithah and A M Sahidu
[+](#) Open abstract [View article](#) [PDF](#)
-
- OPEN ACCESS** 012116
The effect of kappa-carrageenan fortification on the physicochemical and organoleptic properties of milkfish galantin
D Darmawan, L Sulmartiwi and A A Abdillah
[+](#) Open abstract [View article](#) [PDF](#)
-
- OPEN ACCESS** 012117
The application of a high voltage electric field (HVEF) to reduce *Escherichia coli* and *Salmonella thyphimurium* bacteria in red snapper (*Lutjanus* sp.) fillets
D J Subakti, H Pramono and J Triastuti
[+](#) Open abstract [View article](#) [PDF](#)
-
- OPEN ACCESS** 012118
Isolation and identification of fish import consumption bacteria in a fish quarantine center, focusing on the quality control and safety of fishery products at Tanjung Priok, Jakarta
H S Farizky and W H Satyantini
[+](#) Open abstract [View article](#) [PDF](#)
-
- OPEN ACCESS** 012119
Replacement of gum arabic by dry *Spirullina* sp. biomass as a food emulsifier in bread making
D Wulandari, M N G Amin, E D Masithah, M Lamid and M A Alamsjah
[+](#) Open abstract [View article](#) [PDF](#)
-
- OPEN ACCESS** 012120
Nutrient improvement of *Bruguiera gymnorrhiza* peel fruit through fermentation using commercial tempeh (Indonesian fermented soybean) mold
M N G Amin, R A Prastiya, M N Hasan, Zakariya and M A Alamsjah
This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.
[+](#) Open abstract [View article](#) [PDF](#)
-

-
- OPEN ACCESS** 012121
DPPH scavenging property of bioactives from soft corals origin palu bay, Central Sulawesi, Indonesia
W A Tanod, U Yanuhar, Maftuch, D Wahyudi and Y Risjani
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012122
Extraction of bioactive compounds fruit from *Rhizophora mucronata* using sonication method
Ernawati, E Suprayitno, Hardoko and U Yanuhar
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012123
Harvesting *Chaetoceros gracilis* by flocculation using Chitosan
W AA Yamin, S R M Shaleh, F F Ching, R Othman, M Manjaji-Matsumoto, S Mustafa, S Shigeharu and G Kandasamy
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012124
The Use of Rajungan (*Portunus Pelagicus*) Shells as Flour in Wet Noodles Ingredient
A Rahma, A A Abdillah and E Saputra
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012125
The Potential of Lindur Fruit Flour (*Bruguiera Gymnorrhiza*) in Reducing Oil Absorption of Milkfish Nugget during the Deep Frying Process
A Widyastuti, AA Abdillah and Laksmi Sulmartiwi
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012126
Chemical and Sensory Characteristics of Flakes Made from Seaweed (*Euचेuma cottonii*) and Soybean (*Glycine max* (L.) Merrill)
I R Firdarini, Kismiyati and A Manan
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012127
The Effect of Maltodextrin Concentration on the Characteristics of Snappers' (*Lutjanus* sp.) Peptone
R Ningsih, Sudarno and Agustono
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more,
[open our Privacy and Cookies policy.](#) 
-
- OPEN ACCESS** 012128

Physics and Chemical Characteristics of Sargassum Sp. Seaweed with Addition of Sodium Alginate Stabilizer to Different Concentrations

M Via, A A Abdillah and M A Alamsjah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012129

The Effect of Sorbitol Addition on the Characteristic of Carrageenan Edible Film

M Rahmawati, M Arief and W H Satyantini

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012130

The Utilization of Flower Crab (*Portunus Pelagicus*) Shell as Pasta Flavor

R Rahmawati, E Saputra and A A Abdillah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012131

Income analysis of catching fish using dogol (demersal danish seine) in the sub-district of Brondong, Lamongan

A Aulia, A M Sahidu and Agustono

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012132

Area development based on conservation and ecotourism on the Cemara Beach (Pine Trees Beach), Pakis, Banyuwangi, East Java Province, Indonesia

E W Setyaningrum, A T K Dewi, K P Prapti, Z Erwanto and H D Susanti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012133

Socioeconomic and institutional factors affecting the sustainable development for fisheries in Bontang City, Indonesia

B I Gunawan

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012134

Factors influencing the market requirements fulfillment in small medium enterprise fish processors in East Java

W Sulistyowati, Soemarno, A Efani and M Primyastanto

[+ Open abstract](#) [View article](#) [PDF](#)

JOURNAL LINKS

[Journal home](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more,

[see our Privacy and Cookies policy.](#)



Information for organizers

Information for authors

Contact us

Reprint services from Curran Associates

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



PAPER • OPEN ACCESS

The spectrum of light and nutrients required to increase the production of phycocyanin *Spirulina platensis*

To cite this article: H A Wicaksono *et al* 2019 *IOP Conf. Ser.: Earth Environ. Sci.* **236** 012008

View the [article online](#) for updates and enhancements.

You may also like

- [The effect of different temperature on the stability of phycocyanin on microcapsule *Spirulina platensis*](#)
FNW Purnama, TW Agustini and RA Kurniasih
- [Effect of ultrasonication-assisted extraction time on the profile of phycocyanin from wet biomass *Spirulina platensis*](#)
Safrina Dyah Hardiningtyas, Iriani Setyaningsih and Aulia Andhika Radya Susila
- [Improvement stability of phycocyanin from *Spirulina platensis* encapsulated by water soluble chitosan nanoparticles](#)
A Gustiningtyas, I Setyaningsih, S D Hardiningtyas *et al.*



244th Electrochemical Society Meeting

October 8 – 12, 2023 • Gothenburg, Sweden

50 symposia in electrochemistry & solid state science

Abstract submission deadline:
April 7, 2023

Read the call for papers &

submit your abstract!

The spectrum of light and nutrients required to increase the production of phycocyanin *Spirulina platensis*

H A Wicaksono¹, W H Satyantini^{2*} and E D Masithah³

¹Aquaculture, Faculty of Fisheries and Marine Universitas Airlangga, Surabaya 60115, Indonesia

²Department of Fish Health Management and Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya, Indonesia.

³Department of Marine, Faculty of Fisheries and Marine Universitas Airlangga, Surabaya 60115, Indonesia

*Corresponding author: worohastuti79@gmail.com

Abstract. *Spirulina platensis* is one of the microalgae that have several active ingredients that can be used in fisheries, pharmaceuticals, food additives and cosmetics, and one of them contains a high level of phycocyanin. Different light qualities can be used to increase the production of cyanobacterium phycocyanin. Some of the light spectrum can affect pigment synthesis, and besides that, nutrients are also very important to maintain the quantity, quality and production of *Spirulina platensis* in the biosynthesis of phycobiliprotein (phycocyanin and phycoerythrin). This study aims to determine whether there was an influence from a combination of the color spectrum of light and nutrient composition on the phycocyanin production of *Spirulina platensis*. The research method was experimental. The study used a combination of two light spectra (red and blue) as well as two different nutrients (Walneand technical nutritional modification and MT). Each treatment combination was repeated four times. The results showed that the light spectrum and nutrient composition had an effect on *Spirulina platensis* production. The combination of the nutrient MT and red spectrum resulted in the highest production of phycocyanin and a dry weight of *Spirulina* of 5.1078 mgg⁻¹ and 0.6677/0.5 g L⁻¹ respectively. This was significantly different (P<0.05) compared to the other treatment combinations.

1. Introduction

Spirulina is a microalgae with excellent nutritional characteristics. Beside its high protein content, there are several compounds and natural pigments therein. Among them are the phycobiliproteins derived from *Spirulina*, phycoerythrin (PE) and allophycocyanin (AP) which can be found in small amounts. The most abundant is phycocyanin (PC), which is a bright blue pigment [1].

The productivity of the *Spirulina platensis* cells is influenced by the availability of both macro and micronutrients (C, N, P, K, S, Mg, Na, Cl, and Ca, Fe, Zn, Cu, Ni, Co, and Se) [2]. Most of the pigments absorb certain wavelengths of light [3]. Meanwhile, some of the light prompts different cyanobacterium phycocyanin production. According to [4]'s research results, Blue Light (45%) is the most suitable condition, with Red Light (29%) and Green Light (25%) showing second and third place for pigment synthesis respectively. The culturing conditions can influence decisively the growth phases of *Spirulina*, causing changes in its composition and increasing or decreasing the proportion of phycobiliproteins, including phycocyanin [1].

The problem that arises is if the use of different light spectrums and different nutrients affects the production of phycocyanin? Which combination of light spectrum and nutrients provides the optimum phycocyanin production?



Therefore, it is necessary to do research into the use of different light spectrums and nutrients for the production of *Spirulina platensis* phycocyanin. The purpose of this study was to determine the effect of different light spectra on phycocyanin production, to determine the different nutrient compositions involved in phycocyanin production and to find the best combination of light spectrum and nutrients associated with the optimum production of phycocyanin *Spirulina platensis*.

2. Research Method

2.1 Place and time

The research study was implemented on March 1st and lasted until March 31st2015 in the Education Laboratory of the Faculty of Fisheries and Marine, Universitas Airlangga in Surabaya, Indonesia.

2.2 Materials and research

The tools needed in for the pure culture of *Spirulina* are 2 TL 40 watt lamps (red and blue) as the sources of light and energy, aerators, aeration hoses, 16 1 liter-size culture bottles, 16 measuring glass, black plastic as a light barrier, pipette drops, volume pipettes, filter paper, funnels, a Sedgewick rafter, lux meters, thermometers, pH meters, refractometers, nitrate and phosphate test kits and microscopes.

The materials used were a pure stock of microalgae *Spirulina platensis*, a fertilizer with medium technical modifications (MT), a control sample of Walne, fresh water, sea water, alcohol 90%, N-tiosulfate 20 ppm and 60 ppm of chlorine.

2.3 Research design

This research study used an experimental method. The research design used was a Complete Random Factorial Pattern Design with 2 factors (factor A was the color of the lights with levels a1 and a2 and factor B was the nutrients with levels b1 and b2), so we obtained 2 x 2 treatment combinations: a1b1, a1b2, a2b1, and a2b2. Each treatment was repeated 4 times; 4 x 4 = 16 experiments [5] : Combination fertilizer Walne with a red light (a1b1), combination fertilizer Walne with a blue light(a1b2), combination fertilizer MT with a red light (a2b1) and combination fertilizer MT with a blue light (a2b2).

2.4 Work procedures

2.4.1 Tools and materials preparation

The culture equipment to be used was washed thoroughly, rinsed with fresh water and then dried. For the equipment made of heat-resistant glass, it was covered with cotton and gauze and then wrapped in aluminum foil. After that, it was sterilized using an autoclave at 121 ° C for 15 minutes. The heat-resistant equipment was sterilized with a 150 ppm chlorine solution for 24 hours. We then rinsed it with fresh water until clean and until the smell of chlorine was gone.

2.4.2 Culture media preparation

The seawater that will be used for the culture was sterilized using a chlorine solution. The sea water was filtered first using cotton that was placed in a water funnel, before it was sterilized with chlorine 60 ppm for 24 hours and then aerated. The residual chlorine was removed with 20 ppm Na Thiosulfate.

2.4.3 Fertilizer preparation

The fertilizers used as a culture media were the technical medium modification (MT) and Walne medium (control). The liquid fertilizer solutions for nutrients MT and Walne were stored in opaque containers. The liquid fertilizer solutions for the MT and Walne nutrients were then sterilized.

2.4.4 Culture of *Spirulina platensis*

The microalgae pure stock of *Spirulina platensis* was obtained from the Center for Brackish Water Aquaculture Development (BBPBAP) in Jepara. The growth performance of *Spirulina platensis* was assessed in limited light conditions using a 1-liter bottle incubated at a temperature of 30 ° C illuminated from one side with a fluorescent color lamp in a bright cycle and dark cycle (16 hours: 8 hours). The *Spirulina platensis* culture was done using seawater with a salinity of 15 ppt [6]. The *Spirulina platensis* inoculan which was put into the media had a 10⁴units ml⁻¹ density.

2.5 Calculation of phycocyanin

The *Spirulina platensis* cell biomass that was filtered was rinsed with distilled water and dried. The drying of the *Spirulina platensis* was carried out at 40 ° C for 48 hours [6,7]. The dried *Spirulina platensis* cells were then broken down using the sonication method. Phycocyanin extraction was carried out using a 0.1 M Na phosphate buffer (pH 7) with a concentration of 0.04 g of dry *Spirulina platensis* mL⁻¹ solvent. It was then shaken at a speed of 100 rpm at 25 ° C. The samples were collected after 24 hours. It was then centrifused, the supernatant was harvested and its optical density was measured using a spectrophotometer at 615 and 652 nm wavelengths [7].

The calculation of the *Spirulina* extraction results based on the concentration, level of purity and the production of phycocyanin was done, using several formulas. The phycocyanin concentration (PC, mg mL⁻¹) was calculated using the following formula:

$$PC = \frac{(OD_{615} - 0.474(OD_{652}))}{5.34} [6], [7]$$

PC : Phycocyanin concentration (mg mL⁻¹)

OD₆₁₅ : Reading the optical density of the sample at a wavelength of 615 nm

OD₆₅₂ : Reading the optical density of the sample at a wavelength of 652 nm

The calculation of the production of the phycocyanine extraction (mg g⁻¹) was done using the following formula:

$$\text{Phycocyanin} = \frac{PC \times V}{DB} [6]$$

PC : Phycocyanin concentration (mg mL⁻¹),

V : volume (mL) solvent,

DB : dry weight biomass (g).

2.6 Population of *Spirulina platensis*

During the culturing, the counting of the biomass of *Spirulina* population was carried out by taking samples every 24 hours and counting using the Sedgewick rafter counting cell method. The calculation was done using the following equation:

$$\text{Units/ml} = \frac{C \times 1000 \text{ mm}^3}{A \times D \times F}$$

Description:

C = number of organisms calculated

A = Wide, airy viewpoint ($3.14 \times (\frac{d}{2})^2$), mm²

D = depth of fields of view, mm

F = number of fields of view are uncountable.

2.7 Measuring the parameters of the study

The parameters measured in this study included the phycocyanin content of *Spirulina platensis*, the population density, the dry weight of *Spirulina platensis* and water quality.

2.8 Data analysis

The data was analyzed using an ANOVA factorial pattern with an error rate of 5%. If the treatment had a significant effect, then in order to see the treatment with the best results, it was continued with a double comparison of the Honestly Significant Difference [5].

3. Results and discussion

3.1 Results

3.1.1 Phycocyanin *Spirulina platensis*

The measured phycocyanin content of this study ranged from 3.5838 to 5.1078 mg g⁻¹. The data obtained during the study was then analyzed using analysis of variance (ANOVA). In order to find out the differences between all of the treatments, it was necessary to do an honest difference test with a confidence level of 0.05. The results of the test showed significantly different results ($p < 0.05$) and the highest phycocyanin was obtained in treatment a2b1 (nutrient MT combined with a red light) of 5.1078 mg g⁻¹ (Table 1).

Table 1. Phycocyanin content of *Spirulina platensis* (mg g⁻¹) with a combination of nutrients and color spectrums in different lights.

Treatment	Average Production of Phycocyanin
a1b1	4.7590 ^b
a1b2	3.0182 ^d
a2b1	5.1078 ^a
a2b2	3.5838 ^c

^{a,b,c,d} The superscript is different in one column to show the real difference ($p < 0.05$), a1b1: Fertilizer Walne with red light spectrum., a1b2: Fertilizer Walne with spectrum blue light., a2b1: Fertilizer MT with red light spectrum., a2b2: Fertilizer MT with spectrum blue light.

3.1.2 Population density of *Spirulina platensis*

The results of the variance analysis showed that factor a (nutrient type) and factor b (spectrum of light color) were significantly different to the population of *Spirulina platensis*. To find out the difference between the treatments, an Honestly Significant Difference Test (BNJ) was carried out. The data on the population density of *Spirulina platensis* can be seen in Table 2. in treatment a2b1 and a2b2, there was a decrease in the *Spirulina* population on day 8, while treatments a1b1 and a1b2 decreased on day 9.

Table 2. Cell density of *Spirulina platensis* with a combination of nutrients and different light spectrums.

Treatment	Density of <i>Spirulina platensis</i> (10 ⁴ unitml ⁻¹), Days									
	0	1	2	3	4	5	6	7	8	9
a1b1	1	2.31 ^a	4.23 ^a	4.44 ^c	5.48 ^c	9.37 ^b	10.76 ^b	12.26 ^c	13.38 ^a	10.10
a1b2	1	2.48 ^a	3.08 ^c	4.72 ^c	5.95 ^c	8.21 ^c	9.42 ^c	11.71 ^c	12.57 ^b	9.10
a2b1	1	2.58 ^a	3.27 ^c	5.27 ^b	7.04 ^b	10.18 ^a	12.57 ^a	15.10 ^a	13.41 ^a	-
a2b2	1	2.42 ^a	3.70 ^b	5.94 ^a	7.84 ^a	10.31 ^a	12.08 ^a	13.94 ^b	11.36 ^c	-

^{a,b,c} The superscript differs in one column to show the significant difference ($p < 0.05$).




Figure 1. Graph of the average population growth of *Spirulina platensis* (10^4 units/ml). Description: a1b1: Fertilizer Walne with red light., a1b2: Fertilizer Walne with blue light., a2b1: Fertilizer MT with red light., a2b2: Fertilizer MT with blue light.

3.1.3 Dry weight of *Spirulina platensis*

The dry weight of *Spirulina platensis* obtained from this study ranged from 0.4150 to 0.6677/0.5 gL⁻¹. The data obtained during the study was then analyzed using analysis of variance (ANOVA). In order to find out the differences between all of the treatments, it was necessary to do an honest difference test with a confidence level of 0.05. The full statistical test results can be seen in Table 3. The results of the statistical analysis showed significantly different results and the highest dry weight of *Spirulina platensis* was achieved in a2b1 treatment (nutrient MT and red light spectrum) with 0.6677/0.5 gL⁻¹. This was significantly different from the other treatments ($p < 0.05$).

Table 3. The dry weight of *Spirulina platensis* (in 0.5 L⁻¹) with a combination of nutrients and color spectrums with different lights

Treatment	Dry Weight (in 0.5L ⁻¹) g
a1b1	0.4787 ^b
a1b2	0.4150 ^c
a2b1	0.6677 ^a
a2b2	0.5135 ^b

Description: the superscript is different in one column to show the real difference ($p < 0.05$), a1b1: Fertilizer Walne with red light., a1b2: Fertilizer Walne with blue light., a2b1: Fertilizer MT with red light., a2b2: Fertilizer MT with blue light.

3.1.4 Water quality

The water quality and light intensity during the study were measured as supporting data. For seven days, the culture of the water quality and light intensity was measured every 24 hours. The range of water quality and light intensity can be seen in Table 4.

Table 4. Range of water quality and light intensity during the period of the *Spirulina* culture

Parameter	Range
pH	7-9

Temperature (°C)	30-34
Salinity (ppt)	30-32
Nitrate (mg/l)	12.5– 100
Phosphat (mg/l)	0.25– 2
Intensity (Lux)	4232-4257

3.2 Discussion

The results showed that the use of the red spectrum against the production of *Spirulina platensis* phycocyanin was different from the use of the blue spectrum. The highest phycocyanin content was found in the use of the red spectrum compared to the use of the blue spectrum. These things indicate that more disorders can increase the production of phycocyanin. Radiation rays between 380 and 750 nm are enough energy to help the chemistry to absorb the molecules, such as those occurring along the photosynthetic pathway in microalgae [3,8]. Therefore, visible light is the main source of autotrophic microalgae energy to produce organic compounds in photosynthesis. The red light spectrum at a wavelength of 700 nm was supported by the statement by [1] that high absorption occurs in visible light and that phycocyanin pigments absorb the main yellow and orange light which results in high absorbance across the spectrum. Phycocyanin production, which is higher when the red filtered light is used, adapts to the narrow range of light provided by the filter.

The maximum red spectrum absorbed will produce energy which is also optimal in that it can synthesize ATP. The ATP results from photosystem I will be assisted by photons from a wavelength of 700 nm (red spectrum) to produce NADPH[9]. The formed NADPH plays a role in the synthesis process of phycocyanin; it is thought that this causes an increase in the content of phycocyanin *Spirulina platensis*.

The culture growth rates in the MT media were higher than in the Walne media on the same day. This is thought to be the source of the nitrogen used - (NH₄)₂CO (urea) and (NH₄)₂SO₄ (ZA) quickly dissolves in water so then it is quickly utilized by the microalgae for metabolic processes [10]. Nitrogen is needed for the synthesis of amino acids, which form proteins and other cellular components such as phycocyanin. This does not have a significant effect on biomass [11].

Culture growth rates in the MT media were higher than in the Walne media on the same day. This was thought to be due to the source of phosphorus in the MT media, which was more quickly utilized by the microalgae for metabolic processes. The source of the MT phosphate media was Na₂HPO₄, and Walne's medium was NaH₂PO₄.2H₂O. Phosphorus deficiency causes the algae cells to decrease the protein content followed by the degradation of various cell components associated with protein synthesis including phycocyanin.

4 Conclusion

From the research that has been carried out, it can be concluded that the use of the light color spectrum influences the production of phycocyanin *Spirulina*. The nutrient composition has an effect on the production of phycocyanin. There was an interaction between the nutrients and the different color spectra on the production of phycocyanin *Spirulina*. The combination of nutrient MT and the red spectrum produced the highest amount of phycocyanin by 5.1078 mg g⁻¹.

5. References

- [1] Walter A, Júlio C de C, Vanete T S, Ana B B de F, Vanessa G and Carlos R S 2011 *Braz. Arch. Biol. Technol*, **54** 675-682.
- [2] Habib M A B 2008 *Review on culture, production and use of Spirulina as food for humans and feeds for domestic animals and fish*. Food and agriculture organization of the United Nations.

- [3] Carvalho A P, Silva S O, Baptista J M, and Malcata, F X 2011 *Microb. Biotechnol.*, **89**1275-1288.
- [4] Vijaya V and N Anand 2009 *J. Agricul. Biol. Sci.*, **4**: 1-8.
- [5] Kusningrum R S 2008 *Experimental Design*. Surabaya: Pusat Penerbitan dan Percetakan Universitas Airlangga. (Text in Indonesian)
- [6] Silveira S T, J F M Burkert, J A V Costa, C A V Burkert, S J Kalil 2007 *Bioresour. Technol.*, **98** 1629-1634.
- [7] Satyantini W H 2013 *The technology of Spirulina platensis ficocyanin production and its use as an immunostimulant in duck grouper (Cromileptes altivelis)*.
- [8] Kommareddy A and G Anderson 2003 *Study of Light as a Parameter in the Growth of Algae in a Photo-Bio Reactor (PBR) in 2003 ASAE Annual Meeting*. American Society of Agricultural and Biological Engineers.
- [9] Richmond A 2004 *Handbook of microalgal culture: biotechnology and applied psychology*. **577**: Wiley Online Library.
- [10] Barus D A 2013 *The content of ficocyanin, protein, and antioxidants of spirulina platensis grown in the media and the age of cultivation are different*.
- [11] Qasim M, Imran N, Majeeda R, Khawar A S, Abdul A, Zahida F and Zubair A 2012 *Afr. J. Microbiol. Res.*, **6** 2788-2792.