

# Table of contents

Volume 236

2019

◀ Previous issue    Next issue ▶

**The 1st International Conference on Fisheries and Marine Science 6 October 2018, East Java, Indonesia**

Accepted papers received: 16 January 2019

Published online: 01 March 2019

Open all abstracts

## Preface

**OPEN ACCESS** 011001

The 1st International Conference on Fisheries and Marine Science

+ Open abstract     View article     PDF

**OPEN ACCESS** 011002

Organizing Committee

+ Open abstract     View article     PDF

**OPEN ACCESS** 011003

Conference Photos

+ Open abstract     View article     PDF

**OPEN ACCESS** 011004

Peer review statement

+ Open abstract     View article     PDF

## Papers

**OPEN ACCESS** 012001

The Increase in  $\beta$ -carotene Content in *Dunaliella salina* from the Application of Different Light Intensities

N Sugiati, E D Masithah, W Tjahjaningsih and A A Abdillah

+ Open abstract     View article     PDF

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more,

**OPEN ACCESS** and Cookies policy.

012002 

Ammonia-eliminating potential of *Gracilaria* sp. And zeolite: a preliminary study of the efficient ammonia eliminator in aquatic environment

M R Royan, M H Solim and M B Santanumurti

[+ Open abstract](#) [View article](#) [PDF](#)

---

OPEN ACCESS

012003

Identification of extracellular enzyme-producing bacteria (proteolytic, cellulolytic, and amylolytic) in the sediment of extensive ponds in Tanggulangrejo, Gresik

OA Artha, Sudarno, H Pramono and LA Sari

[+ Open abstract](#) [View article](#) [PDF](#)

---

OPEN ACCESS

012004

Addition of water from the treatment pond of pangasius fillet waste (*Pangasius* sp.) with different concentrations in the cultivation medium due to the population growth of *Daphnia* sp.

H P Alvian, E D Masithah and M H Azhar

[+ Open abstract](#) [View article](#) [PDF](#)

---

OPEN ACCESS

012005

The growth and survival rate in lettuce aquaponic systems (*Latuca sativa*) of eels in various stocking densities of eel (*Monopterus albus*)

N K Portalia, L Sulmartiwi and B S Rahardja

[+ Open abstract](#) [View article](#) [PDF](#)

---

OPEN ACCESS

012006

The prevalence of benedeniasis in humpback grouper (*Cromileptes altivelis*) in floating net cages in Situbondo Regency, East Java, Indonesia

S O Wijaya, S Subekti and Kismiyati

[+ Open abstract](#) [View article](#) [PDF](#)

---

OPEN ACCESS

012007

The prevalence of fungi on groupers (*Epinephelus* sp.) in cage mariculture systems of the northern coast of Surabaya, East Java

E Yuliasuti, R Kusdawarti and Sudarno

[+ Open abstract](#) [View article](#) [PDF](#)

---

OPEN ACCESS

012008

The spectrum of light and nutrients required to increase the production of phycocyanin *Spirulina platensis*

H A Wicaksono, W H Satyantini and E D Masithah

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more,

[see our Privacy and Cookies policy.](#)

012009

The growth and survival rate of the larvae of the sunu grouper (*Plectropomus leopardus*) in different temperatures

L Lutfiyah, D S Budi and M F Ulkhaq

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012010

Preservation of common carp (*Cyprinus carpio*) sperm using 0.9% NaCl and ringer's lactate solution

D S Budi, L A Adawiyah and L Lutfiyah

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012011

Comparison of the efficiency (flash point, freezing point, and viscosity test) of biodiesels from *Sargassum* sp.

M B Santanumurti, M R Royan, S H Samara, S Sigit and M A Alamsjah

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012012

Study of patterns in the relationship of ecdysis with the age of freshwater crayfish *Cherax quadricarinatus* aged 76 Days

A H Fasya

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012013

Maximum density in the *Moina macrocopa* culture able to produce parthenogenesis in female offspring

A S Mubarak, D Jusadi, M Z Junior and M A Suprayudi

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012014

The oxygen content and dissolved oxygen consumption level of white shrimp *Litopenaeus vannamei* in the nanobubble cultivation system

D P Galang, A K Ashari, L Sulmatiwi, G Mahasri, Prayogo and LA Sari

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012015

The effect of the epiphytes of *Chaetomorpha crassa* on the total chlorophyll-a and growth of *Gracilaria verrucosa*

A L L Handayani, R J Triastuti and L Sulmartiwi

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.

012016

Growth monitoring of koi fish (*Cyprinus carpio*) in natural hatchery techniques in Umbulan, Pasuruan, East Java

F P Putri and N N Dewi

[+](#) Open abstract [View article](#) [PDF](#)

---

OPEN ACCESS

012017

Dynamic Ratio Correlation of N:P in relation to the Diatom Abundance in the Intensive System of the Vannamei (*Litopenaeus vannamei*) Shrimp Pond

E D Masithah, D D Nindarwi, T Rahma and dan R R Satrya P I

[+](#) Open abstract [View article](#) [PDF](#)

---

OPEN ACCESS

012018

Dynamic ratio correlation of N:P on the abundance of Bluegreen algae in an intensive system in a white shrimp (*Litopenaeous vannamei*) pond

E D Masithah, D D Nindarwi, A L A Suyoso and D Husin

[+](#) Open abstract [View article](#) [PDF](#)

---

OPEN ACCESS

012019

Dynamic ratio correlation of N:P toward phytoplankton explosions in intensive systems of white shrimp pond

E D Masithah, D D Nindarwi, D Husin and T Rahma

[+](#) Open abstract [View article](#) [PDF](#)

---

OPEN ACCESS

012020

Development of water and nutrient management models to improve multitrophic seafarming productivity

J A Surbakti, I A L Dewi, M A Alamsjah and M Lamid

[+](#) Open abstract [View article](#) [PDF](#)

---

OPEN ACCESS

012021

Pond soil characteristic in reclaimed tidal lowlands and its correlation with the water quality for aquaculture

M Fitriani, I Wudtisin, M Kaewnern and R H Susanto

[+](#) Open abstract [View article](#) [PDF](#)

---

OPEN ACCESS

012022

The dynamics of total organic matter (tom) on sangkuriang catfish (*clarias gariepinus*) farming at upt ptpbp2kp and the effectiveness of freshwater bivalve (*anodonta woodiana*) in reducing the total organic matter with varying density

D Arfiati, C D G Putra, A H Tullah, S W A Permanasari and A W Puspitasari

[+](#) Open abstract [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



---

**OPEN ACCESS** 012023

Growth and morphological changes in relation to the maturation of male Japanese eel, *Anguilla japonica* injected with human chorionic gonadotrophin (HCG) in the different interval in the tropical region

Y T Hee, F F Ching and S Senoo

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS** 012024

Genetic diversity of the endangered species *Sphyrna lewini* (Griffith and Smith 1834) in Lombok based on mitochondrial DNA

S Hadi, N P Anggraini, E Muttaqin, B M Simeon, B Subhan and H Madduppa

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS** 012025

Sex ratio and size at first maturity of razor clam *Solen* sp. in Pamekasan and Surabaya coastal area, East Java, Indonesia

N Trisyani, N I Wijaya and I Yuniar

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS** 012026

Improving productivity and water quality of catfish, *Clarias* sp. cultured in an aquaponic ebb-tide system using different filtration

E Setiadi, I Taufik, Y R Widyastuti, I Ardi and D Puspaningsih

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS** 012027

Different substrate of trickling filter on growth, survival rate, and water quality of common carp (*Cyprinus carpio*) cultivation by using an intensive recirculation system

E Setiadi, I Taufik, Y R Widyastuti, I Ardi and A Saputra

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS** 012028

Water quality dynamic, production and profitability of catfish, *Clarias* sp. cultured at different design construction of aquaponic

Y R Widyastuti, E Setiadi, I Taufik and L Setijaningsih

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS** 012029


Effect of C:N ratio on the spore production of *Bacillus* sp. indigenous shrimp pond

A Yuniarti, N B Arifin, M Fakhri and A M Hariati

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



- 
- OPEN ACCESS** 012030  
Comparative Test on Bacteria in the Digestive Tract of Vannamei Shrimp (*Litopenaeus vannamei*) at Intensive and Extensive Ponds in Ujungpangkah, Gresik  
D Ningrum, M Arief and dan K T Pursetyo  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012031  
Growth Performance of Laboratory-Scale *Chaetoceros calcitrans* in Different Containers  
M Jannah, M F Ulkhaq, M H Azhar, Suciyono and dan W Soemarjati  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012032  
Study of the Dynamic Density and Diversity of Plankton at Different Brackishwater Pond Managements in Gresik, East Java  
O Tilahwatih, E D Masithah and dan B S Rahardja  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012033  
The Effect of Demineralization Stage of Agar's Solid Waste on the Characterization of Activated Carbon  
R Febrianto, Sudarno and R Kusdarwati  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012034  
The Dynamic of Density and Diversity of Cyanophyta in Different Pond Bases in Educational Pond of Faculty of Fisheries and Marine Universitas Airlangga  
S Z Cahyani, E D Masithah and Prayogo  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012035  
Effect of Different Salinity Level within Water Against Growth Rate, Survival Rate (FCR) of Catfish (*Clarias* sp.)  
D Prananingtyas, Prayogo and S Rahardja  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012036  
Molecular identification and phylogenetic reconstruction of two fiddler crabs (*Uca forcipata* and *Uca triangularis*)  
S Andriyono, H Pramono and H W Kim  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012037  
See our Privacy and Cookies policy. 

The molecular identification and phylogenetic reconstruction of Palaemonid and Penaeid shrimp from the southern part of Bangladesh

M J Alam, S Andriyono, A T M Eunos and H W Kim

[+ Open abstract](#) [View article](#) [PDF](#)

---

OPEN ACCESS

012038

Morphometric characteristics of Fur Cockles (*Anadara spp.*) in Wonokromo and Juanda Estuary, Surabaya

P B Pamungkas, K T Pursetyo, J Triastuti and N N Dewi

[+ Open abstract](#) [View article](#) [PDF](#)

---

OPEN ACCESS

012039

Stock status of ark clams (*Anadara spp.*) based on dredge fishing of the east coast of Surabaya, Indonesia

N N Dewi, K T Pursetyo, O P Darmono, F R Fachri, F S Puspitasari and A Damora

[+ Open abstract](#) [View article](#) [PDF](#)

---

OPEN ACCESS

012040

The distribution patterns and biomass of bivalves in Segoro Tambak estuary, Sedati, Sidoarjo, East Java

S H Liyana, L A Sari, N N Dewi, E D Masithah, A M Sahidu and K T Pursetyo

[+ Open abstract](#) [View article](#) [PDF](#)

---

OPEN ACCESS

012041

Inventorization of reef fish on Tabuhan Island, Banyuwangi, East Java, Indonesia

Suciyono, M A Azhar, M F Ulkhaq and H Kenconoajati

[+ Open abstract](#) [View article](#) [PDF](#)

---

OPEN ACCESS

012042

Dynamic study on the effect of calcium hydroxide and sodium bicarbonate treatment on the N/P ratio and plankton abundance

M R N Tsany, E D Masithah, B S Rahardjo and D D Nindarwi

[+ Open abstract](#) [View article](#) [PDF](#)

---

OPEN ACCESS

012043

Distribution patterns and the biomass of bivalves at Segoro Tambak estuary, Sedati, Sidoarjo, East Java

S H Liyana, L A Sari, N N Dewi, E D Masithah, A M Sahidu and K T Pursetyo

[+ Open abstract](#) [View article](#) [PDF](#)

---

OPEN ACCESS

012044

Optimization of diatom *Hyaloecclesia ostrearia* cultivation in different mediums and nutrients

See our privacy and cookies policy.



S Arsad, C Stavrakakis, V Turpin, P Rossa, Y Risjani, L A Sari, F S Prasetya and J-L Mouget

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012045

Coastal ecosystem model based on environmental suitability and carrying capacity of the fishpond in Banyuwangi Region, East Java, Indonesia

E W Setyaningrum, Maghdalena, A T K Dewi, M. Yuniartik and E D Masithah

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012046

Development and succession of sessile macrofouling organisms on the artificial structure in the Shallow Coastal Waters of Sabah, Malaysia

M A M Affandy, J Madin, K P Jakobsen and M Auluck

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012047

Fish species difference around the light of metal halide lamps and LED lamps with mini purse seine operation

M A Sofijanto, D Arfiati, T D Lelono and A Muntaha

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012048

Status of coral diseases and compromised health syndromes on Pemuteran shallow reefs, North Bali island

W Karim

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012049

Management options for restoring artificial coral reefs in Indonesia: strengthening in institutional approach

Rudianto and Ahmad Zainul

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012050

Growth of salt-secretor and non-salt secretor mangrove seedlings with varying salinity and their relations to habitat zonation

M Basyuni, Ramayani, A Hayullah, Prayunita, M Hamka, L A Putri and S Baba

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012051

Distribution of *Ctenactis Echinata* and *Fungia Consinna* coral on Mamburit island,

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.





Sawiya, D Arfiati, Guntur and U Zakiyah

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012052

Diversity species and condition of seagrass ecosystem in Teluk Awur and Prawean Jepara

I Riniatsih, A Ambariyanto, E Yudiati, R Hartati, W Widianingsih and R T Mahendrajaya

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012053

Copper (Cu) and Cadmium (Cd) toxicity on growth, chlorophyll-a and carotenoid content of phytoplankton *Nitzschia* sp

D Hindarti and A.W. Larasati

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012054

Vegetation Characteristic and Micro Environment of Mangrove Rehabilitation Forest at Coastal Areas of East Sinjai, South Sulawesi

H Setiawan

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012055

The Use of Water Lettuce (*Pistia stratiotes*) as Phytoremediator for Concentration and Deposits of Heavy Metal Lead (Pb) Tilapia (*Oreochromis niloticus*) Gills

A A D Amalia, B S Rahardja and Rr J Triastuti

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012056

The Effectiveness of Heavy Metals Pb, Cd and Zn Reduction in NPK Fertilizer Waste Combined with Biofilters of Seaweed (*Gracillaria* sp.), Blood Clam (*Anadara* sp.), and Zeolite

A R K Sari, R K Harryes, F A Anggraini, M A Alamsyah and dan A Ahadi

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012057

The Effectiveness of Combining *Gracilaria* Sp. Seaweed Biofilter and *Anadara granosa* Shell with Zeolite in the Decrease in the Level of Mercury (Hg) Heavy Metal

J A Spespatri, B S Rahardja and A A Abdillah

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012058

The Effectiveness of Combination of Seaweed (*Gracillaria* sp.), Blood Clam (*Anadara granosa*), and Zeolite as Biofilter in the Reduction of Heavy Metal Copper (Cu)

A A D Amalia, B S Rahardja and Rr J Triastuti

[+ Open abstract](#) [View article](#) [PDF](#)



[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012059

Bioaccumulation of Cadmium (Cd) Heavy Metal on Seaweed (*Gracilaria* sp.) in Traditional Fishpond of Jabon Subdistrict, Sidoarjo District

O Ardiyansyah, Sudarno and Rosmanida

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012060

Identification of Proteolytic Bacterial Isolates in Sediment Ecosystem of Gunung Anyar Mangrove Forest, Surabaya

P B Utomo, Sudarno and B S Rahardja

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012061

The Analysis of Cockle (*Anadara inaequalvis*) Gonad Maturity Level in the Estuary of Banjar Kemuning River, Sedati, Sidoarjo

R F Saputra, E D Masithah and P D Wulansari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012062

Spatial and Temporal Variation of Biomass Blood Cockle (*Anadara* sp.) in Estuaries Dadapan, Sedati Sub-District, Sidoarjo, East Java

M R Ramadhan, K T Pursetyo, Prayogo and N N Dewi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012063

Distribution Patterns and Biomass of Bivalve in Juanda and Segoro Tambak Estuary in Sedati, Sidoarjo, East Java

S H Liyana, E D Masithah and A M Sahidu

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012064

Analysis of Cadmium (Cd) Heavy Metal on Sediment and Mangrove Leaves *Avicennia marina* at Mangrove Ecotourism Wonorejo, Surabaya

W C Dermawan, Prayogo and B S Rahardja

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012065

Analysis of Lead (Pb) Value Comparison on Seaweed (*Euचेuma cottonii*) in Bluto and Saronggi Sumenep Marine, Madura, East Java

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy](#) and [Cookies](#) policy.



[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012066

The potential addition of lemuru oil to commercial feed to increase the content of EPA and DHA in eels (*Monopterus albus*)

A Imanisa, M B Santanumurti, M Lamid and Agustono

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012067

Effect of lysine in addition to commercial feed on crude protein and the energy digestibility of gourami (*Osphronemus gouramy*)

D Setiyawan, S H Samara, Agustono and M A A Arif

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012068

Addition of the papain enzyme to commercial feed against protein retention and feed efficiency in eels (*Anguilla bicolor*)

DA Liono, M Arief, Prayogo and W Isoni

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012069

Combination of papain enzyme and phytase enzyme in commercial feed and the protein and energy retention of tilapia *Oreochromis niloticus*

A Saifulloh, M B Santanumurti, M Lamid and W P Lokapirnasari

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012070

The effect of giving cake artificial feed on the survival rate, and growth of Common carp (*Cyprinus carpio*) larva in an Installation of Freshwater Culture (IBAT) in Punten, Batu.

I P Zainiyah, Rozi, W H Satyantini and A M Sahidu

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012071

The Utilization of Phytase Enzymes and SEM Analysis in order to increase the Quality of Rice Bran as a Layer and Fish Feed

M Lamid, Anam Al-Arif and S H Warsito

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012072

The dynamic relationship of phytoplankton abundance and diversity in relation to white shrimp (*Litopenaeus vannamei*) feed consumption in intensive ponds

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012073

Effect of partial replacement of fish meal with *Spirulina platensis* meal in practical diets and culture location on growth, survival, and color enhancement of percula clownfish *Amphiprion percula*

S Hudaidah, B Putri, S H Samara and Y T Adiputra

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012074

The Effect of Different level of Probiotic Addition on Commercial Feed against Digestibility and Efficiency of Nile Tilapia Feed (*Oreochromis Niloticus*)

D Taufik, M Arief and H Kenconoajati

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012075

The Effect of Adding Synbiotics Into Commercial Feed Towards Protein Retention and Fat Retention of Dumbo Catfish (*Clarias sp.*)

H Syevidiana, M Arief and I S Hamid

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012076

The Effect of Adding Lysine in Commercial Feed on Growth Rate, Feed Efficiency, and Feed Conversion Ratio to Tambaqui (*Colossoma Macropomum*)

L V D Putra, U Agustono and S H Kenconoajati

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012077

The Effect of Coconut Shell Liquid Smoke in Commercial Feed on Total Bacteria of *Pseudomonas Aeruginosa* in the Tilapia's Kidney (*Oreochromis niloticus*)

M Rahmawati, Sudarno and S Subekti

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012078

The Effect of Coconut Shell Liquid Smoke in Commercial Feed Towards Total *Pseudomonasaeruginosa* Bacteria on Gastrointestinal Tract Tilapia (*Oreochromis Niloticus*)

S Rahmadini, Sudarno and S Subekti

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012079

Antibacterial activity of honey in preserving high-pressure cooked milkfish stored at room temperature

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



D A Hakim, W Tjahjaningsih and Sudarno

[+ Open abstract](#) [View article](#) [PDF](#)

---

OPEN ACCESS

012080

Bacterial composition in the gastrointestinal tract of *Uca* spp crabs fed on *Avicennia marina* leaf litter

M A B Kareho, E D Masithah and W Tjahjaningsih

[+ Open abstract](#) [View article](#) [PDF](#)

---

OPEN ACCESS

012081

The correlation between ectoparasite infestation and the total plate count of *Vibrio* sp. in pacific white shrimp (*Litopenaeus vannamei*) in ponds

G Mahasri, Rozi, A T Mukti, W H Satyantini and N M Usuman

[+ Open abstract](#) [View article](#) [PDF](#)

---

OPEN ACCESS

012082

In vitro study of an ethanolic extract of coffea leaves to inhibit freshwater pathogenic bacteria

H Kenconoajati, MF Ulkhaq, DS Budi and MH Azhar

[+ Open abstract](#) [View article](#) [PDF](#)

---

OPEN ACCESS

012083

The effect of noni fruits (*Morinda citrifolia*) with different ripeness stages against the total erythrocytes and leukocytes of comet goldfish (*Carassius auratus*) infested by *Argulus*

E Setyaningsih, Kismiyati and S Subekti

[+ Open abstract](#) [View article](#) [PDF](#)

---

OPEN ACCESS

012084

The protection capacity of the crude and whole protein spores of *Myxobolus koi* as an immunostimulant material development in goldfish (*Cyprinus carpio*) for preventing Myxobolus

G Mahasri, M Yusuf, R Woro and M B Santanumurti

[+ Open abstract](#) [View article](#) [PDF](#)

---

OPEN ACCESS

012085

Identification of white spot syndrome virus (WSSV) in pacific white shrimps (*Litopenaeus vannamei*) from ponds postexposure to immunogenic membrane proteins (*Zoothamnium penaei*)

P A Wiradana, G Mahasri, R E R Sari, U C Marwiyah and R Prihadhana

[+ Open abstract](#) [View article](#) [PDF](#)

---

OPEN ACCESS

012086

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



Gills and swimming leg histopathologies in pacific white shrimp (*Litopenaeus vannamei*) from ponds exposed to the immunogenic membrane proteins of *Zoothamnium penaei*

R E R Sari, G Mahasri, P A Wiradana and U C Marwiyah

[+](#) [Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012087

Total plate count and identification of vibrio in pacific white shrimp (*Litopenaeus vannamei*) from ponds and in those exposed to immunogenic protein membrane *Zoothamnium penaei*

U C Marwiyah, G Mahasri, R E Ratnasari and P A Wiradana

[+](#) [Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012088

Effectiveness of *Nitrobacter* on the specific growth rate, survival rate and feed conversion ratio of dumbo catfish *Clarias* sp. with density differences in the aquaponic system

A S Taragusti, M B Santanumurti, B S Rahardja and Prayogo

[+](#) [Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012089

The erythrocyte and leucocyte profile of saline tilapia (*Oreochromis Niloticus*) in a cultivation system with nanobubbles

M Gunanti, P D Wulansari and K Kinzella

[+](#) [Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012090

The prevalence and intensity of ectoparasites infecting vanname shrimp (*Litopenaeus vannamei*) reared in different ponds

N Nurlatiffah, Kismiyati and MF Ulkhaq

[+](#) [Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012091

Protozoan parasites of Vannamei Shrimp (*Litopenaeus vannamei*) in farmed fish from Pasuruan, Indonesia

U Hafidloh and P D W Sari

[+](#) [Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012092

Growth of *Bacillus* sp. and *Flavobacterium* sp. in culture media with the addition of liquid whey tofu waste

W H Satyantini, R M Pratiwi, A M Sahidu and D D Nindarwi

[+](#) [Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



OPEN ACCESS

012093

The effect of noni *Morinda citrifolia* L. fruit extracts on the gill histopathological changes of Nile tilapia *Oreochromis niloticus*

A T Mukti, E Dewi, W H Satyantini, L Sulmartiwi, Sudarno and M Hassan

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012094

Prevalence and intensity of ectoparasites in Pacific white shrimp (*Litopenaeus vannamei*) seeds from a pond and hatchery

G Mahasri, T Hidayat and Sudarno

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012095

Prevalence and intensity of ectoparasites in gabus fish (*Channa striata*) at Cangkringan Fishery Cultivation Technology Development Center, Sleman, Yogyakarta

E N Fitriani, Rozi, M Arief and H Suprpto

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012096

Antibacterial activity of bitter melon (*Momordica charantia* L.) leaf extract against *Aeromonas hydrophila*

D A Masithoh, R Kusdarwati and D Handijatno

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012097

Determination of the *aerolysin* gene in *Aeromonas hydrophila* using the polymerase chain reaction (PCR) technique

G Christy, R Kusdawarti and D Handijatno

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012098

Bioaccumulation and histopathological effect on the gills and liver of silver barb (*Barbonymus gonionotus*) exposed to the heavy metal nickel

I Purwanti, W Arroisi, B S Rahardja and L Sulmartiwi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012099

Identification and prevalence of the ectoparasite *Octolasmis* in sand lobster (*Panulirus homarus*) and bamboo lobster (*Panulirus versicolor*) in Floating Net Cages in Sape, Bima Regency, West Nusa Tenggara Province, Indonesia

L. Yusgita, Kismiyati, S. Subekti, P D Wulansari and M K Amiin

[+ Open abstract](#) [View article](#) [PDF](#)  
This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



OPEN ACCESS

012100

Treatment of *Chlorella sp.* extract on heat shock cluster (HSC) response from the tissue and bloodcells proliferation of *Epinephelus fuscoguttatus-lanceolatus* infected by *Viral Nervous Necrosis*

U Yanuhar, I Al-Hamidy and N R Caesar

[+](#) Open abstract [View article](#) [PDF](#)

OPEN ACCESS

012101

Superoxide dismutase (SOD) and metallothionein (MT) *Tubifex tubifex* at the acute mercury exposure

I M Widiastuti, A M S Hertika, M Musa and D Arfiati

[+](#) Open abstract [View article](#) [PDF](#)

OPEN ACCESS

012102

Antifungal activity of marine sponges (Class Demospongiae) collected from Biak, Indonesia

R D Kasitowati, K Wittriansyah, A Trianto, D C Pratiwi and M A P Panjaitan

[+](#) Open abstract [View article](#) [PDF](#)

OPEN ACCESS

012103

Antibacterial activities of *Physalis angulata* herb extract on white feces diseases (WFD) in *Litopenaeus shrimp vannamei*

E Saraswati and AS Wijaya

[+](#) Open abstract [View article](#) [PDF](#)

OPEN ACCESS

012104

The Effects of Mercury Chloride (Hgcl<sub>2</sub>) on the Changes in Hematology and Blood Sugar Level in Carps (*Cyprinus carpio*)

I Setiyowati, H Suprpto and G Mahasri

[+](#) Open abstract [View article](#) [PDF](#)

OPEN ACCESS

012105

Change in Two-Spot Catfish Histopathological Liver (*Mystus nigriceps*) Accumulated with Heavy Metal Cadmium (Cd) in Ketingan Estuary, Sidoarjo - East Java, Indonesia

L Anggitasari, H Suprpto and dan D D Nindarwi

[+](#) Open abstract [View article](#) [PDF](#)

OPEN ACCESS

012106

Effectiveness of Pepaya Leaf Extract (*Carica Papaya* L.) to Control Ectoparasite *Argulus* on Common Carp (*Cyprinus Carpio*)

L S Azizah, Kismiyati and A H Fasya

[+](#) Open abstract [View article](#) [PDF](#)  
This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.







- 
- OPEN ACCESS** 012107  
Bacterial Identification from Marine Ornamental Fish in Fish Quarantine, Quality Control and Fishery Products Safety Class I Denpasar, Bali  
N R Rukmana, G Mahasri, S N Hidayah, M F Ulkhaq and H Kenconoajati  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012108  
Prevalence and Intensity of Ectoparasites of Tilapia (*Oreochromis niloticus*) in Ponds with Low, Medium and High Stocking Density  
M Indahsari, Kismiyati and M F Ulkhaq  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012109  
Texture profile of the bread produced from composite flour *Bruguiera gymnorrhiza* flour (BGF) and wheat flour  
M N G Amin, M N Hasan, Zakariya, S A Pralebda, H Pramono, E Saputra, S Subekti and M A Alamsjah  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012110  
The potential of seaweed waste (*gracilaria* sp. and *eucheuma cottonii*) as a medium density fiberboard (mdf)-based pot material for better water use efficiency in tomato plants  
H Kurnia, R R Rifadi, Agustono, M N G Amin, S A Sudjarwo and M A Alamsjah  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012111  
Characterization of edible coating based on surimi fillet catfish as biodegradable packaging  
E Saputra  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012112  
Characteristics of the fish protein isolate recovered from Sardine by-products using the Isoelectric Solubilization- Precipitation method  
H. Pramono, N T Irawan and M R A Firdaus  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012113  
The potential of peptides derived from the chymotrypsin hydrolysate of soft shelled turtle yolk against the Angiotensin I Converting Enzyme  
D Y Pujiastuti and J L Hsu  
[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



- 
- OPEN ACCESS** 012114  
Kappa and iota carrageenan combination of *Kappaphycus alvarezii* and *Eucheuma spinosum* as a gelatin substitute in ice cream raw material product  
I Suryani, D I Permata Sari, D M Astutik and A Abdillah  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012115  
Detection of antibiotic-resistant *Salmonella* sp. in the seafood products of Surabaya local market  
H Pramono, A Kurniawan, N Andika, T F Putra, M A R Hazwin, S Utari, A P Kurniawan, E D Masithah and A M Sahidu  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012116  
The effect of kappa-carrageenan fortification on the physicochemical and organoleptic properties of milkfish galantin  
D Darmawan, L Sulmartiwi and A A Abdillah  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012117  
The application of a high voltage electric field (HVEF) to reduce *Escherichia coli* and *Salmonella thyphimurium* bacteria in red snapper (*Lutjanus* sp.) fillets  
D J Subakti, H Pramono and J Triastuti  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012118  
Isolation and identification of fish import consumption bacteria in a fish quarantine center, focusing on the quality control and safety of fishery products at Tanjung Priok, Jakarta  
H S Farizky and W H Satyantini  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012119  
Replacement of gum arabic by dry *Spirullina* sp. biomass as a food emulsifier in bread making  
D Wulandari, M N G Amin, E D Masithah, M Lamid and M A Alamsjah  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012120  
Nutrient improvement of *Bruguiera gymnorrhiza* peel fruit through fermentation using commercial tempeh (Indonesian fermented soybean) mold  
M N G Amin, R A Prastiya, M N Hasan, Zakariya and M A Alamsjah  
[+ Open abstract](#) [View article](#) [PDF](#)
- This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).
- 

- 
- OPEN ACCESS** 012121  
DPPH scavenging property of bioactives from soft corals origin palu bay, Central Sulawesi, Indonesia  
W A Tanod, U Yanuhar, Maftuch, D Wahyudi and Y Risjani  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012122  
Extraction of bioactive compounds fruit from *Rhizophora mucronata* using sonication method  
Ernawati, E Suprayitno, Hardoko and U Yanuhar  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012123  
Harvesting *Chaetoceros gracilis* by flocculation using Chitosan  
W AA Yamin, S R M Shaleh, F F Ching, R Othman, M Manjaji-Matsumoto, S Mustafa, S Shigeharu and G Kandasamy  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012124  
The Use of Rajungan (*Portunus Pelagicus*) Shells as Flour in Wet Noodles Ingredient  
A Rahma, A A Abdillah and E Saputra  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012125  
The Potential of Lindur Fruit Flour (*Bruguiera Gymnorhiza*) in Reducing Oil Absorption of Milkfish Nugget during the Deep Frying Process  
A Widyastuti, AA Abdillah and Laksmi Sulmartiwi  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012126  
Chemical and Sensory Characteristics of Flakes Made from Seaweed (*Eucheuma cottonii*) and Soybean (*Glycine max* (L.) Merrill)  
I R Firdarini, Kismiyati and A Manan  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012127  
The Effect of Maltodextrin Concentration on the Characteristics of Snappers' (*Lutjanus* sp.) Peptone  
R Ningsih, Sudarno and Agustono  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012128  
This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more,  
[open our Privacy and Cookies policy.](#) 

## Physics and Chemical Characteristics of Sargassum Sp. Seaweed with Addition of Sodium Alginate Stabilizer to Different Concentrations

M Via, A A Abdillah and M A Alamsjah

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012129

## The Effect of Sorbitol Addition on the Characteristic of Carrageenan Edible Film

M Rahmawati, M Arief and W H Satyantini

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012130

## The Utilization of Flower Crab (*Portunus Pelagicus*) Shell as Pasta Flavor

R Rahmawati, E Saputra and A A Abdillah

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012131

## Income analysis of catching fish using dogol (demersal danish seine) in the sub-district of Brondong, Lamongan

A Aulia, A M Sahidu and Agustono

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012132

## Area development based on conservation and ecotourism on the Cemara Beach (Pine Trees Beach), Pakis, Banyuwangi, East Java Province, Indonesia

E W Setyaningrum, A T K Dewi, K P Prapti, Z Erwanto and H D Susanti

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012133

## Socioeconomic and institutional factors affecting the sustainable development for fisheries in Bontang City, Indonesia

B I Gunawan

[+ Open abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS**

012134

## Factors influencing the market requirements fulfillment in small medium enterprise fish processors in East Java

W Sulistyowati, Soemarno, A Efani and M Primyastanto

[+ Open abstract](#) [View article](#) [PDF](#)

## JOURNAL LINKS

---

Journal home

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more,

see our Privacy and Cookies policy.



---

Information for organizers

---

Information for authors

---

Contact us

---

Reprint services from Curran Associates

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



PAPER • OPEN ACCESS

## Coastal ecosystem model based on environmental suitability and carrying capacity of the fishpond in Banyuwangi Region, East Java, Indonesia

To cite this article: E W Setyaningrum *et al* 2019 *IOP Conf. Ser.: Earth Environ. Sci.* **236** 012045

View the [article online](#) for updates and enhancements.

You may also like

- [Divorce and Family Role Based on Slerek as Traditional Fishing Method in Muncar Banyuwangi Coastal Area](#)  
Burhani Buhani and Sri Surahmi
- [Exotic Heritage from Coastal East Java of Batik Bayuwangi](#)  
Syaiful Hadi, Ikhwanul Qiram and Gatut Rubiono
- [Design of the Needs Model for the Development of Young Generation Interests in the Agricultural Sector in Banyuwangi Regency](#)  
Muksin, I Juanda, N D Wahyono et al.



### 244<sup>th</sup> Electrochemical Society Meeting

October 8 – 12, 2023 • Gothenburg, Sweden

50 symposia in electrochemistry & solid state science

Abstract submission deadline:

**April 7, 2023**

Read the call for papers &

**submit your abstract!**

# Coastal ecosystem model based on environmental suitability and carrying capacity of the fishpond in Banyuwangi Region, East Java, Indonesia

E W Setyaningrum<sup>1,\*</sup>, Maghdalena<sup>2</sup>, A T K Dewi<sup>1</sup>, M. Yuniartik<sup>1</sup>, E D Masithah<sup>3</sup>

<sup>1</sup> Faculty of Agriculture and Fisheries, University of 17 Agustus 1945, Banyuwangi, Indonesia.

<sup>2</sup> Faculty of Science Education, University of 17 Agustus 1945, Banyuwangi, Indonesia.

<sup>3</sup> Faculty of Fisheries and Marine, University of Airlangga, Surabaya, Indonesia.

\*Corresponding author: ervinawahyu@untag-banyuwangi.ac.id

**Abstract.** Banyuwangi coastline (175.8 Km) is the longest coastal area in East Java. There are various potential coastal resources, one of them is aquaculture. Unfortunately, the utilization of the fishpond is not optimal and most of the fishpond areas were abandoned. The main cause is a decline in water quality and/or changes in coastal habitats. This study aims: (1) to identify the suitability and the carrying capacity of the fishpond and (2) to determine the coastal ecosystem model in Banyuwangi based on the spatial analysis. A descriptive method was used and direct observation and interview survey were conducted for collecting primary data. The data were analyzed according to the suitability and carrying capacity of the study area. A GIS analysis was used to determine the coastal ecosystem model. The results showed that Banyuwangi has suitable coastal areas for developing aquaculture is 1,220.535 Ha with a carrying capacity of 898.71Ha. Therefore, Banyuwangi coastal area facing critical problems due to ecological constraints of exceeding its carrying capacity limit for about 26.43% or 322.827 Ha.

## 1. Introduction

Banyuwangi is a regency with the largest area in East Java (around 5,782.50 km<sup>2</sup>) and borders with Jember, Bondowoso, Situbondo, and Bali. In addition, Banyuwangi regency has the longest coastal area in East Java (175.8 Km), which is located in nine different sub-districts; one sub-district faces Indonesian Ocean, seven sub-districts face Bali Strait, and one sub-district faces the Java Sea. The northern part, from Bajulmati to Wongsorejo village, almost 60% fishpond is not producing in the last two years. While the eastern coastal area from Bulusan to Muncar Banyuwangi has around 30% unproductive shrimp fishpond both the traditional or intensive ones due to disease, weather, and other reason. Those fishponds are left by the owners.

Fishpond productivity has been increasing since 1985, however, there was a decrease in the 2000s and started to increase afterward. The decrease in fishpond productivity in a given year needs attention, especially related to fishpond carrying capacity in the Regency, so that fishpond carrying capacity can be known earlier that fishpond area resources allocation can be determined more precisely [1]. The final result is the concept of sustainable aquaculture can be realized in Banyuwangi Regency. In the Government Regulation of the Republic of Indonesia Number 26 of 2008 concerning National Spatial Planning, it has been determined that the policy of developing cultivation area including development control of aquaculture activity so as not to exceed environment carrying capacity. The spatial planning act covers both spatial and environmental issues.



Land suitability is the adaptability of land for a particular purpose, through land value determination (class) and land use patterns linked to the area potential, thus, more directed land use along with its sustainability can be attempted [2]. Ecological changes in the fishpond environment will affect environmental carrying capacity which then affects fishpond production. Environmental carrying capacity for fishpond is the natural ability to provide the existence of tolerable fishpond. This natural ability will ultimately affect fishpond production [3]. Carrying capacity is an important concept for ecosystem-based management which helps define the upper limits of aquaculture without causing “unacceptable change” to both natural ecosystem and social functions and structures [4].

Carrying capacity is a factor with a major influence on development. In this context, carrying capacity is a widely used concept that encompasses four aspects namely physical, production, ecological, and social carrying capacity [5] [6] [7]. One of the natural resources and carrying capacity is a physical environment in which the development carried out. Physical carrying capacity assumes the entire water body is leased for aquaculture, being little more than the total area suitable for aquaculture [3]. In the other hand, ecological carrying capacity as the maximum density of fish an ecosystems can naturally support during the period minimum available habitat that effect to ecological process, species, population, or communities in the environment [4] [8]. From this fact, a harmony between development and physical carrying capacity is needed. To achieve this harmony, it is important to know the physical carrying capacity of the environment. By doing so, suitable development activities to that carrying capacity can be determined [9].

Therefore, the ecosystem model in the development of a coastal area in Banyuwangi is needed to optimize the area to be more beneficial for community welfare, especially in the economic context.

## **2. Method**

### *2.1 Research time and location*

We applied the research in 9 sites of Banyuwangi coastal area (Figure 1). The research was conducted from March to August 2018 at fishery laboratory University of Airlangga Surabaya and fishery laboratory University of 17 August 1945 Banyuwangi.

### *2.2 Data collection*

The data needed include biological parameter, water quality, and substrate quality (Table 1). This parameter measurement was carried out at 9 sites points along the coast of Banyuwangi Regency. Besides primary data, secondary data from a biological parameter, water quality, and substrate were also collected.

### *2.3 Data analysis*

The analysis carried out includes the suitability analysis of coastal area for fishpond designation and the carrying capacity analysis of fishpond area, then spatial analysis was conducted. Spatial data processing in this research was data in the form of maps, satellite remote sensing data, and spatial modeling for environmental suitability analysis.

The suitability analysis of fishpond area was intended to determine land and coastal area suitability (physical, chemical, and biological) of designated aquaculture. This analysis was conducted by measuring several environmental parameters of ecological requirements consisting of a physical substrate, water quality, and hydro-oceanography for aquaculture development (Table 2).



**Table 1.** Parameters of observed aquatic environment and measurement tools/methods


Parameter	Tool/method	Remarks
Biology		
1. Chlorophyll	Filtration	Laboratory Analysis
2. Mangrove	Image & Ground Check Map	Mangrove Mapping
Water Quality		
1. Dissolved Oxygen	DO meter/Winkler	In-Situ/Laboratory
2. Salinity	Salinometer	In-Situ
3. pH	pH meter	In-Situ
4. TSS	Spectrophotometer	Laboratory Analysis
5. Ammonia	Spectrophotometer	Laboratory Analysis
6. Nitrate	Spectrophotometer	Laboratory Analysis
7. Phosphate	Spectrophotometer	Laboratory Analysis
8. Temperature	Thermometer	In-Situ
9. Heavy Metal	Spectrophotometer	Laboratory Analysis
Substrate Quality		
1. pH	pH meter	In-Situ
2. Substrate Texture	Fraction Analysis	Laboratory Analysis

**Table 2.** Parameters of land and water suitability (score) for fishpond

Environmental Characteristics	Suitability			
	S1 (4)	S2 (3)	S3 (2)	N (1)
Soil				
Slope (%)	0 – 3	3 – 6	6 – 8	>8
Depth (cm)	>150	100 – 150	75 – 100	< 75
Texture	Quite Fine	Medium	Fine	Coarse
Distance from the coast (m)	200 – 300	>300 – 4000	<200	>4000
Distance from a river (m)	0 – 1000	>1000 – 2000	>2000 – 3000	>3000
Drainage	Periodically inundated	Rather rarely inundated	Rarely inundated	Uninundated
Water				
Dissolved Oxygen (mg/l)	>5	3 – 5	1 – < 3	<1
Salinity (‰)	12 – 20	>20 – 35	>35 – 50	>50
Temperature (°C)	28 – 30	>30 – 35	12 – <18	<12
pH	7.5 – 8.5	>8.5 – 10	>10 – 11	<4
Ammonia (mg/l)	<0.3	0.3 – 0.4	>0.4 – 0.5	>0.5
Hydro-oceanography				
Tidal Amplitude (m)	1.5 – 2.5	1 – <1.5 >2.5 – 3.0	0.5 – <1.5 >3.0 – 3.5	<0.5 >3.5
Rainfall (mm/th)	2500 – 3000	2000 – <2500	1000 – <2000 >3000 – 3500	<1000 >3500
Dry Season (<60 mm)	1 – 2	>2 – 3	>3 – 5	>5

Source: Modification from Bakosurtanal (1996); Hardjodiwegen and Widiatmaka (2007).

Note: S1: Very suitable, S2: Quite suitable, S3: Conditionally suitable, N: Not suitable



**Figure 1.** Map of the research area showing the sampling sites located in Banyuwangi coastal area

The coastal area carrying capacity for aquaculture in Banyuwangi Regency was calculated by the ideal land suitability approach in that coastal area. This carrying capacity analysis was carried out on land with S1 (very suitable) and S2 (suitable) suitability. While S3 (conditionally suitable) and N (not suitable) were not recommended for aquaculture development. Land carrying capacity for development was influenced by land suitability. The equation used to calculate fishpond area carrying capacity is:

$$DDT = \sum_{i=1}^n \frac{Spi}{4n} \times 100\%$$

Note:

DDT: Fishpond Area Carrying Capacity

Spi: Score of parameter i

n: The number of parameters used

### 3. Results and discussion

#### 3.1 Water quality of Banyuwangi coastal area

Environmental parameter data taken are water quality such as temperature, salinity, pH, DO, NH<sub>4</sub>, NO<sub>3</sub>, PO<sub>4</sub>, Alkalinity (CO<sub>3</sub>, HCO<sub>3</sub>) and TOM in Banyuwangi coastal area with 9 sites of data collection points represented all districts along the Banyuwangi coast with three replication (Table 3).

**Table 3.** Water quality of Banyuwangi coastal area.

Water Quality Research Location		DO	Temperature	pH	Salinity	NH <sub>4</sub> (ppm)	NO <sub>3</sub> (ppm)	PO <sub>4</sub> (ppm)	Alkalinity		TOM
									CO <sub>3</sub> (ppm)	HCO <sub>3</sub> (ppm)	
Alas Buluh	High	6.7	30.3	7.6	26	0	1	0	12	116	59.04
	Low	6.5	29.7	7.3	25	0	0	0	36	112	52.45
Kampe	High	7.1	30.1	7.3	22	0	0	0	24	100	56.7
	Low	7	28.8	7	23	0	0	0.1	12	144	51.19
BP3	High	8	27.3	7.2	20	0	0	0	24	116	55.61
	Low	6.1	27.5	7.1	20	0	0	0.1	16	136	59.04
Cemara Beach	High	7.5	31	9	25	0	0	0.1	80	88	54.98
	Low	6.4	29.3	8	25	0	0	0	32	100	55.61
Pakem Kertosari	High	7.7	31.6	7.2	24	0	0	0	12	112	54.98
	Low	7.4	29.3	7.4	27	0	0	0	16	124	49.92
Santen Island	High	7.2	29.7	7.2	26	0	0	0	36	92	61.93
	Low	6.5	29.2	7.4	26	0	0	0	36	100	63.2
Blimbingsari	High	6.1	30.3	8.9	27	0	0	0.1	12	120	52.45
	Low	6.4	30.1	8.4	27	0	0	0.1	12	140	53.75
Pangpang Bay	High	7.04	30.3	6.9	23	0.7	0	0	24	98	56.88
	Low	6.9	29.17	6	18	0.8	0	0	24	116	56.88
Lampon	High	6.9	30.6	7.1	25	0	0	3	44	84	50.56
	Low	6.8	30.7	6.9	26	0	0	0	44	80	63.2

Data collection of water quality is categorized in 2 conditions, namely at highs and lows. Based on the data above, it can be seen that there is indeed a difference between the value of water quality at high tide, but the difference is not significant.

The research location is in the waters along the coast of Banyuwangi Regency, where coastal waters are a very narrow part of the ocean when compared to the area of the waters. This region is a meeting area between the influence of land and sea and has very diverse properties with changes in the natural environment occurring very quickly in time and space. Not only experience periodic drying and soaking every day, but also the temperature difference is greater both daily and yearly than in other parts of the sea.

Water temperature can affect the life of aquatic biota indirectly, through its influence on the solubility of oxygen in the water. The higher the water temperature, the lower the solubility of oxygen in the water, and conversely the higher the solubility of oxygen, the lower the water temperature. Temperature indirectly affects metabolism, the solubility of gases and various chemical reactions in water [10].

Photosynthesis is not directly proportional to the intensity of light. In the water column 10-15 m upward, the speed of photosynthesis is lower than the 15-30 m layer, because light at sea level is too intensive for most biota that can be injured by ultraviolet light. Photosynthesis occurs up to 100 meters, where light intensity is only 1% of the surface [11]

NH<sub>4</sub> is a form of nitrogen in natural waters. Nitrogen in seawater is absorbed by marine organisms to meet the need for nitrogen as one of the main components of amino acid formation which is the beginning of protein formation. And that happened at the location of the research conducted as stated by Kennish [12]. that nutrient sources were obtained from river water input, through soil washing and rock decay.

Salinity is a very important factor for the growth of plankton. Changes in salinity in the waters cause plankton maintaining the balance of osmotic pressure between protoplasm and water. Therefore salinity can affect plankton abundance and distribution. Naturally, the fluctuations of salinity in tidal areas are caused by two things: heavy rain and large evaporation [13].

### 3.2 Mangrove Distribution in Banyuwangi Regency

Based on the spatial analysis results, the distribution of mangrove in Banyuwangi is almost evenly distributed along the coast from north to south. The area of mangrove along the coast is 2,608.846 Ha, and that area is larger than fishpond area in Banyuwangi.




Figure 2. Mangrove distribution in Banyuwangi Regency

### 3.3 Coastal chlorophyll in Banyuwangi regency

After the sampling collection, it was then taken to Universitas Airlangga Surabaya laboratory to be analyzed. For chlorophyll measurement results, there are three types of chlorophyll in Banyuwangi coastal area, namely chlorophyll-a which produces blue-green color, chlorophyll b which produces yellowish green color, and chlorophyll c which produces brown-green color. Chlorophyll-a is a complex compound between magnesium and porphyrin which contains cyclopentanone ring (ring V). The four nitrogen atoms are bonded together. Chlorophyll-a is a form of chlorophyll which is found in all autotrophic plants. Chlorophyll b is the second chlorophyll found in green plants. Chlorophyll b is also bound to proteins in a cell. It is found in Chlorophyta green algae and terrestrial plants. Chlorophyll a and b are the strongest to absorb red and purple light spectrum, a green spectrum is the least absorbed. Therefore, when white light shines on structures containing chlorophyll such as leaves, a green light will be sent and reflected so that the structure appears green. Carotene is in the chromoplast which is a colored plastide containing pigments besides chlorophyll. Chlorophyll c is found in Phaeophyta brown algae and Bacillariophyta diatoms.

**Table 4.** Data of Chlorophyll Analysis in Banyuwangi Coastal Area in 2018

Chlorophyll type Research location	Chlorophyll -a	Chlorophyll -b	Chlorophyll -c	Total Chlorophyll
Alas Buluh	0.06138	0.07764	0.635417	0.23052
	0.23125	0.30776	0.62979	1.1688
Kampe	1.188194	0.37722	0.43438	0.9827
	0.24125	0.49149	0.50901	1.24175
BP3	0.117361	0.05198	0.16242	0.2313
	0.283333	0.33241	0.16434	0.53755
Cemara Beach	0.11172	0.14986	0.31964	0.31964
	0.4734	0.04749	0.18366	0.70455
Pakem Kertosari	0.22056	0.06712	0.20936	0.49704
	0.26148	0.11888	0.27036	0.65072
Santen Island	0.03768	0.0885	0.09484	0.22102
	0.03223	0.01779	0.05335	0.10337
Blimbingsari	0.00853	0.02865	0.05669	0.09387
	0.00588	0.01583	0.04321	0.08734
Pangpang Bay	0.04451	0.13286	0.19279	0.37034
	0.120622	0.360051	0.522461	1.003621
Lampon	0.11399	0.11865	0.19987	0.43251
	0.03069	0.03882	0.04575	0.11526

### 3.4 Heavy metals (Cu, Hg, Pb, and Sn) in Banyuwangi Coastal Area

Heavy metals analyzed in this research is copper (Cu), mercury (Hg), lead (Pb) and lead (Sn). The following is the results of heavy metal tests conducted in the laboratory of Research Institute and Industry Standardization Surabaya.

In general, the source of heavy metals on the coast can be divided into two, namely sources that enter naturally to the waters and artificial marine waters. While heavy metals that enter the ocean waters can come from three sources, namely:

- Input from the coastal area originating from the river and the results of coastal abrasion by wave activity.
- Inputs from the deep sea include metals released by volcanic activity in deep seas and metals released from particles through chemical processes.
- Inputs from nearshore land environments, including metals originating from the atmosphere as dust particles.

While the source of artificial metals is metal that was released during the metal and rock industry process. Some industries only use certain heavy metals for their production activities. However, in general, most industries use various types of heavy metal elements, making it difficult to trace the origin of sources of pollution.

Of the four heavy metals mentioned above, different concentrations of heavy metals are obtained in seawater. This difference in concentration is possible due to the variability of metals in water caused by currents, adsorption, tides, or deposition [13].

**Table 5.** Test Results of Heavy Metals (Cu, Hg, Pb, and Sn) in Banyuwangi Coastal Area in 2018

Parameter	Unit	Test Results						Test method
		P. 2169 Alas buluh (low tide)	P. 2170 Alas buluh (high tide)	P. 2171 Kampe (high tide)	P. 2172 Kampe (low tide)	P. 2173 BP 3 (high tide)	P. 2174 BP 3 (low tide)	
Copper (Cu)	mg/L	<0.0223	0,026	0,032	<0.0223	0,026	0,026	SNI 6989.6 : 2009
Mercury (Hg)	mg/L	<0.0005	<0.0005	<0.0005	<0.0005	<0.0005	<0.0005	SNI 6989.78 : 2011
Lead (Pb)	mg/L	0.012	0.015	0.015	0.016	0.017	0.015	SNI 6989.46:2009
Lead (Sn)*	mg/L	<0.1050	<0.1050	0.469	<0.1050	<0.1050	<0.1050	APHA Ed.21.311 B,2005

Parameter	Unit	Test Results						Test method
		P. 2175 P. Santen (high tide)	P. 2176 P. Santen (low tide)	P. 2177 P. Pakem (high tide)	P. 2178 P. Pakem (low tide)	P. 2179 P. Cemara (high tide)	P. 2180 P. Cemara (low tide)	
Copper (Cu)	mg/L	<0.0223	<0.0223	0,026	<0.0223	0,03	<0.0223	SNI 6989.6 : 2009
Mercury (Hg)	mg/L	<0.0005	<0.0005	<0.0005	<0.0005	<0.0005	<0.0005	SNI 6989.78 : 2011
Lead (Pb)	mg/L	0.017	0.018	0.017	0.022	0.018	0.018	SNI 6989.46:2009
Lead (Sn)*	mg/L	<0.1050	<0.1050	<0.1050	<0.1050	<0.1050	4.136	APHA Ed.21.311 B,2005


Note:

- Parameters were tested according to the parameter
- Symbol “<” shows Limit Of Quantity value of the tests

### 3.5 Fishpond Land Suitability in Banyuwangi Regency

The area of aquaculture fishpond in Banyuwangi Regency in the last 5 years (2012 – 2016) has been stable as well as the number of fishery households (FH) in the same period, that is 498 RTP. Reviewed from the number of FH compared to the fishpond area, the ratio of fishpond area per FH in Banyuwangi Regency has also been stable, which is 2.77 ha/FH. It shows that the area managed by each FH for the last five years has tended to be stagnant.

The potential of the fishpond in Banyuwangi Regency is relatively high considering the vast fishpond area and relatively good fishpond condition. Due to the problem of fishpond management, shrimp disease, there are many fishponds that are currently not operating.



**Figure 3.** Fishpond area in Banyuwangi Regency

**Table 6.** The Development of Area and Number of Fishery Households (FH) in Banyuwangi Regency from 2012 to 2016

Year	Fishpond Area (ha)	Number of Fishery Households (people)	Business Area Ratio (ha/FH)
2012	1381.34	498	2.77
2013	1381.76	498	2.77
2014	1381.76	498	2.77
2015	1381.76	498	2.77
2016	1381.76	498	2.77

Source: Processed from the Department of Fisheries and Food Security Banyuwangi Regency (2018)

Carrying capacity analysis was carried out on the lands with S1 (very suitable) and S2 (suitable) suitability. While land with conditionally suitable category (S3) and not suitable (N) was not recommended for fishpond land development. Land carrying capacity for development is influenced by land suitability level.

**Table 7.** Suitability Parameter of Area and Water (score) for Fishpond

Environment Characteristics	Suitability				
	Wongsorejo Sub-district Coastal Area	Banyuwangi Sub-district Coastal Area	Blimbingsari Sub-district Coastal Area	Pangpang Bay Coast (Muncar and Tegaldlimo Sub-district)	Lampon Coastal Area (Pesanggaran Sub-district)
<i>Soil</i>					
Slope (%)	13.39 (N)	3.11 (S2)	3.12 (S2)	2 (S1)	15 (N)
Depth (cm)	25 (N)	100 (S2)	25 (N)	20 (N)	150 (S1)
Texture	Medium (S2)	Quite fine (S1)	Quite fine (S1)	Quite fine (S1)	Rough (N)
Distance from the coast (m)	< 200 (S3)	>300 (S2)	>300 (S2)	>300 (S2)	>300 (S2)
Distance from a river (m)	3000 (S3)	500 (S1)	>1000 (S2)	0 – 1000 (S1)	0 – 1000 (S1)
Drainage	Not flooded (N)	Periodically flooded (S1)	Rarely flooded (S2)	Periodically flooded (S1)	Periodically flooded (S1)
<i>Water</i>					
Dissolved oxygen (mg/l)	6.9 (S1)	7.12 (S1)	6.1 (S1)	4.88 (S2)	6.85 (S1)
Salinity (‰)	22.67 (S2)	25.5 (S2)	27 (S2)	20.67 (S2)	25.5 (S2)
Temperature (°C)	28.95 (S1)	30.02 (S2)	30.3 (S2)	30.12 (S2)	30.65 (S2)
pH	7.25 (S2)	9 (S2)	8.9 (S2)	6.9 (S2)	7 (S2)
Ammonia (mg/l)	0 (S1)	0 (S1)	0 (S1)	0.45 (S3)	0 (S1)
<i>Hydro-oceanography</i>					
Tidal Amplitude (m)	3 (S2)	1.08 (S2)	1.08 (S2)	1.08 (S2)	4 (N)
Rainfall (mm/th)	179.21 (N)	179.21 (N)	179.21 (N)	179.21 (N)	179.21 (N)
Dry Season (<60mm/year)	2 (S1)	2 (S1)	2 (S1)	2 (S1)	2 (S1)

Note: S1: Very suitable, S2: Suitable, S3: Conditionally suitable, N: Not suitable

Assessment analysis of fishpond land suitability category was conducted on all coastal land in Banyuwangi Regency. From the total of 175.8 km coastal land with 1,381.76 Ha fishpond area, based



on the suitability category, estimated fishpond area in Banyuwangi with conditionally suitable (S2) category is 1,220.535 Ha.

The recommended suitability of fishpond designated area is the land with S1 (very suitable) and S2 (suitable) category. However, based on the suitability analysis, the fishpond area along the coast fell into S2 (suitable) category is 1,220.535 Ha meaning that the coastal area in Banyuwangi is suitable for developing fishpond.

Until 2016, according to the Department of Fisheries, this figure is different from the fishpond area in Banyuwangi which is 1381.76 Ha. This difference is in accordance with the survey result and ground check carried out in this research that in 2018, there are some fishpond areas are not operating due to the damage, both facilities, and infrastructure. In addition, another serious issue that should be addressed immediately is the occurrence of input channel silting, both in the fishpond irrigation channel and river mouths in some fishpond areas. This problem was also caused by the absence of optimal irrigation arrangements for the fishpond area development. In principle, the designated area for aquaculture must fulfill physical, chemical, biological, technical, social, economic, hygienic, and legal requirements. In order to meet those requirements, according to Fauzi *et al* [2], there are four main aspects that need to be considered as criteria in determining fishpond location, namely ecological, soil, biological, and social aspects. Those four aspects are supporting elements for developing aquaculture in Banyuwangi coastal area and can be used as assessment bases in designing land suitability model.

In addition, another factor affecting the decrease in fishpond production was land-use change, mostly becoming housing. Dwipradnyana [15] explained that land-use change is an issue that can give significant effects on production, as well as social and environmental aspects. Land-use change is a serious threat to food security since the effect is permanent. Since this change affects more on spatial suitability, long-term economic benefit and other alternative measures can be taken to make it more beneficial instead of giving major damage.

### 3.6 Fishpond Carrying Capacity in Banyuwangi Regency

Fishpond development with the concept of sustainability requires resource utilization under the area carrying capacity. Besides, fishpond area has exceeded the carrying capacity of Banyuwangi coastal area.

**Table 8.** Carrying Capacity and Recommendation Fishpond Area in Banyuwangi Regency


Location	Suitable fishpond area (Ha)	Carrying Capacity (%)	Fishpond area in accordance with carrying capacity (Ha)
Wongsorejo	211.039	64.29	135.677
Banyuwangi	350.031	82.14	287.515
Rogojampi	226.202	75	169.652
Pangpang Bay (Muncar and Tegaldlimo Sub-district)	430.503	75	322.877
Lampon (Pesanggaran Sub-district)	3.76	71.43	2.686
<b>Carrying capacity</b>	<b>1,221.535</b>	<b>73.57</b>	<b>898.71</b>

The fishponds in Banyuwangi Regency have exceeded environment carrying capacity showed by mangrove ecological function, irrigation system, and not optimal production. The area carrying capacity for aquaculture development in Banyuwangi Regency is 73.57% of the total area suitable for

developing fishpond. From the suitable fishpond area (1,221.535 Ha) can be developed based on the carrying capacity in Banyuwangi Regency 898.71 Ha.

Based on **Table 8**, it can be seen that from several coastal areas with aquaculture potential, almost all of them have exceeded the carrying capacity in which the fishpond utilization needed to be reduced. For fishpond in Wongsorejo coastal area, since the area is large, the reduction of land utilization was greater too, reaching 64.29% or around 35.71% Ha to make it suitable to carrying capacity and fishpond sustainability. Likewise, land utilization should be reduced up to 70%-80% in the coastal areas in Banyuwangi, Rogojampi, Muncar, Tegaldlimo, and Pesanggaran.

Thus, from 1,221.535 Ha suitable area, around 26.43% or 322,827 Ha area should be restored to support fishpond environment. The buffer area is recommended as a green area which is mangrove vegetation.



**Figure 4.** Map of Fishpond Carrying Capacity in Banyuwangi Regency.

The buffer area needs to be provided in an aquaculture bed. This buffer area is an area borders with the sea or river that is not used for aquaculture, instead of the place of mangrove vegetation which is the native of the area. Mangrove is a natural buffer which can withstand the storm and strong wind as well as being care area and foraging place for an economic commodity such as shrimp, crab, fish, and oysters. This buffer area also functions to trap sediment, protect water quality, hold toxic materials, and slow down surface water flow [1].

Presidential Decree Number 32 of 1990 concerning Management of Protected Areas explains that buffer area is needed as median with the minimum width of 130 x average value of the annual difference between the highest and the lowest in meters measured from the lowest ebb line. This is also in accordance with the Government Regulation of the Republic of Indonesia Number 26 of 2008 concerning National Spatial Plan that is the strategy to prevent negative impacts of human activities may cause environmental damage such as protecting environment ability to absorb substances, energy, and or other components that are disposed of into it.

#### 4. Conclusion

The results showed that the Banyuwangi coastal areas were suitable for developing brackish water pond culture is 1.220,535 Ha, with a carrying capacity of 898,71 Ha. Therefore, the Banyuwangi coastal area being faced with critical problems due to ecological constraints of exceeded its carrying capacity limit for about 26,43% or 322,827 Ha.

#### 5. References

- [1] Mustafa A and Tarunamulia 2009 *Journal Riset Akuakultur* **4** 3 395-406
- [2] Fauzi Y, Susilo B and Mayasari Z M 2009 *Journal Forum Geografi* **23** 2 101-111
- [3] Yulianda F 2008 *Jurnal Ilmu-Ilmu Perairan dan Perikanan Indonesia* **15** 2 157-163
- [4] Byron C J and Costa-Pierce B A 2010 *Carrying capacity tools for use in the implementation of an Ecosystems Approach to Aquaculture* (UK: FAO Expert Workshop University of Stirling UK, 6-8 December)
- [5] English G J, Hayden B J and Ross A H 2000 *An overview of factors affecting the carrying capacity of coastal embayments for mussel culture* (New Zealand: NIWA)
- [6] McKindsey C W, Thetmeyer H, Landry T and Silvert W 2006 *Aquaculture* **261** 451-462
- [7] Guyondet T, Comeau L A, Cedric B, Grant J, Rosland R, Sonier R and Filgueira R 2014 *Estuaries and Coasts* **38** (5) 1593-1618
- [8] Ayllon D, A Almodóvar, GG Nicola, I Parra and B Elvira 2012 *Journal of Fisheries Research* **134-136** 95-103
- [9] Susetyo A D and Santoso E B 2016 *JURNAL TEKNIK ITS* **5** 1 2301-9271 Print
- [10] Gufran MH dan Baso BT 2007 *Pengelolaan Kualitas Air dalam Budidaya Perairan* (Jakarta: Rineka Cipta)
- [11] Romimohtarto K dan Juwana S 2009 *Biologi Laut, Ilmu Pengetahuan tentang Biota Laut* (Jakarta: Penerbit Djambatan)
- [12] Kennish M J 1994 *J. Coastal Res, Special Issue* **12**
- [13] Nybakken J W 1992 *Biologi Laut Suatu Pendekatan Biologis* (Jakarta: PT Gramedia)
- [14] Sagala SL, R Bramawanto ARTD, Kuswardani dan WS Pranowo 2014 *Jurnal Ilmu dan Teknologi Kelautan Tropis* **6** (2) 297 – 310
- [15] Dwipradyana I M M 2014 *Faktor-Faktor yang Mempengaruhi Konversi Lahan Pertanian serta Dampaknya terhadap Kesejahteraan Petani* (Denpasar: Universitas Udayana)

### **Acknowledgments**

We would like to thank The Ministry of Research, Technology, and Higher Education who has provided the opportunity to conduct research through Higher Education Research Collaboration 2018 scheme.