

ISSN: 1412-033X
E-ISSN: 2085-4722

BIODIVERSITAS

Journal of Biological Diversity

Volume 20 - Number 5 - May 2019

[Home](#)

[Content and Archives](#)

[Aims and Scope](#)

[Editorial Board](#)

[Guidance for Authors](#)

[Ethical Guidelines](#)

[Charges](#)

[Membership](#)

[Previously Reviewers](#)

[Conference Events](#)

EDITOR-IN-CHIEF:

Sutarno

EDITORIAL MEMBERS:

English Editors: Graham Eagleton (grahameagleton@gmail.com), Suranto (surantouns@gmail.com); Technical Editor & Banking: Artini Pangastuti (pangastuti_tutut@yahoo.co.id), Solichatun (solichatun_s@yahoo.com); Distribution & Marketing: Rita Rakhmawati (oktia@yahoo.com); Webmaster: Ari Pitoyo (aripitoyo@yahoo.co.id)

MANAGING EDITORS:

Ahmad Dwi Setyawan (unsjournals@gmail.com)

EDITORIAL BOARD (COMMUNICATING EDITORS):

Abd Fattah N. Abd Rabou, Islamic University of Gaza, Palestine
Agnieszka B. Najda, University of Life Sciences in Lublin, Lublin, Poland
Ajay Kumar Gautam, Abhilashi University Mandi, Himachal Pradesh, India
Annisa, Padjadjaran University, Sumedang, Indonesia
Alan J. Lymbery, Murdoch University, Perth, Australia
Bambang Hero Saharjo, Institut Pertanian Bogor, Bogor, Indonesia
Daiane H. Nunes, State University of Londrina, Londrina, Brazil
Darlina Md. Naim, University Sains Malaysia, Penang, Malaysia
Ghulam Hassan Dar, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, Srinagar, India
Hassan Pourbabaee, University of Guilan, Somehsara, Guilan, Iran
Joko Ridho Witono, Center for Plant Conservation-Bogor Botanical Gardens, Indonesian Institute of Sciences, Bogor, Indonesia
Kartika Dewi, Research Center for Biology, Indonesian Institute of Sciences, Cibinong, Bogor, Indonesia
Katsuhiko Kondo, University of Missouri, Columbia, USA
Kusumadewi Sri Yulita, Research Center for Biology, Indonesian Institute of Sciences, Cibinong, Bogor, Indonesia
Livia Wanntorp, Naturhistoriska riksmuseet, Stockholm, Sweden
M. Jayakara Bhandary, Government Arts and Science College, Karwar, Karnataka, India
Mahdi Reyahi-Khoram, Islamic Azad University (Hamadan Branch), Hamadan, Iran
Mahendra Kumar Rai, SGB Amravati University, Maharashtra, India
Mahesh K. Adhikari, Adhikari Niwas, Kathmandu, Nepal
Maria Panitsa, University of Patras, Agrinio, Greece
Mochamad A. Soendjoto, Lambung Mangkurat University, Banjarbaru, Indonesia
Mohamed M.M. Najim, University of Kelaniya, Kelaniya, Sri Lanka
Mohib A. Shah, Nepean Telehealth Technology Centre, Sydney, Australia
Nurhasanah, Mulawarman University, Samarinda, Indonesia
Praptiwi, Research Center for Biology, Indonesian Institute of Sciences, Cibinong, Bogor, Indonesia
Rasool B.Tareen, University of Balochistan, Quetta, Pakistan
Seyed Aliakbar Hedayati, Gorgan University of Agricultural Sciences and Natural Resources, Iran
Seyed Mehdi Talebi, Arak University, Iran
Shahabuddin, Universitas Tadulako, Palu, Indonesia
Shahir Shamsir, Universiti Teknologi Malaysia, Skudai, Malaysia
Shri Kant Tripathi, Mizoram University, Aizawl, India

Sugeng Budiharta, Purwodadi Botanical Gardens, Indonesian Institute of Sciences, Pasuruan, Indonesia

Sugiyarto, Universitas Sebelas Maret, Surakarta, Central Java, Indonesia

Subash C. Santra, University of Kalyani, India

Taufiq Purna Nugraha, Research Center for Biology, Indonesian Institute of Sciences, Cibinong, Bogor, Indonesia

Yosep S. Mau, Nusa Cendana University, Kupang, Indonesia

ISSN 1412-033X | E-ISSN 2085-4722

PUBLISHER: Society for Indonesian Biodiversity

CO-PUBLISHER: Department of Biology, FMNS, Sebelas Maret University Surakarta

FIRST PUBLISHED: 2000

ADDRESS:

Sebelas Maret University

Jl. Ir. Sutami 36A Surakarta 57126. Tel. +62-271-7994097, Tel. & Fax.: +62-271-663375, email: editors@smujo.id

ONLINE: biodiversitas.mipa.uns.ac.id

[Download the Editorial Boards.](#)

[Home](#) | [Archives](#) | [Contact Us](#)

Copyright © Biodiversitas, 2012. All Rights Reserved

[Home](#)[Content and Archives](#)[Aims and Scopes](#)[Editorial Board](#)[Guidance for Authors](#)[Ethical Guidelines](#)[Charges](#)[Membership](#)[Previously Reviewers](#)[Conference Events](#)**Biodiversitas Vol. 20, No. 5, May 2019**

Google Search









 only search Biodiversitas

Table of Content

Cover [\[PDF\]](#)Editorial Board & Guidance for Authors [\[PDF\]](#)

1. Genetic structure and diversity between and within African and American oil palm species based on microsatellite markers
AZIS NATAWIJAYA, SINTHO W ARDIE, MUHAMAD SYUKUR, ISMAIL MASKROMO, ALEX HARTANA, SUDARSONO SUDARSONOO [\[PDF\]](#)
2. Phenotypic plasticity in the ovarium of crested flower of *Hibiscus rosa-sinensis*
SAIFUDIN SAIFUDIN, ANDI SALAMAH [\[PDF\]](#)
3. Phylogenetic study of endophytic fungi associated with *Centella asiatica* from Bengkulu and Malaysian accessions based on the ITS rDNA sequence
NANI RADIASTUTI, DWI NINGSIH SUSILOWATI, HUSHSHILA A BAHALWAN [\[PDF\]](#)
4. Actinomycetes of rhizosphere soil producing antibacterial compounds against Urinary Tract Infection bacteria
PRAMESITA PRAWADIKA APSARI, SRI BUDIARTI, ARIS TRI WAHYUDI [\[PDF\]](#)
5. Species diversity and phylogenetic analysis of *Heliconia* spp. collections of Purwodadi Botanic Garden (East Java, Indonesia) inferred by *rbcL* gene sequences
LIA HAPSARI, TRIMANTO TRIMANTO, DIDIK WAHYUDI [\[PDF\]](#)
6. Changes of ecological wisdom of Sundanese People on conservation of wild animals: A case study in Upper Cisokan Watershed, West Java, Indonesia
SIDIK PERMANA, JOHAN ISKANDAR, PARIKESIT PARIKESIT, TEGUH HUSODO, ERRI N MEGANTARA, RUHYAT PARTASASMITA [\[PDF\]](#)
7. Diversity and inheritance in cowpea (*Vigna unguiculata*) on protein and yield components characters
INDAH PURNAMASARI, SOBIR SOBIR, MUHAMAD SYUKUR [\[PDF\]](#)
8. Population assessment and species distribution modeling of *Paris polyphylla* in Sikkim Himalaya, India
DAWA LHENDUP LEPCHA, ADITYA PRADHAN, DHANI RAJ CHHETRI [\[PDF\]](#)
9. Short Communication: Community structure of prawns (Decapoda: Penaeidae and Palaemonidae) in mangrove inlets of Kuala Langsa, Aceh, Indonesia
TEUKU M FAISAL, HELMY AKBAR, ANDIKA PUTRININGTIAS, SINGGIH AFIFA PUTRA [\[PDF\]](#)
10. Short communication: Endophytic actinobacteria isolated from ginger (*Zingiber officinale*) and its potential as a pancreatic lipase inhibitor and its toxicity
SRI RAHAYU, LENNI FITRI, YULIA SARI ISMAIL [\[PDF\]](#)
11. Growth parameter, mortality, recruitment pattern, and exploitation rate of white shrimp *Penaeus indicus* in northern coastal waters of Western Central Java, Indonesia
SURADI WIJAYA SAPUTRA, ANHAR SOLICHIN, WIWIET TEGUH TAUFANI, SITI RUDIYANTI, NINIEK WIDYORINI [\[PDF\]](#)
12. Genetic diversity in the growth of white jabon (*Neolamarckia cadamba*) provenance-progeny test: Comparing study in the nursery and field
DEDE J SUDRAJAT1, YULIANTI YULIANTI, EVAYUSVITA RUSTAM, ICHSAN SUWANDHI [\[PDF\]](#)
13. The presence of microplastics in the digestive tract of commercial fishes off Pantai Indah Kapuk coast, Jakarta, Indonesia
AYU RAMADHINI HASTUTI, DJAMAR TF LUMBANBATU, YUSLI WARDIATNO [\[PDF\]](#)
14. Correlation among Snpb11 markers, root growth, and physiological characters of upland rice under aluminum stress
MIFTAHUL HUDA FENDIYANTO, RIZKY DWI SATRIO, SUHARSONO SUHARSONO, ARIS TIAHJOLEKSONO, M MIFTAHUDIN [\[PDF\]](#)
15. Agronomical performance of soybean genotypes infected by Cowpea Mild Mottle Virus in various level of nitrogen
ENDRIK NURROHMAN, SITI ZUBAIDAH, HERU KUSWANTORO [\[PDF\]](#)
16. Ethnobotany of traditional medicinal plants used by Dayak Desa Community in Sintang, West Kalimantan, Indonesia
MARKUS IYUS SUPIANDI, SUSRIYATI MAHANAL, SITI ZUBAIDAH, HENDRIKUS JULUNG, BENEDIKTUS EGE [\[PDF\]](#)
17. Diversity of phenotypic characteristics of White Tailed-Yellow Chicken populations reared under free range system in Phitsanulok Province, Thailand
SUPHAWADEE YAEMKONG, TUAN NGUYEN NGOC [\[PDF\]](#)
18. Diversity of mammals in forest patches of Cisokan, Cianjur, West Java, Indonesia
TEGUH HUSODO, PUPUT FEBRIANTO, ERRI NOVIAR MEGANTARA, SYA SYA SHANIDA, MUHAMMAD PAHLA PUJIANTO [\[PDF\]](#)

19. Effect of phytoremediation on the morpho-anatomical characters of some aquatic macrophytes
JONES T NAPALDET, INOCENCIO Jr E BUOT, MACRINA T ZAFARALLA, IRENEO Jr. L LIT, RACHEL C SOTTO
[\[PDF\]](#)
20. Potential of endophytic bacteria from rice root as potassium solvent
WARZATULLISNA WARZATULLISNA, LENNI FITRI, YULIA SARI ISMAIL [\[PDF\]](#)
21. Bioprospecting of polyhydroxyalkanoates-producing bacteria from Indonesian marine environment
WATUMESA A TAN, IRA WIJAYA, TRESNAWATI PURWADARIA [\[PDF\]](#)
22. Hobby and business on trading birds: Case study in bird market of Sukahaji, Bandung, West Java and Splendid, Malang, East Java (Indonesia)
BUDIAWATI S ISKANDAR, JOHAN ISKANDAR, RUHYAT PARTASASMITA [\[PDF\]](#)
23. Some behavior features of a female *Panthera pardus tulliana* during pregnancy and at first parturitions
MAKHAR M ERTUEV, UMAR A SEMENOV [\[PDF\]](#)
24. Short Communication: Identification and characterization of nitrifying bacteria in mud crab (*Scylla serrata*) recirculation aquaculture system by 16S rRNA sequencing
YUNI PUJI HASTUTI, IMAN RUSMANA, KUKUH NIRMALA, RIDWAN AFFANDI, SISKATRIDESEANTI [\[PDF\]](#)
25. Mass production of the entomopathogenic nematode, *Steinernema carpocapsae* on *Tenebrio molitor* and *Spodoptera litura*
HERI PRABOWO, SRI ADIKADARSIH, JANIS DAMAIYANI [\[PDF\]](#)
26. Short Communication: Antagonistic activity of bacteria isolated from coral *Acropora* sp of Karimunjawa Islands, Indonesia against Acroporid White Syndrome
AGUS SABDONO, AGUS TRIANTO, OCKY KARNA RADJASA, DIAH PERMATA WIJAYANTI [\[PDF\]](#)
27. Study of the genus *Curcuma* in Indonesia used as traditional herbal medicines
DYAH SUBOSITI, SLAMET WAHYONO [\[PDF\]](#)
28. Niche partitioning on Sumatran Elephant (*Elephas maximus sumatranus*) and human in space and resources in Riau, Indonesia
YOHANES WISNU SUKMANTORO, HADI SUKADI ALIKODRA, AGUS PRIJONO KARTONO, EFRANSJAH EFRANSJAH [\[PDF\]](#)
29. The potential of indigenous bacteria from oil sludge for biosurfactant production using hydrolysate of agricultural waste
NI'MATUZAHROH, SILVIA KURNIA SARI, IRINE PUSPA NINGRUM, APRILLA DILA PUSFITA, LISA MARJAYANDARI, NASTITI TRIKURNIADEWI, SYAHRIAR NUR MAULANA MALIK IBRAHIM, FATIMAH FATIMAH, TRI NURHARIYATI, TINI SURTININGSIH, HANIF YULIANI [\[PDF\]](#)
30. Isolation of high lipids content microalgae from Wonorejo rivers, Surabaya, Indonesia and its identification using *rbcL* marker gene
TRIONO BAGUS SAPUTRO, KRISTANTI INDAH PURWANI, DINI ERMAVITALINI, ARIF FUAD SAIFULLAH [\[PDF\]](#)
31. Chromosome numbers, RAPD and ISSR profiles of six *Zingiber* species found in Manipur, India
LAISHRAM BIDYALEIMA, RAJKUMAR KISHOR, GURUMAYUM JITENDRA SHARMA [\[PDF\]](#)
32. Evaluation of agronomic performances of rainfed barley double-haploids (DHs) lines under semi-arid conditions
LARBI KARKOUR, MOHAMMED FENNI, DALILA RAMLA, DJOUHER GAAD, ABDELKADER BENBELKACEMH [\[PDF\]](#)
33. The impact of herbivorous insect on leaves of mangrove species *Rhizophora stylosa* and its relation to leaf nutrient level
INDAH TRISNAWATI, MUKHAMMAD MURYONO, ISKA DESMAWATI [\[PDF\]](#)
34. The effect of gamma rays irradiation on diversity of *Musa*
MARIA SERVIANA DUE, ARI SUSILOWATI, AHMAD YUNUS [\[PDF\]](#)
35. Indigenous knowledge of bamboos by Naga community, Tasikmalaya District, West Java, Indonesia
BUDI IRAWAN, RUHYAT PARTASASMITA, NGESTI RAHAYU, TIA SETIAWATI, JOHAN ISKANDAR [\[PDF\]](#)
36. Diversity and characterization of entomopathogenic fungi from rhizosphere of maize plants as potential biological control agents
NOVRI NELLY, MY SYAHRAWATI, HASMIANDY HAMID, TRIMURTI HABAZAR, DWI NASRI GUSNIA [\[PDF\]](#)
37. Short communication: Morphological characteristics of flower and fruit in several rambutan (*Nephelium lappaceum*) cultivars in Serang City, Banten, Indonesia
GUT WINDARSIH, MUHAMMAD EFENDI [\[PDF\]](#)
38. Differences among clinical isolates of *Pseudomonas aeruginosa* in their capability of forming biofilms and their susceptibility to antibiotics
DIDIK WAHYUDI, ABU THOLIB AMAN, NIKEN SATUTI NUR HANDAYANI, ENDANG SUTARININGSIH SOETARTO [\[PDF\]](#)
39. Diversity and distribution of medicinal plants in the Universitas Sumatera Utara Arboretum of Deli Serdang, North Sumatra, Indonesia
RAHMAWATI RAHMAWATI, J B SAMOSIR, R BATUBARA, A RAUF [\[PDF\]](#)
40. Latitudinal species diversity and density of cryptic crustacean (Brachyura and Anomura) in micro-habitat Autonomous Reef Monitoring Structures across Kepulauan Seribu, Indonesia
GESTEN HAZERI, DWI LISTYO RAHAYU, BEGINER SUBHAN, ANDRIANUS SEMBIRING, AJI WAHYU ANGGORO, AHMAD TAUFIK GHOZALI, HAWIS H MADDUPPA [\[PDF\]](#)
41. Land cover change and land use suitability analyses of coastal area in Bantul District, Yogyakarta, Indonesia
SARAH AFSHOLNISSA, ENDANG HERNAWAN, TIEN LASTINI [\[PDF\]](#)
42. Using the modern morphometric approach to determine sexual dimorphism of three medically important flies (Order: Diptera) in Thailand
TANAWAT CHAIPHONGPACHARA, SEDTHAPONG LAOJUN [\[PDF\]](#)



Biodiversitas

Q3 Animal Science and Zoology
best quartile

SJR 2019
0.27

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com" style="border: 1px solid #ccc; padding: 2px 5px; display: inline-block;">https://www.scimagojr.com
```

Human Behaviour and Brain

HumanBaB is a peer-reviewed, open access, interdisciplinary and international journal.

Human Behaviour and Brain

OPEN

Biodiversitas

11

H Index

Country	Indonesia - IIII SIR Ranking of Indonesia
Subject Area and Category	Agricultural and Biological Sciences Animal Science and Zoology Plant Science Biochemistry, Genetics and Molecular Biology Molecular Biology
Publisher	Biology department, Sebelas Maret University Surakarta
Publication type	Journals
ISSN	1412033X, 20854722
Coverage	2014-2020
Scope	"Biodiversitas, Journal of Biological Diversity" or Biodiversitas encourages submission of manuscripts dealing with all biological aspects of plants, animals and microbes at the level of gene, species, and ecosystem.
	Homepage
	How to publish in this journal
	Contact
	Join the conversation about this journal

Human Behaviour and Brain

HumanBaB is a peer-reviewed, open access, interdisciplinary and international journal.

Human Behaviour and Brain

The potential of indigenous bacteria from oil sludge for biosurfactant production using hydrolysate of agricultural waste

NI'MATUZHROH^{1,*}, SILVIA KURNIA SARI¹, IRINE PUSPA NINGRUM¹, APRILLA DILA PUSFITA¹, LISA MARJAYANDARI¹, NASTITI TRIKURNIADEWI¹, SYAHRIAR NUR MAULANA MALIK IBRAHIM¹, FATIMAH¹, TRI NURHARIYATI¹, TINI SURTININGSIH¹, HANIF YULIANI²

¹Department of Biology, Faculty of Science and Technology, Universitas Airlangga, Kampus C Mulyorejo Surabaya 60115, East Java, Indonesia. Tel.: +62-31-5936501, Fax.: +62-31-5926804. *email: nimatuzahroh@fst.unair.ac.id

²Agency for The Assessment and Application of Technology. Jl. Raya Puspitpek, Serpong, Tangerang Selatan 15311, Banten, Indonesia.

Manuscript received: 31 January 2019. Revision accepted: 26 April 2019.

Abstract. Ni'matuzahroh, Sari SK, Ningrum IP, Pusfita AD, Marjayandari L, Trikurniadewi N, Ibrahim SNMM, Fatimah, Nurhariyati T, Surtiningsih T, Yuliani H. 2019. The potential of indigenous bacteria from oil sludge for biosurfactant production using hydrolysate of agricultural waste. *Biodiversitas* 20: 1374-1379. Biosurfactants are amphipathic compounds which are useful in various fields of health, industry, and remediation. Biosurfactants are produced by bacteria that grow in hydrocarbon or sugar substrates. Hydrolysis product of agricultural waste can be used as a biosurfactant production medium. This research aims to obtain biosurfactant producing bacteria from Balongan oil sludge, Indonesia. The ability to grow and produce biosurfactant by indigenous bacteria was tested using a medium of Synthetic Mineral Water (SMW) added by 209.3 ppm of rice straw hydrolysis product (RSHP). The growth of bacteria was evaluated through Total Plate Count (TPC) and biosurfactant production was evaluated through measurement of emulsification activity and surface tension. Six indigenous bacteria were capable to produce biosurfactants in the RSHP. Emulsification activity was not detected, but surface tension reduction was founded. The best biosurfactant was indicated by surface tension value of 53.56 mN/m with TPC value of 20.07 CFU/mL at the 5th day of incubation by BP (1) 5. The indigenous bacteria were identified as *Propionibacterium* BP (1) 1, *Propionibacterium* BP (1) 3, *Bacillus* BP (1) 4, *Corynebacterium* BP (1) 5, *Corynebacterium* BP (1) 8, and *Rothia* BP (1) 6. Utilization of sugar as hydrolysis product of agricultural waste is an innovation of raw materials for biosurfactant production.

Keywords: Agricultural waste, biosurfactant, indigenous bacteria, oil sludge, rice straw

INTRODUCTION

Biosurfactants are amphipathic compounds which are useful in various fields of health, industry, and remediation. Biosurfactants have low toxicity, easy manufacture, and wide application (Elazzazy et al. 2015). Biosurfactants are produced by bacteria that grow in hydrocarbon environments and can be secreted on the surface or outside cells in growth media with noticeable by surface tension reduction and emulsifying activities. (Kosaric 1993). Biosurfactant contains hydrophobic and hydrophilic molecules. Biosurfactant production by bacteria requires carbon nutrients, one of which is a sugar substrate.

The use of agricultural waste as a supplier of carbon sources in the form of sugar substrates as a result of hydrolysis has not been further exposed. The results of hydrolysis of agricultural waste can be used as a medium for biosurfactant production. Organic material in organic waste is a source of micronutrients (Wahyono 2011). The use of agricultural waste has been groundbreaking as an alternative material for biosurfactant production. Utilization of agricultural wastes (barley bran, trimming vine shoots, corn cobs, and *Eucalyptus globulus* chips) for biosurfactant production by bacteria have been done to reduce production costs (Moldes et al. 2007). Rice straw decomposing also has been successfully used for biosurfactant production by *Pseudomonas aeruginosa* BSZ-07 (Quizhuo et al. 2008).

One of agricultural waste that can produce hydrolyzing sugar is rice straw. According to the Indonesian Center for Rice Research, in the East Java region for March 2018, rice yields reached 2,026,739 tons (BB Padi 2018) and produced 12-15 tons per hectare of rice straw. Rice straw contains polysaccharides in lignocellulose. Lignocellulose is an abundant material, renewable resources, and less attention (Peralta et al. 2012). In rice straw, lignocellulose contains 32% cellulose, 24% hemicellulose, and 14% lignin (Chandel et al. 2007 in Novalina, 2014). Rice straw also produces 40-43% of carbons. The high cellulose and hemicellulose content can be used as a result of hydrolysis of rice straw and carbon for biosurfactant-producing bacteria. In this study, *Penicillium* sp. H9 was used to hydrolysis rice straw and its hydrolysis product called rice straw hydrolysis product (RSHP) was used as substrate for biosurfactant production.

Indonesia has a diversity of microorganisms that have the potential to be explored, one of which is a group of bacteria. Hydrocarbonoclastic bacteria in oil sludge are known to produce biosurfactants. Biosurfactant production by indigenous or soil bacteria has been carried out (Amani et al. 2011; Ni'matuzahroh et al. 2017; Lee et al. 2018). This study reveals the ability of indigenous bacteria to grow and produce biosurfactant and knowing the prospect of utilizing RSHP for alternative substrate for biosurfactant production.

MATERIALS AND METHODS

Procedures

Sample collection and bacteria isolation

Oil sludge was obtained from PT. Pertamina, Balongan, West Java, Indonesia. Ten grams of the oil sludge was added to 250 mL Erlenmeyer flasks containing 90 mL distilled water and homogenized with a shaker for 15 minutes. One mL of supernatants were put into a sterile Petri dish and then added 15 mL of Nutrient Agar. This Petri dish was incubated at 30°C for 24 hours on incubator. The bacteria isolates that have been obtained then characterized and purified.

Morphological and physiological characterization of bacteria

All isolated bacteria were characterized macroscopically, microscopically, and physiologically. Macroscopic characterization was observed by the morphological colony of form, color, elevation, edges, and consistency. Microscopic observation of the isolates was performed before 18 hours and after 24 hours to make sure the shape and the presence of Gram. Bacteria biochemical test was using Microbact Kit GNB 12A and 12B according to the protocol. Almost bacteria identified were included in the Gram-positive bacteria groups so that the species name was compared to the Bergey's Manual of Determinative Bacteriology 9th and Cowan and Steel's Manual for Identification of Medical Bacteria Third Edition.

Hydrolysis of rice straw by Penicillium sp.H9

The hydrolysis process was carried out enzymatically by *Penicillium* sp. H9, a microbial collection of Microbiology Laboratory of Universitas Airlangga which was obtained from Alas Purwo National Park. Organic agriculture waste, rice straw, was dried for two days to reduce the water content then delignified mechanically by the grinding process. Rice straw powder and dilute NaOH were mixed in a solid-liquid (1:10) for 60 minutes at 100°C. The mixture was cooled to room temperature and washed to pH 7.0 with distilled water then dried to remove the water. Two gram of pretreatment rice straw was added to 100 mL media Mendel-Sternberg's and autoclaved at 121°C for 15 minutes. The media was cooled to room temperature and added 4% (v/v) spores suspension of *Penicillium* sp. H9 then incubated at shaker incubator for 6 days. Sample was filtrated through sterile filter unit (Minisart NML syringe filter 0.2 µm). The final filtrate was served as RSHP in the following experiments. The reducing sugar content was measured by the Somogyi-Nelson method.

Biosurfactant production

Media Synthetic mineral water used for biosurfactant production in this study was modified from Pruthi and Cameotra (1997). The composition of medium was the following (g/L): (NH₄)₂SO₄ (3.0 g/L), NaCl (10 g/L), MgSO₄.7H₂O (0.2g/L), CaCl₂ (0.01g/L), MnSO₄.H₂O (0.001 g/L), H₃BO₃ (0.001 g/L), ZnSO₄.7H₂O (0.001 g/L), CuSO₄.5H₂O (0.001 g/L), CoCl₂.6H₂O (0.005g/L) dan NaMoO₄.2H₂O (0,001 g/L). AMS buffer consists of (g/50mL): KH₂PO₄ (5 g), K₂HPO₄ (2.62047 g) and Fe (g/50mL) Fe₃O₄ (0.0006 g). The total culture volume was 20 mL and added 1.5 mL of rice straw sugar hydrolysis with the concentration 209.3 ppm. 2% (v/v) bacteria suspension was used for the culture. Microbial culture was incubated for 0, 1, 3, and 5 days in a rotary shaker 120 rpm at room temperature.

Detection of growth and biosurfactant product

Bacteria cultures that incubated during the incubation time were measured for TPC, surface tension (ST), and emulsification activity (EA). Samples of culture medium were centrifuged for 15 minutes at 6000 rpm. 10 ml supernatants were taken for surface tension analysis using Tensiometer Du-Nuoy, Surface tension value was stated in mN/m or dyne/cm. The emulsification activity was obtained from 1 ml of supernatant which added by 1 ml of kerosene, the mixture was homogenized with vortex for 2 minutes, and the emulsification activity can be observed after 1 hour and 24 hours.

RESULTS AND DISCUSSION

Morphological and microscopical characteristics of bacteria

Six morphologically distinct bacteria colonies were isolated from Balongan's oil sludge. Isolates were identified macroscopically, microscopically, and physiologically. Macroscopic and microscopic characters are presented in table 1. Every colony displayed different characters (Figure 1), but microscopic characters of some bacteria showed the similarity of characters. Only one bacterium has rod-shaped and Gram-negative, BP (1) 6 isolates, and other bacteria were identified with rod-shaped and Gram-positive (Figure 2). Isolate BP (1) 4 have a longer rod-shaped compared with the others.

Table 1. Macroscopic and microscopic characters of oil sludge indigenous bacteria from Balongan

Isolates	Macroscopic characters of colony				Microscopic characters of cell		
	Color	Elevation	Edge	Shape	Consistency	Shape	Gram
BP (1) 1	White	Raised	Entire	Circular	Opaque	Rod	Positive
BP (1) 3	White yellowish	Raised	Entire	Circular	Opaque	Rod	Positive
BP (1) 4	White	Flat	Undulate	Irregular	Translucent	Rod	Positive
BP (1) 5	White	Flat	Entire	Circular	Transparent	Rod	Positive
BP (1) 6	White Bone	Flat	Undulate	Irregular	Translucent	Rod	Negative
BP (1) 8	Orange	Flat	Entire	Circular	Transparent	Rod	Positive




Figure 1. Macroscopic morphology of indigenous bacteria from Balongan's oil sludge. A is isolate of BP (1) 1; B is isolate of BP (1) 3; C is isolate of BP (1) 4; D is isolate of BP (1) 5; E is isolate of BP (1) 6; and F is isolate of BP (1) 8




Figure 2. Cell of indigenous bacteria from Balongan's oil sludge. A. Isolate of BP (1) 1; B. Isolate of BP (1) 3; C. Isolate of BP (1) 4; D. Isolate of BP (1) 5; E. Isolate of BP (1) 6; and F is isolate of BP (1) 8

Physiological characteristic of bacteria

Based on differences in microscopic and macroscopic character, physiological characteristics were carried out to obtain the genus name. The physiological test of the bacteria is shown in Table 2.

According to Jaccard's index (Real et al. 1996), the isolates were identified using similarity index (J) values. The isolates were identified as *Propionibacterium* sp. (isolate BP (1) 1) (J = 60%), *Propionibacterium* sp. (isolate BP (1) 3) (J = 60%), *Bacillus* sp. (isolate BP (1) 4) (J = 80%), *Corynebacterium* sp. (isolate BP (1) 5) (J = 75%), *Rothia* sp. (isolate BP (1) 6) (J = 75%), and *Corynebacterium* sp. (isolate BP (1) 8) (J = 51%).

Biosurfactant production

All isolated bacteria were able to grow and produce biosurfactants when grown on synthetic mineral water containing RSHP as the main carbon. This is evidenced by the growth response shown by the TPC are and the decrease in ST of the supernatant culture which indicates the production of biosurfactant. Value of TPC and ST are displayed in figure 2.

Growth response of bacteria had formed after first-day incubation and increased continuously until the end of the incubation. This data shows that the addition of RSHP as a carbon source can be used by indigenous bacteria to grow. Biosurfactant production of isolates BP (1) 1, BP (1) 3, and BP (1) 5 tends increased from the first day of incubation to the end of incubation where all three bacteria are in the exponential phase.

Table 2. The biochemical test of bacteria using microbact GNB 12A and 12B

Type of test	Bacteria isolate code					
	1	3	4	5	6	8
Oxidase	+	+	+	+	+	+
Motility	-	-	-	-	-	-
Nitrate	+	+	-	+	-	-
Lysine	-	+	+	+	+	-
Ornithine	-	-	-	-	-	-
H2S	-	-	-	-	-	-
Glucose	-	-	+	-	-	-
Mannitol	-	-	-	-	-	-
Xylose	-	-	+	-	-	-
ONPG	-	-	-	-	-	-
Indole	-	-	-	-	-	-
Urease	+	-	-	-	-	-
VP	-	-	-	-	-	-
Citrate	-	+	-	+	+	-
TDA	-	-	-	-	-	-
Gelatin	-	-	-	-	-	-
Malonate	-	-	-	-	+	-
Inositol	-	-	-	-	-	-
Sorbitol	-	-	-	-	-	-
Rhamnose	-	-	-	-	-	-
Sucrose	-	-	-	-	-	-
Lactose	-	-	-	-	-	-
Arabinose	-	-	-	-	-	-
Adonitol	-	-	-	-	-	-
Raffinose	-	-	-	-	-	-
Salicin	-	-	-	-	-	-
Catalase	+	+	+	+	+	+
Arginine	-	-	-	-	-	-
After 24 hours	-	-	-	-	-	-
After 48 hours	-	-	-	-	-	-




Figure 2. Total Plate Count and Surface Tension result of culture supernatant during 0, 1, 3 and 5 days incubation by: A. *Propionibacterium* sp. BP (1) 1; B. *Propionibacterium* sp. BP (1) 3; C. *Bacillus* sp. BP (1) 4; D. *Corynebacterium* sp. BP (1) 5; E. *Rothia* BP (1) 6; F. *Corynebacterium* sp. BP (1) 8 was using RSHP as carbon source

Table 3. Growth rate, generation time, TPC and ST result in fifth day incubation time

Isolates	Growth rate (cell/hour)	Generation time (hour)	TPC (CFU/mL)	ST (mN/m)
BP (1) 1	0.41	2.47	19.42±0.02	62.30±4.31
BP (1) 3	0.32	3.15	19.48±0.02	54.15±0.60
BP (1) 4	0.44	2.28	20.12±0.05	66.52±4.61
BP (1) 5	0.33	3.03	20.07±0.04	53.56±2.66
BP (1) 6	0.44	2.30	20.08±0.02	69.72±3.09
BP (1) 8	0.43	2.34	20.22±0.46	65.80±6.62

The highest biosurfactant production was obtained on the fifth-day incubation by isolate BP (1) 5 with a surface tension value of 53.56 ± 2.66 mN/m and 54.15 ± 0.60 mN/m. The pattern of biosurfactant production by BP (1) 5 and BP (1) 3 shows that bacterial growth is in line with the quantity of biosurfactant product. The high value of surface tension is in line with the result of TPC values, but inversely proportional to the generation time of bacteria. Isolate BP (1) 5 takes 0.61 hours to replicate while isolates with the highest generation time only need 0.46 hours to replicate (Table 3).

Discussion

All isolated bacteria from Balongan's oil sludge were identified into four genera as *Propionibacterium* sp., *Bacillus* sp., *Corynebacterium* sp., and *Rothia* sp. based on the results of physiological tests using microbact 12A and 12B. Four genera were able to grow and produced biosurfactants when grown on sugar hydrolysis of rice straw. Generally, the type of biosurfactant produced by *Propionibacterium* sp., *Bacillus* sp., *Corynebacterium* sp., and *Rothia* sp have potential in the process of hydrocarbon degradation. Several studies have reported similar results where the bacterial isolated from hydrocarbon contaminated areas was obtained *Bacillus*, *Corynebacterium*, *Propionibacterium*, and *Rothia* (Calvo et al. 2004; Sette et al. 2007; Sathishkumar et al. 2008; Yang et al. 2017). *Bacillus* produced surfactin, the type of biosurfactant which capable to emulsify the hydrocarbon. In addition, biosurfactant also has the potential as anti-microbial agents and inhibitors biofilm formation. The lipopeptide, biosurfactant types produced by *Bacillus* sp. strain SW9 that was capable of inhibiting biofilm formation in a wide range of bacteria. While, lipopeptide produced by *Propionibacterium freudenreichii* was known to inhibit *Pseudomonas aeruginosa* biofilm production (Wu et al. 2013; Hajfarajollah et al. 2014). The ability to inhibit biofilm formation is owned by Corixyn, the type of biosurfactant, produced by *Corynebacterium xerosis* strain NS. This biosurfactant disrupted biofilm preform of *S. aureus*, *S. mutans*, *P. aeruginosa*, and *E. coli* beside of that, it is also reported this ability as an anti-microbial agent (Dalili et al. 2015). Meanwhile, the type of biosurfactant from the genus *Rothia* has not been reported.

All of indigenous bacteria from oil sludge were able to grow and produce biosurfactant in RSHP. Reducing sugar in hydrolysate of rice straw has the following composition

of 41-43% glucose, 14.8-20.2% xylose, 2.7-4.5% arabinose, 1.8% mannose, and 0.4% galactose (Maiorella 1985; Roberto et al. 2003). Biosurfactant production associated with growth response has been reported for isolate AB-Cr1, where the bacterial growth showed a parallel relationship with biosurfactant production and correlated with the glucose utilization (Ruzniza 2005). The result of this study is similar to Mulligan and Gibbs (1993), biosurfactant compound was a bacterial bioproduct produced during exponential and stationer phases. Biosurfactant production was found parallel to the highest reduction of surface tension which observed during the exponential growth of bacteria (Ruzniza 2005).

Six indigenous bacteria can grow and produce biosurfactants in the RSHP. The most potential isolate was *Corynebacterium* BP (1) 5 with TPC value of 20.07 CFU/mL and ST value of 53.56 mN/m. RSHP have the prospect to be an alternative substrate for biosurfactant production. Moreover, utilization of RSHP can reduce organic waste in the environment.

ACKNOWLEDGEMENTS

Authors would like to thank Faculty of Science and Technology, Universitas Airlangga, Surabaya, Indonesia which has funded the research through the *Penelitian Unggulan Fakultas* grant scheme in 2018 and Suwarni for his aid during this research.

REFERENCES

- Amani H, Haghghi M, Sarrafzadeh MH, Mehrnia MR, Shahmirzaee F. 2011. Optimization of the production of biosurfactant from Iranian indigenous bacteria or the reduction of surface tension and enhanced oil recovery. *Petroleum Sci Technol* 29: 301-311.
- BB Padi. 2018. Luas Panen Padi di beberapa Wilayah. Balai Besar Penelitian Tanaman Padi, Subang. <http://bbpadi.litbang.pertanian.go.id>.
- Barrow GI, Feltham RKA. 1993. *Cowan and Steel's Manual for Identification of Medical Bacteria*. 3rd ed. Cambridge University Press, UK.
- Bergey DH, Holt JG. 1994. *Bergey's Manual of Determinative Bacteriology*. William & Wilkins, Baltimore.
- Calvo C, Toledo FL, González-López J. 2004. Surfactant activity of a naphthalene degrading *Bacillus pumilus* strain isolated from oil sludge. *J Biotechnol* 109 (3): 255-262.
- Dalili D, Amini M., Faramarzi MA, Fazeli MR, Khoshayand MR, Samadi N. 2015. Isolation and structural characterization of Coryxin, a novel cyclic lipopeptide from *Corynebacterium xerosis* NS5 having emulsifying and anti-biofilm activity. *Colloids and Surfaces B: Biointerfaces* 135: 425-432.
- Elazzazy AM, Abdelmoneim TS, Almaghrabi OA. 2015. Isolation and characterization of biosurfactant production under extreme environmental conditions by alkali-halo-thermophilic bacteria from Saudi Arabia. *Saudi J Biol Sci* 22: 466-475.
- Hajfarajollah H, Mokhtarani B, Noghabi KA. 2014. Newly antibacteria and antiadhesive lipopeptide biosurfactant secreted by a probiotic strain, *Propionibacterium freudenreichii*. *Appl Biochem Biotechnol* 174: 2725-2740.
- Real R, Vargas J M. 1996. The probabilistic basis of Jaccard's index of similarity. *Syst Biol* 45 (3): 380-385.
- Kosaric N. 1993. *Biosurfactants: Production, Properties, Application*. Marcel Dekker, Inc, New York.
- Maiorella BL. Ethanol. In: Young M, editor. 1985. *Comprehensive Biotechnology*, Vol. 3. Pergamon Press, Oxford.

- Moldes AB, Torrado AM, Barral MT, Domiáñez JM. 2007. Evaluation of biosurfactant production from various agricultural residues by *Lactobacillus pentosus*. *J Agric Food Chem* 55: 4481-4486.
- Mulligan CN and Gibbs BF. 1993. Factors influencing the economics of biosurfactant. In: Cosaric N (ed.). *Biosurfactant Production, Properties, and Application*. Marcel Dekker Inc. New York
- Ni'matuzahroh, Yuliawatin ET, Kumalasari DP, Trikurniadewi N, Pratiwi IA, Salamun, Fatimah, Sumarsih S, Yuliani H, 2017. Potency of Oil Sludge Indigenous Bacteria from Dumai-Riau in Producing Biosurfactant on Variation of Saccharide Substrates. *Proceeding of International Conference on Green Technology* 8: 339-340.
- Novalina I. 2014. Hydrolysis Enzymatic of Rice Straw by *Penicillium* sp. H9 on Variation of pH and Temperature. [Thesis]. Universitas Airlangga, Surabaya. [Indonesian].
- Peralta-Yahya PP, Zhang F, del Cardayre SB, Keasling JD. 2012. Microbial engineering for the production of advanced biofuels. *Nature* 488: 320-328.
- Pruthi V, Cameotra. 1997. Rapid identification of biosurfactant producing bacterial strain using cell surface hydrophobicity techniques. *Biotechnol Tech.* 11: 671-674.
- Qiuzhuo Z, Weimin C, Juan W. 2008. Stimulatory effects of biosurfactant produced by *Pseudomonas aeruginosa* BSZ-07 on rice straw decomposing. *J Environ Sci* 20: 975-980.
- Roberto IC, Mussatto SI, Rodrigues RCLB. 2003. Dilute-acid hydrolysis for optimization of xylose recovery from rice straw in a semi-pilot reactor. *Industr Crops Prod* 7 (3):171-176.
- Ruzniza, Binti Mohd Zawawi. 2005. Production of Biosurfactant by Locally Isolated Bacteria from Petrochemical Waste. [Thesis]. Faculty of Science. Universiti Teknologi Malaysia. [Malaysian]
- Sathishkumar M, Binupriya AR, Baik, SH, Yun SE. 2008. Biodegradation of crude oil by individual bacterial strains and a mixed bacterial consortium isolated from hydrocarbon contaminated areas. *CLEAN-Soil, Air, Water* 36 (1): 92-96.
- Sette LD, Simioni KC, Vasconcellos SP, Dussan LJ, Neto EV, Oliveira VM. 2007. Analysis of the composition of bacterial communities in oil reservoirs from a southern offshore Brazilian basin. *Antonie Van Leeuwenhoek* 91 (3): 253-266.
- Wahyono S. 2011. Pengolahan Sampah dan Aspek Sanitasi. *Jurnal Teknologi Lingkungan* 2: 113-118. [Indonesian]
- Lee DW, Lee H, Kwon BO, Khim JS, Yim UH, Kim BS, Kim JJ. 2018. Biosurfactant-assisted bioremediation of crude oil by indigenous bacteria isolated from Taean beach sediment. *Environ Pollut* 241: 254-64.
- Wu ZY, Ye CS, Guo F, Zhang SH, Yu X. 2013. Evidence for Broad-Spectrum Biofilm Inhibition by the Bacterium *Bacillus* sp Strain SW9. *Appl Environ Microbiol* 79:1735-1738.
- Yang Q, Zhao H, Du B. 2017. Bacteria and bacteriophage communities in bulking and non-bulking activated sludge in full-scale municipal wastewater treatment systems. *Biochem Eng J* 119: 101-111.