



# Biocatalysis and Agricultural Biotechnology

The Official Journal of the  
International Society of  
Biocatalysis and Agricultural  
Biotechnology (ISBAB)





ScienceDirect

# Biocatalysis and Agricultural Biotechnology

Supports *open access*

[Latest issue](#)   [All issues](#)   ●●●

[Search in this journal](#)

---

## About the journal

[Aims and scope](#)   [Editorial board](#)

---

### Editor-in-Chief

Ching T. Hou

USDA-ARS National Center for Agricultural Utilization Research, Peoria, Illinois, United States

### Associate Editor-in-Chief

Yung-Sheng (Vic) Huang

National Chung Hsing University, Taichung, Taiwan

### Editors

Yeh-Jin Ahn

Sangmyung University Department of Biotechnology, Seoul, Korea, Republic of

Tsunehiro Aki

Hiroshima University, Higashi-Hiroshima, Japan

Richard Ashby

USDA-ARS Eastern Regional Research Center, Wyndmoor, Pennsylvania, United States

Aydin Berenjian

University of Waikato, Hamilton, New Zealand

Milan Certik

Slovak University of Technology, Bratislava, Slovakia

Guanqun (Gavin) Chen

University of Alberta, Edmonton, Alberta, Canada

Yu-Ting Chen

National Chung Hsing University, Taichung, Taiwan

MubarakAli Davoodbasha

B S Abdur Rahman Crescent Institute of Science and Technology School of Life Sciences, Chennai, Tamil Nadu, India

Hooi Ling Foo

Putra Malaysia University Faculty of Biotechnology and Biomolecular Sciences, Serdang, Malaysia

Matthew K. Gilbert

U.S. Department of Agriculture (USDA), Agricultural Research Service (ARS), Food and Feed Safety Research Unit, New Orleans, Louisiana, United States

Abd El-Latif Hesham

Assiut University, Genetics Department, Assiut, Egypt

David Hildebrand

University of Kentucky, Lexington, Kentucky, United States

Masashi Hosokawa

Hokkaido University Faculty of Fisheries Sciences Graduate School of Fisheries Sciences School of Fisheries Sciences, Hakodate, Japan

Lu-Kwang (Luke) Ju

University of Akron, Akron, Ohio, United States

Beom Soo Kim

Chungbuk National University, Cheongju, Korea, Republic of

Igor Kovalchuk

University of Lethbridge, Lethbridge, Alberta, Canada

Adepu Kiran Kumar

Sardar Patel Renewable Energy Research Institute, Vallabh Vidhyanagar, India

Siqing Liu

USDA-ARS National Center for Agricultural Utilization Research, Peoria, Illinois, United States

Thomas A. McKeon

USDA-ARS Western Regional Research Center, Albany, California, United States

Kazuo Miyashita

Hokkaido University, Faculty of Fisheries Sciences, Department Bioresources Chemistry, Hakodate, Japan

Toshihiro Nagao

Osaka Research Institute of Industrial Science and Technology, Osaka, Japan

Madhavan Nampoothiri

National Institute for Interdisciplinary Science and Technology CSIR, Trivandrum, Kerala, India

Jun Ogawa

Kyoto University, Kyoto, Japan

Ramesh N. Patel

SLRP Associates, Biotechnology Consultation, Bridgewater, NJ, United States

Xiao Qiu

University of Saskatchewan, Saskatoon, Saskatchewan, Canada

Ulrich Schörken

University of Applied Sciences Cologne, Koln, Germany

Sengottayan Senthil-Nathan

Manonmaniam Sundaranar University Department of Environmental Science, Tirunelveli, India

Yuji Shimada

Osaka Research Institute of Industrial Science and Technology, Osaka, Japan

Jay Shockey

USDA-ARS Southern Regional Research Center, New Orleans, Louisiana, United States

Pau Loke Show

University of Nottingham - Malaysia Campus Department of Chemical and Environmental Engineering, Semenyih, Malaysia

Lie-Fen Shyur

Agricultural Biotechnology Research Centre Academia Sinica, Taipei, Taiwan

Rajinder Singh

Malaysian Palm Oil Board, Kajang, Malaysia

Vijay Pratap Singh

Chowdhary Mahadev Prasad Degree College, Allahabad, India

Anita Slavica

University Hospital Centre Zagreb, Zagreb, Croatia

Daniel Solaiman

USDA-ARS Eastern Regional Research Center, Wyndmoor, Pennsylvania, United States

Koretaro Takahashi

Hokkaido University Faculty of Fisheries Sciences Graduate School of Fisheries Sciences School of Fisheries Sciences, Hakodate, Japan

Sheng-Yang (David) Wang

National Chung Hsing University, Taichung, Taiwan

Randall Weselake

University of Alberta, Edmonton, Alberta, Canada

Suk Hoo Yoon

Korea Food Research Institute, Wanju-gun, Korea, Republic of

Arash Zibae

University of Guilan, Department of Plant Protection, Rasht, Iran, Islamic Republic of

## Editorial Advisory Board

Casimir Akoh

University of Georgia Department of Food Science & Technology, Athens, Georgia, United States

Yuen May Choo

Malaysian Palm Oil Board, Kajang, Malaysia

Mee-Len Chye

University of Hong Kong, Hong Kong, Hong Kong

Sevim Erhan

USDA-ARS Eastern Regional Research Center, Wyndmoor, Pennsylvania, United States

John Harwood

Cardiff University, Cardiff, United Kingdom

Kiyoshi Hayashi

National Agriculture and Food Research Organization, Tsukuba, Japan

Chieh-Chen Huang

National Chung Hsing University, Taichung, Taiwan

Jei-Fu Shaw

I-Shou University, Kaohsiung City, Taiwan

Ming-Che Shih

Academia Sinica, Taipei, Taiwan

Hajime Taniguchi

National Agriculture and Food Research Organization, Tsukuba, Japan

Andrew Hwei-Jiung Wang

Academia Sinica, Taipei, Taiwan

Teruyoshi Yanagita

Saga University, Saga, Japan

Chang-Hsien Yang

National Chung Hsing University, Taichung, Taiwan

Shyi-Dong Yeh

National Chung Hsing University, Taichung, Taiwan

Gow-Chin Yen, PhD

National Chung Hsing University Department of Food Science, Taichung, Taiwan



[About ScienceDirect](#)

[Remote access](#)

[Shopping cart](#)

[Advertise](#)

[Contact and support](#)

[Terms and conditions](#)

[Privacy policy](#)

We use cookies to help provide and enhance our service and tailor content and ads. By continuing you agree to the **use of cookies**.

Copyright © 2020 Elsevier B.V. or its licensors or contributors. ScienceDirect® is a registered trademark of Elsevier B.V.

ScienceDirect® is a registered trademark of Elsevier B.V.





ScienceDirect

## Biocatalysis and Agricultural Biotechnology

Supports *open access*

[Latest issue](#)   [All issues](#)   ●●●

[Search in this journal](#)

---

### Volume 24

March 2020

[↓](#) [Download full issue](#)

[← Previous vol/issue](#)

[Next vol/issue →](#)

Receive an update when the latest issues in this journal are published

[Sign in to set up alerts](#)

*Review*

---

Review article ○ Abstract only

**Medicinal plants: Treasure trove for green synthesis of metallic nanoparticles and their biomedical applications**

Harish Chandra, Pragati Kumari, Elza Bontempi, Saurabh Yadav

Article 101518



[Purchase PDF](#)    [Article preview](#) 

## Agricultural Biotechnology

---

Research article  Abstract only

Mechanisms of acid-resistant *Rhodopseudomonas palustris* strains to ameliorate acidic stress and promote plant growth

Nguyen Quoc Khuong, Duangporn Kantachote, Phitthaya Nookongbut, Jumpen Onthong, ... Ampaitip Sukhoom

Article 101520

[Purchase PDF](#)    [Article preview](#) 

Research article  Abstract only

Physicochemical factors modulate regeneration and *Agrobacterium*-mediated genetic transformation of recalcitrant indica rice cultivars - ASD16 and IR64

Sathish Sundararajan, Venkatesh Rajendran, Safia Nayeem, Sathishkumar Ramalingam

Article 101519

[Purchase PDF](#)    [Article preview](#) 

Research article  Abstract only

Eco-physiological responses of desert and riverain legume plant species to extreme environmental stress

Zainab G. Ahmed, Usama Radwan, Magdi A. El-Sayed

Article 101531

[Purchase PDF](#)    [Article preview](#) 

Research article  Abstract only

Potential insecticidal activity of *Sarocladium strictum*, an endophyte of *Cynanchum acutum*, against *Spodoptera littoralis*, a polyphagous insect pest

Ashraf S.A. El-Sayed, Ahmed H. Moustafa, Hussein A. Hussein, Aly A. El-Sheikh, ... Gamal A. Enan

Article 101524

[Purchase PDF](#)    [Article preview](#) 

Research article  Abstract only

Synthesis, phytotoxic evaluation and *in silico* studies for the development of novel natural products-inspired herbicides

Ricardo P. Rodrigues, Adriano C.M. Baroni, Carlos A. Carollo, Daniel P. Demarque, ... João M. de Siqueira

Article 101559

[Purchase PDF](#) Article preview 

Research article  Abstract only

*Pseudomonas taiwanensis* (MTCC11631) mediated induction of systemic resistance in *Anthurium andreanum* L against blight disease and visualisation of defence related secondary metabolites using confocal laser scanning microscopy

S. Dhanya, Varghese Sherin, K. Divya, J. Sreekumar, M.S. Jisha

Article 101561

[Purchase PDF](#) Article preview 

Research article  Abstract only

Development of a novel probiotic milk product with enhanced antioxidant properties using mango peel as a fermentation substrate

Gabriela Mourad Vicenssuto, Ruann Janser Soares de Castro

Article 101564

[Purchase PDF](#) Article preview 

*Biocatalysis and Bioprocess*

---

Research article  Abstract only

Composition, thermal behavior and antioxidant activity of pracaxi (*Pentaclethra macroloba*) seed oil obtained by supercritical CO<sub>2</sub>

Gerson Lopes Teixeira, Laércio Galvão Maciel, Simone Mazzutti, Cintia Bernardo Gonçalves, ... Jane Mara Block

Article 101521

[Purchase PDF](#) Article preview 

Research article  Abstract only

*In silico* and *in vitro* comparison of nicotinamide adenine dinucleotide phosphate dependent xylose reductase rossmaan fold in *Debaryomycetaceae* yeast family

Nagarajan Arumugam, Thulasinathan Boobalan, Soorangkattan Saravanan, Muthuramalingam Jothi Basu, ... Thangavel Kavitha

Article 101508

[Purchase PDF](#)    [Article preview](#) 

Research article  Abstract only

Biocatalytic reduction of 5-hydroxymethylfurfural to 2,5-furandimethanol using coconut (*Cocos nucifera* L.) water

Ananda S. Amarasekara, Cristian D. Gutierrez Reyes, Rocio Garcia Obregon

Article 101551

[Purchase PDF](#)    [Article preview](#) 

Research article  Abstract only

A specific, rapid and high-throughput cascade catalytic method for determination of plasma uric acid by using uricase and trivalent peroxidase-mimicking DNAzyme

Zahra Karami, Nasrin Sohrabi, Arastoo Badoei-dalfard

Article 101549

[Purchase PDF](#)    [Article preview](#) 

Research article  Abstract only

Optimization of various process parameters for biodegradation of 4-chlorophenol using Taguchi methodology

Sudhansu Sandhibigraha, Subhasis Mandal, Mayank Awasthi, Tarun kanti Bandyopadhyay, Biswanath Bhunia

Article 101568

[Purchase PDF](#)    [Article preview](#) 

### *Bioactive Agents*

---

Research article  Abstract only

The combined effect of formic acid and Nisin on potato spoilage

Ya'u Sabo Ajingi, Songsirin Ruengvisesh, Pongsak Khunrae, Triwit Rattanarojpong, Nujarin Jongruja

Article 101523

[Purchase PDF](#)    [Article preview](#) 

Research article  Abstract only

Green route for the synthesis of zinc oxide nanoparticles from *Melia azedarach* leaf extract and evaluation of their antioxidant and antibacterial activities

Kayal Vizhi Dhandapani, Devipriya Anbumani, Arumugam Dhanesh Gandhi, Purandaradas Annamalai, ...  
Babujanarthanam Ranganathan

Article 101517

[Purchase PDF](#)    Article preview 

Research article  Abstract only

Mycosynthesis of anticancer drug taxol by *Aspergillus oryzae*, an endophyte of *Tarennasiatica*, characterization, and its activity against a human lung cancer cell line

Gopal Suresh, Dhanasegaran Kokila, Thirunavukarasu Chitrikha Suresh, Subramanian Kumaran, ...  
Arumugam Veera Ravi

Article 101525

[Purchase PDF](#)    Article preview 

Research article  Abstract only

The impact of nitrogen concentrations on production and quality of food and feed supplements from three cyanobacteria and potential application in biotechnology

Ahmed Issa, Esmat Ali, R. Abdel-Basset, M.F. Awad, ... S.A. Hassan

Article 101533

[Purchase PDF](#)    Article preview 

Research article  Abstract only

Metabolite profiling of *Artemisia carvifolia* Buch transgenic plants and estimation of their anticancer and antidiabetic potential

Erum Dilshad, Hammad Ismail, Mubarak Ali Khan, Rosa Maria Cusido, Bushra Mirza

Article 101539

[Purchase PDF](#)    Article preview 

Research article  Abstract only

Combination treatment of MCF-7 spheroids by *Pseudomonas aeruginosa* HI1 levan and cisplatin

Asmaa Ezzat, Walid Fayad, Ahmed Ibrahim, Zeinat Kamel, ... Mona A. Esawy

Article 101526

[Purchase PDF](#)    Article preview 

Research article  Abstract only

Termiticidal activity of chitinase enzyme of *Bacillus licheniformis*, a symbiont isolated from the gut of *Globitermes sulphureus* worker

Nurul Akmar Hussin, Abdul Hafiz Ab Majid

Article 101548

[Purchase PDF](#)    Article preview 

Short communication  Abstract only

Antiparasitic activity of *Eichhornia crassipes* leaves extract

Somia M. Elagib

Article 101556

[Purchase PDF](#)    Article preview 

Research article  Abstract only

Improving oxidative and microbial stability of beef using Shahri Balangu seed mucilage loaded with Cumin essential oil as a bioactive edible coating

Behrooz Alizadeh Behbahani, Mohammad Noshad, Hossein Jooyandeh

Article 101563

[Purchase PDF](#)    Article preview 

Research article  Abstract only

Cytological effects of herbicide alachlor in somatic cells of maize (*Zea mays* L.) and soybean (*Glycine max* Merrill.)

N.K. Hemanth Kumar, Shobha Jagannath

Article 101560

[Purchase PDF](#)    Article preview 

## Industrial Enzyme


---

Research article  Abstract only

Statistical modelling and optimization of protease production by an autochthonous *Bacillus aryabhatai* Ab15-ES: A response surface methodology approach

Adegoke Isiaka Adetunji, Ademola Olufolahan Olaniran

Article 101528

[Purchase PDF](#)    Article preview 

Research article  Abstract only

## The production of $\beta$ -mannanase from *Kitasatospora* sp. strain using submerged fermentation: Purification, characterization and its potential in mannoooligosaccharides production

Yopi, Nanik Rahmani, Siti Amanah, Pugoh Santoso, Puspita Lisdiyanti

Article 101532

[↓ Purchase PDF](#)    Article preview

Research article  Abstract only

## Enzymatic production of methyl esters from low-cost feedstocks

Guilherme Martinez Mibielli, Ana Paula Fagundes, Letícia Renata Bohn, Matheus Cavali, ... José Vladimir Oliveira

Article 101558

[↓ Purchase PDF](#)    Article preview

Research article  Abstract only

## A novel lipolytic yeast *Meyerozyma guilliermondii*: Efficient and low-cost production of acid and promising feed lipase using cheese whey

Adriana Knob, Simone Cristine Izidoro, Lorena Tigre Lacerda, André Rodrigues, Vanderlei Aparecido de Lima

Article 101565

[↓ Purchase PDF](#)    Article preview

### Bioenergy

---

Research article  Abstract only

## Rubber seed oil extraction: Effects of solvent polarity, extraction time and solid-solvent ratio on its yield and quality

Chiazor Faustina Jisieike, Eriola Betiku

Article 101522

[↓ Purchase PDF](#)    Article preview

Research article  Abstract only

## Exploring *Pongamia* seed cake hydrolysate as a medium for microbial lipid production by *Aspergillus ochraceus*

Harshitha Madhusoodan Jathanna, Chandrayan Vaman Rao, Louella Concepta Goveas

Article 101543

[Purchase PDF](#)    Article preview 

*Environmental*

---

Research article  Abstract only

### Enhanced textile wastewater treatment by a novel biofilm carrier with adsorbed nutrients

Jéssica Mulinari, Cristiano José de Andrade, Heloísa de Lima Brandão, Adriano da Silva, ... Antônio Augusto Ulson de Souza

Article 101527

[Purchase PDF](#)    Article preview 

Research article  Abstract only

### Antioxidant capacity and phytochemical characterization of *Spathodea campanulata* growing in different climatic zones in Brazil

Valter H.M. Santos, Igor O. Minatel, Giuseppina P.P. Lima, Regildo M.G. Silva, Chung-Yen O. Chen

Article 101536

[Purchase PDF](#)    Article preview 

Research article  Abstract only

### L-arginine amidinohydrolase by a new *Streptomyces* isolate: Screening and statistical optimized production using response surface methodology

Mohamed Abdelraof, Mostafa M. Abo Elsoud, Mohsen Helmy Selim, Amany Ahmed Hassabo

Article 101538

[Purchase PDF](#)    Article preview 

Research article  Abstract only

### Larvicidal activity of the methanolic, hydroethanolic and hexanic extracts from *Acmella oleracea*, solubilized with silk fibroin, against *Aedes aegypti*

Inana F. Araújo, Hellen A. Loureiro, Victor H.S. Marinho, Fernando B. Neves, ... Irlon M. Ferreira

Article 101550

[Purchase PDF](#)    Article preview 

Research article  Abstract only

## A feasible scale-up production of *Sporosarcina pasteurii* using custom-built stirred tank reactor for in-situ soil biocementation

Armstrong I. Omoregie, Enzo A. Palombo, Dominic E.L. Ong, Peter M. Nissom

Article 101544

[↓ Purchase PDF](#)    Article preview [✓](#)

Research article  Abstract only

## Development of regression model for bacteriocin production from local isolate of *Lactobacillus acidophilus* MS1 using Box-Behnken design

Mahwish Salman, Muhammad Shahid, Tanzila Sahar, Shazia Naheed, ... Arif Nazir

Article 101542

[↓ Purchase PDF](#)    Article preview [✓](#)

Research article  Abstract only

## Analysis of remediation potential of whole bacterial cells on wastewater decolourisation and detoxification

Rabia Nisar, Burarah Arooj, Bushra Muneer, Roquyya Gul, Mahjabeen Saleem

Article 101557

[↓ Purchase PDF](#)    Article preview [✓](#)

Research article  Abstract only

## Biosynthesis of silver nanoparticles using *phyllanthus emblica* fruit extract for antimicrobial application

R. Renuka, K. Renuka Devi, M. Sivakami, T. Thilagavathi, ... K. Kaviyarasu

Article 101567

[↓ Purchase PDF](#)    Article preview [✓](#)

*Lignocellulases : from Isolation to Structure-Function Mechanism and Industrial Application; edited by Ni Nyoman Tri Puspaningsih, Kazuo Sakka, Muhammad Mukram Mahamed Mackeen and Rosli Md Illias*

---

Research article  Abstract only

## Extracellular hydrolytic enzyme activities of indigenous actinomycetes on pretreated bagasse using choline acetate ionic liquid

Heri Satria, Yandri, Nurhasanah, Suripto Dwi Yuwono, Dian Herasari

Article 101503



[Purchase PDF](#)    [Article preview](#) 

Research article  Abstract only

### Supplementation of fermented coffee-peel flour to increase high-density lipoprotein (HDL) cholesterol, docosahexaenoic acids (DHA) and eicosapentaenoic acids (EPA) deposition in tilapia fillet

Pingky D. Fitria, Muhamad Amin, Widya P. Lokapirnasari, Mirni Lamid

Article 101502

[Purchase PDF](#)    [Article preview](#) 

Research article  Abstract only

### Antimicrobial activity of biosurfactants produced by actinomycetes isolated from rhizosphere of Sidoarjo mud region

Achmad Arifiyanto, Tini Surtiningsih, Ni'matuzahroh, Fatimah, ... Nur Hidayatul Alami

Article 101513

[Purchase PDF](#)    [Article preview](#) 

Research article  Abstract only

### Application of enzyme cocktails from Indonesian isolates to corncob (*Zea mays*) waste saccharification

One Asmarani, Artati Dian Pertiwi, Ni Nyoman Tri Puspaningsih

Article 101537

[Purchase PDF](#)    [Article preview](#) 

Research article  Abstract only

### Substitution of commercial feed with phytase-fermented rice bran and turmeric flour to increase EPA, DHA, and protein depositions in broiler meat

Muhammad A. Al-Arif, Sunaryo H. Warsito, Muhamad Amin, Mirni Lamid

Article 101535

[Purchase PDF](#)    [Article preview](#) 

Research article  Abstract only

### Bioconversion of agricultural waste hydrolysate from lignocellulolytic mold into biosurfactant by *Achromobacter* sp. BP(1)5

Ni'matuzahroh, Silvia Kurnia Sari, Nastiti Trikurniadewi, Syahriar Nur Maulana Malik Ibrahim, ... Fatimah

Article 101534

[Purchase PDF](#)    [Article preview](#) 

## Research

---

Research article  Abstract only

### Separation and partial purification of collagenolytic protease from peacock bass (*Cichla ocellaris*) using different protocol: Precipitation and partitioning approaches

Vagne de Melo Oliveira, Márcia Nieves Carneiro da Cunha, Caio Rodrigo Dias de Assis, Juanize Matias da Silva Batista, ... Ana Lúcia Figueiredo Porto

Article 101509

[Purchase PDF](#)    [Article preview](#) 

Research article  Abstract only

### Bioprocess for hydrolysis of galacto-oligosaccharides in soy molasses and tofu whey by recombinant *Pseudomonas chlororaphis*

Daniel K.Y. Solaiman, Richard D. Ashby, Nicole V. Crocker

Article 101529

[Purchase PDF](#)    [Article preview](#) 

## *Application of environmental biotechnology for sustainable future : edited by Habibollah Younesi*

---

Research article  Abstract only

### Evaluation on biodiesel cold flow properties, oxidative stability and enhancement strategies: A review

Chee Bing Sia, Jibrail Kansedo, Yie Hua Tan, Keat Teong Lee

Article 101514

[Purchase PDF](#)    [Article preview](#) 

Research article  Abstract only

### Optimization of enzymatic hydrolysis for acid pretreated date seeds into fermentable sugars

Hekma Salem Hasan Ba Hamid, Ku Syahidah Ku Ismail

Article 101530

[Purchase PDF](#)    [Article preview](#) 

Research article  Abstract only

## Bioadsorption and enzymatic biodecolorization of effluents from ethanol production plants

Zhila Ziaei-Rad, Mojdeh Nickpour, Mehrdad Adl, Mohammad Pazouki

Article 101555

[Purchase PDF](#) Article preview

---

*Marine Natural Products as Pharmaceutical Agents; edited by Ganesh Kumar Vijayakumar, Stalin Dhas T, Praveen Ramasamy, Francis Borgio J and Antony Samrot Vincent*

Research article  Abstract only

## Survey and isolation of marine cyanobacteria from eastern coast of India as a biodiesel feedstock

V.S. Uma, Dineshbabu Gnanasekaran, Uma Lakshmanan, Prabaharan Dharmar

Article 101541

[Purchase PDF](#) Article preview

---

*Functional Lipids for human; edited by In-Hwan Kim, Byung Hee Kim & Hak-Ryul Kim*

Research article  Abstract only

## Optimal production of 7,10-epoxy-octadeca-7,9-dienoic acid from 7,10-dihydroxy-8(*E*)-octadecenoic acid by heat treatment

Joel B. Ellamar, In-Hwan Kim, Ching T. Hou, Hak-Ryul Kim

Article 101545

[Purchase PDF](#) Article preview

Research article  Abstract only

## Peroxidase of *Cedreia fissilis* leaves: Biochemical characterization and toxicity of enzymatically decolorized solution of textile dye Brilliant Sky-Blue G

Wikeff Fritzke, Elian Gabriel Salla, Margarete Dulce Bagatini, Beatriz da Silva Rosa Bonadiman, ... Aniela Pinto Kempka

Article 101553

[Purchase PDF](#) Article preview

Erratum Full text access

Corrigendum to facile biofabrication, characterization, evaluation of photocatalytic, antipathogenic activity and *in vitro* cytotoxicity of zinc oxide nanoparticles

Krishnasamy Sekar Rajkumar, Sridhar Arun, Manikandan Dinesh Babu, Perumalsamy Balaji, ...

Ramasamy Thirumurugan

Article 101477

[Download PDF](#)

---

[< Previous vol/issue](#)

[Next vol/issue >](#)

ISSN: 1878-8181

Copyright © 2020 Elsevier Ltd. All rights reserved



[About ScienceDirect](#)

[Remote access](#)

[Shopping cart](#)

[Advertise](#)

[Contact and support](#)

[Terms and conditions](#)

[Privacy policy](#)

We use cookies to help provide and enhance our service and tailor content and ads. By continuing you agree to the **use of cookies**.

Copyright © 2020 Elsevier B.V. or its licensors or contributors. ScienceDirect® is a registered trademark of Elsevier B.V.

ScienceDirect® is a registered trademark of Elsevier B.V.

RELX™

× ⓘ

**Free English Writing Tool**  
 Grammarly makes sure everything you type is effective and mistake-free. Try now

DOWNLOAD Grammarly

# Biocatalysis and Agricultural Biotechnology

# 30

H Index

**Country** Netherlands -  SIR Ranking of Netherlands

**Subject Area and Category** [Agricultural and Biological Sciences](#)  
[Agronomy and Crop Science](#)  
[Food Science](#)

[Biochemistry, Genetics and Molecular Biology](#)  
[Biotechnology](#)

[Chemical Engineering](#)  
[Bioengineering](#)

[Immunology and Microbiology](#)  
[Applied Microbiology and Biotechnology](#)



**Publisher** Elsevier BV

**Publication type** Journals

**ISSN** 18788181

**Coverage** 2012-2020

**Scope** Biocatalysis and Agricultural Biotechnology is the official journal of the International Society of Biocatalysis and Agricultural Biotechnology (ISBAB). The journal publishes high quality articles especially in the science and technology of biocatalysis, bioprocesses, agricultural biotechnology, biomedical biotechnology, and, if appropriate, from other related areas of biotechnology. The journal will publish peer-reviewed basic and applied research papers, authoritative reviews, and feature articles. The scope of the journal encompasses the research, industrial, and commercial aspects of biotechnology, including the areas of: biocatalysis; bioprocesses; food and agriculture; genetic engineering; molecular biology; healthcare and pharmaceuticals; biofuels; genomics; nanotechnology; environment and biodiversity; and bioremediation.


-  [Homepage](#)
- [How to publish in this journal](#)
-  [Join the conversation about this journal](#)

× ⓘ


**Free English Writing Tool**  
 Grammarly makes sure everything you type is effective and mistake-free. Try now

DOWNLOAD Grammarly


Quartiles




SJR




Citations per document




Total Cites Self-Cites




External Cites per Doc Cites per Doc




% International Collaboration




Citable documents Non-citable documents



Cited documents Uncited documents



Biocatalysis and Agricultural Biotechnology



← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com" style="display: inline-block; border: 1px solid black; padding: 2px 5px;">https://www.scimagojr.com
```

# Journal Pre-proof

Antimicrobial activity of biosurfactants produced by actinomycetes isolated from rhizosphere of Sidoarjo mud region

Achmad Arifiyanto, Tini Surtiningsih, Ni'matuzahroh, Fatimah, Dyah Agustina, Nur Hidayatul Alami



PII: S1878-8181(19)31290-3

DOI: <https://doi.org/10.1016/j.bcab.2020.101513>

Reference: BCAB 101513

To appear in: *Biocatalysis and Agricultural Biotechnology*

Received Date: 15 September 2019

Revised Date: 22 December 2019

Accepted Date: 22 January 2020

Please cite this article as: Arifiyanto, A., Surtiningsih, T., Ni'matuzahroh, , Fatimah, , Agustina, D., Alami, N.H., Antimicrobial activity of biosurfactants produced by actinomycetes isolated from rhizosphere of Sidoarjo mud region, *Biocatalysis and Agricultural Biotechnology* (2020), doi: <https://doi.org/10.1016/j.bcab.2020.101513>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Ltd.

### Credit Author Statement

**Achmad Arifiyanto:** Conceptualization, Writing-Original draft preparation, Software. **Tini Surtiningsih:** Data curation and Visualization. **Ni'matuzahroh:** Methodology and Investigation. **Fatimah:** Supervision and Validation **Dyah Agustina:** Project administration. **Nur Hidayatul Alami:** Resources.

Journal Pre-proof



1 Green Industry Research Symposium for Sustainable Development 2019

2 Lignocellulases: from Isolation to Structure-Function Mechanism and Industrial Application

3 In Collaboration with Biocatalys and Agricultural Biotechnology

4 (Scimago Q2, Indexed by Scopus)

5  
6 **Antimicrobial Activity of Biosurfactants Produced by Actinomycetes**  
7 **Isolated from Rhizosphere of Sidoarjo Mud Region**

8 Achmad Arifiyanto<sup>a\*</sup>, Tini Surtiningsih<sup>b</sup>, Ni'matuzahroh<sup>bc</sup>, Fatimah<sup>bc</sup>, Dyah Agustina<sup>c</sup>, Nur  
9 Hidayatul Alami<sup>d</sup>,

10 <sup>a</sup>Department of Biology, Universitas Lampung, Jl. Prof. Sumantri Borojonegoro No. 1, Bandarlampung, 35145, Indonesia

11 <sup>b</sup>Department of Biology, Universitas Airlangga, Kampus C Mulyorejo, Surabaya, 60115, Indonesia

12 <sup>c</sup>Research Center for Bio-Molecule Engineering, Universitas Airlangga, Kampus C Mulyorejo Surabaya, 60115, Indonesia

13 <sup>d</sup>Department of Biology, Institut Teknologi Sepuluh Nopember, Sukolilo, Surabaya, 60115, Indonesia  
14

---

15 **Abstract**

16 A unique location with a unique environment attracted many researchers to explore bacteria as an isolate collection especially for  
17 actinomycetes group because of the potential of their antimicrobial metabolites. Information on antimicrobial activity by  
18 Streptomyces from the Sidoarjo mud area is limited. Biosurfactant was also one of the metabolites produced by the group of  
19 actinomycetes. Biosurfactant screening was carried out by growing isolates in a series of tests in the hemolytic, tributyrin agar,  
20 surface tension, and emulsification activity. They fermented on glycerol, n-hexane and molasses substrates for 7 days. The  
21 antimicrobial activity of organic extract evaluated using a paper disc diffusion method against E. coli ATCC 25922 and S. aureus  
22 ATCC 6538P. The results of biosurfactant screening tests showed differences in surface tension and varying emulsification activities  
23 in the isolates obtained. Supernatants produced by biosurfactants have been shown to be able to show antimicrobial activity as  
24 indicated by the formation of inhibitory zones associated with pathogenic growth. AF1 as a selected isolates identified as  
25 *Streptomyces* sp. MN394821 using 16S rDNA gene analysis. It has 100% similarity with accession-coded strain KY236015.1 and  
26 98% with accession coded *Streptomyces rochei* AB1 strain GU434672.1.

27  
28 *Keywords:* Antibiotic; Active compound; Biosurfactant; Streptomyces

---

29 **1. Introduction**

30 Microbes are found to live freely or stick to the surface of soil particles, but most soil bacteria interact with plant  
31 roots known as the rhizosphere. The main reason was the increasing rate of availability of dissolved organic compounds  
32 derived from exudation of plant roots [1]. The location around Sidoarjo mudflow was once a fertile agricultural land, the  
33 environmental conditions at that location are also interesting to be investigated further about the potential of  
34 microorganisms in these habitats. A number of studies using the microflora of these habitats have been widely reported.  
35 Poernomo [2] interested in proteolytic enzymes, while Habibie [3] examined the potential of xylanase, Prachayo [4]

\*Corresponding author.

E-mail address: achmad.arifiyanto@fmipa.unila.ac.id

36 identified potential bacteria to inhibit pathogenic molds in rice, and overall informs the thermophilic ability of bacteria  
37 isolated, as the research focus by Dagdag [5].

38 Actinomycetes or Order Actinomycetales are members of the Actinobacteria group. Actinomycetes belong to the  
39 group of Gram positive bacteria, with filaments characteristic. He plays an important role in various industries because  
40 of his ability produced a number of metabolite compounds that vary in structure and function. Many metabolites by  
41 actinomycetes are reported to have antagonistic abilities against bacteria, yeast, and mold. For this reason, many  
42 actinomycetes are developed and used as medicinal ingredients in the prevention of various diseases. The search for  
43 actinomycetes metabolites develops from exploration of certain regions, unique environments, variations in isolation  
44 methods, innovation to manipulation of genetic material [6]. Most microorganisms, including actinomycetes, are known  
45 to produce active metabolites which can reduce surface tension. This compound itself is better known as surfactant. A  
46 number of biosurfactant groups are known to have antimicrobial activity, including glycolipids, lipopeptides &  
47 lipoproteins [7]. On the other hand, actinomycetes bacteria are spread over a wide habitat. It can be found in various  
48 types of soil samples, fresh and marine waters, mangrove ecosystems [8], thermophilic and alkaliphilic [9], acidophilic  
49 [10] and halophilic [11] or even though at Antarctic region [12]. Therefore it is also called the extremophilic group.  
50 Different ecological environments, with the surrounding physical, chemical and biological conditions often play a role  
51 in the discovery of new types of bacteria and metabolites.

52 The actinomycetes study in Indonesia itself is being carried out. Susilowati [13] collected actinomycetes in various  
53 agricultural and plantation areas on the islands of Java, Sumatra, Sulawesi, and NTT by focusing isolation in the soil's  
54 rhizosphere. Nurkanto [14] explored land actinomycetes in Ternate. Meanwhile there is no information about  
55 actinomycetes isolated from Sidoarjo mud.  
56

## 57 **2. Materials and Methods**

### 58 *2.1. Sampling and isolation of actinomycetes*

59 Soil samples taken using soil borers at a depth of 30 cm through line transect method around six burst  
60 embankments (112.70487,-7.52393) (112.70325,-7.52912) (112.70326,-7.53282) (112.70355,-7.53272) (112.70734,-  
61 7.51744) (112.71107,-7.51237). Isolation of actinomycetes was carried out using the dry-heating method. The soil dried  
62 in the oven for approximately 8 days at 70°C. Soil samples crushed and filtered with a 3-5 mm sieve. Then 1 g of soil  
63 was taken, suspended in 9 mL of sterile water, shaken in 2 minutes, and left for 1 minute. They diluted fully for 10-8  
64 dilutions. The last 3 series of dilutions are 10-6, 10-7, and 10-8 also the first dilution is taken as much as 0.1 mL and  
65 spread on a Petri dish containing Starch Casein Agar or SCA (Himedia®) medium which is added with griseofulvin and  
66 chloramphenicol 0.05 ppm. The surface of the medium is flattened using a spatula. The Petri dish was then incubated  
67 for 1-3 weeks at 70°C.

68 Isolates were observed microscopically and accompanied by Gram staining to determine whether or not the  
69 actinomycetes group was present. Storage of isolates was carried out on sloping SCA media and as stock culture and  
70 work culture. Long-term storage of o them stored using liquid drying method with cryoprotective media whose  
71 composition consisted of ribitol or adenitol 1.5 g; phosphate buffer pH 7.2 100 mL, and monosodium glutamate 3 g.

### 72 *2.2. Hemolytic test*

73 Isolate is grown on Blood agar medium. A total of 1.5 mL of fresh sterile lamb blood was pipetted into an  
74 Erlenmeyer flask containing 100 ml of mineral-agar media and then homogenized. The blood medium poured and kept  
75 sterile until it got frozen and solid. Biosurfactant potential bacterial strains were scratched on fresh blood media. It was  
76 incubated for 1-2 days at 37°C. It was checked for the zone of hemolysis and color changes formed from the activity of  
77 the test bacteria

### 78 *2.3. Lipase test*

79 Lipase activity was measured by growing bacterial isolates on agar tributyrin medium. This medium is made by  
80 adding 10 mL of tributyrin (Himedia®) to 990 ml of agar solution (5 g of peptone, 3 g of yeast extract, and 15 g of  
81 agar) and then heated and sterilized in an autoclave at 121°C. This medium should not be poured on disposable Petri  
82 dishes or other tools made of plastic. It affected the objectivity in assessing the ability of bacteria in lipase activity tests,  
83 because plastic included as a hydrocarbon group that was sensitive to the lipase enzyme.  
84  
85  
86

#### 87 2.4. Surface tension test

88 Surface tension measurement conducted using the Kruss K100 tensiometer (Kruss GmbH, Hamburg, Germany)  
89 using the Du Nouy ring method. Measurements were made using crude surfactant previously obtained. It tested at room  
90 temperature (3x replication used to improve measurement accuracy and take an average). The formula below referred to  
91 [15],

$$92 \gamma_2 = \gamma_1 \frac{n_1 \rho_1}{n_2 \rho_2} \quad (1)$$

93  
94  
95  
96  
97  $\gamma_1$  = water surface tension, 72 dyne / cm,  $\gamma_2$  = biosurfactant surface tension,  $\rho_1$  = water density,  $\rho_2$  = biosurfactant  
98 density,  $n_1$  = number of drops of water, and  $n_2$  = number of drops of biosurfactant solution.

99  
100  
101 <sup>1</sup> Chakraborty *et al.*, 2015.  
102

#### 103 2.5. Index of Emulsification

104 The ability of biosurfactant emulsifies liquid of hydrocarbons determined using kerosene. Bacterial cultures were  
105 inoculated on a liquid production medium and incubated for 24 hours. Bacterial cell-free supernatant had obtained  
106 through centrifugation at 20.000 G for 10 minutes and homogenized with hydrocarbons on the vortex machine at high  
107 speed for 2 minutes. Index of emulsification counted from height of fluid layer emulsion divided by the entire system  
108 height multiplied by 100. The stability measured after one hour. The emulsification activity of biosurfactant was  
109 determined by measuring the emulsification index (E24 or EI). It can be calculated using the following formula:

$$110 EI = \frac{\text{Total height of the emulsion layer}}{\text{Total height of the system}} \times 100 \quad (2)$$

111  
112  
113 <sup>2</sup> Chakraborty *et al.*, 2015.  
114

#### 115 2.6. Antimicrobial activity test

116 Antimicrobial activity tested using agar disc diffusion test at Mueller-Hinton agar (MHA; Himedia®). *E. Coli*  
117 ATTCC (-), and *Staphylococcus aureus* ATTCC (+) cultured overnight in Mueller-Hinton broth at 37° C. Optical  
118 Density broth cultures were adjusted to 0.1 or equivalent to 108 CFU mL<sup>-1</sup> inoculum (according to McFarland turbidity  
119 standards). About 500 µL inoculated on Petri dish then the MHA medium (10mL) was poured on the surface and spread  
120 uniformly. It air dried for 10 minutes under laminar sterile air flow. Biosurfactant extracted using chloroform: methanol  
121 (3:1/v:v). Sterile Whatman disc filter (5 mm in diameter) dripped with 10 µL of a different solution of biosurfactant in  
122 methanol aseptically on the surface of the MHA agar. A disc dipped by methanol used as a negative control. The dish is  
123 incubated in 37°C for 24 hours. The inhibition zone (mm) around the well and discs measured triplicate re regarding the  
124 antibiotic zone scale (Himedia®).

#### 125 2.7. Fourier Transform Infrared Spectroscopy (FTIR)

126 FTIR spectroscopy (8201PC Shimadzu, Japan) was carried out using crude biosurfactant (extracted using  
127 chloroform: methanol (3:1/v:v)). The FTIR spectrum, with a resolution of 4 cm<sup>-1</sup>, was collected from 500 to 4000 cm<sup>-1</sup>.  
128 The basic functional groups of the biosurfactant were analyzed

#### 129 2.8. Isolate identification

130 Suspected isolates of actinomycetes were identified by morphological, physiological, biochemical [16] and  
131 molecular approaches. Molecular identification of bacteria suspected of actinomycetes tested using primers 27F-  
132 5'AGAGTTTGATCCTGGCTCAG-3 'and 149R-5'GGTTACCTTGTTACGACTT3'. PCR was carried out by mixing  
133 2.5 µL 10X PCR buffer, 0.5 µL mixtures of dNTP, 1.0 µL of each PCR primer, 0.5 µL of Taq DNA polymerase, and  
134 1.0 µL of each template DNA. Initial denaturation at 94°C for 25 seconds, followed by denaturation at 94°C for 10  
135 seconds for 35 cycles, annealing at 46°C for 30 seconds, elongation at 72°C for 1.5 minutes, then final extension at

136 72°C for 10 minute. The PCR results were then further sequenced and the results were compared with other bacterial  
 137 sequences at Genbank.  
 138

### 139 3. Results and Discussion

#### 140 3.1. Biosurfactant screening

141 Biosurfactant is a variety of groups of active compounds produced by various types of microorganisms. Has a polar  
 142 and non-polar group that is able to separate water from oil or air surface with water so that it can reduce interface  
 143 tension or surface tension. Because of this ability, biosurfactants are widely offered as a solution to overcome waste oil  
 144 pollution [17]. A number of biosurfactants are known as antibacterial, antifungal and antiviral agents. Actinomycetes  
 145 are a large group of actinobacteria members known for their ability in antibiotic activity and many are reported to be  
 146 able to produce biosurfactants.

##### 147 3.1.1. Hemolytic activity

148 Hemolytic reaction on the Blood agar plate was observed by lifting Petri to the light source, the results of hemolytic  
 149 activity can be seen in table 1. There are 3 types of hemolysis namely beta hemolysis ( $\beta$ ), alpha hemolysis ( $\alpha$ ), and  
 150 gamma hemolysis ( $\gamma$ ). Beta hemolysis ( $\beta$ ) or commonly called total hemolysis is defined as the lysis of all red blood  
 151 cells. A clear zone, close to the color of the transparency of the base media and surround the colony. Many species of  
 152 bacteria produce toxins or poisons that can destroy red blood cells. Some species display various levels of beta  
 153 hemolysis. The hemolysis alpha ( $\alpha$ ) also called partial hemolysis is a decrease in red blood cell hemoglobin for  
 154 methemoglobin in the medium around the colonies. This causes changes in green or brown in the medium. Color can be  
 155 likened to "bruising" cells. Microscopic examination of alpha-hemolyzed red blood cells shows that the cell membrane  
 156 is intact, so the cell is not really lysis. Gamma hemolysis ( $\gamma$ ) is also called non hemolysis. Gamma shows a lack of  
 157 hemolysis, because there is no reaction in the surrounding medium.

158 Table 1: Hydrolysis results for actinomycetes grown in blood agar medium  
 159


No	Isolate	Hydrolysis results	
1	AA1	+	$\alpha$
2	AA2	-	$\gamma$
3	AB1	+	$\alpha$
4	AB8	+	$\alpha$
5	AF1	+	$\beta$
6	AF2	+	$\beta$
7	AF3	+	$\alpha$
8	AF4	+	$\alpha$
9	AF5	-	$\gamma$
10	AF6	+	$\alpha$
11	AG1	-	$\gamma$
12	AG2	+	$\alpha$

160 From the results of the study as presented in Table 1, there are 4 isolates that are negative for hemolytic activity, namely  
 161 AA2, AF3, AF5, and AG1. While 8 other isolates were categorized positively hydrolyzing the blood agar medium  
 162 namely AA1, AB1, AB8, AF1, AF2, AF4, AF6, and AG2. There are only two isolates namely AF1 and AF2 which  
 163 show the results of  $\beta$  hemolysis. Suthindhiran [18] suspected that the ability of actinomycetes in hemolytic activity is  
 164 likely to be used as an anticancer agent because of its cytotoxic properties in human and mouse blood cells.

##### 165 3.1.2. Lipolytic activity

166 The tributyrin medium is widely used to test the lipase enzyme, which is exoenzyme, which is able to hydrolyze the  
 167 medium containing tributyrin oil. Lipase plays a role in breaking the lipid complex into simple fragments. Tributyrin oil  
 168 is a type of lipid group called triglycerides. Triglycerides are composed of glycerol and fatty acids. The tributyrin oil  
 169 suspension will make the medium become opaque or look thick. Lipase enzymes produced by microorganisms will


170 break down tributyrins, so that a halo zone (clear) formed around the area of microbial growth. Microbes including  
 171 actinomycetes will break the bonds of glycerol and fatty acids so that they can be converted into final products that will  
 172 be used to produce energy and other metabolic processes. According Fig.1 AF3 and AF1 was top producer of lipase.



173 Fig. 1. Comparison of colony growth and formation of clear zones in actinomycetes isolates through lipolytic test  
 174  
 175

### 176 3.1.3. Surface tension test and Emulsification index


177 The resulting free cell supernatant is measured by its surface tension value using a Du Nouy tensiometer at room  
 178 temperature. The results of surface tension measurements and a decrease in the value of the surface tension to the  
 179 control can be seen in Fig 2. The value of the surface tension measured at day 0 (control) and day 7). The decrease in  
 180 surface tension is caused by the presence of biosurfactants in the supernatant produced by bacterial isolates during the  
 181 growth process.



182 Fig. 2. Surface tension on actinomycetes isolates growing on glycerol, n-hexane and molase substrate media  
 183

184 Based on the results of surface tension measurements (Fig. 2), it was found that most of the isolates were able to reduce  
 185 surface tension except AF2, AF3 and AF4 isolates on glycerol fermentation medium, AA2 and AF4 isolates on n-  
 186 hexane medium and AA2, AF3, and AF4 isolates on molasses medium. Three isolates that were characterized as  
 187 actinomycetes had surface tension values ranging from  $\Delta$  56.22 mN / m for AF6,  $\Delta$  57.27 mN / m for AB8, and  $\Delta$  57.33  
 188 mN / m for AA1.

189 AF3 isolates ranked first in reducing surface stress on fermentation using n-hexane substrate (Fig. 2) with a value of  $\Delta$   
 190 41.28 mN / m followed by AA1 with a reduced surface tension of  $\Delta$  41.28 mN / m. On molasses substrate (Fig 2) AA1  
 191 isolate reduced the surface tension by  $\Delta$  60.39 mN / m, at once being the highest among other isolates.



192

193 Fig. 3. Emulsification activity conducted in the glycerol, n-hexane and molasses fermentation medium. AE1G:  
 194 Emulsification activity of 1<sup>st</sup> hour at glycerol media, AE2G: Emulsification activity after a day at glycerol  
 195 media, AE1H: Emulsification activity of 1<sup>st</sup> hour at n-hexane media, AE2H: Emulsification activity after a  
 196 day at n-hexane media, AE1M: Emulsification activity of 1<sup>st</sup> hour at molase media, AE2M: Emulsification  
 197 activity after a day at molase media.

198 The emulsification activity (AE) of biosurfactants determined by measuring the emulsification index produced (Fig. 3).  
 199 The presence of emulsification activity in the media can be an indicator that biosurfactant compounds have been  
 200 produced by microorganisms. AE data indicate that there was an emulsion forming which appeared to be a gas bubble  
 201 enveloping the fermentation medium. Bacteria use an emulsion as a mechanism of interaction with the substrate and it  
 202 is one of the biosurfactant characters produced by microorganisms. Bacteria expand surface contact with the substrate  
 203 by producing biosurfactants, as well as facilitating substrate transport into the cell. AE was observed in the first hour of  
 204 contact of the supernatant and hydrocarbons namely kerosene and it subsequently observed again after 24 hours.

### 205 3.2. Antimicrobial activity test


206 Media fermented until 7<sup>th</sup> day was continued by testing the antimicrobial activity. The results of the antimicrobial  
 207 activity test (figure 4) was show that AF1 isolates have the largest average inhibition zone diameter in inhibiting the  
 208 growth of *S. aureus* ATCC 6538P which around 12 mm  $\pm$  3.75 mm especially in the molasses medium. AA1 isolates as  
 209 isolates with the highest inhibitory zone in the molasses medium were 7.7 mm  $\pm$  2.15 mm. AB8 with inhibition zone of  
 210 6.65 mm  $\pm$  2.28 mm in the glycerol medium. Molasses medium contributes positively compared to other media in terms  
 211 of producing antimicrobial biosurfactants. Isolate AA1, AA2, AB8, AF1, AF2 and AF6 produced biosurfactants that are  
 212 more antagonistic to *S. aureus* ATCC 6538P when they had grown on molasses medium. AB8, AA1 and AF2 isolates  
 213 are able to produce biosurfactants which inhibit the growth of pathogens both *S. aureus* ATCC 6538P and *E. coli*  
 214 ATCC 25922 (see Fig. 4).

215 *Streptomyces* VITSDK1 spp., [18] are more antagonistic to Gram negative bacteria, with a inhibition zone of 10.3 mm  
 216 in *Klebsiella pneumoniae*. *Streptomyces* VITSDK1 spp., inhibits *Staphylococcus aureus* by 5.3 mm. This shows that the  
 217 antimicrobial activity obtained in this study was reach above it in terms of inhibition zones in Gram positive or negative  
 218 Gram bacteria.

219 Antimicrobial mechanisms might be occur due to disruption of cell wall synthesis, damage to membrane permeability,  
 220 inhibition of protein synthesis, DNA / RNA synthesis and metabolism. The ability to reduce surface tension in  
 221 actinomycetes isolates was thought to play a role in damaging cell walls and permeability of pathogenic membranes.  
 222 The integrity of the disturbed wall and membrane will inhibit various target cell metabolisms. Biosurfactants also acted



223 as fungicidal, bactericidal, insecticidal and anti-viral ingredients because of their anti-adhesive agents and enzyme  
 224 inhibitors [19]. Rhamnolipid was one of the biosurfactant groups which has antibacterial activity [20], and anticapacity  
 225 produced from *P. aeruginosa* AT10 [21]. Shoforolipids from *C. bombicola* yeast are broad-spectrum as antibacterial in  
 226 Gram positive or negative [22]. Lipopeptide was also exploited and gave positive result against methicillin-  
 227 resistant *Staphylococcus aureus* [23].



228

229 Fig. 4. The clear zone formation (mm) indicates the inhibition of bacterial growth due to exposure of the fermented  
 230 supernatant after 7 days on the glycerol, n-hexane and molasses substrate media; SG: *S. aureus* ATCC 6538P  
 231 as pathogen in medium glycerol, SH: *S. aureus* ATCC 6538P as pathogen in medium n-hexane, SM: *S.*  
 232 *aureus* ATCC 6538P as pathogen in medium molase, EG: *E. coli* ATCC 25922 as pathogen in medium  
 233 glycerol, EH: *E. coli* ATCC 25922 as pathogen in medium n-hexane, EM: *E. coli* ATCC 25922 as pathogen  
 234 in medium molase.

### 235 3.3. Fourier Transform Infrared Spectroscopy (FTIR)

236 Result of IR spectral analysis shown different peak at Fig. 5, the sharp at  $3410.15\text{ cm}^{-1}$  was expressed as O-H group.  
 237 The appearance of this peak was confirmed that the sample contains phenol or alcohol group. Wave numbers ranging  
 238 from  $3600\text{ cm}^{-1}$  to  $3100\text{ cm}^{-1}$  was observed. This corresponded to compound that contains amine (N-H) group. Sharp  
 239 absorbance peaks are observed at  $2924.09\text{ cm}^{-1}$ ,  $1157.29\text{ cm}^{-1}$ , and  $1026.13\text{ cm}^{-1}$ . They were indicated aliphatic chains  
 240 with ether and amines. The peak  $1651.07\text{ cm}^{-1}$  was double bonds (alkenes). Spectrum  $1419.61\text{ cm}^{-1}$  that is a bending  
 241 vibration of C-H from  $\text{CH}_2$  while  $1373.32\text{ cm}^{-1}$  is contains nitro compounds. Amine group based on FTIR spectrum  
 242 with positive lipolytic activity regarding halo zone at tributyrin media was allegedly the presence of a lipopeptide  
 243 biosurfactant.

244

245

246

247

248

249

250

251

252

253

254


255

256

257

258

259




260 Fig. 5. FTIR spectrum of the biosurfactant.

## 261 3.4. Identification selected isolate

262 According to Fig. 4, AF1 was the highest isolate that has ability inhibit pathogen proven by clear zone formation.  
 263 It sequenced using 16S rRNA and tabulation data from BLAST® results indicated that isolate AF1 belongs to the genus  
 264 *Streptomyces* spp., and it deposited at GenBank under MN394821 accession number. The percentage of identity  
 265 reached 100% similarity with the *Streptomyces* sp. strain of actinomycetes 16S ribosomal RNA with accession code  
 266 KY236015.1. (Fig 6)

267



268

269 Fig.6. (a) *Streptomyces* sp. strain AF1 (left, Author collection) and (b) *Streptomyces rochei* strain AB1 (right, compared  
 270 to [24])

## 271 4. Conclusions

272 Lapindo mud volcano in Sidoarjo proven as a unique environment that now *Streptomyces* sp., collected from there.  
 273 *Streptomyces* sp. strain AF1 has a unique character. It's able to grow in high temperature around 70° C and produce  
 274 biosurfactant with antimicrobial activity. It potentially developed as candidate for enzymatic and antibiotic producer for  
 275 further research.

## 276 5. References

- 277 1 Söderberg K.H. and Bååth E. 1998. Bacterial Activity Along a Youngbarley Root Measured by The  
 278 Thymidine and Leucine Incorporation techniques. *Soil Biology and Biochemistry*. vol.30, no.10-11 pp.1259-  
 279 1268.
- 280 2 Poernomo A.T, Suryagama D, and Purwanto D.A. 2008. The Influence of pH on The Proteolytic Enzyme  
 281 Activity Thermophiles Bacteria from Lapindo Mud. *Majalah farmasi airlangga*. vol.6, no.2, pp.51-52.
- 282 3 Habibie F.M, Sigres D.P, and Isilami S.L.N. 2014. Isolasi dan Karakterisasi Mikroba Termofilik Penghasil  
 283 Xilanase Dari Lumpur Panas Lapindo Sebagai Alternative Pengganti Klorin Pada Industry Kertas. *Jurnal*  
 284 *pangan dan agroindustry*. vol.2, no.4.
- 285 4 Prachayo R, Khalimi K, and Wijana, G. 2015. Kajian Potensi Bakteri Lumpur Lapindo Sebagai Agen Hayati  
 286 Terhadap *Pyricularia Oryzae* Dan Agen Biostimulan Pada Tanaman Padi. *E-jurnal agroekoteknologi tropika*,  
 287 vol.4, no.1.
- 288 5 Dagdag E.E.A, and Sukoso. 2015. Isolation and Characterization of A3 and S3 Isolate Thermophilic Bacteria  
 289 from Lapindo Sidoarjo Mud, East Java. *International journal of ChemTech research*. vol.8, no.2, pp.541-548.
- 290 6 Nurkanto A, Listyaningsih F, Julistiono H and Agusta A. 2010. Eksplorasi Keanekaragaman Aktinomisetes  
 291 Tanah Ternate Sebagai Sumber Antibiotik. *Jurnal Biologi Indonesia*, vol.6, no.3, pp. 325-339.
- 292 7 Kiran G.S, Thomas T.A, Selvin J.B, Sabarathnam and Lipton A.P. 2010. Optimization and Characterization of  
 293 New Lipopeptide Biosurfactant Produced by Marine *Brevibacterium aureum* MSA13 in Solid State Culture.  
 294 *Bioresources Thechnology*. vol.101, pp.2389-2396.
- 295 8 Sengupta S, Pramanik S, Ghosh A, and Bhattacharyya M. 2015. Antimicrobial activities of actinomycetes  
 296 isolated from unexplored regions of Sundarbans mangrove ecosystem. *BMC Microbiol*, vol.15, no.170,  
 297 <https://doi.org/10.1186/s12866-015-0495-4>.
- 298 9 Shivlata L and Satyanarayana T. 2015. Thermophilic and alkaliphilic *Actinobacteria*: biology and potential  
 299 applications. *Front Microbiol*. vol.6, no.1014. <https://doi.org/10.3389/fmicb.2015.01014>.
- 300 10 Poomthongdee N, Duangmal K, and Wasu P.A. 2015. Acidophilic actinomycetes from rhizosphere soil:  
 301 diversity and properties beneficial to plants. *The Journal of Antibiotics*. vol.68, pp.106-114.
- 302 11 Ballav S, Kerkar S, Thomas S, and Augustine N. 2015. Halophilic and Halotolerant Actinomycete from A  
 303 Marine Saltern Of Goa, India Producing Anti-Bacterial Metabolites. *Journal of Bioscience and*  
 304 *Bioengineering*. vol.119, no.3.pp. 323-330.



- 305 12 Montero K.N, Lamilla C, Abanto M, Maruyama F et al. 2019. Antarctic *Streptomyces fidesensis* So13.3 strain  
306 as a promising source for antimicrobials discovery. *Nature Scientific Reports*. vol.9, pp.7488,  
307 <https://doi.org/10.1038/s41598-019-43960-7>.
- 308 13 Susilowati D.N, Hastuti R.D, Yuniarti E. 2007. Isolasi dan Karakterisasi Aktinomisetes Penghasil Antibakteri  
309 Enteropatogen *E.coli* k1.1, *P.pseudomodallei* 02 05 dan *L. monocitogenes* 5407. *Jurnal AgroBiogen*. vol.3,  
310 no.1, pp.15-23.
- 311 14 Nurkanto A and Julistiono H. 2014. Screening and Study of Antifungal Activity of leaf litter Actinomycetes  
312 Isolated from Ternate Island, Indonesia. *Asian Pacific Journal of Tropical Medicine*. vol.7, pp.S238-S243.
- 313 15 Chakraborty S, Ghosh M, Chakraborti S, Jana S et al. 2015. Biosurfactant Produced from Actinomycete  
314 Nocardiosis A17: Characterization and Its Biological Evaluation. *International Journal of Biological*  
315 *Macromolecules*. vol.79, pp.405-12. c
- 316 16 Sharma M, Dangi P and Choudhary M. 2014. Actinomycetes: Source, Identification, and Their Applications,  
317 *International Journal of Current Microbiology and Applied Science*. vol.3, no.2, pp.801-832.
- 318 17 Ward P and Owen. 2010. *Microbial Biosurfactant and Biodegradation : Biosurfactants*. Lands Bioscience and  
319 Springer Science Business Media.
- 320 18 Suthindhiran K and Kannabiran K. 2009. Hemolytic Activity of *Streptomyces* VITSDK1 spp. Isolated from  
321 Marine Sediments in Southern India. *Journal de Mycologie Médicale*. vol. 19, pp.77-86.
- 322 19 Rangarajan V and Sem R. 2013. An inexpensive strategy for facilitated recovery of metals and fermentation  
323 products by foam fractionation process. *Coll. Surf. B Biointerfaces*. vol.104, pp.99-106.
- 324 20 Santos D.K.F, Rufino R.D, Luna J.M, Santos V.A, Sarubbo L.A. 2016. Biosurfactants: Multifunctional  
325 Biomolecules of the 21st Century. *International Journal of Molecular Sciences*. vol. 17, no.3, pp.401.
- 326 21 Abalos A, Pinazo A, Infante M.R, Casals M, García F, Manresa A. 2001. Physicochemical and antimicrobial  
327 properties of new rhamnolipids produced by *Pseudomonas aeruginosa* AT10 from soybean oil refinery wastes.  
328 *Langmuir*. vol.17, pp.1367-1371.
- 329 22 Joshi K. N, Prabhune A. 2013. A biosurfactant sophorolipid acts in synergy with antibiotics to enhance their  
330 efficiency. *BioMed. Res*. pp.1-8.
- 331 23 Sharma D, Mandal S.M, Manhas R.K. 2014. Purification and characterization of a novel lipopeptide from  
332 *Streptomyces amritsarensis* sp. nov. active against methicillin-resistant *Staphylococcus aureus*. *AMB Express*.  
333 vol.4:50. -y
- 334 24 Aouar L, Lerat S, Ouffroukh A, Boulahrouf A, and Beaulieu C. 2012. Taxonomic identification of  
335 rhizospheric actinobacteria isolated from Algerian semi-arid soil exhibiting antagonistic activities against plant  
336 fungal pathogens. *CAN J PLANT PATHOL*. vol.34, pp.1-12.