

## Ecology, Environment and Conservation Editorial Advisory Board

### Chief Editor

**Prof.(Dr.) R.K.Trivedy, Pune, India**

#### EDITORIAL ADVISORY BOARD

- |  |  |
|--|--|
| 1. Dr. Teresa Ferreira, Portugal             | 19. Dr. A. Olawale, Nigeria                          |
| 2. Dr. Michael Ukwuru, Nigeria               | 20. Dr. Ing. Agr. Mario Ridardo Sabbatini, Argentina |
| 3. Dr. Moses Inbaraj, Chennai                | 21. Dr. Philip C. Reid, U.K.                         |
| 4. Dr. D.J. Lee, Taiwan                      | 22. Dr. Mohd. Yusuf, Malaysia                        |
| 5. Dr. Christial Paul P.delacruz, Phillipnes | 23. Dr. Oswaldo A. Feernandez, Argentina             |
| 6. Dr. T. Bahorun, Mauritius                 | 24. Dr. Ms. Mirela Tulik, Warsaw, Poland             |
| 7. Dr. Linda Blackwell, Australia            | 25. Dr. L.L. Chukwu, Nigeria                         |
| 8. Dr. G. Zellner, Netherlands               | 26. Dr. Azni Idris, UPM, Malaysia                    |
| 9. Dr. Wilson S. Tisera, Kupang, Indonesia   | 27. Dr. Vikas Sharma, J&K, India                     |
| 10. Dr. M.F. Hamoda, Kuwait                  | 28. Dr. Amresh Chandra Pandey, Jharkhand, India      |
| 11. Dr. H.A.Abrahamse, South Africa          | 29. Prof.(Dr.) Agoes Soegianto, Indonesia            |
| 12. Dr. Arulmozhiyal R., Salem               | 30. Dr. A.K. Panigrahi, Berhampur, India             |
| 13. Dr. Hassan Ibrahim Ali, Sudan            | 31. Dr. Ahmed El Mahmoudi, Saudi Arabia              |
| 14. Dr. A.R.Ghosh, Burdwan, India            | 32. Dr. Seyed Mohammad Tajbakhsh, Iran               |
| 15. Prof. M. Zaman, Bangladesh               | 33. Dr. Amin L. Setyo, Indonesia                     |
| 16. Dr. Marcantonio Bragdin, Venice, Italy   | 34. Dr. Francis Gbogbo, Ghana                        |
| 17. Dr. Z. Fuat Topark, Turkey               | 35. Dr. S. Shabanlou, Iran                           |
| 18. Dr. Z. Li. Bonn, Germany                 |  |

[Back to EEC Journal Details](#)

## Ecology, Environment and Conservation Journal Papers

Issue: Vol 26, April Suppl. Issue, 2020

### [PATHOGENIC FOOD-BORNE BACTERIA IN SHELLFISH AND SHRIMP FROM THE LARGEST TRADITIONAL SEAFOOD MARKET IN SURABAYA, INDONESIA](#)

Sin War Naw, Putu Angga Wiradana, Agoes Soegianto and Aondohemba Samuel Nege

[Get Abstract](#)[Get Paper](#)

### [DNA BAR CODING: STUDY OF BANANAS \(MUSA SPP.\) WILD AND CULTIVARS GROUP FROM EAST JAVA INFERRED BY RBC L GENE SEQUENCES](#)

Rojaunnajah Kartika Ainiyah, Verina Wahyunindita, Windi Nur Pratama, Intan Ayu Pratiwi, Edy Setiti Wida Utami and Sucipto Hariyanto

[Get Abstract](#)[Get Paper](#)

### [ROLE OF EDUCATION IN CONSERVING “KALEKA” FOREST IN DAYAK PEOPLE OF CENTRAL KALIMANTAN, INDONESIA](#)

Holten Sion, Muhamad Affandi and Febrina Natalia

[Get Abstract](#)[Get Paper](#)

### [DIVERSITY OF INDIGENOUS ENTOMOPATHOGENIC BACILLI FROM DOMESTICS BREEDING SITES OF DENGUE HEMORRHAGIC FEVER VECTOR BASED ON THE TOXICITY AGAINST AEDESAEGYPTI LARVAE](#)

Salamun, Ni'matuzahroh, Fatimah, M.I.F. Maswantari, M.U. Rizka, T. Nurhariyati and A. Supriyanto

[Get Abstract](#)[Get Paper](#)

### [PARASITIZATION OF PARASITIC PLANTS ON FRUIT PLANTS IN BANGKALAN REGENCY AND MALANG CITY, INDONESIA](#)

Solikin

[Get Abstract](#)[Get Paper](#)

### [CRUDE METHANOL EXTRACT OF BROTOWALI LEAVES \(TINOSPORA CRISPA\) AS BIOLARVICIDE AGAINST DENGUE VECTOR AEDES AEGYPTI](#)

Manikya Pramudya, Rosmanida, Nabilah Istighfari Zuraidassanaaz, Nadyatul Ilma Indah Savira, Elma Sakinatussajdah and Intan Permata Putri

[Get Abstract](#)[Get Paper](#)

### [THE EFFECT OF OKRA PODS \(ABELMOSCHUS ESCULENTUS L.\) METHANOL EXTRACT ON WHITE BLOOD CELL COUNT, PHAGOCYtic ACTIVITY, AND IFN-GAMMA LEVEL IN MUS MUSCULUS EXPOSED SODIUM NITRITE](#)

Sri Puji Astuti Wahyuningsih, Siti Richa Isnaini, Dwi Winarni, Win Darmanto, Manikya Pramudya, Anni Mar'atun Nafi'ah, Khoirunnisa, Elma Sakinatus Sajdah and Baiq Naili Dewi Atika

## Search Articles

### Journal Issues

[Vol 26, Issue 2, 2020](#)[Vol 26, June Suppl. Issue, 2020](#)[Vol 26, April Suppl. Issue, 2020](#)[Vol 26, Issue 1, 2020](#)[Vol 26, Feb Suppl. Issue, 2020](#)[Vol 25, Issue 4 2019](#)[Vol 25, Issue 3 2019](#)[Vol 25, Nov Suppl. Issue, 2019](#)[Vol 25, Sept Suppl. Issue, 2019](#)[Vol 25, Aug Suppl. Issue, 2019](#)[Vol 25, July Suppl. Issue, 2019](#)[Vol 25, Issue 2 2019](#)[Vol 25, May Suppl. Issue, 2019](#)[Vol 25, April Suppl. Issue, 2019](#)[Vol 25, Issue 1 2019](#)[Vol 24, Issue 4 2018](#)[Vol 24, Issue 3 2018](#)[Vol 24, Issue 2 2018](#)[Vol 24, Issue 1 2018](#)[Vol 24, March Suppl. Issue 2018](#)[Vol 24, Feb. Suppl. Issue 2018](#)[Vol 23, Issue 4, 2017](#)[Vol 23, Nov. Suppl. Issue 2017](#)[Vol 23, Sept. Suppl. Issue 2017](#)[Vol 23, Issue 2, 2017](#)[Vol 23, Issue 3, 2017](#)[Vol 23, Issue 1, 2017](#)[Vol 23, Feb 2017 Suppl. Issue](#)[Vol 22, Dec 2016 Suppl. Issue](#)[Vol 22, Issue 4, 2016](#)[Vol 22, Sept. Suppl. Issue, 2016](#)[Vol 22, Issue 3, 2016](#)[Vol. 22, June Suppl. Issue 2016](#)[Vol 22, Issue 2, 2016](#)[Vol. 22, April Suppl. Issue 2016](#)[Vol 22, Issue 1, 2016](#)[Vol 21, Issue 4, 2015](#)[Vol. 21 Dec. 2015 Suppl. Issue](#)[Vol. 21 Nov. 2015 Suppl. Issue](#)[Vol 21, Issue 3, 2015](#)[Vol 21, Issue 2, 2015](#)[Vol. 21 Suppl. Issue August 2015](#)[Vol 21, Suppl. Issue June 2015](#)[Vol 21, Issue 1, 2015](#)[Supplement Issue, Dec. 2014](#)[Special Issue-2014](#)

[Get Abstract](#)[Get Paper](#)

**SPINY LOBSTERS SPECIES COMPOSITION BETWEEN NORTH AND SOUTH SEA OF EAST JAVA, INDONESIA**

Arief Setyanto, Soemarno, D.G.R. Wiadnya, Cahyo Prayogo, R.J. West, M. Tsamenyi, S. Halimah and N.A. Rahman

[Get Abstract](#)[Get Paper](#)

**ECOLOGICAL STUDY OF PRIMITIVE BRACHIOPODS LINGULA SP. IN PROBOLINGGO, EAST JAVA, INDONESIA**

Rakmawati and Sucipto Hariyanto

[Get Abstract](#)[Get Paper](#)

**IDENTIFICATION AND TOXICITY TEST OF ORTHOSIPHONSTAMINEUS LEAVES EXTRACT AGAINST HOUSEFLIES(MUSCA DOMESTICA)**

Surahmaida, Umarudin and Junairiah

[Get Abstract](#)[Get Paper](#)

**THE EFFECTS OF LEAD ACETATE EXPOSURE ON BLOOD COMPONENT AND KIDNEY: THE MECHANISM OF OXIDATIVE STRESS**

Sugiharto, Win Darmanto, Sri Puji Astuti W., Kevita Putri, Nur Khoirunnisa, Dea B. Mulyawan, and Arin N. Shanti

[Get Abstract](#)[Get Paper](#)

**THE EFFECTS OF LEAD ACETATE EXPOSURE ON BLOOD COMPONENT AND KIDNEY: THE MECHANISM OF OXIDATIVE STRESS**

Sugiharto, Win Darmanto, Sri Puji Astuti W., Kevita Putri, Nur Khoirunnisa, Dea B. Mulyawan, and Arin N. Shanti

[Get Abstract](#)[Get Paper](#)

**AN EVIDENCE OF THE MISSING COMPLETE PECTORAL FINS TRAIT IN CLARIAS GARIEPINUS REARED IN POND IS HERITABLE**

Farikhah Farikhah, Sukoso Sukoso, Uun Yanuhar, Feni Iranawati and Muchlis Zainudin Arif

[Get Abstract](#)[Get Paper](#)

**DIVERSITY OF MODERATELY THERMO-ALKALIPHILE BACTERIA PRODUCING AMYLASE ORIGINATED FROM PARIANGAN HOT SPRING, WEST SUMATRA, INDONESIA**

Gustina Indriati, Ruth Rize Pass Megahati and Dwi Hilda Putri

[Get Abstract](#)[Get Paper](#)

**LIPID ACCUMULATION OF SKLETONEMA COSTATUM UNDER SILICA, NITROGEN AND AERATION STRESS**

Tutik Nurhidayati, Aini Ika Rahmawati, Endang Purwanti Setyaningsih, Kristanti Indah Purwani, Edwin Setyawan and Nurul Jadid

[Get Abstract](#)[Get Paper](#)

**TOXICITY EFFECT OF COPPER ON HISTOPATHOLOGY OF OREOCHROMIS NILOTICUS LIVER WITH DIET OF LACTIC ACID BACTERIA**

Alfiah Hayati, Rosana Dyah Umami, Aken Puti Wanguyun, Primadika Ashar Dwi Putra, Afinia Primananda Taufiq and Agus Supriyanto

[Get Abstract](#)[Get Paper](#)[Vol.20, Issue 4, 2014](#)[Vol.20, Issue 3, 2014](#)[Vol.20, Issue 2, 2014](#)[Vol.20 Issue 01, 2014](#)[Vol.19 Issue 04, 2013](#)[Vol.19 Issue 03, 2013](#)[Vol.19, Issue 02, 2013](#)[Vol.19, Issue 01, 2013](#)[Vol.18, Issue 04, 2012](#)[Vol.18, Issue 3, 2012](#)[Vol.18, Issue 2, 2012](#)[Vol.18, Issue 1, 2012](#)[Vol.17, Issue 4, 2011](#)[Vol.17, Issue 3, 2011](#)[Vol.17, Issue 2, 2011](#)[Vol.17, Issue 1, 2011](#)[Vol.16, Issue 4, 2010](#)[Vol.16, Issue 3, 2010](#)[Vol.16, Issue 2, 2010](#)[Vol.16, Issue 1, 2010](#)[Vol.15, Issue 04, 2009](#)[Vol.15, Issue 03, 2009](#)[Vol.15, Issue 02, 2009](#)[Vol.15, Issue 1, 2009](#)[Vol.14, Issue 04, 2008](#)[Vol.14, Issue 2-3, 2008](#)[Vol.14, Issue 2-3, 2008](#)[Vol.14, Issue 1, 2008](#)[Vol.14, Issue 2-3, 2008](#)[Vol.13, Issue 04, 2007](#)[Vol.13, Issue 2, 2007](#)[Vol.13, Issue 1, 2007](#)[Vol.12, Issue 4, 2006](#)[Vol.12, Issue 3, 2006](#)[Vol.12, Issue 2, 2006](#)[Vol.12, Issue 1, 2006](#)[Vol.12, Issue 01, 2006](#)[Vol.11, Issue 3,4, 2005](#)[Vol.11, Issue 2, 2005](#)[Vol.11, Issue 1, 2005](#)[Vol.10, Issue 04, 2004](#)[Vol.10, Issue 03, 2004](#)[Vol.10, Issue 02, 2004](#)[Vol.10, Issue 01, 2004](#)[Vol.09, Issue 04, 2003](#)[Vol.09, Issue 03, 2003](#)[Vol.09, Issue 02, 2003](#)[Vol.08, Issue 04, 2002](#)[Vol.08, Issue 03, 2002](#)[Vol.08, Issue 01, 2002](#)[Vol.07, Issue 04, 2001](#)[Vol.07, Issue 03, 2001](#)[Vol.07, Issue 02, 2001](#)[Vol.07, Issue 01, 2001](#)[Vol.06, Issue 04, 2000](#)[Vol.06, Issue 03, 2000](#)[Vol.06, Issue 02, 2000](#)[Vol.06, Issue 01, 2000](#)[Vol.05, Issue 04, 1999](#)[Vol.05, Issue 03, 1999](#)[Vol.05, Issue 02, 1999](#)[Vol.05, Issue 01, 1999](#)[Vol.04, Issue 1,2, 1998](#)[Vol.03, Issue 3,4, 1997](#)[Vol.03, Issue 01, 1997](#)

**RISK ASSESSMENT OF INVASIVE ALIEN PLANT SPECIES IN PANGANDARAN NATURE RESERVE, WEST JAVA, INDONESIA**

Arif Mukhtar Hasbi and Dian Rosleine

[Get Abstract](#)

[Get Paper](#)

**PHENOTYPIC DIVERSITY OF FIVE GOAT POPULATIONS IN TROPICAL ENVIRONMENT**

Arsyad Alawiansyah, Richi Yuliavian Kusminanto, Nuzul Widyas, Ahmad Pramono, Muhammad Cahyadi and Sutarno

[Get Abstract](#)

[Get Paper](#)

**NEW ISOLATION OF HYDRO CARBONOCLASTIC BACTERIA FROM LOCAL LIMESTONE MINING AS PROMISING CONCRETE STRENGTH ENHANCER**

Enny Zulaika, M. Andry Prio Utomo, Nur Hidayatul Alami, Endry Nugroho Prasetyo, Tri Wulan and Agoes Soegianto

[Get Abstract](#)

[Get Paper](#)

**BIRD DIVERSITY OF RESORT RANU DARUNGAN, BROMO TENGGER SEMERU NATIONAL PARK, INDONESIA**

Boni Herdiawan, Reni Ambarwati and Toni Artaka

[Get Abstract](#)

[Get Paper](#)

**CHARACTER VISUALIZATION OF PYCNOSCELUS SP. (BLATTODEA: COCKROACH) IN HOUSEHOLD ORGANIC WASTE COMPOSTER IN SURABAYA**

Moch.Affandi, Emilia Anjar Prastiwi, Nindya Ayu Damayanti, Chairunisa Firdaus, Azizah Kusumadewi, Anisa Rahmawati, Puspanjali Prahasto, Audry Putri Kristanti, Alifatus Alkurnia Sukma Firdaus, Hardiansah Pramana Abdilah, Muhammad Arifudin Mubarak, Syahriar

[Get Abstract](#)

[Get Paper](#)

**ISOLATION AND CHARACTERIZATION OF COCKROACH ENDOSYMBIONT BACTERIA WITH POTENTIAL TO PRODUCE HYDROLYTIC ENZYME OF ORGANIC MATERIAL**

Ni'matuzahroh, Nastiti Trikurniadewi, Syahriar Nur Maulana Malik Ibrahim, Achmad Zainal Abidin, Ana Mariatul Khiftiyah, Silvia Kurnia Sari, Eka Narendra Nuswantara, Fawwas Nurmansyah, Muhammad Aufar Rizki Wildan Rahman, Habibah Lailatul Maghfirah, Miftahul

[Get Abstract](#)

[Get Paper](#)

**MORPHOPHYSIOLOGICAL CHARACTERISTIC RESPONSES OF SOYBEAN (GLYCINE MAX L.) GROBOGAN VARIETY IN WATER LOGGING STRESS**

Vita Siti Fatimah and Tutik Nurhidayati

[Get Abstract](#)

[Get Paper](#)

**THE NUMBERS OF LYMPHOCYTES AND CAPILLARIES IN SILICA EXPOSED LUNG TISSUES OF BALB/C MICE (MUSMUSCULUS) TREATED WITH MORINGAOLEIFERA LEAVES EXTRACT**

Kusmiyati, Soedjajadi Keman, Muhammad Amin, Suwarno and Agoes Soegianto

[Get Abstract](#)

[Get Paper](#)

**BIODIVERSITY RICHNESS OF TERRESTRIAL ORCHIDS IN BROMO TENGGER SEMERU NATIONAL PARK, INDONESIA**

Lita Soetopo and Muhamad Jauhari

[Get Abstract](#)

[Get Paper](#)

**CHARACTERISTICS OF RICE PLANT WITH LOW METHANE EMISSION**

Noer Rahmi Ardiarini, Finsa Dwi Arisandi and Prihasto Setyanto

[Vol.02, Issue 1,2, 1996](#)

[Vol.01, Issue 14, 1995](#)

Looking for Past Issues?

[Click here to get them!!](#)

[Get Abstract](#)[Get Paper](#)

**EXPLORING THE LANDRACE RICE (ORYZA SATIVA L) FOR SALINITY TOLERANT AND AGRONOMIC TRAITS**

Nafisah, Celia Roza, Nani Yunani, Rahmini and Priatna Sasmita

[Get Abstract](#)[Get Paper](#)

**BIODIVERSITY OF PLANT GROWTH PROMOTING RHIZO BACTERIA FROM RHIZOSPHERE OF PINUS SP AND CASUARINA SP AT RADEN SOERJO FOREST PARK**

Brigita Nur Diyan Agustiana, Indra Gumilar, RizqiaNurul Alfiyanita, Alifah Nastiti, Tesalonika Tetya Virginia, Siti Rizqiyatul Mukarromah, Annisa Dwi Savitri, Lunace Rahma Kinayoh, Erna Erviana Nadhifa, Khafid Jallaludhin, Ni'matuzahroh and Almando Gerald

[Get Abstract](#)[Get Paper](#)

**CORN COB HYDROLYZATE FROM PENICILLIUM CITRINUM H9 AS AN ALTERNATIVE SUBSTRATE FOR BIO SURFACTANT PRODUCTION BY HYDROCARBONOClastic BACTERIA**

Fatimah, Silvia Kurnia Sari, Nastiti Trikurniadewi, Syahriar Nur Maulana Malik Ibrahim, Ana Mariatul Khiftiyah, Khudrotul Nisa Indriyasari, Tri Nurhariyati, Tini Surtiningsih and Hanif Yuliani, Ni'matuzahroh

[Get Abstract](#)[Get Paper](#)

**INFESTATION OF BIO FOULINGS (BALANUS SP. AND LITTORARIASp.) ON MANGROVE RHIZOPHORA SP. AND AVICENNIA SP. AT GADINGREJO, PASURUAN, INDONESIA**

Oktiyas Muzaky Luthfi, Evi Safitri and Agoes Soegianto

[Get Abstract](#)[Get Paper](#)

**LENGTH-WEIGHT RELATIONSHIP AND CONDITION FACTOR OF SYGNATHOIDES BIACULEATUS (BLOCH, 1785) CAUGHT FROM SEA GRASS AREA OF TANAKEKE ISLANDS, TAKALAR DISTRICT, SOUTH SULAWESI, INDONESIA**

Ardi Eko Mulyawan, Yeni Savitri Andi Lawi, Syaifiuddin, Farid Samawi and Agoes Soegianto

[Get Abstract](#)[Get Paper](#)

**COMPARISON OF WATER MASS CHARACTERISTICS BETWEEN THE INDONESIAN THROUGH FLOW EXIT PASSAGE AND SOUTH JAVA SEA, INDONESIA**

B. Semedi, A. Purnomosakti, D. K. Saputra, I.B. Wahyono and A. Soegianto

[Get Abstract](#)[Get Paper](#)

**COMPARISON OF WATER MASS CHARACTERISTICS BETWEEN THE INDONESIAN THROUGH FLOW EXIT PASSAGE AND SOUTH JAVA SEA, INDONESIA**

B. Semedi, A. Purnomosakti, D. K. Saputra, I.B. Wahyono and A. Soegianto

[Get Abstract](#)[Get Paper](#)

**MORPHOLOGY-BASED GENETIC DIVERSITY OF EARLY MATURITY RICE GERM PLASM, THE RELATION TO YIELD COMPONENT**

Indria W. Mulsanti, Nani Yunani and Trias Sitaresmi

[Get Abstract](#)[Get Paper](#)

**THE POPULATION OF SOLEN SP. (BIVALVIA: SOLENIDAE) FROM PAMEKASAN, INDONESIA**

Reni Ambarwati and Bambang Irawan

[Get Abstract](#)[Get Paper](#)

**LARVICIDAL ACTIVITY OF METHANOL FRACTIONS FROM CARICAPAPAYA LEAVES EXTRACT AGAINST AEDES AEGYPTI**

Tri Puji Lestari Sudarwati, M.A.H. Ferry Fernanda, Hilya Nur Imtihani and Intan A. Pratiwi

[Get Abstract](#)

[Get Paper](#)

**MOLECULAR DETECTION AND MORPHOLOGICAL VARIED OF BLASTOCYSTIS SP IN BEEF CATTLE IN SIAK SRI INDRAPURA RIAU INDONESIA**

Lucia Tri Suwanti, Mufasirin, Nunuk Dyah Retno Lastuti, Yuli Susana and Suci M. Rizki

[Get Abstract](#)

[Get Paper](#)

**REMOVAL OF LEAD (II) FROM AQUEOUS SOLUTION USING COMPOSITE ADSORBENT OF CASSAVA PEEL WASTE/MAHOGANY SAWDUST/BENTONITE**

Eko Prasetyo Kuncoro, Ganden Supriyanto, Thin Soedarti, Nilaturrohmah and Gabriel Putra Utama

[Get Abstract](#)

[Get Paper](#)

---

[Home](#) | [International Journals](#) | [Books](#) | [Environmental Consulting](#) | [About Us](#) | [Contact Us](#) | [Submit Paper](#) | [Search Journal Article](#) |

[Become a fan](#) on Facebook

[Follow us](#) on Twitter



---

© EM International 2012-2019 | **Developed by Eneblur Consulting**

---

[i](#) [x](#)

## molecutrack

We Focus on Clean Energy Solutions That Replace Hazardous and Carbon Intensive Fuel.


molecutrack.com OPEN

# Ecology, Environment and Conservation

<b>Country</b>	India - <a href="#">IIII</a> <a href="#">SJR Ranking of India</a>
<b>Subject Area and Category</b>	<a href="#">Agricultural and Biological Sciences</a> <a href="#">Ecology, Evolution, Behavior and Systematics</a>  <a href="#">Environmental Science</a> <a href="#">Ecology</a> <a href="#">Nature and Landscape Conservation</a>
<b>Publisher</b>	<a href="#">EM International</a>
<b>Publication type</b>	Journals
<b>ISSN</b>	0971765X
<b>Coverage</b>	1997-2019
<b>Scope</b>	Published Quarterly Since 1995. Ecology, Environment and Conservation is published in March, June, September and December every year. ECOLOGY, ENVIRONMENT AND CONSERVATION is one of the leading International environmental journal. It is widely subscribed in India and abroad by Institutions and Individuals in education and research as well as by Industries, Govt. Departments and Research Institutes.
<a href="#">?</a>	<a href="#">Homepage</a> <a href="#">How to publish in this journal</a> <a href="#">Contact</a>  <a href="#">Join the conversation about this journal</a>








# 12

H Index



SJR +

Citations per document +



**Ecology, Environment and Conservation**

Q4 Ecology  
best quartile

SJR 2019  
0.14

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```

L **Leta Rafael Levis** 2 weeks ago

I have published my article on this journal in 2017, no 23 Issue 2 with the title of the article was :  
Farmers attitude .....  
How ever, I have got the URL for full text.  
So, please let me know the URL for the article full text.

Change my email address: in 2017 when I sent my article to this journal I still use email address:



# Isolation and characterization of cockroach endosymbiont bacteria with potential to produce hydrolytic enzyme of organic material

Ni'matuzahroh<sup>12\*</sup>, Nastiti Trikurniadewi<sup>12</sup>, Syahriar Nur Maulana Malik Ibrahim<sup>12</sup>, Achmad Zainal Abidin<sup>1</sup>, Ana Mariatul Khiftiyah<sup>12</sup>, Silvia Kurnia Sari<sup>12</sup>, Eka Narendra Nuswantara<sup>1</sup>, Fawwas Nurmansyah<sup>1</sup>, Muhammad AUFAR Rizki Wildan Rahman<sup>1</sup>, Habibah Lailatul Maghfirah<sup>1</sup>, Miftahul Jannah<sup>1</sup>, Anna Raisa Masrurin<sup>1</sup>, Lailatus Saidah<sup>1</sup>, Risza Lailiana Makrifah<sup>1</sup>, Fatimah<sup>12</sup> and Moch Affandi<sup>12</sup>

<sup>1</sup>*Department of Biology, Faculty of Science and Technology, Universitas Airlangga, Surabaya, Indonesia*

<sup>2</sup>*Research Center for Bio-Molecule Engineering, Universitas Airlangga, Surabaya, Indonesia*

(Received 27 September, 2019; Accepted 10 January, 2020)

## ABSTRACT

Cockroaches are one of the most popular decomposers in organic matter processing. Its ability to decompose organic matter is supported by the association with the digestive tract microbiota. Microbial exploration of the cockroach gastrointestinal tract is still rarely studied, especially in Indonesia. This study aims to obtain potential bacterial isolates that produce enzymes that can decompose of organic matter derived from cockroach midgut. Cockroach samples were obtained from household organic wastes composts. Midguts of cockroach sample were surgically removed, mashed aseptically, and inoculated into nutrient agar medium. Endosymbiont bacteria grown in the media then purified, characterized, and tested for enzymatic abilities such as amylase, protease, lipase and cellulase. The abundance of cockroach midgut endosymbiont bacteria was detected as much as  $1.1 \times 10^4$  CFU/mL. A total of 20 endosymbiont bacterial isolates had different characters and had the ability to hydrolyze starch, cellulose, protein, and lipid. Three potential bacterial isolates in hydrolyzing starch, cellulose, protein, and lipid were EKA 4, EKA 8, and EKA 20. Cockroach endosymbiont isolates are potential to be applied for organic waste treatment.

*Key words:* Cockroach, Endosymbiont, Enzyme, Midgut, Organic waste

## Introduction

Increasing human population, urbanization, economic growth, biomass production and consumption patterns enhance the amount of solid wastes (Shaoli and Debnath, 2019). Treatment technologies for solid waste can be done by direct use, physico-chemical, thermochemical, and biological treatment (Lohri *et al.*, 2017). Biological treatment applied to solid organic waste on a household scale is com-

monly done by composting (Warunasinghe and Yapa, 2016; Bakari *et al.*, 2017). Composting is the process of decaying organic matter aerobically to produce humus (Epstein, 1996; Sikora, 1998; Lohri *et al.*, 2017). Decomposition is a process of physical or chemical changes in organic matter of animals and plants into simple organic compounds (Susanti and Halwany, 2017). Lohri *et al.*, (2016) revealed that, composting in a household scale is generally done by stacking organic waste in barrels and relying on

Corresponding author's email: [nimatuzahroh@fst.unair.ac.id](mailto:nimatuzahroh@fst.unair.ac.id)

the passive aeration process. The main actors in composting organic material are microorganisms such as fungi, yeasts, bacteria and macro invertebrates such as insects (Morales and Wolff, 2010; Lohri *et al.*, 2017). Decomposed organic matter becomes a natural habitat for these organisms (Tuomela *et al.*, 2000). The association of insects in the process of composting organic waste is able to increase the speed of physical and chemical changes of organic waste during decomposition as well as causing a decrease in the molecular weights of compounds (Morales and Wolff, 2010). This makes insects as reliable organisms that can recycle organic matter (Morales and Wolff, 2010; Tarli *et al.*, 2014). Among all insects, cockroaches, member of order Blattodea, are important detritivore in the ecosystem (Evangelista *et al.*, 2013; Mullins, 2015). As detritivores, cockroaches not only consume fresh food types, but even rotten human leftover, and excrement from other cockroaches (Rozendaal, 1997; Moges, *et al.* 2016). The wide variety of organic materials in biological waste products makes the Blattodea order very important for ecosystem functioning (Weiiser and Sieman, 2008; Ifeanyi and Odunayo, 2015).

Cockroaches share their habitat with various types of bacteria which makes association in the form of symbiotic relation is not uncommon. Symbiotic bacteria are transmitted vertically between generations and horizontally from the environment (Bright and Bulgheresi, 2010). Horizontal transmission mainly sourced from organic materials in the habitat (Zhang and Zhang, 2018). Bacteria from the habitat enter the digestive tract of cockroaches and stay for a short time or transient (allochthonous bacteria) and others are able to attach and form colonization of the digestive tract mucus epithelium (autochthonous bacteria). The endosymbiont bacteria showed beneficial effects for the host. In the case of *Blattella germanica*, among others, the endosymbionts increased the host resistant to insecticides (Pietri, *et al.*, 2018), pathogenic fungi (Zhang, 2018), helped nitrogen fixation (Patino-Navarrete, 2014), and helped degrade the complex organic components by producing extracellular enzymes (Tinker and Ottesen, 2016).

Endosymbiont bacteria of cockroach digestive tract work together to alter food (Zhang and Zhang, 2018). Each bacterium can produce different hydrolytic enzymes such as cellulase, amylase, lipase, and protease (Singh *et al.*, 2016). Their enzyme products

have the potential to be utilized for various fields in industry. Thus, in this research, we isolated cockroach's endosymbiont and tested their potencies to produce hydrolytic enzymes which can be utilized in various industries.

Endosymbiont diversity in the digestion track of cockroaches is greatly influenced by their diet (Schauer *et al.*, 2012). Omnivorous cockroaches *Periplaneta americana* has a higher endosymbiont diversity compared to wood-feeding cockroach *Cryptocercus punctulatus* (Colman *et al.*, 2012; Pérez-Cobas *et al.*, 2015). Some exploratory research of endosymbiont bacteria in the digestive tract of cockroaches has been carried out including on *Blattella germanica* (Pérez-Cobas, *et al.*, 2015), *Shelfordella lateralis* (Schauer, 2012), *Periplaneta americana* (Tinker and Ottesen, 2016), *Panchlora* (Gontang *et al.*, 2017), and litter-feeding cockroach *Pycnoselus surinamensis* (Richards *et al.*, 2017). However, there is still rare information about the hydrolytic enzymes produced by the endosymbiont bacteria of *Pycnoselus surinamensis*.

In this study, endosymbiont bacteria were isolated from the gut of *Pycnoselus surinamensis*. The diversity of bacteria was detected by culture-dependent methods. This study will add the data about diversity of endosymbiont bacteria from gut of *Pycnoselus surinamensis* that have ability to produce hydrolytic enzymes of starch, cellulose, protein, and lipid.

## Materials and Methods

### Cockroach collection

Cockroaches were obtained from Surabaya urban household waste treatment plants which had been acclimated for 2 weeks at the Faculty of Science and Technology, Universitas Airlangga, Surabaya, Indonesia. The most abundant cockroach that was chosen as the object of this study was *Pycnoselus surinamensis* according to Roth, 1998.

### Dissection and endosymbiontbacteria isolation of cockroach

A Cockroach obtained by hand sorting was anesthetized with chloroform, and then the legs were removed. The side of the cockroach was slashed from the posterior to anterior to obtain the intact digestive tract. The digestive tract organ was separated from the body. The digestive tract organ was steril-

ized according to Wynants *et al.* (2018) with modification: digestive tract organ was washed with 10 mL of 70% alcohol then followed by two washes of 10 mL of sterile distilled water. Each washing step was carried out for 15 seconds on a vortex with a speed of 2000 rpm. The gut was cut aseptically, crushed, and dissolved in a physiological saline solution (NaCl 0.85%). One mL of the suspension was diluted at stratified dilution until it reaches a certain bacterial density. Bacteria were grown on Nutrient Agar (NA) (Oxoid), Strach Casein Agar (SCA), and de Man, Rogosa, and Shape Agar (MRSA) with pour plate technique and incubated for 24-48 hours at temperatures of approximately 29 °C.

#### Quantification and characterization of cockroach endosymbiont bacteria

Bacteria that grow in each medium were counted using the Total Plate Count (TPC) principle. Colonies that have different characters were purified and rejuvenated for hydrolytic assay. Macroscopic characterizations were done by observing the character of bacterial colonies consisting of shape, size, color, margin, elevation, and transparency, while microscopic characterizations were done by Gram staining and observing at shape of bacterial cells.

#### Hydrolytic assay

Amylolytic, proteolytic, cellulolytic, and lipolytic activities of each isolates were assessed on solid media that had been modified with the test substrates. Hydrolytic activity by amylase-producing potential bacteria was carried out on NA which was added by starch 2% (w/v) (Banerjee and Ghosh, 2014), hydrolytic activity by protease-producing potential bacteria was carried out on NA which was added by skim milk 2% (v/v) (Bhowmik, *et al.*, 2015), hydrolytic activity by cellulase-producing potential bacteria was carried out on Carboxy Methyl Cellulose (CMC), hydrolytic activity by potential lipase-producing bacteria was carried out on NA which was added by glycerol 2% (v/v). Incubation was carried out for 24 hours at a temperature of approximately 29 °C. Visualization of hydrolytic activity of each substrate was different. To determine the presence of amylase was done by adding a 2% iodine indicator solution to the test media, to determine the presence of proteases without the addition of any reagents, to determine the presence of cellulase was done by adding Congo red indicator solution then washed with NaCl 1% solution for 20 min-

utes, to determine the presence of lipase by adding rhodamine indicator solution which was then observed under UV light. Positive test results were indicated by the presence of a clear zone around the bacterial colony. The hydrolytic activity index was calculated using the formula (Febriyanto *et al.*, 2015):

$$\text{Hydrolytic Activity Index} = \frac{\text{Bacteria Colony Zones Diameter (mm)}}{\text{Bacteria Colony Diameter (mm)}}$$

#### Identification of potential bacteria based on physiological character

The three enzymatic potential bacteria were identified using Microbact kit GNB 12A and 12 B (Oxoid) based on manufacturer's instructions. Identification of bacteria isolates obtained based on Bergey's Manual of Determinative Bacteriology, Holt (1994).

### Results

#### Quantification and characterization of cockroach endosymbiont bacteria

The abundance of endosymbiont bacteria in the cockroach gut obtained was  $1.1 \times 10^4$  CFU/mL. A total of 20 bacterial isolates have different characteristics of macroscopic and microscopic. One of twenty isolate was difficult to rejuvenate (EKA 14) (data not shown). Macroscopic and microscopic characters of bacteria are shown in Table 1.

#### Hydrolytic activity

From a total of 20 bacterial isolates, 16 bacterial isolates had amylolytic activity, 15 bacterial isolates had proteolytic activity, 16 bacterial isolates had cellulolytic activity, and 11 bacterial isolates had lipolytic activity. The data is shown at Table 2. Observation of amylase, protease, and cellulase enzymes were carried out quantitatively, while observation the activity of lipase enzymes was carried out qualitatively by observing the presence of zones and fluorescence visible when exposed to UV light.

The results of the hydrolytic activity index showed the average index value of potential isolates was more than 1, with an average standard deviation of  $\pm 10\%$  (Table 2). It shows that the bacteria potential to produce amylase, protease, and cellulase. In the hydrolytic assay of amylase, 80% of bacterial isolates showed amylolytic activity with the highest index of 3.84 (EKA 20) and the lowest of 1.03 (EKA 9). As many as 75% of bacterial isolates

**Table 1.** Macroscopic and microscopic characteristic of cockroach endosymbiont bacteria

Bacteria Codes	Macroscopic Character					Microscopic Character		
	Color of Colony	Shape	Margin	Elevation	Structure	Shape of Cell	Existence of Spore	Gram Staining
EKA 1	Cream	Irregular	Undulate	Raised	Opaque	Rod	+	+
EKA 2	Yellow	Irregular	Undulate	Crateriform	Opaque	Shape Variable	-	-
EKA 3	Cream	Irregular	Entire	Flat	Translucent	Rod	+	+
EKA 4	Cream	Circular	Entire	Crateriform	Opaque	Rod	+	+
EKA 5	Cream	Irregular	Filiform	Raised	Opaque	Rod	+	+
EKA 6	Cream	Irregular	Undulate	Umbonate	Translucent	Rod	+	+
EKA 7	Cream	Irregular	Filiform	Flat	Translucent	Rod	+	+
EKA 8	Cream	Irregular	Undulate	Flat	Translucent	Rod	+	+
EKA 9	Cream	Irregular	Undulate	Flat	Translucent	Rod	+	+
EKA 10	Cream	Irregular	Entire	Convex	Opaque	Rod	+	+
EKA 11	Cream	Irregular	Curled	Flat	Opaque	Rod	+	+
EKA 12	Cream	Circular	Entire	Convex	Opaque	Rod	+	+
EKA 13	Cream	Irregular	Curled	Flat	Opaque	Rod	+	+
EKA 15	Orange	Irregular	Undulate	Raised	Opaque	Rod	+	+
EKA 16	Orange	Irregular	Undulate	Flat	Opaque	Rod	+	+
EKA 17	Cream	Irregular	Undulate	Raised	Opaque	Rod	+	+
EKA 18	Cream	Irregular	Undulate	Raised	Translucent	Rod	+	+
EKA 19	Cream	Irregular	Lobate	Flat	Translucent	Rod	+	+
EKA 20	Cream	Irregular	Curled	Flat	Opaque	Rod	+	+

**Table 2.** Hydrolytic activity index of endosymbiont bacterial isolates

Isolate code	Amylase Index	Protease Index	Celullase Index	Lipase Activity
EKA 1	2.39 ± 0.21	2.16 ± 0.17	1.67 ± 0.00	+++
EKA 2	2.66 ± 0.05	-	2.53 ± 0.05	-
EKA 3	1.74 ± 0.07	2.53 ± 0.29	1.58 ± 0.12	++
EKA 4	2.91 ± 0.04	5.94 ± 0.02	2.42 ± 0.34	+++
EKA 5	1.31 ± 0.04	2.16 ± 0.20	1.32 ± 0.13	-
EKA 6	2.10 ± 0.05	1.41 ± 0.02	1.44 ± 0.04	++
EKA 7	2.18 ± 0.12	1.97 ± 0.00	1.84 ± 0.20	+++
EKA 8	2.48 ± 0.14	2.95 ± 0.16	2.63 ± 0.13	+++
EKA 9	1.03 ± 0.05	-	1.47 ± 0.13	-
EKA 10	-	-	-	++
EKA 11	1.88 ± 0.04	-	2.17 ± 0.05	-
EKA 12	-	2.33 ± 0.19	-	+++
EKA 13	1.21 ± 0.03	1.09 ± 0.13	1.23 ± 0.10	-
EKA 15	1.15 ± 0.05	3.21 ± 0.35	1.01 ± 0.07	-
EKA 16	2.02 ± 0.17	4.02 ± 0.09	1.16 ± 0.13	-
EKA 17	1.05 ± 0.04	3.61 ± 0.00	1.11 ± 0.09	+
EKA 18	-	3.18 ± 0.29	-	+
EKA 19	3.30 ± 0.15	2.31 ± 0.08	3.99 ± 0.46	-
EKA 20	3.84 ± 0.01	2.64 ± 0.25	2.43 ± 0.05	+

(-) not detected, (+) fluorescent colony, (++) fluorescent zone, (+++) fluorescent colony and zone

showed proteolytic activity with the highest index of 5.94 (EKA 4) and the lowest of 1.09 (EKA 13). In the hydrolytic assay of cellulose, 80% of bacterial isolates showed cellulolytic activity with the highest index of 3.99 (EKA 19) and the lowest 1.01 (EKA 15).

Whereas 55% of bacterial isolates showed lipolytic activity, with potential isolates of 5 isolates (EKA 1, EKA 4, EKA 7, EKA 8, and EKA 12).

Several bacterial isolates were found to have the ability to multi-enzyme. Isolates capable of produc-

ing amylase, protease, cellulase, and lipase enzymes together were 8 isolates; they were EKA 1, EKA 3, EKA 4, EKA 6, EKA 7, EKA 8, EKA 17 and EKA 20. Bacterial isolates that were able to produce amylase, protease, and cellulase are 5 isolates, namely EKA 5, EKA 13, EKA 15, EKA 16, and EKA 19. Isolates that could produce amylase and cellulase are 3 isolates namely EKA 2, EKA 3, and EKA 11. Isolates that were able to produce protease and lipase are 2 isolates namely EKA 12, and EKA 18. Bacteria that have the most potential enzyme multi-activity were EKA 4, EKA 8 and EKA 20. Enzyme multi-activity of cockroach gut endosymbiont bacterial is shown in Fig. 1.




Fig 1. Summary of cocroach endosymbiont bacteria enzymaticpotency

**Identification of potential bacteria based on physiological character**

Physiological characters of EKA 4, EKA 8, and EKA 20 are shown in Table 3. Based on the book of Bergey’s Manual of Determinative Bacteriology, EKA 4, EKA 8, and EKA 20 were *Bacillus brevis* (EKA 4), *Bacillus badius* (EKA 8), and *Bacillus pantothenicus* (EKA20) with percent of probability 91.3% and 85.0%, 95.2% respectively.

**Discussion**

Anatomically, the digestive tract consists of crop, foregut, midgut, and hindgut (Cruden and Markovetz, 1987). Enzymatic digestion generally

Table 3. Physiological character of EKA 4, EKA 8, and EKA 20

Characteristic	EKA 4	EKA 8	EKA 20
Oxidase	+	+	-
Motility	+	+	+
Nitrate	+	+	+
Lysine	-	-	-
Ornithine	-	-	-
H <sub>2</sub> S	-	-	-
Glucose	-	-	-
Mannitol	-	-	-
Xylose	-	-	-
ONPG	+	-	+
Indole	-	-	-
Urease	-	-	-
VP	+	+	+
Citrate	-	-	-
TDA	-	-	-
Gelatin	+	+	+
Malonate	-	-	-
Inositol	-	-	-
Sorbitol	-	-	-
Rhamnose	-	-	-
Sucrose	-	-	-
Lactose	-	-	-
Arabinose	-	-	-
Adonitol	-	-	-
Raffinose	-	-	-
Salicin	-	-	-
Catalase	+	+	+
Arginine	-	-	-

occurs in the midgut intestine which produces degradation enzymes (Bagde *et al.*, 2013). Besides being produced by midgut, enzymes are also produced by endosymbiont microbes that live in the digestive tract. Each part of intestine has a different microbial community. The results of previous studies on the visualization of digestive tract microorganisms *Periplaneta americana* and *Shelfrodella lateral* use culture-dependent or culture-independent based method shows that the anterior colon has the highest density/ abundance (Bignell, 1976; Schauer *et al.*, 2012). Different cockroach diets greatly affect the diversity of bacteria. Omnivorous cockroaches *Periplaneta americana* has a higher endosymbiont diversity compared to wood-feeding cockroach *Cryptocercus punctulatus* (Colman *et al.*, 2012; Pérez-Cobas *et al.*, 2015).

Bacteria as providers a source of carbohydrate by increasing the efficiency of polymers such as lignin, hemicellulose and cellulose, xylan and pectin, can also contribute to the digestion of lipids and pro-



teins (Douglas, 2009; Visotto *et al.*, 2009). The role of endosymbiont bacteria in the digestive tract, among others, produces hydrolytic enzymes that decompose organic matter. The ability of cockroach endosymbiont bacteria to produce amylase enzymes was triggered by decomposer animal food. Bacteria could be present in large quantities in the digestive tract along with amylase and bacteria were involved in product digestion (Anand *et al.*, 2010). This is also supported by Supriyatna and Ukit's research (2016) that the presence of some cellulolytic bacteria in the intestines of black soldier fly larvae helped the digestion process in organic matter. Endogenous cellulase was present in some insects and termites (Taggar, 2015).

Some microbes that play a role in the digestive track of *Pycnoscelus surinamensis* were the group of Planctomycetaceae, Candidatus, Proteobacteria, Sulfurospirillum, Actinomyces, Lactobacillales, Betaproteobacteria, Desulfobivibrionaceae, Ruminococcaceae, Micrococcales, and Weissella (Richards *et al.*, 2017). This study found a group of rod-shaped Gram-positive spore forming bacteria endorsed by the Firmicutes group. This study confirm the results of previous studies conducted by Richards *et al.* (2017), using a culture-independent method namely Next-Generation Sequencing (NGS), found that the bacterial community of *Pycnoscelus surinamensis* was dominated by Bacteroidetes, Firmicutes and Proteobacteria. Predominantions of the Bacteroidetes and Firmicutes group were common in omnivorous groups including *Blatellagermanica*, *Shelfrodella lateralis* (Schauer *et al.*, 2012; Carrasco *et al.*, 2014; Perez-Cobas, 2015). Based on Correia *et al.*, (2018), Enterobacter found in intestinal track of *Pycnoscelus surinamensis*. Some species of the genus Enterobacter can produce cellulase enzyme (Sari *et al.*, 2017).

The results of the enzymatic test of the isolates shelter in the digestive tract of *Pycnoscelus surinamensis* showed that the eight isolates were also able to produce the digestive enzymes, amylase, protease, cellulase, and lipase. Cellulose-decomposing microorganisms become inseparable parts of the digestive tract microflora of cockroaches and other insects that eat material consisted of cellulose (Bagde *et al.*, 2013). Sharma *et al.* (2018), successfully isolated eight endosymbiont bacteria (14K, 16K, 22K, 24K, 28K, 29K, 30K, 31K) from the *Periplaneta americana* intestinal track. All of these isolates were

known to be able to excrete protease and cellulase enzymes. Isolate 29K which was genetically similarly with *Bacillus* spp. was able to excrete cellulase, protease, and keratinase. Cruden and Markovetz (1979), found in *Periplaneta americana* and *Euderes posticus* that there were a number of obligate anaerobic bacteria that can degrade carboxymethyl cellulose. In this study the enzymatic index method was used to compare the enzyme excretion produced by each bacterium. The enzymatic index method is a fast and simple way to select isolates that are most potential for enzyme excretion (Castro *et al.*, 2014). An index value above 1.0 is an indication of enzyme secretion (Carrim *et al.*, 2006). Based on research Florencio *et al.* (2012), strains that show an enzyme index higher than 1.50 were considered as potential enzyme producers. Exploration of bacteria that produce hydrolytic enzymes from the digestive tract of potential decomposers would then be continued.

## Conclusion

Eighty percent of isolated bacteria from cockroach's gut have potency to produce hydrolytic enzymes; amylase, protease, cellulase, and lipase. The potential bacteria producing these four enzymes are EKA 4, EKA 8 and EKA 20.

## Acknowledgement

The authors are thankful to Indonesian Ministry of Research, Technology, and Higher Education for funding in this research through the Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT) program

## References

- Anand, A.A.P.A., Vennison, S.J., Sankar, S.G., Prabhu, D.I.G., Vasan, P. T., Raghuraman, T., Geoffrey, C.J. and Vendan, S.E. 2010. Isolation and characterization of bacteria from the gut of *Bombyx mori* that degrade cellulose, xylan, pectin, and starch and their impact on digestion. *Journal of Insect Science*. 10:1-20. <https://doi.org/10.1673/031.010.10701>
- Bagde, U.S., Gopi, U. and Prasad, R. 2013. Isolation and characterization of gut-associated microbes in cockroach. *African journal of Microbiology Research*. 7 : 2034-2039. <https://doi.org/10.5897/AJMR12.844>
- Bakari, S.S., Laila, M.M., Maalim, M.K., Aboubakari, Z.M., Salim, L.A. and Ali, H.R. 2016. Characterization of household solid waste compost inoculated with ef-

- fective microorganism. *Modern Environmental Science and Engineering*. 2 : 194-200. [https://doi.org/10.15341/mese\(2333-2581\)/03.02.2016/007](https://doi.org/10.15341/mese(2333-2581)/03.02.2016/007)
- Banerjee, S. and Ghosh, K. 2014. Enumeration of gut associated extracellular enzyme-producing yeasts in some freshwater fishes. *Journal of Applied Ichthyology*. 30: 986-993. <https://doi.org/10.1111/jai.12457>
- Bergey, D.H. and Holt, J.G. 1994. *Bergey's Manual of Determinative Bacteriology*. 9th Edition. Williams & Wilkins: Maryland.
- Bhowmik, S., Islam, S., Ahmed, M.M., Hossain, M.B. and Hossain, M.A. 2015. Protease Producing Bacteria and Activity in Gut of Tiger Shrimp (*Penaeus monodon*). *Journal of Fisheries and Aquatic Science*. 10: 489-500. <https://doi.org/10.3923/jfas.2015.489.500>
- Bignell, D.E. 1976. Some observations on the distribution of gut flora in the American cockroach, *Periplaneta americana*. *Journal of Invertebrate Pathology*. 29:338-343. [https://doi.org/10.1016/S0022-2011\(77\)80040-2](https://doi.org/10.1016/S0022-2011(77)80040-2)
- Bright, M. and Bulgheresi, S. 2010. A complex journey: transmission of microbial symbionts. *Nature Reviews Microbiology*. 8 : 218-230. <https://doi.org/10.1038/nrmicro2262>
- Carrasco, P., Pérez-Cobas, A.E., van de Pol, C., Baixeras, J., Moya, A. and Latorre, A. Succession of the gut microbiota in the cockroach *Blattella germanica*. *International Microbiology*. 17 : 99-109. <https://doi.org/10.2436/20.1501.01.212>
- Carrim, A.J.I., Barbosa, E.C. and Vieira, M. 2006. Enzymatic activity of endophytic bacterial isolates of *Jacaranda decurrens* Cham. (Carobinha-do-campo). *Brazilian Archives of Biology and Technology*. 49 : 353-359. <https://doi.org/10.1590/S1516-89132006000400001>
- Castro, R.A., Quecine, M.C., Lacava, P.T., Batista, B.D., Luvizotto, D.M., Marcon, J., Ferreira, A., Melo, I.S. and Azevedo, J.L. 2014. Isolation and enzyme bioprospection of endophytic bacteria associated with plants of *Brazilian mangrove* ecosystem. *Springer Plus*. 3 : 1-9. <https://doi.org/10.1186/2193-1801-3-382>
- Colman, D.R., Toolson, E.C. and Takacs-Vesbach, C.D. 2012. Do diet and taxonomy influence insect gut bacterial communities?. *Molecular Ecology*. 21:5124-5137. <https://doi.org/10.1111/j.1365-294X.2012.05752.x>
- Correia, D. da S., Samuel, R.P., Diogo, N.P., Paula, V.M., Gustavo, R.X. and Maria, E.F.C. 2018. Microbial diversity associated to the intestinal tract of soil invertebrates. *Applied Soil Ecology*. 131 : 38-46. <https://doi.org/10.1016/j.apsoil.2018.07.009>
- Cruden, D.L. and Markovetz, A.J. 1979. Carboxymethyl cellulose decomposition by intestinal bacteria of cockroaches. *Applied and Environmental Microbiology*. 38:372-379. <https://aem.asm.org/content/38/3/369>
- Cruden, D.L. and Markovetz, A.J. 1987. Microbiology ecology of the cockroach gut. *Annual Reviews*. 41 : 617-643. *Applied and Environmental Microbiology*. 38 : 369-372. <https://doi.org/10.1146/annurev.mi.41.100187.003153>
- Douglas, A.E. 2009. The microbial dimension in insect nutritional ecology. *Functional Ecology*. 23 : 38-47. <https://doi.org/10.1111/j.1365-2435.2008.01442.x>
- Epstein, E. 1996. *The Science of Composting*. CRC Press LLC: Florida.
- Evangelista, D.A., Bourne, G. and Ware, J.L. 2013. Species richness estimates of Blattodeas.s. (Insecta: Dictyoptera) from Northern Guyana vary depending upon methods of species delimitation. *Systematic Entomology*. 39 : 150-158. <https://doi.org/10.1111/syen.12043>
- Febriyanto, A., Rusmana, I. and Rafiuddin, R. 2015. Characterization and identification of cellulolytic bacteria from gut worker *Macrotermes gilvus*. *HAYATI Journal Biosciences*. 22 : 197-200. <https://doi.org/10.1016/j.hjb.2015.07.001>
- Florencio, C., Couri, S. and Farinas, C.S. 2012. Correlation between agar plate screening and solid-state fermentation for the prediction of cellulase production by *Trichoderma* strains. *Enzyme Research*. 1-7. <https://doi.org/10.1155/2012/793708>
- Gontang, E.A., Aylward, F.O., Carlos, C., Glavina del Rio, T., Chovatia, M, Fern, A, Lo, C.C., Malfatti, S. A., Tringe, S.G., Currie, C.R. and Kolter, R. 2017. Major changes in microbial diversity and community composition across gut sections of a juvenile *Panlhora* cockroach. *Plos ONE*. 12:1-21. <https://doi.org/10.1371/journal.pone.0177189>
- Ifeanyi, O.T. and Odunayo, O.O. 2015. Microbiology of cockroach - A public health concern. *International Journal of Scientetic Research*. 4 : 485-488. <https://doi.org/10.36106/ijsr>
- Lohri, C.R., Diener, S., Zabaleta, I., Mertenat, A. and Zurbrugg, C. 2017. Treatment technologies for urban solid biowaste to create value products: are view with focus on low- and middle-income settings. *Reviews in Environmental Science and Bio-Technology*. 16 : 81-130. <https://doi.org/10.1007/s11157-017-9422-5>
- Moges, F., Eshetie, S., Endris, M., Huruy, K., Muluye, D., Feleke, T., Silassie, F.G., Ayalew, G. and Nagappan, R. 2016. Cockroaches as a sources of high bacterial pathogens with multidrugs resistant strains in Gondar Town, Ethiopia. *Biomed Research International*. 4:1-6. <https://doi.org/10.1155/2016/2825056>
- Morales, G.E. and Wolff, M. 2010. Insects associated with the composting process of solid urban waste separated at the source. *Revista Brasileira de Entomologia*. 54 : 645-653. <https://doi.org/10.1590/S0085-56262010000400017>
- Mullins, D.E. 2015. Physiology of environmental adaptations and resource acquisition in cockroaches. *An-*

- nual Review of Entomology*. 60 : 473-492. <https://doi.org/10.1146/annurev-ento-011613-162036>
- Patiño-Navarrete, R., Piulachs, M., Belles, X., Moya, A., Latorre, A. and Peretó, J. 2014. The cockroach *Blattella germanica* obtains nitrogen from uric acid through a metabolic pathway shared with its bacterial endosymbiont. *Biology Letter*. 10 : 384-396. <https://doi.org/10.1098/rsbl.2014.0407>
- Pérez-Cobas, A. E., Maiques, E., Angelova, A., Carrasco, P., Moya, A. and Latorre, A. 2015. Diets shapes the gut microbiota of the omnivorous cockroach *Blattella germanica*. *Federation of European Microbiological Societies*. 91 : 1-14. <https://doi.org/10.1093/femsec/fiv022>
- Pietri, J.E., Tiffany, C. and Liang, D. 2018. Disruption of the microbiota affects physiological and evolutionary aspects of insecticide resistance in the German cockroach, an important urban pest. *Plos ONE*. 13 : 1-17. <https://doi.org/10.1371/journal.pone.0207985>
- Richards, C., Otani, S., Mikaelyan, A. and Poulsen, M. 2017. *Pycnoscelus surinamensis* cockroach gut microbiota respond consistently to a fungal diet with out mirroring those of fungus-farming termites. *Plos ONE*. 12 : 1-19. <https://doi.org/10.1371/journal.pone.0185745>
- Roth, L.M. 1998. The cockroach genus *Pycnoscelus* scudder, with a description of *Pycnoscelus femapterus*, sp. nov. (Blattaria: Blaberidae: Pycnoscelinae). *Oriental Insects*. 32 : 93-130. <https://doi.org/10.1080/00305316.1998.10433770>
- Rozendaal, A. 1997. Vector control: Methods for use by individuals and communities. World Health Organization: England.
- Sari, W.N., Safika, Darmawi, and Yudha, F. Isolation and identification of a cellulolytic Enterobacter from rumen of Aceh cattle. *Veterinary World*. 2231-0961. <https://doi.org/10.14202/vetworld.2017.1515-1520>
- Schauer C., Thompson, C.L. and Brune, A. 2012. The bacterial community in the gut of the cockroach *Shelfordella lateralis* reflects the close evolutionary relatedness of cockroaches and termites. *Applied and Environmental Microbiology*. 78 : 2758-2767. <https://doi.org/10.1128/AEM.07788-11>
- Shaoli D. and Debnath, B. 2016. Prevalence of health hazards associated with solid waste disposal - A case study of Kolkata, India. *Procedia Environmental Sciences*. 35 : 201-208. <https://doi.org/10.1016/j.proenv.2016.07.081>
- Sharma, S., Prasad, R.K., Chatterjee, S., Sharma, A., Vairale, M.G. and Yadav, K.K. 2018. Characterization of *Bacillus* species with keratinase and cellulase properties isolated from feather dumping and cockroach gut. *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences*. 89 : 1079-1086. <https://doi.org/10.1007/s40011-018-1026-5>
- Sikora, L.J. 1998. Benefits and drawbacks to composting organic by-products. In: *Beneficial Co-Utilization of Agricultural, Municipal, and Industrial by-Products*. Sally Brown J. Scott Angle Lee Jacobs (Editors). Springer: Dordrecht. [https://doi.org/10.1007/978-94-011-5068-2\\_6](https://doi.org/10.1007/978-94-011-5068-2_6)
- Singh, R., Kumar, M., Mittal, A. and Mehta, P.K. 2016. Microbial enzymes: industrial progress in 21st century. 3 *Biotech*. 6(174): 1-5. <https://doi.org/10.1007/s13205-016-0485-8>
- Supriyatna, A. and Ukit, U. 2016. Screening and isolation of cellulolytic bacteria from gut of black soldier fly larvae (*Hermetia illucens*) feeding with rice straw. *Biosaintifika*. 8 : 314-320. <https://doi.org/10.15294/biosaintifika.v8i3.6762>
- Susanti, P.D. and Halwany, W. 2017. Dekomposisi serasah dan keanekaragaman mikroba di areal perikanan di Kabupaten Niyawai. *Jurnal Ilmu Kehutanan*. 11 : 212-223. <https://doi.org/10.22146/jik.28285>
- Taggar, M.S. 2015. Insect cellulolytic enzymes: Novel sources for degradation of lignocellulosic biomass. *Journal of Applied and Natural Science*. 7 : 625-. <https://doi.org/10.31018/jans.v7i2.656>
- Tarli, V.D., Pequeno, P.A.C.L., Franklin, E., W. De Morais, J., Zouza, J.L.P., Oliveira, A.H.C. and Guilherme, D.R. 2014. Multiple environmental controls on cockroach assemblage structure in a tropical rain forest. *Biotropica*. 46 : 1-10. <https://doi.org/10.1111/btp.12138>
- Tinker, K. A. and Ottesen, E. A. 2016. The core gut microbiome of the American cockroach, *Periplaneta americana*, is stable and resilient to dietary shifts. *Applied and Environmental Microbiology*. 82 : 6603-6610. <https://doi.org/10.1128/AEM.01837-16>
- Tuomela, M., Vikman, M., Hatakka, A. and Itavaara, M. 2000. Biodegradation of lignin in a compost environment: A review. *Bioresour Technol*. 72 : 169-183. [https://doi.org/10.1016/S0960-8524\(99\)00104-2](https://doi.org/10.1016/S0960-8524(99)00104-2)
- Visotto, L.E., Oliveira, M.G., Guedes, R.N., Ribon, A.O. and Good-God, P.I. 2009. Contribution of gut bacteria to digestion and development of the velvetbean caterpillar, *Anticarsia gemmatilis*. *Journal Insect Physiol*. 55 : 185-191. <https://doi.org/10.1016/j.jinsphys.2008.10.017>
- Warunasinghe, W.A.A.I. and Yapa, P.I. 2016. A survey on household solid waste management (SWM) with social reference to peri-urban area (Kottawa) in Colombo. *Procedia Food Science*. 12 : 718-729. <https://doi.org/10.1016/j.profoo.2016.02.038>
- Weisser, W.W. and Siemann, E. 2008. The various effects of insects on ecosystem functioning. In: Weisser, W. W. and Siemann, E. (Editors). *Insect and Ecosystem Function*. 173 : 3-24. Springer: Verlag Berlin Heidelberg. [https://doi.org/10.1007/978-3-540-74004-9\\_1](https://doi.org/10.1007/978-3-540-74004-9_1)



- Wynants, E., Froominckx, L., Crauwels, S., Verreth, C., De Sme, J., Sandrock, C., Wohlfahrt, J., Van Schelt, J., Depraetere, S., Lievens, B., Van Miert, S., Claes, J. and Van Campenhout, L. 2019. Assessing the microbiota of black soldier fly larvae (*Hermetia illucens*) reared on organic waste streams on four different locations at laboratory and large scale. *Environmental Microbiology*. 77: 913-930. <https://doi.org/10.1007/s00248-018-1286-x>
- Zhang, X.C. and Zhang, F. 2018. The potential control strategies based on the interaction between indoor cockroaches and their symbionts in china. Chapter Four. *Advances in Insect Physiology*. 55 : 55-122. <https://doi.org/10.1016/bs.aip.2018.07.001>
-