

Ecology, Environment and Conservation Editorial Advisory Board

Chief Editor

Prof.(Dr.) R.K.Trivedy, Pune, India

EDITORIAL ADVISORY BOARD

- | | |
|--|--|
| 1. Dr. Teresa Ferreira, Portugal | 19. Dr. A. Olawale, Nigeria |
| 2. Dr. Michael Ukwuru, Nigeria | 20. Dr. Ing. Agr. Mario Ridardo Sabbatini, Argentina |
| 3. Dr. Moses Inbaraj, Chennai | 21. Dr. Philip C. Reid, U.K. |
| 4. Dr. D.J. Lee, Taiwan | 22. Dr. Mohd. Yusuf, Malaysia |
| 5. Dr. Christial Paul P.delacruz, Phillipnes | 23. Dr. Oswaldo A. Feernandez, Argentina |
| 6. Dr. T. Bahorun, Mauritius | 24. Dr. Ms. Mirela Tulik, Warsaw, Poland |
| 7. Dr. Linda Blackwell, Australia | 25. Dr. L.L. Chukwu, Nigeria |
| 8. Dr. G. Zellner, Netherlands | 26. Dr. Azni Idris, UPM, Malaysia |
| 9. Dr. Wilson S. Tisera, Kupang, Indonesia | 27. Dr. Vikas Sharma, J&K, India |
| 10. Dr. M.F. Hamoda, Kuwait | 28. Dr. Amresh Chandra Pandey, Jharkhand, India |
| 11. Dr. H.A.Abrahamse, South Africa | 29. Prof.(Dr.) Agoes Soegianto, Indonesia |
| 12. Dr. Arulmozhiyal R., Salem | 30. Dr. A.K. Panigrahi, Berhampur, India |
| 13. Dr. Hassan Ibrahim Ali, Sudan | 31. Dr. Ahmed El Mahmoudi, Saudi Arabia |
| 14. Dr. A.R.Ghosh, Burdwan, India | 32. Dr. Seyed Mohammad Tajbakhsh, Iran |
| 15. Prof. M. Zaman, Bangladesh | 33. Dr. Amin L. Setyo, Indonesia |
| 16. Dr. Marcantonio Bragdin, Venice, Italy | 34. Dr. Francis Gbogbo, Ghana |
| 17. Dr. Z. Fuat Topark, Turkey | 35. Dr. S. Shabanlou, Iran |
| 18. Dr. Z. Li. Bonn, Germany | |

[Back to EEC Journal Details](#)

Ecology, Environment and Conservation Journal Papers

Issue: Vol 26, April Suppl. Issue, 2020

[PATHOGENIC FOOD-BORNE BACTERIA IN SHELLFISH AND SHRIMP FROM THE LARGEST TRADITIONAL SEAFOOD MARKET IN SURABAYA, INDONESIA](#)

Sin War Naw, Putu Angga Wiradana, Agoes Soegianto and Aondohemba Samuel Nege

[Get Abstract](#)

[Get Paper](#)

[DNA BAR CODING: STUDY OF BANANAS \(MUSA SPP.\) WILD AND CULTIVARS GROUP FROM EAST JAVA INFERRED BY RBC L GENE SEQUENCES](#)

Rojaunnajah Kartika Ainiyah, Verina Wahyunindita, Windi Nur Pratama, Intan Ayu Pratiwi, Edy Setiti Wida Utami and Sucipto Hariyanto

[Get Abstract](#)

[Get Paper](#)

[ROLE OF EDUCATION IN CONSERVING "KALEKA" FOREST IN DAYAK PEOPLE OF CENTRAL KALIMANTAN, INDONESIA](#)

Holten Sion, Muhamad Affandi and Febrina Natalia

[Get Abstract](#)

[Get Paper](#)

[DIVERSITY OF INDIGENOUS ENTOMOPATHOGENIC BACILLI FROM DOMESTICS BREEDING SITES OF DENGUE HEMORRHAGIC FEVER VECTOR BASED ON THE TOXICITY AGAINST AEDESAEGYPTI LARVAE](#)

Salamun, Ni'matuzahroh, Fatimah, M.I.F. Maswantari, M.U. Rizka, T. Nurhariyati and A. Supriyanto

[Get Abstract](#)

[Get Paper](#)

[PARASITIZATION OF PARASITIC PLANTS ON FRUIT PLANTS IN BANGKALAN REGENCY AND MALANG CITY, INDONESIA](#)

Solikin

[Get Abstract](#)

[Get Paper](#)

[CRUDE METHANOL EXTRACT OF BROTOWALI LEAVES \(TINOSPORA CRISPA\) AS BIOLARVICIDE AGAINST DENGUE VECTOR AEDES AEGYPTI](#)

Manikya Pramudya, Rosmanida, Nabilah Istighfari Zuraidassanaaz, Nadyatul Ilma Indah Savira, Elma Sakinatussajidah and Intan Permata Putri

[Get Abstract](#)

[Get Paper](#)

[THE EFFECT OF OKRA PODS \(ABELMOSCHUS ESCULENTUS L.\) METHANOL EXTRACT ON WHITE BLOOD CELL COUNT, PHAGOCYTTIC ACTIVITY, AND IFN-GAMMA LEVEL IN MUS MUSCULUS EXPOSED SODIUM NITRITE](#)

Sri Puji Astuti Wahyuningsih, Siti Richa Isnaini, Dwi Winarni, Win Darmanto, Manikya Pramudya, Anni Mar'atun Nafi'ah, Khoirunnisa, Elma Sakinatus Sajidah and Baiq Naili Dewi Atika

Search Articles

Journal Issues

[Vol 26, Issue 2, 2020](#)

[Vol 26, June Suppl. Issue, 2020](#)

[Vol 26, April Suppl. Issue, 2020](#)

[Vol 26, Issue 1, 2020](#)

[Vol 26, Feb Suppl. Issue, 2020](#)

[Vol 25, Issue 4 2019](#)

[Vol 25, Issue 3 2019](#)

[Vol 25, Nov Suppl. Issue, 2019](#)

[Vol 25, Sept Suppl. Issue, 2019](#)

[Vol 25, Aug Suppl. Issue, 2019](#)

[Vol 25, July Suppl. Issue, 2019](#)

[Vol 25, Issue 2 2019](#)

[Vol 25, May Suppl. Issue, 2019](#)

[Vol 25, April Suppl. Issue, 2019](#)

[Vol 25, Issue 1 2019](#)

[Vol 24, Issue 4 2018](#)

[Vol 24, Issue 3 2018](#)

[Vol 24, Issue 2 2018](#)

[Vol 24, Issue 1 2018](#)

[Vol 24, March Suppl. Issue 2018](#)

[Vol 24, Feb. Suppl. Issue 2018](#)

[Vol 23, Issue 4, 2017](#)

[Vol 23, Nov. Suppl. Issue 2017](#)

[Vol 23, Sept. Suppl. Issue 2017](#)

[Vol 23, Issue 2, 2017](#)

[Vol 23, Issue 3, 2017](#)

[Vol 23, Issue 1, 2017](#)

[Vol 23, Feb 2017 Suppl. Issue](#)

[Vol 22, Dec 2016 Suppl. Issue](#)

[Vol 22, Issue 4, 2016](#)

[Vol 22, Sept. Suppl. Issue, 2016](#)

[Vol 22, Issue 3, 2016](#)

[Vol. 22, June Suppl. Issue 2016](#)

[Vol 22, Issue 2, 2016](#)

[Vol. 22, April Suppl. Issue 2016](#)

[Vol 22, Issue 1, 2016](#)

[Vol 21, Issue 4, 2015](#)

[Vol. 21 Dec. 2015 Suppl. Issue](#)

[Vol. 21 Nov. 2015 Suppl. Issue](#)

[Vol 21, Issue 3, 2015](#)

[Vol 21, Issue 2, 2015](#)

[Vol. 21 Suppl. Issue August 2015](#)

[Vol 21, Suppl. Issue June 2015](#)

[Vol 21, Issue 1, 2015](#)

[Supplement Issue, Dec. 2014](#)

[Special Issue-2014](#)

[Get Abstract](#)[Get Paper](#)

SPINY LOBSTERS SPECIES COMPOSITION BETWEEN NORTH AND SOUTH SEA OF EAST JAVA, INDONESIA

Arief Setyanto, Soemarno, D.G.R. Wiadnya, Cahyo Prayogo, R.J. West, M. Tsamenyi, S. Halimah and N.A. Rahman

[Get Abstract](#)[Get Paper](#)

ECOLOGICAL STUDY OF PRIMITIVE BRACHIOPODS LINGULA SP. IN PROBOLINGGO, EAST JAVA, INDONESIA

Rakmawati and Sucipto Hariyanto

[Get Abstract](#)[Get Paper](#)

IDENTIFICATION AND TOXICITY TEST OF ORTHOSIPHONSTAMINEUS LEAVES EXTRACT AGAINST HOUSEFLIES(MUSCA DOMESTICA)

Surahmaida, Umarudin and Junairiah

[Get Abstract](#)[Get Paper](#)

THE EFFECTS OF LEAD ACETATE EXPOSURE ON BLOOD COMPONENT AND KIDNEY: THE MECHANISM OF OXIDATIVE STRESS

Sugiharto, Win Darmanto, Sri Puji Astuti W., Kevita Putri, Nur Khoirunnisa, Dea B. Mulyawan, and Arin N. Shanti

[Get Abstract](#)[Get Paper](#)

THE EFFECTS OF LEAD ACETATE EXPOSURE ON BLOOD COMPONENT AND KIDNEY: THE MECHANISM OF OXIDATIVE STRESS

Sugiharto, Win Darmanto, Sri Puji Astuti W., Kevita Putri, Nur Khoirunnisa, Dea B. Mulyawan, and Arin N. Shanti

[Get Abstract](#)[Get Paper](#)

AN EVIDENCE OF THE MISSING COMPLETE PECTORAL FINS TRAIT IN CLARIAS GARIEPINUS REARED IN POND IS HERITABLE

Farikhah Farikhah, Sukoso Sukoso, Uun Yanuhar, Feni Iranawati and Muchlis Zainudin Arif

[Get Abstract](#)[Get Paper](#)

DIVERSITY OF MODERATELY THERMO-ALKALIPHILE BACTERIA PRODUCING AMYLASE ORIGINATED FROM PARIANGAN HOT SPRING, WEST SUMATRA, INDONESIA

Gustina Indriati, Ruth Rize Pass Megahati and Dwi Hilda Putri

[Get Abstract](#)[Get Paper](#)

LIPID ACCUMULATION OF SKLETONEMA COSTATUM UNDER SILICA, NITROGEN AND AERATION STRESS

Tutik Nurhidayati, Aini Ika Rahmawati, Endang Purwanti Setyaningsih, Kristanti Indah Purwani, Edwin Setyawan and Nurul Jadid

[Get Abstract](#)[Get Paper](#)

TOXICITY EFFECT OF COPPER ON HISTOPATHOLOGY OF OREOCHROMIS NILOTICUS LIVER WITH DIET OF LACTIC ACID BACTERIA

Alfiah Hayati, Rosana Dyah Umami, Aken Puti Wanguyun, Primadika Ashar Dwi Putra, Afinia Primananda Taufiq and Agus Supriyanto

[Get Abstract](#)[Get Paper](#)[Vol.20, Issue 4, 2014](#)[Vol.20, Issue 3, 2014](#)[Vol.20, Issue 2, 2014](#)[Vol.20 Issue 01, 2014](#)[Vol.19 Issue 04, 2013](#)[Vol.19 Issue 03, 2013](#)[Vol.19, Issue 02, 2013](#)[Vol.19, Issue 01, 2013](#)[Vol.18, Issue 04, 2012](#)[Vol.18, Issue 3, 2012](#)[Vol.18, Issue 2, 2012](#)[Vol.18, Issue 1, 2012](#)[Vol.17, Issue 4, 2011](#)[Vol.17, Issue 3, 2011](#)[Vol.17, Issue 2, 2011](#)[Vol.17, Issue 1, 2011](#)[Vol.16, Issue 4, 2010](#)[Vol.16, Issue 3, 2010](#)[Vol.16, Issue 2, 2010](#)[Vol.16, Issue 1, 2010](#)[Vol.15, Issue 04, 2009](#)[Vol.15, Issue 03, 2009](#)[Vol.15, Issue 02, 2009](#)[Vol.15, Issue 1, 2009](#)[Vol.14, Issue 04, 2008](#)[Vol.14, Issue 2-3, 2008](#)[Vol.14, Issue 2-3, 2008](#)[Vol.14, Issue 1, 2008](#)[Vol.14, Issue 2-3, 2008](#)[Vol.13, Issue 04, 2007](#)[Vol.13, Issue 2, 2007](#)[Vol.13, Issue 1, 2007](#)[Vol.12, Issue 4, 2006](#)[Vol.12, Issue 3, 2006](#)[Vol.12, Issue 2, 2006](#)[Vol.12, Issue 1, 2006](#)[Vol.12, Issue 01, 2006](#)[Vol.11, Issue 3,4, 2005](#)[Vol.11, Issue 2, 2005](#)[Vol.11, Issue 1, 2005](#)[Vol.10, Issue 04, 2004](#)[Vol.10, Issue 03, 2004](#)[Vol.10, Issue 02, 2004](#)[Vol.10, Issue 01, 2004](#)[Vol.09, Issue 04, 2003](#)[Vol.09, Issue 03, 2003](#)[Vol.09, Issue 02, 2003](#)[Vol.08, Issue 04, 2002](#)[Vol.08, Issue 03, 2002](#)[Vol.08, Issue 01, 2002](#)[Vol.07, Issue 04, 2001](#)[Vol.07, Issue 03, 2001](#)[Vol.07, Issue 02, 2001](#)[Vol.07, Issue 01, 2001](#)[Vol.06, Issue 04, 2000](#)[Vol.06, Issue 03, 2000](#)[Vol.06, Issue 02, 2000](#)[Vol.06, Issue 01, 2000](#)[Vol.05, Issue 04, 1999](#)[Vol.05, Issue 03, 1999](#)[Vol.05, Issue 02, 1999](#)[Vol.05, Issue 01, 1999](#)[Vol.04, Issue 1,2, 1998](#)[Vol.03, Issue 3,4, 1997](#)[Vol.03, Issue 01, 1997](#)

RISK ASSESSMENT OF INVASIVE ALIEN PLANT SPECIES IN PANGANDARAN NATURE RESERVE, WEST JAVA, INDONESIA

Arif Mukhtar Hasbi and Dian Rosleine

[Get Abstract](#)

[Get Paper](#)

PHENOTYPIC DIVERSITY OF FIVE GOAT POPULATIONS IN TROPICAL ENVIRONMENT

Arsyad Alawiansyah, Richi Yuliavian Kusminanto, Nuzul Widyas, Ahmad Pramono, Muhammad Cahyadi and Sutarno

[Get Abstract](#)

[Get Paper](#)

NEW ISOLATION OF HYDRO CARBONOCLASTIC BACTERIA FROM LOCAL LIMESTONE MINING AS PROMISING CONCRETE STRENGTH ENHANCER

Enny Zulaika, M. Andry Prio Utomo, Nur Hidayatul Alami, Endry Nugroho Prasetyo, Tri Wulan and Agoes Soegianto

[Get Abstract](#)

[Get Paper](#)

BIRD DIVERSITY OF RESORT RANU DARUNGAN, BROMO TENGGER SEMERU NATIONAL PARK, INDONESIA

Boni Herdiawan, Reni Ambarwati and Toni Artaka

[Get Abstract](#)

[Get Paper](#)

CHARACTER VISUALIZATION OF PYCNOSCELUS SP. (BLATTODEA: COCKROACH) IN HOUSEHOLD ORGANIC WASTE COMPOSTER IN SURABAYA

Moch.Affandi, Emilia Anjar Prastiwi, Nindya Ayu Damayanti, Chairunisa Firdaus, Azizah Kusumadewi, Anisa Rahmawati, Puspanjali Prahasto, Audry Putri Kristanti, Alifatus Alkurnia Sukma Firdaus, Hardiansah Pramana Abdilah, Muhammad Arifudin Mubarak, Syahriar

[Get Abstract](#)

[Get Paper](#)

ISOLATION AND CHARACTERIZATION OF COCKROACH ENDOSYMBIONT BACTERIA WITH POTENTIAL TO PRODUCE HYDROLYTIC ENZYME OF ORGANIC MATERIAL

Ni'matuzahroh, Nastiti Trikurniadewi, Syahriar Nur Maulana Malik Ibrahim, Achmad Zainal Abidin, Ana Mariatul Khiftiyah, Silvia Kurnia Sari, Eka Narendra Nuswantara, Fawwas Nurmansyah, Muhammad Aufar Rizki Wildan Rahman, Habibah Lailatul Maghfirah, Miftahul

[Get Abstract](#)

[Get Paper](#)

MORPHOPHYSIOLOGICAL CHARACTERISTIC RESPONSES OF SOYBEAN (GLYCINE MAX L.) GROBOGAN VARIETY IN WATER LOGGING STRESS

Vita Siti Fatimah and Tutik Nurhidayati

[Get Abstract](#)

[Get Paper](#)

THE NUMBERS OF LYMPHOCYTES AND CAPILLARIES IN SILICA EXPOSED LUNG TISSUES OF BALB/C MICE (MUSMUSCULUS) TREATED WITH MORINGAOLEIFERA LEAVES EXTRACT

Kusmiyati, Soedjajadi Keman, Muhammad Amin, Suwarno and Agoes Soegianto

[Get Abstract](#)

[Get Paper](#)

BIODIVERSITY RICHNESS OF TERRESTRIAL ORCHIDS IN BROMO TENGGER SEMERU NATIONAL PARK, INDONESIA

Lita Soetopo and Muhamad Jauhari

[Get Abstract](#)

[Get Paper](#)

CHARACTERISTICS OF RICE PLANT WITH LOW METHANE EMISSION

Noer Rahmi Ardiarini, Finsa Dwi Arisandi and Prihasto Setyanto

[Vol.02, Issue 1,2, 1996](#)

[Vol.01, Issue 14, 1995](#)

Looking for Past Issues?

[Click here to get them!!](#)

[Get Abstract](#)[Get Paper](#)

EXPLORING THE LANDRACE RICE (ORYZA SATIVA L) FOR SALINITY TOLERANT AND AGRONOMIC TRAITS

Nafisah, Celia Roza, Nani Yunani, Rahmini and Priatna Sasmita

[Get Abstract](#)[Get Paper](#)

BIODIVERSITY OF PLANT GROWTH PROMOTING RHIZO BACTERIA FROM RHIZOSPHERE OF PINUS SP AND CASUARINA SP AT RADEN SOERJO FOREST PARK

Brigita Nur Diyan Agustiana, Indra Gumilar, RizqiaNurul Alfiyanita, Alifah Nastiti, Tesalonika Tetya Virginia, Siti Rizqiyatul Mukarromah, Annisa Dwi Savitri, Lunace Rahma Kinayoh, Erna Erviana Nadhifa, Khafid Jallaludhin, Ni'matuzahroh and Almando Gerald

[Get Abstract](#)[Get Paper](#)

CORN COB HYDROLYZATE FROM PENICILLIUM CITRINUM H9 AS AN ALTERNATIVE SUBSTRATE FOR BIO SURFACTANT PRODUCTION BY HYDROCARBONOClastic BACTERIA

Fatimah, Silvia Kurnia Sari, Nastiti Trikurniadewi, Syahriar Nur Maulana Malik Ibrahim, Ana Mariatul Khiftiyah, Khudrotul Nisa Indriyasari, Tri Nurhariyati, Tini Surtiningsih and Hanif Yuliani, Ni'matuzahroh

[Get Abstract](#)[Get Paper](#)

INFESTATION OF BIO FOULINGS (BALANUS SP. AND LITTORARIASp.) ON MANGROVE RHIZOPHORA SP. AND AVICENNIA SP. AT GADINGREJO, PASURUAN, INDONESIA

Oktiyas Muzaky Luthfi, Evi Safitri and Agoes Soegianto

[Get Abstract](#)[Get Paper](#)

LENGTH-WEIGHT RELATIONSHIP AND CONDITION FACTOR OF SYGNATHOIDES BIACULEATUS (BLOCH, 1785) CAUGHT FROM SEA GRASS AREA OF TANAKEKE ISLANDS, TAKALAR DISTRICT, SOUTH SULAWESI, INDONESIA

Ardi Eko Mulyawan, Yeni Savitri Andi Lawi, Syaifiuddin, Farid Samawi and Agoes Soegianto

[Get Abstract](#)[Get Paper](#)

COMPARISON OF WATER MASS CHARACTERISTICS BETWEEN THE INDONESIAN THROUGH FLOW EXIT PASSAGE AND SOUTH JAVA SEA, INDONESIA

B. Semedi, A. Purnomosakti, D. K. Saputra, I.B. Wahyono and A. Soegianto

[Get Abstract](#)[Get Paper](#)

COMPARISON OF WATER MASS CHARACTERISTICS BETWEEN THE INDONESIAN THROUGH FLOW EXIT PASSAGE AND SOUTH JAVA SEA, INDONESIA

B. Semedi, A. Purnomosakti, D. K. Saputra, I.B. Wahyono and A. Soegianto

[Get Abstract](#)[Get Paper](#)

MORPHOLOGY-BASED GENETIC DIVERSITY OF EARLY MATURITY RICE GERM PLASM, THE RELATION TO YIELD COMPONENT

Indria W. Mulsanti, Nani Yunani and Trias Sitaresmi

[Get Abstract](#)[Get Paper](#)

THE POPULATION OF SOLEN SP. (BIVALVIA: SOLENIDAE) FROM PAMEKASAN, INDONESIA

Reni Ambarwati and Bambang Irawan

[Get Abstract](#)[Get Paper](#)

LARVICIDAL ACTIVITY OF METHANOL FRACTIONS FROM CARICAPAPAYA LEAVES EXTRACT AGAINST AEDES AEGYPTI

Tri Puji Lestari Sudarwati, M.A.H. Ferry Fernanda, Hilya Nur Imtihani and Intan A. Pratiwi

[Get Abstract](#)

[Get Paper](#)

MOLECULAR DETECTION AND MORPHOLOGICAL VARIED OF BLASTOCYSTIS SP IN BEEF CATTLE IN SIAK SRI INDRAPURA RIAU INDONESIA

Lucia Tri Suwanti, Mufasirin, Nunuk Dyah Retno Lastuti, Yuli Susana and Suci M. Rizki

[Get Abstract](#)

[Get Paper](#)

REMOVAL OF LEAD (II) FROM AQUEOUS SOLUTION USING COMPOSITE ADSORBENT OF CASSAVA PEEL WASTE/MAHOGANY SAWDUST/BENTONITE

Eko Prasetyo Kuncoro, Ganden Supriyanto, Thin Soedarti, Nilaturrohmah and Gabriel Putra Utama

[Get Abstract](#)

[Get Paper](#)

[Home](#) | [International Journals](#) | [Books](#) | [Environmental Consulting](#) | [About Us](#) | [Contact Us](#) | [Submit Paper](#) | [Search Journal Article](#) |

[Become a fan](#) on Facebook

[Follow us](#) on Twitter





© EM International 2012-2019 | **Developed by Eneblur Consulting**

molecutrack

We Focus on Clean Energy Solutions That Replace Hazardous and Carbon Intensive Fuel.


molecutrack.com OPEN

Ecology, Environment and Conservation

Country	India -  SJR Ranking of India
Subject Area and Category	Agricultural and Biological Sciences Ecology, Evolution, Behavior and Systematics Environmental Science Ecology Nature and Landscape Conservation
Publisher	EM International
Publication type	Journals
ISSN	0971765X
Coverage	1997-2019
Scope	Published Quarterly Since 1995. Ecology, Environment and Conservation is published in March, June, September and December every year. ECOLOGY, ENVIRONMENT AND CONSERVATION is one of the leading International environmental journal. It is widely subscribed in India and abroad by Institutions and Individuals in education and research as well as by Industries, Govt. Departments and Research Institutes.
	Homepage How to publish in this journal Contact  Join the conversation about this journal








12

H Index



SJR +

Citations per document +



Ecology, Environment and Conservation

Q4 Ecology
best quartile

SJR 2019
0.14

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```

L **Leta Rafael Levis** 2 weeks ago

I have published my article on this journal in 2017, no 23 Issue 2 with the title of the article was :
Farmers attitude
How ever, I have got the URL for full text.
So, please let me know the URL for the article full text.

Change my email address: in 2017 when I sent my article to this journal I still use email address:

Corn Cob Hydrolyzate from *Penicillium citrinum* H9 as an alternative substrate for Biosurfactant production by Hydrocarbonoclastic Bacteria

Fatimah¹², Silvia Kurnia Sari¹², Nastiti Trikurniadewi¹², Syahriar Nur Maulana Malik Ibrahim¹², Ana Mariatul Khiftiyah¹², Khudrotul Nisa Indriyarsari¹, Tri Nurhariyati¹², Tini Surtiningsih¹² and Hanif Yuliani³, Ni'matuzahroh^{12*}

¹Department of Biology, Faculty of Science and Technology, Universitas Airlangga, Surabaya, Indonesia

²Research Center for Bio-Molecule Engineering, Universitas Airlangga, Surabaya, Indonesia

³Agency for the Assessment and Application of Technology, BPPT, South Tangerang, Indonesia

(Received 27 September, 2019; Accepted 10 January, 2020)

ABSTRACT

Corn cob is a potential agricultural waste that can be utilized as raw materials in bioindustries. Commonly, corn cob is regarded as agricultural waste and would be eliminated by burning, however this elimination method would produce a toxic fume that polluted the atmosphere. So, new utilization methods of corn cob need to be developed. This research aimed to reveal the potency of corn cob hydrolyzate (CCH) that produced enzymatically using *Penicillium citrinum* H9 as substrate for biosurfactant production. The CCH was used as growth substrate for biosurfactant production by seven indigenous bacteria of Balongan oil sludge. All isolates were tested the growth and the capability of biosurfactant production through surface tension and emulsification activity measurement. One potential isolate was obtained from this observation. BP1(5) isolate has a lowering surface tension at 56.64 ± 4.63 mN/m in five days incubation and the growth rate was 0.52 "cell/hour. The best biosurfactant product was obtained at concentration 200 ppm of CCH in five days incubation by 23.93 ± 3.56 mN/m in surface tension reductions but the emulsification activity was not detected. This research showed that the CCH be utilized as substrate for biosurfactant production by hydrocarbonoclastic bacteria.

Key words: Biosurfactant, Corn Cob Hydrolyzate, Hydrocarbonoclastic Bacteria

Introduction

Agricultural waste especially corn cob in Indonesia, are increasing very year, and to reduce their accumulation, most of them are eliminated by burning. Some Indonesian researchers have observed their utilization to decrease the waste and increase the value by conversion, for instance, to feedstock with

high protein (Marhaeni, 2002), as well as, to xylose and glucose (Oktaviani *et al.*, 2019). Furthermore, corn cob has been reported that it is promising substrate for biosurfactant production due to their low cost (Ni'matuzahroh *et al.*, 2019a).

The world's interest in biosurfactants has increased considerably in recent years due to their potential application in various industrial fields. In

the pharmaceutical sector biosurfactants were used due to their anti-bacterial properties, anti-biofilm and anti-cancer activities (Rangarajan and Sen, 2013; Karlapudi *et al.*, 2018). In the food industry, biosurfactants were used as emulsifiers of the bread making process. Furthermore, biosurfactants can also be employed for bioremediation of environments polluted by petroleum and heavy metals (Banat *et al.*, 2000; Mulligan and Wang, 2006).

Based on global market insights, the biosurfactant market in 2017 reached USD 1.8 billion and predicted to be more than USD 2.7 billion in 2024. The high biosurfactants demand is caused by their low toxicity, biodegradability, and stable in temperature, pH and salinity with broad range compared to synthetic surfactants (Zambry *et al.*, 2017; Varjani *et al.*, 2017). Despite people's interest in biosurfactants, the molecules are still unable to compete with synthetic surfactants in the world market because of high production cost, inefficient production methods, and expensive substrates requirement (Makkar and Cameotra, 2002). The solution for this problem is to explore the potential of isolates, optimize production, and use inexpensive raw materials and substrates from waste. Several previous studies have published the ability of organic waste such as rice straw, wheat straw, and sugar cane-bagasse to hydrolyze to produce sugar as a substrate for biosurfactant production (Joy *et al.*, 2019; Ni'matuzahroh *et al.*, 2019b).

This study used corn cob as a substrate for biosurfactant production by hydrocarbonoclastic bacteria. Corn cob consist of mainly three types of polymers cellulose, hemicellulose, and lignin, so that it is potentially hydrolyzed (Ni'matuzahroh *et al.*, 2018; Saha, 2004). Previous studies have shown that corn cob can be hydrolyzed into simple sugars and used for the production of biosurfactants by *Bacillus subtilis* 3KP with a reduction in surface tension of $12.55 \pm \pm 3,95\text{mN/m}$ (Ni'matuzahroh *et al.*, 2018). This paper explored the potency of corn cobhydrolysate as a substrate for biosurfactant production by hydrocarbonoclastic bacteria isolates with variations in substrate concentration and incubation time.

Materials and Methods

Microorganism

Hydrocarbonoclastic bacteria, which were isolated from Balongan oil sludge in the previous study

(Ni'matuzahroh *et al.*, 2019a) and collection of microbiology laboratory of Universitas Airlangga, were used in this study. The seven phenotypically different colonies obtained, namely BP(1)1, BP(1)2, BP(1)3, BP(1)4, BP(1)5, BP(1)6, and BP(1)8, were initially transferred onto Nutrient Agar (NA) slant for re-culture before used as an inoculum for screening biosurfactant production using corn cob hydrolyzate (CCH) as main substrate.

Enzymatic hydrolysis of corn cob by *Penicillium citrinum*. H9

Corn cob was collected from plantation in the Bojonegoroarea, East Java. Sample was dried and ground roughly into 1-5 mm pieces. Chemical delignification process was carried out by mixing corn cob powder with NaOH 1% solution with a ratio of 1:10 (w/v) for an hour at 100 °C. The mixture was filtered and washed to adjust the pH into 7 then dried in an oven for 24 hours at 60 °C. The hydrolysis process was carried out enzymatically by *Penicillium citrinum* H9. Two grams of pretreatment corn cob as the sole carbon source was added to 100 mL of Mandel-Sternberg's medium which consisted of KH_2PO_4 (2 g/L), $(\text{NH}_4)_2\text{SO}_4$ (1.4 g/L), CaCl_2 (0.3 g/L), $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (0.3 mg/L), $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ (5 mg/L), $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ (1.6 mg/L), $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ (1.4 mg/L), and CoCl_2 (2 mg/L) and arranged at pH 5 using citrate buffer. The media was cooled to room temperature then it was add 4% (v/v) of spore suspension of *Penicillium citrinum* H9 and was incubated in shaker incubator for 6 days. The concentration of corn cob hydrolysate sugar was measured by Somogyi-Nelson's method after the glucose standard equation was made by using the same method.

Screening for an efficient isolate for biosurfactant production

A loopful of each purified isolate was separately inoculated into 100 mL Erlenmeyer flasks containing 20 mL nutrient broth and incubated for 24 hours. This culture was used for starter. Screening which was taken for potentials isolates in biosurfactant production was carried out on mineral water synthetic medium with a total volume of 20 mL. The media was added with corn con hydrolysate (CCH) with a concentration of 209.3 ppm as much as 7.5% (v/v) and bacterial starter 2% (v/v) OD_{650} 0.5. The cultures were agitated at 120 rpm, at 30 °C for 0,1,3, and 5 days. Potential isolate was screened on the basis of emulsification, reduction of

surface tensions of fermented broth, and total plate count. The potential isolate then was used on the optimization of biosurfactant production process.

Optimization of biosurfactant production

The optimization of biosurfactant production was conducted by varying the concentration of corn cob hydrolysate sugar and the incubation time. The concentration of corn cob hydrolysate sugar was varied (0, 100, and 200 ppm) with 0, 1, 3, and 5 incubation time. Become these factors were chosen to obtain higher productivity of the biosurfactant. Mineral water synthetic medium utilized for biosurfactant production in this study was modified from Pruthi and Cameotra (1997) with the composition of: $(\text{NH}_4)_2\text{SO}_4$ (3 g/L), NaCl (10 g/L), $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (0.2 g/L), CaCl_2 (0.01 g/L), $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ (0.001 g/L), H_3BO_3 (0.001 g/L), $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ (0.001 g/L), $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ (0.001 g/L), $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ (0.005 g/L), and $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$ (0.001 g/L). Mineral water synthetic medium's buffer consisted of KH_2PO_4 (5 g/50 mL), K_2HPO_4 (2.6207 g/50 mL), and Fe_3O_4 (0.0006 g/50 mL). The total culture volume was 20 mL which consisted of mineral water synthetic medium, 7.5 % (v/v) of corn cob hydrolysate sugar, and 2% (v/v) of bacteria suspension. Microbial culture was incubated in a shaker at room temperature with the agitation rate of 120 rpm.

Biosurfactant productivity test

Before the bacteria cultures were tested for their surface tension (ST) and emulsification activity (EA), the optical density was measured by spectrophotometry. Samples of the culture medium were centrifuged for 15 minutes with an agitation rate of 3000 rpm. As many as 10 mL of supernatant was taken for surface tension analysis using Tensiometer Du-Nuoy ring method. The value of its surface tension was stated in mN/m or equivalent to dynes/cm. Its surface tension was calculated as follows:

$$r = r_0 \frac{\theta}{\theta_0} \quad (1)$$

where r_0 was water surface tension at $t^\circ\text{C}$, θ was the sample values shown on the Tensiometer, and θ_0 was distilled water values shown on the Tensiometer Du-Nuoy. The emulsification activity was obtained by mixing 1 mL of supernatant and 1 mL of kerosene then homogenized with vortex for 2 minutes. The emulsification activity was observed after 1 hour and 24 hours and calculated using the fol-

lowing formula:

$$EA(\%) = \frac{\text{Total height of the emulsified layer}}{\text{Total height of the liquid layer}} \times 100\% \quad (2)$$

Results and Discussion

Enzymatic hydrolysis of corn cob

Corn cob was composed by heterogeneous complexes of carbohydrate polymers such as lignin, cellulose, and hemicellulose that can be hydrolyzed into simple sugar. However cellulose and hemicellulose are coated with lignin, which protects them against enzymatic hydrolysis. Physical and chemical pretreatment (delignification) were performed to break the lignin seal to expose cellulose and hemicellulose for enzymatic action. Corn cob was physically treated by grinding and milling to reduce particle size and increase the surface area that will be accessed by *Penicillium citrinum* H9. In addition to further optimizing the hydrolysis process, chemical delignification is also carried out by using NaOH. Alkali delignification using NaOH has been proven to be more effective in breaking ester bonds between hemicellulose, cellulose, and lignin compared to acidic or oxidative pretreatment, while it also being able to avoid hemicellulose fragmentation polymer, thus making the biomass increasing in the porosity (Tarkow and Feist, 1969; Gaspar *et al.*, 2007).

Zhang and Cai (2008) stated that delignification of rice straw using 2% NaOH for 1 hour at 85 °C was able to reduce lignin content up to 36%. While in this research, it was showed that corn cob delignification using 1% NaOH at 100 °C for 1 hour was able to increase the hemicellulose content until 10.48% and cellulose 17.22%, and reduce the lignin up to 7.07%.

According to this result, it indicates that delignification process plays an important role in decreasing lignin content and increasing cellulose and hemicellulose content, so it is easy to be hydrolyzed enzymatically. Enzymatic hydrolysis is the second step in the production of biosurfactants from lignocellulosic waste. It involves the breakdown of compound cellulose and hemicellulose using enzymes. The hemicellulose contains several sugars such as xylan, mannan, glucan, arabinan, and galactan, while cellulose is only composed of glucans. Therefore, the main hydrolysis product of the hemicellulose was pentoses and hexoses while cellulose was glucose (Taherzadeh and Niklasson, 2004).

Screening for an efficient bacteria using CCH as carbon source

Seven indigenous bacteria of Balongan oil sludge had been successfully isolated in previous studies (Ni'matuzahroh *et al.*, 2019a). All isolates were tested for growth and their ability to produce biosurfactants on the CCH substrate. The results of the test are shown in the Table 1.

Based on the Table 1, it can be seen that bacteria can grow in the substrate. Based on the growth rate and generation time of all bacterial isolates, show that all isolates are able to use CCH as a carbon source for their growth. This is due to the hydrolysis of corn cob contains xylose, glucose, glycerol, arabinose and acetic acid (Chen *et al.*, 2019 Moldes *et al.*, 2007).

In addition to be used for growth substrate, CCH has also been proven capable of being used as a substrate for biosurfactant production. This is consistent with the study of Moldes *et al.*, (2007) which

states that *Lactobacillus pentosus* grown on reducing sugar of corn cob and able to reduce surface tension by 18 mN/m. Biosurfactant production were indicated by presence of surface tension and emulsification activity (El-Sheshtawy *et al.*, 2015). Based on this research, all of the isolates didn't show emulsification activity, while the lowest surface tension were shown by BP(1)5 after 5 days incubation (Table 1). Surface tension of BP(1)5 was also lower than uninoculated media, as many as 72.69 mN/m. Furthermore BP(1)5 isolate showed surface tension reduction after 1, 3, and 5 days of incubation, in contrast to other isolates which showed fluctuated value of surface tension reduction (Fig. 1). From this screening proses we obtained the potential isolate for biosurfactant production using CCH which is BP(1)5. BP(1)5 is used in the optimization of biosurfactant production with variations in substrate concentration and incubation time.

Table 1. The growth rate, generation time, and surface tension of BP(1)5 bacteria on CCH substrate

Isolates	Log TPC at 5 th day (CFU/mL)	Growth Rate (cell/hour)	Generation Time (hour)	Surface Tension (mN/m) 5 th day
BP(1)1	19.57±0.07	0.42	2.38	57.06±1.17
BP(1)2	14.83±0.00	0.24	4.18	72.17±0.00
BP(1)3	17.80±2.91	0.36	2.82	60.01±3.55
BP(1)4	19.89±0.05	0.42	2.40	70.16±1.03
BP(1)5	19.82±0.09	0.52	1.94	56.64±4.63
BP(1)6	19.85±0.47	0.42	2.40	68.06±1.27
BP(1)8	18.89±0.41	0.40	2.47	66.50±5.29




Fig. 1. Surface tension value of a BP(1)5 in various incubation time

Optimization of biosurfactan production by BP(1)5

The research was proven that sugar from corn cob hydrolysis can be used in biosurfactant production. Moreover, the optimal concentration of sugar that gave the best yield of biosurfactant was 200 ppm in 5 days incubation (Fig. 2). The addition of 200 ppm CCH on the culture of BP(1)5 could hold stationary phase longer than the others concentrations. In the culture with 200 ppm of CCH, in 5 days incubation, the density of microbes (OD_{650}) was 0.137. On the other hand concentration of sugar 200 ppm could decrease the surface tension from 62.996 mN/m at day 0 to 48.866 mN/m at day 5, it was the lowest surface tension compared to other cultures in some sugar concentrations which were 56.728 mN/m and 53.351 mN/m for 0 and 100 ppm sugar respectively. Microbial ability to decrease surface tension is an indicator of biosurfactant production by bacteria (El-Sheshtawy *et al.*, 2015). Based on the Picture 2C, the lowest surface tensions were showed at stationary phase, so biosurfactant was mostly produced at stationary phase. The same results were also shown in the research conducted by El-Sheshtawy *et al.* (2015) and Thavasi *et al.* (2011). Patowary *et al.* (2017) stated that biosurfactant production was started at stationary phase until death phase.

Conclusion

Based on the research, CCH can be used as substrate for growth of bacteria and biosurfactant production by hydrocarbonoclastic bacteria BP(1)5. The optimum concentration of CCH that could be used as substrate for growth and production of biosurfactant was 200 ppm in five days incubation. By using 200 ppm of the hydrolyzate, the surface tension could be decreased to 48.866 mN/m and the bacterial optical density was 0.137 (OD_{650}) after five days incubation

Acknowledgement

We gratefully thank to Faculty of Science and Technology, Universitas Airlangga for funding this research through Penelitian Unggulan Fakultas (PUF) 2019.

References

Banat, I.M., Makkar, R.S. and Cameotra, S.S. 2000. Potential Commercial Applications of Microbial Surfactants




Fig. 2. Optical density and surface tension of the BP(1)5 on CCH strain with various sugar concentration (a) 0 ppm, (b) 100 ppm, (c) 200 ppm

tants A review article. *Applied Microbiology and Biotechnology*. 53 : 495-508. <https://doi.org/10.1007/s002530051648>

Chen C., Lin, J., Wang, W., Huang, H. and Li, S. 2019. Cost-Effective Production of Surfactin from Xylose-Rich Corn Cob Hydrolysate Using *Bacillus subtilis* BS-37. *Waste and Biomass Valorization*. 10(2) : 341-347. <https://doi.org/10.1007/s12649-017-0052-5>

El-Sheshtawy, H.S., Aiad, I., Osman, M.E., Abo-ELnasr, A.A. and Kobisy, A.S. 2015. Production of Biosurfactant from *Bacillus licheniformis* for Microbial Enhanced Oil Recovery and Inhibition the Growth of Sulfate Reducing Bacteria. *Egyptian Journal of Petroleum*. 24(2) : 155-162. <https://doi.org/10.1016/j.ejpe.2015.05.005>

Gaspar, M., Kalman, G. and Reczey, K. 2007. Corn Fiber

- as A Raw Material for Hemicellulose and Ethanol Production. *Process Biochemistry*. 42(7) : 1135–1139. <https://doi.org/10.1016/j.procbio.2007.04.003>
- Global market insights Research, Biosurfactants Market worth over \$2.7 bn by 2024. [Online]. Global Market Insights, Inc.(2018). (Accessed 8 Agustus 2019). <https://www.globenewswire.com/news-release/2018/10/03/1600564/0/en/Biosurfactants-Market-will-register-5-5-CAGR-to-surpass-2-7-billion-by-2024-Global-Market-Insights-Inc.html>
- Joy, S., Rahman, P.K.S. M, Khare, S.K. and Sharma, S. 2019. Production and Characterization of Glycolipid Biosurfactant from *Achromobacter* sp.(PS1) Isolate using One-Factor-at-A-Time (OFAT) Approach with Feasible Utilization of Ammonia-Soaked Lignocellulosic Pretreated Residues. *Bioprocess and Biosystems Engineering*. 42(8) : 1301-1315. <https://doi.org/10.1007/s00449-019-02128-3>
- Karlapudi, A.P., Venkateswarulu, T.C., Srirama, K., Kota, R.K., Mikkili, I. and Kodali, V.P. 2018. Evaluation of Anti-Cancer, Anti-Microbial and Anti-Biofilm Potential of Biosurfactant Extracted from An *Acinetobacter* M6 strain. *Journal of King Saud University-Science* (In Press). <https://doi.org/10.1016/j.jksus.2018.04.007>
- Makkar, R.S. and Cameotra, S.S. 2002. Effect of Various Nutritional Supplements on Biosurfactant Production by a Strain of *Bacillus subtilis* at 45 °C. *Journal of Surfactants and Detergents*. 5(1) : 11-17. <https://doi.org/10.1007/s11743-002-0199-8>
- Marhaeni, 2002. Bioconversion of Waste of Corn Cobs to Diet as Protein Sources. *AGRIS*. 2 (2): 182-192. <http://agris.fao.org/agrissearch/search.do?recordID=ID2005000185>
- Moldes, A.B., Torrado, A.M., Barral, M.T. and Domínguez, J.M. 2007. Evaluation of Biosurfactant Production from Various Agricultural Residues by *Lactobacillus pentosus*. *Journal of Agricultural and Food Chemistry*. 55(11) : 4481-4486. <https://doi.org/10.1021/jf063075g>
- Mulligan, C.N. and Wang, S. 2006. Remediation of a heavy metal-contaminated soil by a rhamnolipid foam. *Engineering Geology*. 85(1-2) : 75-81. <https://doi.org/10.1016/j.enggeo.2005.09.029>
- Ni'matuzahroh, Sari, S.K., Ningrum, I.P., Pusfita, A.D., Marjayandari, L., Trikurniadewi, N., Ibrahim, S.N.M.M, Fatimah, Surtiningsih T, Nurhayati, T. and Yuliani, H. 2019a. The potential of Indigenous bacteria from oil sludge for biosurfactant production using hydrolysate of agricultural waste. *Biodiversitas*. 20(5) : 1374-1379. <https://doi.org/10.13057/biodiv/d200529>
- Ni'matuzahroh, Sari, S.K., Trikurniadewi, N., Pusfita, A.D., Ningrum, I.P., Ibrahim, S.N.M.M., Nurhariyati, T., Fatimah and Surtiningsih, T. 2019b. Utilization of Rice Straw Hydrolysis Product of *Penicillium* sp. H9 as A Substrate of Biosurfactant Production by LII61 Hydrocarbonoclastic Bacteria. *IOP Conf. Series: Earth and Environmental Science*. 217(1). <https://doi.org/10.1088/1755-1315/217/1/012028>
- Ni'matuzahroh, Fatimah, Surtiningsih, T. 2018. Pemanfaatan Limbah Organik Sebagai Media Produksi Biosurfaktanoleh Bakteri Hidrokarbonoklastik. *Research Report*. Fakultas Sains dan Teknologi, Universitas Airlangga: Surabaya.
- Oktaviani, M., Hermiati, E., Thontowi, A., Laksana, R.P.B., Kholida, L.N., Andriani, A., Yopi, Mungunwardoyo, W. 2019. Production of xylose, glucose, and other products from tropical lignocellulose biomass by using maleic acid pretreatment. *IOP Conf. Series: Earth and Environmental Science*. 251. <https://doi.org/10.1088/1755-1315/251/1/012013>
- Patowary, K., Patowary, R., Kalita, M.C. and Deka, S. 2017. Characterization of Biosurfactant Produced during Degradation of Hydrocarbons Using Crude Oil as Sole Source of Carbon. *Frontiers in Microbiology*. 8 : 279. <https://doi.org/10.3389/fmicb.2017.00279>
- Pruthi, V. and Cameotra, S.S. 1997. Rapid Identification of Biosurfactant-Producing Bacterial Strains Using a Cell Surface Hydrophobicity Technique. *Biotechnology Techniques*. 11(9): 671-674. <https://doi.org/10.1023/A:1018411427192>
- Rangarajan, V. and Sen, R. 2013. An Inexpensive Strategy for Facilitated Recovery of Metals and Fermentation Products by Foam Fractionation Process. *Colloids and Surfaces B: Biointerfaces*. 104 : 99–106. <https://doi.org/10.1016/j.colsurfb.2012.12.007>
- Saha, B.S. 2004. Lignocellulose Biodegradation and Applications in Biotechnology. In Saha, B. C. and Hayashi, K. (Editors). 2004. *Lignocellulose Biodegradation*. American Chemical Society. 889 : 2-34 : Washington DC, USA. <https://doi.org/10.1021/bk-2004-0889.ch001>
- Taherzadeh, M.J. and Niklasson, C. 2004. Ethanol from Lignocellulosic Materials: Pretreatment, Acid and Enzymatic Hydrolyses, and Fermentation. In Saha, B.C. and Hayashi, K (Editors). 2004. *Lignocellulose Biodegradation*. American Chemical Society. 889 : 49-68. Washington DC, USA. <https://doi.org/10.1021/bk-2004-0889.ch003>.
- Tarkow, H. and Feist, W.C. 1969. A Mechanism for Improving the Digestibility of Lignocellulosic Materials with Dilute Alkali and Liquid Ammonia. In Hajny G. J and Reese E. T (Editors). 1969. *Cellulases and Their Applications*. *Advances in Chemistry Series*; American Chemical Society. 95: 197–218. Washington DC, USA. <https://doi.org/10.1021/ba-1969-0095.ch012>
- Thavasi, R., Nambaru, V.R.M.S., Jayalakshmi, S., Balasubramanian, T. and Banat, I.M. 2011. Biosurfactant Production by *Pseudomonas aeruginosa* from Renewable Resources. *Indian Journal of Microbiology*. 51(1) : 30-36. <https://doi.org/10.1007/>

- s12088-011-0076-7
- Varjani, S.J. and Upasani, V.N. 2017. Critical Review on Biosurfactant Analysis, Purification and Characterization using Rhamnolipid as A Model Biosurfactant. *Bioresource Technology*. 232 : 389–397. <https://doi.org/10.1016/j.biortech.2017.02.047>
- Zambry, N.S., Ayoib, A., Noh, N.A.M. and Yahya, A.R.M. 2017. Production and Partial Characterization of Biosurfactant Produced by *Streptomyces* sp.R1. *Bioprocess and Biosystems Engineering*. 40(7): 1007–1016. <https://doi.org/10.1007/s00449-017-1764-4>
- Zhang, Q.Z. and Cai, W.M. 2008. Enzymatic Hydrolysis of Alkali-Pretreated Rice Straw by *Trichoderma reesei* ZM4-F3. *Biomass and Bioenergy*. 32(12) : 1130–1135. <https://doi.org/10.1016/j.biombioe.2008.02.006>
-