

Antifungal and antibacterial activity of black betel (*Piper betle* L. var *Nigra*) extract

by Ni'matuzahroh Ni'matuzahroh

Submission date: 27-Apr-2021 10:31AM (UTC+0800)

Submission ID: 1570943926

File name: 4._Antifungal_antibacterila.pdf (463.27K)

Word count: 3485

Character count: 19119

9

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/322492230>

Antifungal and antibacterial activity of black betel (Piper betle L. var Nigra) extract

Article in Bioscience Research · October 2017

CITATIONS

2

READS

236

4 authors, including:



Ni'matuzahroh Ni'matuzahroh
Airlangga University

52 PUBLICATIONS 143 CITATIONS

SEE PROFILE



Nabilah Istighfari Zuraidassanaaz
Airlangga University

3 PUBLICATIONS 3 CITATIONS

SEE PROFILE



Lilis Sulistyorini
Airlangga University

14 PUBLICATIONS 12 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Screening of Cellulolytic Bacteria From Alas Purwo National Park to Produce Cellulase [View project](#)



Utilization of Organic Waste for Biosurfactant Production Substrate by Hydrocarbonoclastic Bacteria [View project](#)

All content following this page was uploaded by Nabilah Istighfari Zuraidassanaaz on 08 February 2019.

The user has requested enhancement of the downloaded file.



Available online freely at www.isisn.org

Bioscience Research

Print ISSN: 1811-9506 Online ISSN: 2218-3973

Journal by Innovative Scientific Information & Services Network



RESEARCH ARTICLE

BIOSCIENCE RESEARCH, 2017 14(4): 750-755.

OPEN ACCESS

Antifungal and antibacterial activity of black betel (*Piper betle* L. var Nigra) extract

Junairiah^{1*}, Ni' matuzahroh¹, Nabilah Istighfari Zuraidassanaaz¹ and lilis Sulistyorini²

¹Department of Biology, Faculty of Science and Technology, University of Airlangga, St. Mulyorejo, Sukolilo, Mulyorejo, Surabaya 60115, East Java, **Indonesia**,

²Faculty of Public Health, University of Airlangga, St. Mulyorejo, Sukolilo, Mulyorejo, Surabaya 60115, East Java, **Indonesia**.

*Correspondence: alip.jun1@gmail.com Accepted: 10 Sep. 2017 Published online: 07 Nov. 2017

Black betel (*Piper betle* L. var Nigra) belong to Piperaceae family and one of endemic species of Indonesia. This study was aimed to determine antifungal and antibacterial activity of n-hexane, ethyl acetate, and methanol extracts of *Piper betle* L. var Nigra toward pathogenic microbes *Candida albicans* ATCC 10231, *Staphylococcus aureus* ATCC 25923, and *Escherichia coli* ATCC 25922. Leaf powder of black betel was macerated using n-hexane, ethyl acetate, and methanol solvents. Antifungal and antibacterial activity was determined using diffusion and dilution tests. Data collected from both tests including diameter of inhibition zone (mm) and Minimal Inhibitory Concentration (MIC), Minimal Bactericidal Concentration (MBC), Minimal Fungicidal Concentration. Data was then analyzed statistically. Result showed that methanol extract of black betel had the highest inhibition activity compared to n-hexane and ethyl acetate extract.

Keywords: antifungal, antibacterial, *Piper betle* L. var Nigra

INTRODUCTION

Piperaceae belong to Magnoliopsida class, sub ordo Nymphaeiflorae and Piperales ordo. This plant family have been widely known as source of bioactive compound (Santhakumari et al. 2003; Lei et al. 2003; Majumdar et al. 2002; Kato and Furlan et al. 2007; Regasini et al. 2009). Piperaceae consisted of 10 genera and 2000 species. *Piper* genus is known to contain secondary metabolites of polyphenol, alkaloid, steroid, saponin, and tannin (Navickiene et al. 2006; Regasini et al. 2008).

Water, methanol, ethyl acetate, and ether extracts of *Piper betle* L. were found to be able to inhibit growth of *Streptococcus pyrogenes*, *Staphylococcus aureus*, *Proteus vulgaris*, and *Escherichia coli* (Chakraborty and Shah, 2011), while ethanol, hexane, ethyl acetate,

hydromethanol extracts of leaf and stem from both *Piper arboretum* and *Piper tuberculatum* could inhibit growth of *Candida albicans*, *Candida krusei*, *Candida parasilopsis*, and *Cryptococcus neoformans* (Regasini et al. 2009). In addition, ethanol and chloroform extracts of *Piper nigrum* were also able to inhibit *Escherichia coli*, *Salmonella typhi*, *Pseudomonas sp.*, and *Staphylococcus aureus* (Ganesh et al. 2014). Ethanol, hexane, chloroform, ethyl acetate, methanol, and water extracts of *Piper hayneanum* could suppress growth of *Staphylococcus aureus* and *Candida albicans* (Bastos et al. 2011), while ethanol extract of black pepper (*Piper nigrum* L.) was able to suppress *Staphylococcus aureus*, *Bacillus subtilis*, *Pseudomonas aeruginosa*, *Escherichia coli*, *Alternaria alternata*, *Aspergillus niger*, *Aspergillus flavus*, and *Fusarium*

oxysporum (Rani et al. 2013).

Betel variety currently recorded including green, red, and black betel. Black betel (*Piper betle* L. var *Nigra*) is one of Piperaceae family contained various secondary metabolites, such as alkaloid, flavonoid, saponin, terpenoid, and steroid. Thus, this plant has potential as antibacterial, antifungal, anti-diabetic, antiulcer, antiplatelet, antifertility, antitumor, anti-mutagenic, and anti-helminthic agents (Lei et al. 2003; Majumdar et al. 2002). Based on previous study, few study has been conducted to determine the activities of n-hexane, ethyl acetate, and methanol extracts of *Piper betle* L. var *Nigra* towards pathogenic microbes *Candida albicans* ATCC 10231, *Staphylococcus aureus* ATCC 25923, and *Escherichia coli* ATCC 25922.

MATERIALS AND METHODS

Plant Collection

Black betel (*Piper betle* L. var *Nigra*) was obtained from Kayon Flower Mart, Surabaya, and East Java, Indonesia. This plant was identified in Plant Physiology Laboratory, Faculty of Science and Technology, University of Airlangga.

Pathogenic Microbes

Pathogenic microbes used were *Candida albicans* ATCC 10231, *Staphylococcus aureus* ATCC 25923, and *Escherichia coli* ATCC 25922. The three microbes were collection of Microbiology Laboratory, Department of Biology, Faculty of Science and Technology, University of Airlangga.

Extraction Procedure

Leaves of *Piper betle* L. var *Nigra* were firstly washed and air-dried, then crushed into powder. Betel powder weighed to 29.5 grams then extracted using n-hexane, ethyl acetate, and methanol solvents respectively. Extraction was performed using maceration method for three days and repeated for three times. Crude extract

filtered using filter paper, then extract was made into a series of concentration; 0 ppm; 250 ppm; 500 ppm; 750 ppm; and 1000 ppm.

Microbe Culture

Examined microbe was made into stock solution. Microbe suspension was produced by mixing several ose of bacteria from slanted agar into 20 mL saline water. Suspension was homogenized, then taken several mL for measuring its optical density using spectrophotometer. Stock used had OD 0.1 at 600 nm wavelength for fungi and 625 nm for bacteria.

Diffusion Test

Medium used for diffusion test was Mueller Hinton agar (MHA). 1 mL microbe suspension was placed in petri dish, then 15 mL MHA medium was added, mixture was homogenized and left to be solidified. On the surface of medium, 3 paper discs (6 mm diameter) saturated with 20 μ l extract of respective solvent at each concentration of 0, 250, 500, 750, and 1000 ppm. Diameter of inhibition zone was measured using caliper.

Dilution test

1 mL microbe suspension in Mueller Hinton Broth (MHB) medium was placed into reaction tube filled with 1 mL betel extract of each solvent type and concentration. Each culture was incubated for 24 hours. Next, 0.1 mL of each culture was placed into petri dish and added 15 mL medium, then incubated for 24 hours for bacteria and 48 hours for fungi. If microbe growth was observed in medium, then concentration used was MIC value. If microbe growth was not observed, then concentration used was MBC/MFC value.

RESULTS

Diffusion test was a method commonly used for determining microbe activity, due to its inexpensive cost. Result of diffusion test was presented in Table 1.

Table 1. Antifungal and antibacterial activities of three extracts from *Piper betle* L. var *Nigra*

Microbe strain	Diameter of inhibition zone (mm)														
	n-hexane (μ g/disc)					ethyl acetate (μ g/disc)					methanol (μ g/disc)				
	2	5	10	15	20	2	5	10	15	20	2	5	10	15	20
<i>E. coli</i> ATCC 25922	8,88	8,92	7,20	7,53	6,00	9,57	9,56	6,00	6,00	6,00	7,02	10,20	9,80	7,47	8,22
<i>S. aureus</i> ATCC 25923	6,00	6,00	6,00	7,00	6,00	6,00	6,00	10,63	7,53	10,03	6,00	6,00	12,69	12,00	8,52
<i>C. albicans</i> ATCC 10231	6,00	6,00	6,00	6,00	6,00	6,00	7,42	6,00	6,00	6,00	6,82	7,87	7,68	6,00	6,00

39

Table 2. MIC and MBC of each extract from *Piper betle* var *Nigra* towards *E. coli*, *S. aureus* and *C. albicans*.

Microbe strain	Extract solvent					
	n-hexane		ethyl acetate		methanol	
	MIC (mg/mL)	MBC (mg/mL)	MIC (mg/mL)	MBC (mg/mL)	MIC (mg/mL)	MBC (mg/mL)
<i>E. coli</i> ATCC 25922	0,05	0,1	0,015	0,02	0,0275	0,03
<i>S. aureus</i> ATCC 25923	0,05	0,1	0,02	0,04	0,035	0,02
<i>C. albicans</i> ATCC 10231	-	-	0,02	0,015	0,0275	0,03

It was showed that n-hexane, ethyl acetate, and methanol extracts was able to inhibit the three species of pathogenic microbes. However, each extract was found to have different sensitivity towards specific microbe. N-hexane extract of black betel was found to inhibit *E. coli* ATCC 25922 better at 5 µg/disc, with inhibition zone diameter of 8.92 mm. Extract of ethyl acetate inhibited *S. aureus* ATCC 25923 at 10 µg/disc, with zone diameter of 10.63 mm, while methanol extract was able to inhibit *S. aureus* ATCC 25923 at 10 µg/disc concentration, with zone diameter of 12.69 mm.

For determining MIC and MBC, dilution test was performed for each solvent and concentration used. MIC and MBC was presented in Table 2.

DISCUSSION

Based on result of current study, methanol extract was found as the best extract solvent to be used compared to other solvent n-hexane and ethyl acetate. This was due to methanol as universal solvent, thus was able to bound various compound or secondary metabolites. Methanol extract was found to contain terpenoid, steroid, flavonoid, polyphenol, tannin, and alkaloid. Steroid was found to have function as antibacterial agent (Silvia et al. 2003). Steroid was able to act as antibacterial by preventing spore germination process (Subisha and Subramoniam, 2005). Other than them, alkaloid also had cytotoxic property against bacteria (Ezekiel et al. 2009). Indoquinolone alkaloid specifically previously found to induce cell lysis and morphological alteration on *S. aureus* (Sawer et al. 2005). Flavonoid inhibited function of cytoplasm membrane and energy metabolism (Cushnie and Lamb, 2009). Flavonoid possessed ability as antimicrobe (Ogundipe et al. 2001; Silvia et al. 2003) and was able to suppress growth of *S. aureus* and

E. coli (Chattopadhyay et al. 2001). Flavonoid was currently applied as new compound in antimicrobial therapy (Ozcelik et al. 2008). Polyphenol had antibacterial activity (Taguri et al. 2004), while tannin had previously hypothesized as antibacterial agent (Akiyama, 2001; Futanogawa, 2004; Guittat et al. 2003; Lisgarten et al. 2002).

Previous study had shown that water, methanol, petroleum ether, diethyl acetate, chloroform, and methanol extract of *Tamarindus indica* leaves at 100 mg/mL concentration were found to be able to inhibit growth of various Gram positive (*Bacillus subtilis* ATCC 11774, MRSA ATCC 977, *Staphylococcus aureus* ATCC 29213, and *Micrococcus luteus* ATCC 4698) and Gram negative bacteria species (*Escherichia coli* ATCC 8739, *Klebsiella pneumoniae* ATCC 700603, and *Pseudomonas aeruginosa* ATCC 27853). Lowest inhibition was found from petroleum ether extract towards *Bacillus subtilis* with 11.67 mm zone diameter, while highest inhibition was recorded from methanol extract against *Micrococcus luteus*, with zone diameter of 43.33 mm. In addition, ethanol, petroleum ether, diethyl acetate, ethyl acetate, and chloroform extracts were also found to be able to inhibit fungi species *Aspergillus flavus* ATCC 2000226, *Aspergillus fumigatus* ATCC 204305, *Aspergillus niger* ATCC 1015, and *Candida albicans* ATCC 10231. Highest inhibition was reported from ethanol extract of *Tamarindus indica* towards *Aspergillus fumigatus*, while lowest inhibition was reported from diethyl acetate extract towards *Aspergillus niger* (Gumgumjee et al. 2012).

Piperin isolated from *Piper nigrum* at 25 µl volume was found to be able to inhibit *S. aureus* with zone diameter of 4 mm, however was not able to inhibit *Escherichia coli* (Rani et al. 2013). Water, methanol, ethyl acetate, and ether extracts of *Piper betle* leaves at 5 mg/ml

25 concentration was able to inhibit growth of *S. aureus* and *E. coli* (Chakraborty and Shah, 2011). N-hexane, ethyl acetate, and methanol extract was found unable to suppress *S. aureus*, but able to inhibit *Candida albicans* with inhibition zone diameter of 1.0 up to concentration of 350 µg/disc. MIC of n-hexane, ethyl acetate, and methanol extract were 350 µg/disc, 350 µg/disc, and 1.0 µg/disc respectively (Bastos et al. 2011). Black pepper extract at 1 mg/mL up to 4 mg/mL concentration could suppress *E. coli*, *S. typhi*, *Pseudomonas sp.*, and *Proteus sp.* (Ganesh et al. 2014).

MIC of hexane extract from fruits and leaves of *Piper arboretum* was 250 µg/mL, while MIC of hexane and ethyl acetate extracts of *Piper tuberculatum* was >1000 µg/mL (Regasini et al. 2009). Ethanol extract of *Tamarindus indica* fruits was able to suppress *S. aureus* and *E. coli* have inhibition zone of 18.0 mm and 13.0 mm. MIC of ethanol extract against both bacteria were 500 mg/mL (Gupta et al. 2014). Methanol extract of *Origanum vulgare* could suppress *E. coli* and *S. aureus* at 250 µg/ml concentration (Ashraf et al. 2011). Ethanol extract of leaf, stem, and fruit of *Tamarindus indica* were able to inhibit *E. coli* and *S. aureus* (Nwodo et al. 2011). Water, acetone, and ethanol extract of stem and leaf of *Tamarindus indica* could inhibit *E. coli* and *S. aureus*, but unable to suppress *C. albicans*. MIC of ethanol extract against *E. coli* and *S. aureus* were 15.5 mg/mL and 8 mg/mL respectively. MBC of both extract against *E. coli* and *S. aureus* were 18 mg/mL and 20 mg/mL (Doughari, 2006).

MIC and MBC of ethanol leaf extract from *Tamarindus indica* against *E. coli* were 15.5 mg/mL and 18 mg/mL respectively, while against *S. aureus* was 8 mg/mL and 20 mg/mL (Doughari, 2006). MIC and MBC of hexane and ethyl acetate leaf extracts of *Piper arboretum* against *C. albicans* were respectively 250 µg/mL and >1000 µg/mL. MIC and MBC of leaf extract from *Piper tuberculatum* against *C. albicans* was >1000 µg/mL (Regasini et al. 2009). MIC and MBC ethanol and water extract of *Tamarindus indica* towards *E. coli* ATCC 11775 were 31.25 and 125 mg/mL respectively (Nwodo et al. 2011). Based on data in Table 2, MIC and MBC of ethyl acetate and methanol extracts of black betel was higher compared to extracts from *Tamarindus indica*, *Piper arboretum*, and *Piper tuberculatum*. Based on result of current study, three types of extract using n-hexane, ethyl acetate, and methanol extracts had antibacterial and antifungal

activities. From those varieties of extracts examined, methanol extract had highest antibacterial and antifungal activities.

CONCLUSION

Methanol extract of black betel had the highest inhibition activity compared to n-hexane and ethyl acetate extract.

CONFLICT OF INTEREST

The authors declared that present study was performed in absence of any conflict of interest.

ACKNOWLEDGEMENT

Author would like to thank Ministry of Research, Technology, and Higher Education of Republic Indonesia, who funded this research via Distinguished Research Grant of Higher Education 2017, research contract No. 597/UN3.14/LT/2017

AUTHOR CONTRIBUTIONS

24 Junairiah and Ni'matuzahroh designed and performed the experiments and also wrote the manuscript. Nabilah Istighfari Zuraidassanaaz performed plant collection, diffusion and dilution test, data analysis. Lilis Sulistyorini designed experiments and reviewed the manuscript. All authors read and approved the final version. All authors read and approved the final version.

Copyrights: © 2017 @ author (s).

This is an open access article distributed under the terms of the [Creative Commons Attribution License \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

REFERENCES

- Akiyama H, Kazuyasu F, Yamasaki O, Oono T, Iwatsuki K, 2001. Antibacterial Action of Several Tannins against *Staphylococcus aureus*. *Journal Antimicrobial Chemotherapy* 48 (48): 487-491.
- Ashraf Z, Muhammad A, Imran M, Tareq AH, 2011. In vitro Antibacterial and Antifungal Activity of Methanol, Chloroform and Aqueous Extracts of *Origanum vulgare* and Their Comparative Analysis. *International Journal of*

- Organic Chemistry 1 (04): 257.
- Bastos M, Houly R, Conserva L, Andrade V, Rocha E, Lemos R, 2011. Antimicrobial and Wound Healing Activities of Piper hayneanum. Journal of Chemical and Pharmaceutical Research 3: 213-222.
- Chakraborty D, Shah B, 2011. Antimicrobial, Antioxidative and Antihemolytic Activity of Piper betel Leaf Extracts. Int. J. Pharma. Pharmaceut. Sci 3 : 192-199.
- Chattopadhyay D, Maiti K, Kundu AO, Chakraborty MS, Bhadra R, Maudal SC, Maudal AB, 2001. Antimicrobial Activity of alstoniamacrophylla: A folklore of Bay Islands. Journal of Ethnopharmacol. Lausanne. 77: 49-55.
- Cushnie TPT, Lamb AJ, 2011. Recent Advances in Understanding The Antibacterial Properties of Flavonoids. International Journal of Antimicrobial Agents 38: 99-107.
- Doughari JH, 2006. Antimicrobial Activity of Tamarindus indica Linn. Tropical Journal of Pharmaceutical Research, 5 (2): 597-603.
- Ezekiel OO, Onyeoziri NF, 2009. Preliminary Studies on The Antimicrobial Properties of Buchholzia coriacea (Wonderful Kola). Afr. J. Biotechnol. 8 (3): 472-474.
- Funatogawa K, Hayashi S, Shimomura H, Yoshida T, Hatano T, Ito H, Iria Y, 2004. Antibacterial Activity of Hydrolysable Tannins Derived from Medicinal Plants Against Helicobacter pylori. Microbiol Immunol. 48 (4): 251-261.
- Ganesh P, Kumar P, Suresh R, Saranraj, P, 2014. Phytochemical Analysis and Antibacterial Activity of Pepper (Piper nigrum L.) Against Some Human Pathogens, 3 (2):36-41.
- Guitat L, Alberti P, Rosu F, Van MS, Thetiot E, Pieters L, Gabelica V, De PE, Ottaviani A, Roiu JF, Mergny JL, 2003. Interaction of cryptolepinedanneocryptolepine with Unusual DNA Structures. Bioch. 85: 535-541.
- Gumgumjee NM, Khedr A, Haja AS, 2012. Antimicrobial Activities and Chemical Properties of Tamarindus indica L. Leaves Extract. African Journal of Microbiology Research 6(32): 6172-6181.
- Gupta C, Prakash D, Gupta S, 2014. Studies on The Antimicrobial Activity of Tamarind (Tamarindus indica) and Its Potential as Food Bio-preservative, International Food Research Journal 21(6): 2437-2441.
- Kato MJ, Furlan M, 2007. Chemistry and Evolution of Piperaceae. Pure Appl. Chem. 79: 529-538.
- Lei D, Chan CP, Wang YJ, Wang TM, Lin BR, Huang CH, Lee JJ, Chen, HM, Jeng JH, Chang MC, 2003. Antioxidative and Anti-platelet Effects of Aqueous Inflorescence Piper betel Extract. J Agric Food Chem, 51: 2083-2088.
- Lisgarten JN, Coll M, Portugal J, Wright CW, Aymami J, 2002. The Antimalarial and Cytotoxic Drug Cryptolepine Intercalates into DNA at Cytosine-cytosine Sites. Nature Structural Biol. 9: 57-60.
- Majumdar B, Chaudhuri SR, Roy A, 2002. Potent Antiulcerogenic Activity of Ethanol Extract of Leaf of Piper betel Linn by Antioxidative Mechanism. Ind Jour Clin Bio Chem. 17:49-57.
- Navickiene HMD, Morandim AA, Alécio AC, Regasini LO, Bergamo DC, Telascra M, Cavalheiro AJ, Lopes MN, Bolzani VS, Marques MO, Young MCM, Kato MJ, 2006. Composition and antifungal activity of essential oil from Piper aduncum, Piper arboretum and Piper tuberculatum. Quim. Nova, 29: 467-470.
- Nwodo, UU, Obiyeye GE, Chigor VN, Okoh AI, 2011. Assessment of Tamarindus indica Extracts for Antibacterial Activity. International journal of molecular sciences, 12(10): 6385-6396.
- Ogundipe OO, Moody JO, Houghton PJ, Odelola HA. Bioactive Chemical Constituents from Alchornealaxiflora (benth) pax and Hoffman. Journal of Pharmacol., Lausanne 74: 275-280.
- Özçelik B, Deliorman OD, Özgen S, Ergun F, 2008. Antimicrobial Activity of Flavonoids Against Extended-spectrum β -lactamase (ES β L)-producing *Klebsiella pneumoniae*. Trop J Pharm Res, 7: 1151-1157.
- Rani PU, Hymavathi A, Babu KS, Rao AS, 2013. Bioactivity Evaluation of Prenylated Isoflavones Derived from Derris scandens Benth Against Two Stored Pest Larvae. J. Biopesticides, 6: 14-21.
- Rani SM, Dayanand CD, Shetty J, Vegi PK, Kutty AM, 2013. Evaluation of Antibacterial Activity of Pongamia pinnata linn on Pathogens of Clinical Isolates. American Journal of Phytomedicine and Clinical Therapeutics 1(8): 645-651.
- Regasini LO, Cotinguiba F, Passerini GD, Bolzani VS, Cicarelli RMB, Kato, MJ, Furlan M, 2009. Trypanocidal Activity of Piper arboretum and Piper tuberculatum (Piperaceae). Rev. Bras. Farmacogn. 19:199-203.
- Regasini LO, Cotinguiba F, Siqueira JR, Bolzani

- VS, Silva DHS, Furlan M, Kato MJ, 2008. Radical Scavenging Activity of Piper arboretum and Piper tuberculatum (Piperaceae). *Lat. Am. J. Pharm.* 27: 900-903.
- Santhakumari P, Prakasam A, Puglendi KV, 2003. Modulation of Oxidative Stress Parameters by Treatment with Piper betel Leaf in Streptozotacin Induced Diabetic Rats. *Ind J of Pharmacol.* 35: 373–378.
- Sawer IK, Berry MI, Ford JL, 2005. The Killing Effect on Staphylococcus aureus. *Lett. Appl. Microbiol.* 40: 24-29.
- Silvia HTC, Marcos JS, Evandro W, Izabel YI, 2003. Antimicrobial Activity of Flavonoids and Steroids Isolated from Two Chromolaena Species. *Brazilian Journal of Pharmaceutical Sciences.* 39 (4).
- Subhisha A, Subramoniam A, 2005. Antifungal Activities of A Steroid from Pallavicinia lyellii, *Indian Journal of Pharmacology*, 37(5): 304-308.
- Taguri T, Tanaka T, Kouno I, 2004. Antimicrobial Activity of 10 Different Plant Polyphenols Against Bacteria Causing Food-borne disease. *Biol Pharm Bull* 27(12): 1965-1969.

Antifungal and antibacterial activity of black betel (Piper betle L. var Nigra) extract

ORIGINALITY REPORT

19%

SIMILARITY INDEX

13%

INTERNET SOURCES

16%

PUBLICATIONS

8%

STUDENT PAPERS

PRIMARY SOURCES

1	www.innspub.net Internet Source	1%
2	lppm.ub.ac.id Internet Source	1%
3	Banik Avijit, Tamanna Zerin, Sultana Rajia. "Comparative Phytochemical and Antibacterial Properties of Piper betle Leave Extracts from Barguna and Moheshkhali, Bangladesh", Iranian Journal of Medical Microbiology, 2020 Publication	1%
4	cs.wellesley.edu Internet Source	1%
5	Submitted to iGroup Student Paper	1%
6	www.socialresearchfoundation.com Internet Source	1%
7	Edible Medicinal and Non Medicinal Plants, 2014.	1%

8 [medcraveonline.com](https://www.medcraveonline.com) 1 %
Internet Source

9 Ramy Mohamed Ghazy, Abdallah Elmaghraby, Ramy Shaaban, Ahmed Kamal et al. 1 %
"Effectiveness and Safety of Chloroquine or Hydroxychloroquine as a mono-therapy or in combination with Azithromycin in the treatment of COVID-19 patients: Systematic Review and Meta-Analysis", Cold Spring Harbor Laboratory, 2020
Publication

10 heritagesciencejournal.springeropen.com 1 %
Internet Source

11 Idha Kusumawati, Kresma Oky Kurniawan, Subhan Rullyansyah, Tri Anggono Prijo et al. 1 %
"Anti-aging properties of Curcuma heyneana Valetton & Zipj: A scientific approach to its use in Javanese tradition", Journal of Ethnopharmacology, 2018
Publication

12 E.E. Mgbeahuruike, T. Yrjönen, H. Vuorela, Y. Holm. 1 %
"Bioactive compounds from medicinal plants: Focus on Piper species", South African Journal of Botany, 2017
Publication

13 [worldwidescience.org](https://www.worldwidescience.org)
Internet Source

<1 %

14

www.jpsscscientificpublications.com

Internet Source

<1 %

15

www.mdpi.com

Internet Source

<1 %

16

www.omicsonline.org

Internet Source

<1 %

17

Elumalai K. "Antioxidant Activity and Phytochemical Screening of Different Solvent Extracts *Cluasena excavata burm F.* (Rutaceae)", Antioxidant Activity and Phytochemical Screening of Different Solvent Extracts *Cluasena excavata burm F.* (Rutaceae), 2016

Publication

<1 %

18

eprints.usm.my

Internet Source

<1 %

19

link.springer.com

Internet Source

<1 %

20

mdpi.com

Internet Source

<1 %

21

M.J. Salvador, O.L.A.D. Zucchi, R.C. Candido, I.Y. Ito, D.A. Dias. " Antimicrobial Activity of Crude Extracts and Isolated Constituents of ", *Pharmaceutical Biology*, 2008

<1 %

22

O. C Enechi,, O. F. C. Nwodo,. "Anti-ulcer and gastric anti-secretory activities of seed extract of Buchholzia coriacea in Wistar Albino rats", African Journal of Biotechnology, 2014

Publication

<1 %

23

Rosmiati Rosmiati, Najiah Musa, Habsah Mohammad, Tengku Sifzizul Tengku Muhammad, Andi Parenrengi, Wahyuni Wahyuni. "THE EFFECT OF MARINE SPONGE Aaptos aaptos EXTRACT IN VIBRIOSIS TREATMENT OF BLACK TIGER SHRIMP Penaeus monodon LARVAE", Indonesian Aquaculture Journal, 2014

Publication

<1 %

24

bmccancer.biomedcentral.com

Internet Source

<1 %

25

indonesianjpharm.farmasi.ugm.ac.id

Internet Source

<1 %

26

Aldilene da Silva Lima, José Gracione do Nascimento Sousa Filho, Sinval Garcia Pereira, Giselle Maria Skelding Pinheiro Guillon et al. "Acaricide activity of different extracts from Piper tuberculatum fruits against Rhipicephalus microplus", Parasitology Research, 2013

Publication

<1 %

27

Aline C. Pereira, Hudson W. P. Carvalho, Geraldo H. Silva, Denilson F. Oliveira et al. "Purification of an antibacterial compound from Lantana lilacina", Revista Brasileira de Farmacognosia, 2008

Publication

<1 %

28

Arina Zulfah Primananda, Laksmi Anggun, Rika Murhayati. "Test of Antibacterial Activity 70% Ethanol Extract of Seeds Black Pepper (Piper Nigrum L) Against Bacteria Escherichia Coli Atcc 25922", Journal of Physics: Conference Series, 2021

Publication

<1 %

29

Md Rahman, Md Wadud, Tarequl Islam, Md Hussain, Esfat Bristy, Ashraful Tuhin. "Evaluation of Antibacterial Activity of Piper betel Leaves and Nigella sativa Seeds against Multidrug Resistant Food and Water Borne Pathogenic Bacteria: An in vitro Study Model", Microbiology Research Journal International, 2018

Publication

<1 %

30

Mohammad Omar Faruque, Umme Ruman Ankhi, M. Kamaruzzaman, James W. Barlow, Bo Zhou, Ji Hao, Xinzhou Yang, Xuebo Hu. "Chemical composition and antimicrobial activity of *Congea tomentosa*, an

<1 %

ethnomedicinal plant from Bangladesh",
Industrial Crops and Products, 2019

Publication

31

Muhammad Mutassim Billah, Rafikul Islam, Hajera Khatun, Shahnaj Parvin, Ekramul Islam, SM Anisul Islam, Akbar Ali Mia. "Antibacterial, antidiarrhoeal, and cytotoxic activities of methanol extract and its fractions of *Caesalpinia bonducella* (L.) Roxb leaves", BMC Complementary and Alternative Medicine, 2013

Publication

<1 %

32

kb.psu.ac.th

Internet Source

<1 %

33

moam.info

Internet Source

<1 %

34

mural.maynoothuniversity.ie

Internet Source

<1 %

35

www.academicjournals.org

Internet Source

<1 %

36

www.iums2008.org

Internet Source

<1 %

37

www.scielo.br

Internet Source

<1 %

38

"Microbial Control and Food Preservation", Springer Science and Business Media LLC,

<1 %

2017

Publication

39

Ban, S.H.. "Effects of a bio-assay guided fraction from *Polygonum cuspidatum* root on the viability, acid production and glucosyltransferase of mutans streptococci", *Fitoterapia*, 201001

Publication

<1 %

40

Luis Octávio Regasini, Fernando Cotinguiba, Gabriela Duó Passerini, Vanderlan da Silva Bolzani et al. "Trypanocidal activity of *Piper arboreum* and *Piper tuberculatum* (Piperaceae)", *Revista Brasileira de Farmacognosia*, 2009

Publication

<1 %

41

Md. Anowar Khasru Parvez, Karabi Saha, Juairia Rahman, Rahath Ara Munmun et al. "Antibacterial activities of green tea crude extracts and synergistic effects of epigallocatechingallate (EGCG) with gentamicin against MDR pathogens", *Heliyon*, 2019

Publication

<1 %

42

Muhammad Irfan, Faizah Shahudin, VINCENT HOOPER, Waqar Akram, Rosmaiza Ghani. "The psychological impact of coronavirus on university students and its socio-economic

<1 %

determinants in Malaysia", Cold Spring Harbor Laboratory, 2020

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Antifungal and antibacterial activity of black betel (Piper betle L. var Nigra) extract

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
