

# Antimicrobial Activities and In silico Analysis of Methoxy Amino Chalcone Derivatives

*by Ni'matuzahroh Ni'matuzahroh*

---

**Submission date:** 27-Apr-2021 10:57AM (UTC+0800)

**Submission ID:** 1570971782

**File name:** 3.\_Antimicrobial\_Activities\_and\_In\_silico\_Analysis.pdf (752.69K)

**Word count:** 3963

**Character count:** 21911



Molecular and Cellular Life Sciences: Infectious Diseases, Biochemistry and Structural Biology  
2015 Conference, MCLS 2015

## Antimicrobial Activities and In silico Analysis of Methoxy Amino Chalcone Derivatives

Hery Suwito<sup>a,\*</sup>, Ni'matuzahroh<sup>b</sup>, Alfinda Novi Kristanti<sup>a</sup>, Salwa Hayati<sup>b</sup>,  
Selva Rosyta Dewi<sup>b</sup>, Ilma Amalina<sup>c</sup>, Ni Nyoman Tri Puspaningsih<sup>a,d</sup>

<sup>a</sup>Department of Chemistry, Faculty of Science and Technology – Universitas Airlangga, Surabaya – Indonesia

<sup>b</sup>Department of Biology, Faculty of Science and Technology – Universitas Airlangga, Surabaya – Indonesia

<sup>c</sup>Faculty of Medical Science - University of Maarif Hasyim Latif, Sidoarjo - Indonesia

<sup>d</sup>Proteomic Study Group, Institute of Tropical Disease, Universitas Airlangga, Surabaya - Indonesia

### Abstract

A series of methoxy-4'-amino chalcone derivatives were tested for their antimicrobial activities against *Escherichia coli* ATCC 25923, *Staphylococcus aureus* ATCC 25922 and *Candida albicans* ATCC 10231. Furthermore, their molecular interactions with dihydropteroate synthase (DHPS) of *E. coli* and *S. aureus* were studied with a docking experiment. Compound 4 ((*E*)-1-(4-aminophenyl)-3-(2,3-dimethoxyphenyl)prop-2-en-1-one) exhibited the strongest activity, in which its activity was equal to sulfamerazine and sulfadiazine used as positive controls. In addition, it showed a good potential to be used as a wide spectrum antimicrobial agent. The in silico experiment showed that the prepared compounds had higher affinity to DHPS of *S. aureus* than to DHPS of *E. coli*. The tested compounds showed high similarity interaction with hydroxymethylpterin pyrophosphate (natural substrate of DHPS) in building intermolecular interactions.

© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer-review under responsibility of the organizing committee of the Molecular and Cellular Life Sciences: Infectious Diseases, Biochemistry and Structural Biology 2015 (MCLS 2015)

**Keywords:** Methoxy amino chalcones; antimicrobial; dihydropteroate synthase

\* Corresponding author. Tel.: -; fax: -.  
E-mail address: [herys08032002@yahoo.com](mailto:herys08032002@yahoo.com)

## 1. Introduction

Besides their simple preparation procedures, the increasing interest in chalcones is due to their various pharmacological potentials, such as antitumor<sup>1</sup>, anticancer<sup>2,3</sup>, antimalarial<sup>4</sup>, antihepatotoxic<sup>5</sup>, topoisomerase I inhibitor<sup>6</sup>, antiinflammation<sup>7</sup>, antioxidant<sup>8</sup> and antimicrobial activities<sup>9</sup>.

Overuse and misuse of antimicrobial agents increases cases of antimicrobial drug resistance<sup>10</sup>. Sulfanilamide derivatives as the first generation of antibiotics and penicillin as one of the next generation (Patrick, 2001) have been replaced gradually by new generations of antibiotics<sup>11</sup>. However, the needs for more potential antibiotics are still to be explored due to the global health problems worldwide caused by antimicrobial resistance, especially in the developing countries. Suwito *et al.*<sup>14</sup> have designed and synthesized a series of methoxy-4'-amino chalcone derivatives as antimicrobial agents mimicking the structure of PABA (*p*-amino benzoic acid) in the 4-amino benzoyl moiety<sup>12</sup>. PABA is a substrate of dihydropteroate synthase (DHPS) in the biosynthesis of 7,8-dihydropteroate, which is very important in folic acid biosynthesis needed for cell proliferation<sup>13</sup>. The rationale of the design was that the synthesized compounds could act as a competitive inhibitor of PABA.

In this work we report the study of intermolecular interactions between the prepared compounds with DHPS to develop better understanding of the molecular interactions, so that in the future we can design compounds possessing better anti-microbial activities.

## 2. Methods

### 2.1. Synthesis of chalcone derivatives

The target molecules (compound 1-7) were synthesized using the Claisen-Schmidt reaction as reported by Suwito *et al.*<sup>12</sup> Structure characterization of the prepared compounds was based on spectroscopic evidence.

### 2.2. Antimicrobial activity assay

Seven prepared compounds were tested for their antimicrobial activities against *Escherichia coli* ATCC 25923, *Staphylococcus aureus* ATCC 25922 and *Candida albicans* ATCC 10231. The antimicrobial test was performed using the disc diffusion method and the diameter of the inhibition zone was observed. The data obtained were then analyzed statistically using the Kolmogorov-Smirnov test and the Kruskal-Wallis test. Sulfadiazine and sulfamerazine were used as positive controls.

### 2.3 Software and program

The ligand structures were drawn in ChemBioDraw Ultra 11.0. DS visualizer 2.5 (Accelrys, Inc., USA) and PyMOL (DeLano Scientific LLC, USA) were used to modify the ligand and to visualize the receptor structure and docking results. The preparation of the DHPS *pdbqt* file and determination of the grid box size and position were carried out using AutoDock Tools version 1.5.6. AutoDock 4 was the sole docking program employed in this work<sup>15</sup>.

### 2.4 Preparation of the receptors and ligand structure

The three-dimensional structure of DHPS of *E. coli* complexed with sulfanilamide was retrieved from the Protein Data Bank [PDB:1AJ0], while the three-dimensional structure of DHPS of *S. aureus* complexed with 6-hydroxymethylpterin-diphosphate was downloaded from Protein Data Bank [PDB:1AD4]. The DHPS structure of *E. coli* was prepared for molecular docking by removing the sulphate ions and saved as a *pdb* file for the docking experiment. The DHPS three-dimensional structure of *S. aureus* was prepared for the docking experiment by removing K<sup>+</sup> and Mn<sup>2+</sup> ions and then saved as *pdb* file for molecular docking. The ligand structures were drawn using ChemBioDraw Ultra 11.0, subsequently modified using DS visualizer 2.5 and saved as a *pdb* file.

34

## 2.5 Docking experiment

Molecular docking was carried out using AutoDock 4<sup>15</sup>. AutoDock Tools 1.5.6 was utilized to prepare the input *pdbqt* files of DHPS of *E. coli* and *S. aureus* and to set the size and center of the grid box. The *E. coli* DHPS binding site was set at 41.903 x 8.143 x 2.045 Å, while the *S. aureus* binding site was set at 33.106 x 8.125 x 41.335 Å in the dimensions of x, y, z using 1.000 Å spacing for *E. coli* and 3.75 Å spacing for *S. aureus*. The *pdbqt* input file required for AutoDock4 was prepared by AutoDock Tools 1.5.6. The predicted binding affinity (kcal/mol), which describes the binding strength between ligand and receptor, is calculated based on the scoring function employed in AutoDock4. A more negative binding affinity indicates stronger binding. The docking experiment was performed according to the procedures provided by the AutoDock4 protocol.

29

## 3. Results and discussion

### 3.1. Antimicrobial activity

The compounds used in this work were methoxy-4'-amino chalcone derivatives, the molecular structures of which are displayed in Table 1. Their synthesis and structure characterization have been reported by Suwito *et al.*<sup>12</sup>




Table 1. Structure of the tested chalcones

Comp	Ring B					
	1	2,4	3	4	5	6
1	-	OMe	-	-	-	-
2	-	-	OMe	-	-	-
3	-	-	-	OMe	-	-
4	-	OMe	OMe	-	-	-
5	-	OMe	-	OMe	-	-
6	-	OMe	-	-	OMe	-
7	-	-	-	-	-	-

The data of antimicrobial tests of the entire compounds against *E. coli* ATCC 25923, *S. aureus* ATCC 25922, and *C. albicans* ATCC 10231 have been reported.<sup>14</sup> Dose dependence of the antimicrobial activity of the tested compound was observed. The higher the concentration of the tested compound, the larger the inhibition zone diameter.<sup>14</sup> The effect of the substituent of the prepared chalcone derivatives without considering the variation of concentration toward its activity was studied and is reported in this article. Based on the Kolmogorov-Smirnov test, it was shown that the diameters of the inhibition zones were not normally distributed ( $\alpha = 0.05$ ). The test was then followed with the Kruskal-Wallis test. The results of the tests are presented in Fig. 1.

Based on the analysis using the Kruskal-Wallis test toward *E. coli*, sulfadiazine showed the strongest inhibition activity, followed by sulfamerazine, compound 7 and then compound 4. However, the inhibition activity of compound 7 did not differ significantly from sulfamerazine and compound 4. Statistical analysis of the antimicrobial test toward *S. aureus* showed that compound 4 exhibited the strongest activity. However its activity did not differ significantly from sulfamerazine. The inhibition activity of compounds 7, 1, 5, 3 and 2 also did not differ significantly from sulfadiazine. The inhibition activity analysis toward *C. albicans* showed that sulfadiazine possessed the strongest activity, although its activity did not differ significantly from compounds 5, 3 and 4. Structure-activity relationship analysis employing Kruskal-Wallis test gave us information that the amino group played an important role in the antimicrobial activity, while the methoxy group on ring B played a less significant role. Among the tested compounds, compound 4, (*E*)-1-(4-aminophenyl)-3-(2,3-dimethoxyphenyl)prop-2-en-1-one, exhibited a promising wide spectrum of antimicrobial activity as strong as the positive controls.




Fig. 1. Histogram of the average of inhibition zone diameter and notation of the seven prepared compounds and two positive controls against *E. coli*, *S. aureus*, and *C. albicans* according to the Kruskal-Wallis test. SD = sulfadiazine, SM = sulfamerazine

### 3.2. In silico analysis

This article discusses the molecular interaction between DHPS of *E. coli* and *S. aureus* with the title compounds, as determined by in silico analysis, while the interaction with DHPS of *C. albicans* could not be discussed because the three dimensional structure of DHPS of *C. albicans* was not available in the PDB.

### 3.3. Validation of the docking

In the molecular docking experiment, to ensure that ligand conformations bound correctly with the binding pocket of the target protein, the size and position of the grid box had to be validated. In this work, docking validation was carried out by redocking the co-crystallized DHPS-sulphanilamide complex (PDB:1AJ0)<sup>16</sup> for *E. coli*, and the co-crystallized DHPS-6-hydroxymethylpterin diphosphate complex for *S. aureus* (PDB:1AD4).<sup>17</sup> We found that the binding conformations of both redocked complexes reproduced the binding modes of the co-crystallized complexes with binding affinities  $-2.91$  kcal/mol (RMSD =  $1.58$  Å) for *E. coli*, and  $-7.33$  kcal/mol (RMSD =  $0.83$  Å) for *S. aureus*.

According to Achary *et al.*<sup>16</sup>, the 7,8-dihydropterin pyrophosphate substrate bound in a deep cleft in the barrel, while sulphanilamide bound closer to the surface. Precisely, the sulphanilamide in the *E. coli* DHPS binding site was sandwiched between the main chain of Arg220 and the side chain of Lys221 on one side and the side chain of Arg63 on the other. The sulphonamide NH<sub>2</sub> formed a hydrogen bond to the carbonyl of Ser219, while sulphonamide oxygen accepted a hydrogen bond from the guanidinium of Arg63. Our redocking experiment provided the following observation: the sulphanilamide was flanked between the side chain of Lys221 and the main chain of Thr62. The hydrogen of aminophenyl moiety of sulphanilamide NH<sub>2</sub> donated a hydrogen bond with the carbonyl of Thr62, while the hydrogen of sulphonamide donated a hydrogen bond with the carbonyl of Ser219. Hydrophobic interactions were observed between sulphanilamide and Thr62, Phe190, Arg220, and Lys221.

According to Hampele *et al.*<sup>17</sup>, the interactions of hydroxymethylpterin pyrophosphate with amino acid residues of the *S. aureus* binding site occurred through polar and hydrophobic interactions. Amino acid residues involved in the polar interactions with the pterin moiety of the ligand were Asp167, Asn103, Lys203 and Asp84. The primary amine on C-2 and the protonated nitrogen at N-3 donated hydrogen bonds to the carboxylate oxygen atom of Asp167. Again the 2-NH<sub>2</sub> group donated a hydrogen bond with the side-chain oxygen of Asn103, while the NH group of Asn103 built a hydrogen bond with N-1 of the pterin. Lys203 donated its hydrogen atom to form a hydrogen bond with the oxygen atom of the carbonyl group of the pterin and with the N-5. Asp84 donated a hydrogen bond to the N-8 of the pterin moiety. Our redocking experiment conferred the following observation: the interactions observed between *S. aureus* DHPS binding site with 6-hydroxymethylpterin diphosphate are explained as follows: 6-hydroxymethylpterin diphosphate built hydrophobic interactions with Ile9, Asn11, Arg52, Asp84, Asn103, Met128, Asp167, Phe172, Lys203, Arg239 and His241, while hydrogen bonds existed between the pyrophosphate moiety and four amino acid residues, which were Arg52, Lys203, Arg239, and His241; Arg52 with 4-OH; Asp167 with NH<sub>2</sub>, Asn103 with 1-N-pterin; Arg52 with 4-N-pterin; and Asp84 with H of the N-8-pterin moiety.




Fig. 2. Redocking of sulphanilamide into *E. coli* DHPS (a) and 6-hydroxymethylpterin diphosphate into *S. aureus* DHPS (b). The docking poses of the ligand are shown in green carbons. Residues with hydrophobic contacts with sulphanilamide and 6-hydroxymethylpterin diphosphate are labelled in grey while hydrogen bonds are shown in yellow with dashed lines.

#### 3.4. Docking of the title compounds

In this work we performed molecular docking on 9 compounds, which were 7 derivatives of methoxy-4'-amino chalcone, sulfadiazine, and sulfamerazine. The results presenting the docking poses are displayed in Fig. 2, while the docking results and molecular interactions between tested compounds with amino acid residues of DHPS binding site are tabulated in the Table 2.

General observation of the docking results provided that the tested chalcone derivatives exhibited preponderant affinity with DHPS of *S. aureus* than with *E. coli* (based on the data of binding affinity). Three dimensional structural complementarities between the protein binding site and the ligands is also one of the important factors determining the binding affinity. Based on the sequence alignment of DHPS of *E. coli* and *S. aureus*, only 36% of homology was identified. This results provided that DHPS of both microbes were similar but not identical (Fig. 3), and the three dimensional structures of their binding sites differed consequently. In addition, all the tested compounds were bound in the deep cleft of the binding site, which was in accordance with the binding location of hydroxymethylpterin pyrophosphate. This information gave us an understanding that the prepared compounds, which were designed as competitive inhibitors of PABA, were more suitable to dock into a hydroxymethylpterin pyrophosphate binding location than into a sulphanilamide binding location.




Fig. 3. Docking poses of the tested compounds in the DHPS binding site, (a) *E. coli*, (b) *S. aureus*.




Fig. 4. Superimposed 3D-structure between DHPS of *E. coli* (pink) and *S. aureus* (blue).

In this article we discuss the docking results of compounds 4, 2 and 7. The reasons for the choice of these compounds were shown by the antimicrobial test, in which compound 4 (a dimethoxy-4'-amino chalcone derivative) displayed good potential to be used as a wide spectrum antimicrobial agent, compound 2 representing a monomethoxy-4'-amino chalcone derivative, and compound 7 representing 4'-amino chalcone derivative.

The factors considered for the scoring function in AutoDock4 were van der Waals interactions, electrostatic interactions, hydrogen bonds, desolvation, and rotations<sup>16</sup>. Besides hydrogen bonds, from the docking experiment we also obtained data about van der Waals and electrostatic interaction, which are not displayed in the Table 2.

Table 2. Docking results and molecular interactions between tested compounds with amino acid residues of DHPS.

Comp	DHPS <i>E. coli</i>		DHPS <i>S. aureus</i>	
	$\Delta G$ (kcal/mol)	Hydrogen bond interactions	$\Delta G$ (kcal/mol)	Hydrogen bond interactions
NS	-2.91	Thr62, Ser219	-7.33	Asn11, Asp84, Asn103, Lys203, Arg239, His241
1	-3.13	Pro145	-6.23	Asn103, Asp167, Lys203
2	-2.19	Thr62, Pro145	-6.48	Asn103, Asp167, Lys203
3	-1.05	Thr62, Pro145	-6.19	Asn103, Asp167, Lys203
4	-2.05	Thr62, Pro145	-6.10	Asn103, Asp167, Lys203
5	-1.09	Thr62, Pro145	-6.11	Asn103, Asp167, Lys203, Arg239
6	-0.66	---	-6.47	Asn103, Asp167, Lys203
7	-3.57	Pro145	-6.23	Asn103, Asp167, Lys203
SD	-3.25	Thr62, Arg63	-6.31	Val49, Asp84, Lys203
SM	-3.12	Thr62, Arg63	-6.08	Asn103, Asp167, Arg239

SD = sulfadiazine, SM = sulfamerazine, NS = natural substrate (as in PDB).




Fig. 5. Docking poses and intermolecular interactions of compound 2, 4 and 7 in DHPS binding sites. (a) *E. coli*, (b) *S. aureus*. The docking poses of the ligand are shown in green carbons. Residues with hydrophobic contacts with tested compounds are labelled in grey while hydrogen bonds are shown in grey with dashed lines.

### 3.4.1. Docking into *E. coli* DHPS (Figure 4a)

Compound 2: The location of ring A of compound 2 was flanked by the side chain of Gly189 and Arg63, whilst the ring B was located near the side chain of Lys221. The primary amino group of compound 2 donated hydrogen bonds to the carbonyl groups of Thr62 and Pro145. The position of the methoxy group was pointing to the



guanidinium moiety of Arg220. The amino acid residue Thr62, Arg63, Pro64, Pro145, Gly189, Arg220, Lys221, Ser222, and His257 contributed to hydrophobic interactions.

Compound 4: The position of ring A of compound 4 was flanked by the side chain of Arg63 and Pro64, whilst ring B was near the side chain of Lys221. The primary amino group of compound 4 made a bifurcated hydrogen bond with the carbonyl groups of Thr62 and Pro145. One methoxy group pointed to the guanidinium moiety of Arg63 and another pointed to the guanidinium moiety of Arg220. Amino acid residues Thr62, Arg63, Pro64, Pro145, Gly189, Arg220, Lys221, Ser222 and His257 contributed to hydrophobic interactions.

Compound 7: The side chain of Phe190 and Arg63 flanked the ring A of compound 7, while ring B was flanked by the side chains of Pro232 and Arg220. The primary amino group of compound 7 donated a hydrogen bond to the carbonyl group of Pro145. The amino acid residues involved in the formation of hydrophobic interactions were Thr62, Arg63, Pro64, Pro145, Phe190, Arg220, Lys221, Pro232 and His257.

### 3.4.2. Docking into *S. aureus* DHPS (Figure 4b)

Compound 2: The ring A of compound 2 was in a twisted conformation relative to ring B and was flanked by the side chains of three amino acid residues, Gln105, Asp167, and Ala199, whilst ring B was flanked by Asn103 and Arg239. The methoxy group pointed to the side chain of Asn11. The primary amino group of compound 2 made a bifurcated hydrogen bond with the carbonyl group of side chains of Asp167 and Asn103, while the carbonyl group of compound 2 acted as a hydrogen acceptor in the hydrogen bond with the amino group of the side chain of Lys203. Non-polar interactions were contributed by Asn11, Gly47, Val49, Asp84, Asn103, Gln105, Met128, Asp167, Phe172, Ala199, Lys203 and Arg239.

Compound 4: The ring A of compound 4 was flanked by the side chains of the amino acid residues Gln105, Arg239, and Ala199, whilst the ring B was flanked by Val49 and Ser40. The position of the two methoxy groups was pointing to Asn11 and His55. The amino group of compound 4 donated its two hydrogen atoms to the formation of hydrogen bonds with the side chain carbonyl groups of Asn103 and Asp167, while the carbonyl group of compound 4 acted as a hydrogen acceptor of the side chain amino group of Lys203 in hydrogen bond formation. The following amino acid residues Asn11, Val49, Ser50, His55, Asp84, Asn103, Gln105, Met128, Asp167, Phe172, Ala199, Lys203 and Arg239 contributed to hydrophobic interactions.

Compound 7: The ring A of compound 7 was located in the area flanked by Phe172, Asn103, and Ala199, while ring B was flanked by Val19 and Arg239. The amino group of compound 7 donated hydrogen bonds with the side chain carbonyl group of Asn103 and Asp167. The carbonyl group of compound 7 acted as a hydrogen acceptor of the side chain amino group of Lys203 in hydrogen bond formation. The hydrophobic interactions were contributed by the following amino acid residues: Asn11, Val49, Asp84, Asn103, Gln105, Met128, Asp167, Phe172, Ala199, Lys203 and Arg239.

The docking results showed that the amino acid residues involved actively in the molecular interaction from the docking experiment were in accordance with the results of protein-ligand complexes that were co-crystallized.

## 4. Conclusions

The antimicrobial activities of the methoxy amino chalcone derivatives have been assayed and their molecular interaction with the *E. coli* and *S. aureus* DHPS were studied with a docking experiment using the AutoDock4 program. Compound 4, (*E*)-1-(4-aminophenyl)-3-(2,3-dimethoxyphenyl)prop-2-en-1-one, exhibited a promising wide spectrum antimicrobial activity. The docking experiment gave us information that the prepared compounds were docked in the binding location of hydroxymethylpterin pyrophosphate in the DHPS binding site.

## Acknowledgment

The authors acknowledge the Directorate General of Higher Education of the Indonesian Ministry of Education and Culture for the research funding.

## References

1. Shih H, Deng L, Carrera CJ, Adachi S, Cottam HB, Carson DA. Rational design, synthesis and structure-activity relationship of antitumor (E)-2-benzylidene-1-tetralones and (E)-2-benzylidene-1-indanones. *Bioorg Med Chem Lett*, 2000; 10, 487-490.
2. Yun J-M, Kweon M-H, Kwon H, Hwang, J-K, Mukhtar H. Induction of apoptosis and cell cycle arrest by chalcone panduratin A isolated from *kaempferia pandurata* in androgen-independent human prostate cancer cells PC3 and CU145. *Carcinogenesis*, 2006; 27(7), 1454-1464.
3. Achanta G, Modzelewska A, Feng L, Khan SR, Huang P. A Boronic-chalcone derivative exhibits potent anticancer activity through inhibition of the proteasome. *Mol Pharmacol* 2006; 70, 426-433.
4. Chen M, Christensen SB, Zhai L, Rasmussen MH, Theander TG, Frokjaer S, Steffansen B, Davidsen J, Kharazmi A. The novel oxygenated chalcone, 2,4-dimethoxy-4'-butoxychalcone, exhibits potent activity against human malaria parasite *Plasmodium falciparum* in vitro and rodent parasites *Plasmodium berghei* and *Plasmodium yoelii* in vivo. *JID* 1997; 176:1327-1333.
5. Khan AA, Ahmed B, Alam T. Synthesis and antihepatotoxic activity of some new chalcones containing 1,4-dioxane ring system, *Pak J Pharm Sci*, 2006; 19(4), 290-294.
6. Yoon G, Kang BY, Cheon SH. Topoisomerase I inhibition and cytotoxicity of licochalcones A and E from *Glycyrrhiza inflata*. *Arch Pharm Res*, 2007, 30(3), 313-316.
7. Jin YL, Jin XY, Jin F, Sohn DH, Kim HS. Structure activity relationship studies of anti-inflammatory TMMC derivatives: 4-dimethylamino Group on the B ring responsible for lowering the potency. *Arch Pharm Res*, 2008; 31(9), 1145-1152.
8. Kim B-T, O K-J, Chun J-C, Hwang K-J. Synthesis of dihydroxylated chalcone derivatives with diverse substitution patterns and their radical scavenging ability toward DPPH free radicals. *Bull Korean Chem Soc*, 2008; 29(6), 1125-1130.
9. Choudhary AN, Juyal V. Synthesis of chalcone and their derivatives as antimicrobial agents. *Int J Pharm Pharm Sci* 2011; 3(3), 125-128.
10. Aly MEA, Essam TM, Amin MA. Antibiotic resistance profile of *E. coli* strains isolated from clinical specimens and food sample in Egypt. *Intl. J Microbiol Res* 2012; 3 (3): 176-182.
11. Patrick GL. *Medicinal Chemistry*, 1<sup>st</sup> ed., Oxford: Oxford University Press; 2004
12. Suwito H, Jumina, Mustofa, Pudjastuti P, Fanani MZ, Kimata-Arigo Y, Katahira R, Kawakami, T, Fujiwara T, Hase T, Sirat HM, Puspaningsih NNT. Design and synthesis of chalcone derivatives as inhibitors of the ferredoxin – ferredoxin-NADP<sup>+</sup> reductase interaction of *Plasmodium falciparum*: Pursuing new antimalarial agents. *Molecules*, 2014;19, 21473-21488, doi: 10.3390/molecules191221473.
13. Birmingham A, Derrick JP. The folic acid biosynthesis pathway in bacteria: evaluation of potential for antibacterial drug discovery, *BioEssay* 2002; 24, 637-648.
14. Suwito H, Jumina, Mustofa, Ni'matuzahroh, Puspaningsih NNT. Anticancer and Antimicrobial Activity of Methoxy Amino Chalcone Derivatives. *Der Pharma Chemica*, 2015; 7(3): 89-94.
15. Morris GM, Huey R, Olson AJ. Using AutoDock for ligand-receptor docking. *Curr Protoc Bioinformatics*, 2008; Chapter 8, Unit 8.14, doi: 10.1002/0471250953.bi0814s24.
16. Achary A, Somers DO, Champness JN, Bryant PK, Rosemond J, Stammers DK. Crystal structure of the anti-bacterial sulfonamide drug target dihydropteroate synthase. *Nat Struct Biol* 1997; 4: 490-497.
17. Hampele IC, D'Arcy A, Dale GE, Kostrewa D, Nielsen J, Oefner C, Page MG, Schonfeld HJ, Stuber D, Then RL., Structure and function of the dihydropteroate synthase from *Staphylococcus aureus*. *J Mol Biol* 1997; 268: 21-30.

# Antimicrobial Activities and In silico Analysis of Methoxy Amino Chalcone Derivatives

## ORIGINALITY REPORT

19%

SIMILARITY INDEX

11%

INTERNET SOURCES

14%

PUBLICATIONS

7%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="https://www.mdpi.com">mdpi.com</a> Internet Source	2%
2	Submitted to Higher Education Commission Pakistan Student Paper	1%
3	<a href="https://www.docplayer.net">docplayer.net</a> Internet Source	1%
4	Submitted to Purdue University Student Paper	1%
5	<a href="https://pubs.acs.org">pubs.acs.org</a> Internet Source	1%
6	<a href="https://academic.oup.com">academic.oup.com</a> Internet Source	1%
7	Asif Mir, Humaira Ismatullah, Sobiah Rauf, Umar H.K. Niazi. "Identification of bioflavonoid as fusion inhibitor of dengue virus using molecular docking approach", Informatics in Medicine Unlocked, 2016 Publication	1%
8	Submitted to University of Durham Student Paper	1%

9	Submitted to Thomas Edison State College	1 %
Student Paper		
10	<a href="http://www.portalgaruda.org">www.portalgaruda.org</a>	1 %
Internet Source		
11	Niltui, Piyaporn, Sirichai Kanking, Savaeng Techangamvong, Ekachai Wimolmala, Narongrit Sombatsompop, and E. Wimolmala. "Antibacterial Efficacy and Mechanical Properties of Silica Reinforced Natural Rubber (NR) with HPQM Based Neusilin", Advanced Materials Research, 2013.	1 %
Publication		
12	Suwito, Hery, EmptyYN="Y" Jumina, EmptyYN="Y" Mustofa, Pratiwi Pudjiastuti, Much Fanani, Yoko Kimata-Ariga, Ritsuko Katahira, Toru Kawakami, Toshimichi Fujiwara, Toshiharu Hase, Hasnah Sirat, and Ni Puspaningsih. "Design and Synthesis of Chalcone Derivatives as Inhibitors of the Ferredoxin — Ferredoxin-NADP+ Reductase Interaction of Plasmodium falciparum: Pursuing New Antimalarial Agents", Molecules, 2014.	1 %
Publication		
13	Submitted to Excelsior College	1 %
Student Paper		
14	Submitted to Heriot-Watt University	1 %
Student Paper		

15

Nitya Anand, William A. Remers. "Synthetic Antibacterial Agents", Wiley, 2010

Publication

1 %

16

[link.springer.com](http://link.springer.com)

Internet Source

1 %

17

M. M. Ghorab, M. S. A. El Gaby, N. E. Amin, N. M. H. Taha, M. A. Shehab, I. M. I. Faker. "Dapson in Heterocyclic Chemistry, Part IV: Synthesis of Some Novel Diphenylsulfones Containing Acetamide, Pyrrolidine, Piperazine, and Thiomorpholine Moieties as Antimicrobial and Antitumor Agents", Phosphorus, Sulfur, and Silicon and the Related Elements, 2008

Publication

<1 %

18

"Abstracts accepted for publication only", Clinical Microbiology and Infection, 4/2007

Publication

<1 %

19

Hampele, I.C.. "Structure and function of the dihydropteroate synthase from staphylococcus aureus", Journal of Molecular Biology, 19970425

Publication

<1 %

20

Prashant Chaudhari, Sanjay Bari. "In silico exploration of c-KIT inhibitors by pharmacoinformatics methodology: pharmacophore modeling, 3D QSAR, docking studies, and virtual screening", Molecular Diversity, 2015

Publication

<1 %

21	<a href="http://download.atlantis-press.com">download.atlantis-press.com</a> Internet Source	<1 %
22	<a href="http://www.gssrr.org">www.gssrr.org</a> Internet Source	<1 %
23	<a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a> Internet Source	<1 %
24	Valdameri, Glaucio, Luciana Pereira Rangel, Carmela Spatafora, Jérôme Guitton, Charlotte Gauthier, Ophélie Arnaud, Antonio Ferreira-Pereira, Pierre Falson, Sheila M. B. Winnischofer, Maria E. M. Rocha, Corrado Tringali, and Attilio Di Pietro. "Methoxy Stilbenes as Potent, Specific, Untransported, and Noncytotoxic Inhibitors of Breast Cancer Resistance Protein", ACS Chemical Biology, 2012. Publication	<1 %
25	<a href="http://res.mdpi.com">res.mdpi.com</a> Internet Source	<1 %
26	<a href="http://www.ajmb.org">www.ajmb.org</a> Internet Source	<1 %
27	<a href="http://www.freepatentsonline.com">www.freepatentsonline.com</a> Internet Source	<1 %
28	<a href="http://itd.unair.ac.id">itd.unair.ac.id</a> Internet Source	<1 %
29	Arrieta, M. P., M. A. Peltzer, M. C. Garrigos, and A. Jiménez. "Antibacterial biofilms	<1 %

based on calcium caseinate incorporated with carvacrol", *Microbes In Applied Research Current Advances and Challenges*, 2012.

Publication

---

30

R Novianty, S Ananta, M A Karim. "Herbal plants from Riau Province as inhibitors of COVID-19 binding to ACE2 receptor by computer aided molecular design an in-silico method", *Journal of Physics: Conference Series*, 2021

Publication

---

<1 %

31

[www.embopress.org](http://www.embopress.org)

Internet Source

---

<1 %

32

Alka N. Choudhary, Arun Kumar, Vijay Juyal. "Synthesis, Structure Activity Relationship (SAR), and Biological Activities of Benzylideneacetophenones Derivatives", *Anti-Inflammatory & Anti-Allergy Agents in Medicinal Chemistry*, 2016

Publication

---

<1 %

33

Mohammad H. Baig, Vishal M. Balaramnavar, Gulshan Wadhwa, Asad U. Khan. "Homology modeling and virtual screening of inhibitors against TEM- and SHV-type-resistant mutants: A multilayer filtering approach", *Biotechnology and Applied Biochemistry*, 2015

Publication

---

<1 %

34

Nhut Tran, Linh Tran, Ly Le. "Strategy in structure-based drug design for influenza A virus targeting M2 channel proteins", *Medicinal Chemistry Research*, 2013

Publication

<1 %

35

Okan Zafer Yeşilel, İnci İlker, Mustafa Serkan Soylu, Cihan Darcan, Yasemin Süzen. "Synthesis, crystal structures and antimicrobial properties of copper(II)-thiophene-2,5-dicarboxylate complexes with N-donor ligands", *Polyhedron*, 2012

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On



# Antimicrobial Activities and In silico Analysis of Methoxy Amino Chalcone Derivatives

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---

PAGE 9

---