

# POTENCY OF OIL SLUDGE INDIGENOUS BACTERIA FROM DUMAI-RIAU IN PRODUCING BIOSURFACTANT ON VARIATION OF SACCHARIDE SUBSTRATES

*by Ni'matuzahroh Ni'matuzahroh*

---

**Submission date:** 27-Apr-2021 10:56AM (UTC+0800)

**Submission ID:** 1570969964

**File name:** 2.\_POTENCY\_OF\_OIL\_SLUDGE\_INDIGENOUS.pdf (522.34K)

**Word count:** 4547

**Character count:** 25047

**2**  
**POTENCY OF OIL SLUDGE INDIGENOUS BACTERIA FROM DUMAI-RIAU IN  
PRODUCING BIOSURFACTANT ON VARIATION OF SACCHARIDE  
SUBSTRATES**

Ni'matuzahroh<sup>1</sup>, Erta Tri Yuliawatin<sup>1</sup>, Ditta Putri Kumalasari<sup>1</sup>, Nastiti Trikurniadewi<sup>1</sup>, Intan Ayu Pratiwi, Salamun<sup>1</sup>, Fatimah<sup>1</sup>, Sri Sumarsih<sup>2</sup>, and Hanif Yuliani<sup>3</sup>

<sup>1</sup>Department of Biology, Faculty of Science and Technology, Universitas Airlangga, Surabaya 60115;

<sup>2</sup>Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Surabaya 60115;

<sup>3</sup>Agency for The Assessment and Application of Technology, BPPT, South Tangerang 15314.

---

**Article Info**

**Article history:**

Received Jul 12<sup>th</sup>, 2017

Revised Aug 20<sup>th</sup>, 2017

Accepted Oct 26<sup>th</sup>, 2017

**Keyword:**

Biosurfactant  
D-glucose  
molasses  
oil sludge indigenous bacteria  
sucrose

---

**ABSTRACT**

This study aims to detect the ability of four species of Dumai-Riau oil sludge indigenous bacteria to produce biosurfactants using 3 types of substrate (D-glucose, sucrose and molasses) and to determine the relationship between the five biosurfactant detection methods. The bacteria were grown on liquid mineral medium with addition of 2% saccharide substrates. The culture was incubated for 4 days at 30°C. Production of biosurfactants by bacteria is detected in various ways by indirectly evaluating the presence of biosurfactant in the culture supernatant by measuring the surface tension (mN / m), emulsification (%), oil spreading, drop collapse and knowing the bacteria's ability to lysis blood agar media. The study used a completely randomized design with 4 treatments and 3 replications. The data were analyzed descriptively and statistically using Anova and continued with Duncan test and Pearson correlation test was performed to test the correlation between methods. The results showed the four bacteria gave different growth responses when grown on different substrates. The ability of bacteria to produce biosurfactants also varies depending on the type of substrates. The result of surface tension method has correlation with result of oil spreading method, blood agar test and emulsification activity.

**1**  
Copyright © 2017 Green Technology.  
All rights reserved.

---

**Corresponding Author:**

**Ni'matuzahroh,**  
Department of Biology,  
Faculty of Science and Technology,  
Universitas Airlangga, Surabaya 60115  
Email: [nimatuzahroh@fst.unair.ac.id](mailto:nimatuzahroh@fst.unair.ac.id)

---

**1. INTRODUCTION**

Oil sludge is one of the waste or by products from petroleum exploration and processing. Oil sludge is a hazardous and toxic substance, because of its nature, concentration and quantity that can harm the environment, and the survival of humans and other living things (Katz and Dawson, 1997). The oil sludge components are generally toxic, mutagenic and carcinogenic (Liu, et al., 2010).

Oil mining and processing industries such as PT. Chevron Indonesia and PT. Pertamina contained in Dumai-Riau will produce sludge waste with oil pollutants in considerable amounts, where sludge is obtained from oil drilling and deposits contained in temporary oil storage tanks (Kurniasari, 2005). Oil sludge in oil storage tanks and oil processing pipelines creates mud and crust deposits which can lead to pipe blockage and affect the total volume of oil tanks.

In the oil sludge, there are indigenous microorganisms that can adapt to extreme conditions and lack of oxygen. Indigenous bacteria came from soil and sea water which able to live by utilizing petroleum hydrocarbon compounds as their carbon sources. The capabilities of these bacteria, making indigenous bacteria have an influence in the effort to clean up the oil sludge deposits on processing pipes and oil storage tanks. One of the efforts is utilize biosurfactant as cleaner of tank and oil processing pipe.

Biosurfactant is an amphiphilic compound produced from extra cells and parts of bacterial cell membranes with various substrates including sugar, oil, alkane and waste substrate which are highly potential to be produced in various industrial scales (Banat, 1999; Lin, 1996; Mulligan, 2005). A microorganism that capable to producing biosurfactant compound and a substrate are required to produce biosurfactant. Substrate as a source of nutrients for the growth of microorganisms. The saccharide substrate for the production of biosurfactants may utilize a group of water-soluble monosaccharides and disaccharides such as glucose, sucrose (Desai and Banat, 1997). Molasses is also a water-soluble complex compound containing several compounds of the saccharide group such as glucose, sucrose, fructose and other compounds (Paturau, 1982 in Suastuti, 1998).

2  
To know the potency of indigenous oil sludge bacteria from Dumai-Riau to produce biosurfactant on variation saccharide substrate (D-glucose, sucrose and molasses), product biosurfactant is evaluated through several method, there are hemolytic activity, drop collapse, oil spreading, measurement of surface tension, and emulsification activity. This study is expected to provide information about potency of indigenous oil sludge bacteria from Dumai-Riau to produce biosurfactant on saccharide substrate for the future prospect as an oil tank cleaner.

## 2. RESEARCH METHOD

### Materials

Microorganisms that involved in this study are the oil sludge indigenous bacteria from Dumai-Riau, those are: D1, D2, D3 and D4. Isolates are collection of Microbiology Laboratory of Airlangga University. The bacterial growth medium consisted of Nutrient Agar (NA) (Oxoid), NB (Oxoid) and Mineral Synthetic (MS) Water (Putri and Cameotra, 1997) that modified with (g/L):  $(\text{NH}_4)_2\text{SO}_4$  3g;  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  0,2g; NaCl 10g;  $\text{CaCl}_2$  0.01g;  $\text{MnSO}_4 \cdot \text{H}_2\text{O}$  0.001g;  $\text{H}_3\text{BO}_3$  0.001g;  $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$  0.001g;  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$  0.001g;  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  0.005g;  $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$  0.001g. The buffer material (g/50mL) are  $\text{KH}_2\text{PO}_4$  1g;  $\text{K}_2\text{HPO}_4$  1g and  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$  0,0006g. The growth substrate were molasses, D-glucose (Phyto Technology Laboratories), sucrose (HIMEDIA). Then, the ingredients used for the detection of biosurfactants were agar (Oxoid), crude oil, and kerosene.

### Methods

#### Production of Biosurfactants

Each bacterial culture of the inclined NA aged 24-48 hours was inoculated into 50 mL of NB medium. Then, it was incubated in the incubator shaker for 24 hours at a temperature of 27-30°C to obtain  $\text{OD}_{\lambda 650\text{nm}} = 0.5$ . Bacterial culture is then referred to as a bacterial starter.

Biosurfactant production was performed by 4% (v/v) bacterial starter inoculated on 50 mL sterile MS medium with 2% substrate addition of D-glucose (w/v), sucrose (w/v) and molasses (v/v) on a 100 mL culture bottle separately. The culture was incubated in the incubator shaker with 120 rpm at 30°C for four days (Ni'matuzahroh et al., 2015).

### Growth Assay

Quantification of bacteria was carried out from day zero to day four of incubation time using the Total Plate Count (TPC) method with a series of dilutions. The growth test is accompanied by monitoring bacterial daily pH until the end of the incubation time. The pH measurements were performed using pH indicator and performed three replications in each bacterial culture.

### Biosurfactant Detection Method

#### Hemolytic activity

16  
Hemolytic activity can be done by scraping each strain of pure bacteria on blood agar medium then incubated for 48 hours at 37°C (Youssef et al., 2004). Positive test of the method of hemolytic activity is the formation of inhibitory zone around bacterial colonies. Inhibitory zone classified as  $\alpha$  hemolysis can be observed when around the colonies of bacteria formed greenish zone, zone  $\beta$  haemolysis can be observed to form a zone of white or clear around the bacterial colony. While the inhibitory zone classified as  $\gamma$  hemolysis has a noticeable feature there is no change in other colors around the bacterial colony (Shah et al., 2016).

#### Drop collapse

8  
Drop collapse test is used to determine the ability of oil distribution by biosurfactant activity. The drop collapse test was performed on the fourth day of biosurfactant production by dripping 135  $\mu$ L biosurfactant supernatant over 100  $\mu$ L of oil on a plate having a diameter of 8 mm and a height of 0.25 mm. The presence of biosurfactant is characterized by the formation of clear zone after 1-2 minutes (Jain et al., 1991).

#### Oil spreading

13  
The method of oil spreading follows the procedure of Morikawa et al. (2000). Measurement of spreading oil activity was done by 40 mL of distilled water into a Petri dish (150 mm), then inserting 10  $\mu$ L of crude oil on the surface of the aquades. 10  $\mu$ L of biosurfactant culture was added to the center of the surface of the crude oil. The diameter of the clear zone on the oil surface is measured using a sliding range.

#### 33 Measurement of surface tension

Measurements surface tension (TP) were performed using a Du-Nouy tensiometer. The surface tension value is expressed in units of mN/m or dyne/cm. Surface tension measurements were made on the fourth day of incubation time (Walter, 2010).

Calculation of surface tension using the formula:

$$r = r_0 \frac{\theta}{\theta_0}$$

Information:

r = sample surface tension

$r_0$  = surface tension aquades at  $t^{\circ}\text{C}$

$\theta$  = scale of sample measurement results

$\theta_0$  = scale of the result of aquatic size

#### Emulsification activity test


12  
Emulsification activity test was done according to Suryatmana et al. (2006) by mixing 1 mL of supernatant with 1 mL of test oil (kerosene) into the test tube. The mixture was mixed for 2 minutes, then emulsification activity (AE) (%) was measured after 1 hour and 24 hours. Emulsification activity was observed on the fourth day of incubation time.

Here is the formula for calculating % emulsification activity:


$$\% \text{emulsification} = \frac{\text{height of emulsion}}{\text{total fluid volume}} \times 100\%$$

### 3. RESULTS AND DISCUSSION


The relationship of growth response and pH of indigenous oil sludge isolate on D-glucose, sucrose, and molasses substrate is presented in Figures 1, 2 and 3.



17  
**Figure 1.** The growth response curve and pH of bacterial isolates (a) D1, (b) D2, (c) D3, and (d) D4 on D-glucose substrate .



6  
**Figure 2.** The growth response curve and pH of bacterial isolates (a) D1, (b) D2, (c) D3, and (d) D4 on the sucrose substrate



**Figure 3.** The growth and growth response curves of bacterial pH (a) D1, (b) D2, (c) D3, and (d) D4 on the molasses substrate

#### Test of hemolytic activity

The results of the hemolytic activity test of D1, D2, D3 and D4 bacteria in blood are listed in table 1.

**Table 1.** Results of hemolytic bacteria indigenous oil sludge Dumai-Riau bacterial activity test

No.	Bacterial Isolat	Clear Zone (cm)	Indeks of Clear Zone	Type of Clear Zone
1.	D1	1,97	1,26	$\alpha$
2.	D2	0,74	0,49	$\gamma$ - $\alpha$
3.	D3	0	0	$\gamma$
4.	D4	2,41	1,29	$\beta$

D3 bacterial isolates did not have the ability to lyse blood cells as showed from the measurements of hemolytic zones as listed in table 1. Isolate D3 is classified in hemolysis zone  $\gamma$  because no hemolytic zone (clear zone) is formed around the bacterial colony. D1 produces a zone of  $\alpha$  haemolysis, whereas D4 produces a zone of  $\beta$  hemolysis.

#### Drop collapse test

Drop collapse test results from D1, D2, D3 and D4 bacteria on D-glucose, sucrose, and molasses substrate are presented in Table 2.

**Table 2.** Drop collapse supernatant cultures of the fourth day of bacteria D1, D2, D3 and D4 on D-glucose, sucrose, and molasses substrate

Substrates	Clear Zone (cm)			
	D1	D2	D3	D4
<b>D-glucose</b>	<b>0,34 ±0,03</b>	0,35 ±0,09	0,26 <sup>a</sup> ±0,02	0,22 <sup>a</sup> ±0,04
<b>Sucrose</b>	0,31 ±0,01	0,28 ±0,06	0,20 <sup>a</sup> ±0,03	0,24 <sup>ab</sup> ±0,03
<b>Molasse</b>	0,21 ±0,08	0,39 ±0,03	0,36 <sup>b</sup> ±0,06	<b>0,30<sup>c</sup> ±0,04</b>

The highest drop collapse value produced by D3 bacterial supernatant is  $0.37 \pm 0.05$  cm, D1 bacteria on D-glucose substrate of  $0.34 \pm 0.03$  cm, D2 bacteria on molasses substrate of  $0.41 \pm 0.05$  cm and D4 bacteria on the molasses substrate of  $0.30 \pm 0.04$  cm.

#### Oil spreading test

Tests of spreading oil supernatant bacteria D1, D2, D3 and D4 on sucrose substrate, D-glucose, molasses are presented in table 3.


**Table 3.** Result of oil spreading supernatant test of fourth day culture from bacteria D1, D2, D3 and D4 on D-glucose, sucrose, and molasses substrate

Substrates	Clear Zone (cm)			
	D1	D2	D3	D4
<b>D-glucose</b>	0,24 ±0,07	<b>0,55 ±0,15</b>	0,18 <sup>a</sup> ±0,04	0,23 ±0,03
<b>Sucrose</b>	<b>0,41 ±0,13</b>	0,37 ±0,19	0,34 <sup>a</sup> ±0,08	<b>0,28 ±0,09</b>
<b>Molasse</b>	0,29 ±0,03	0,26 ±0,05	<b>0,71<sup>b</sup> ±0,04</b>	0,22 ±0,00

The largest oil spreading value among the three other bacteria came from D3 bacteria on the molasses substrate of  $0.71 \pm 0.04$  cm. Meanwhile, the largest oil distribution value in D1 and D4 bacteria occurred on sucrose substrate with value of  $0.41 \pm 0.13$  cm and  $0.28 \pm 0.09$  cm respectively, while the largest oil distribution value of D2 bacteria occurred at D-glucose substrate of  $0.55 \pm 0.15$  cm.

#### Surface tension

Values and graphs of surface tension of bacteria D1, D2, D3 and D4 are shown in Fig. 4.




**Figure 4.** Diagrams of supernatant tissue surface degradation value of the fourth day of bacteria D1, D2, D3 and D4 on D-glucose, sucrose, and molasses substrate

In Figure 4, it is known that D1, D2, D3 and D4 bacteria grown on the D-glucose, sucrose, and molasses substrate have the ability to reduce surface tension to media and aquades. The average of surface tension reduction of four isolates on the variationsaccharide substrate is more than 10 dyne/cm.

#### Emulsification activity

Diagram of the value of 1-hour emulsification activity (E1) of bacteria D1, D2, D3 and D4 on D-glucose, sucrose, molasses is presented in figure 5.



**Figure 5.**Diagram of 1 (E1) and 24 hour (E24) emulsification activity values of D1, D2, D3 and D4 bacteria on D-glucose,sucrose, molasses substrate with kerosene test compound.

In the D-glucose and sucrose substrate, the highest emulsification activity value was achieved by the same D1 and D4 bacteria with 24.53± 9.0% and 23.43±5.52% respectively. Meanwhile, the highest emulsification



activity value among the three other bacteria of  $28.85 \pm 5.55\%$  occurred in D3 bacteria with molasses growth substrate. The average decrease in value of 1-hour emulsification activity to the value of the 24 hours emulsification activity is 8.56%.

The statistical test used to show the relationship between biosurfactant detection methods is Pearson correlation. Statistical test using SPSS software (version 21.0) Pearson correlation coefficient ( $\rho$ ) can show the value between -1 (strong negative relationship) and 1 (strong positive relationship). Table 4 shows that on each substrate yields correlation coefficient values between different methods. In the glucose substrate, there is a relationship between oil spreading and surface tension with Pearson correlation coefficient  $\rho = -0,893$ . In the sucrose substrate, the strongest relationship (Pearson correlation coefficient  $\rho = 0,736$ ) occurs between the emulsification activity and the blood agar. In the molasses substrate, several biosurfactant detection methods that show a relationship between oil spreading and blood agar ( $\rho = -0,776$ ), surface tension and drop collapse ( $\rho = 0,738$ ), and between surface tension and emulsification activity ( $\rho = 0,856$ ).

#### Comparative statistics of five screening methods

Table 4. Pearson's correlation results from five screening methods

		Pearson Correlation Coefficient ( $\rho$ )				
		OS	DC	ST	EA	BA
Glucose	OS	1	0,565	-0,893**	0,223	-0,179
	DC	0,565	1	-0,538	-0,318	-0,115
	ST	-0,893**	-0,538	1	-0,390	0,253
	EA	0,223	-0,318	-0,390	1	0,194
	BA	-0,179	-0,115	0,253	0,194	1
Sucrose	OS	1	0,043	-0,421	-0,272	-0,48
	DC	0,043	1	-0,485	0,509	0,415
	ST	-0,421	-0,485	1	-0,130	0,182
	EA	0,509	0,509	-0,130	1	0,736**
	BA	-0,48	0,415	0,182	0,736**	1
Molasse	OS	1	0,232	0,029	0,274	-0,776**
	DC	0,232	1	0,738**	0,547	-0,590*
	ST	0,029	0,738**	1	0,856**	-0,395
	EA	0,274	0,547	0,856**	1	-0,421
	BA	-0,776**	-0,590*	-0,395	-0,421	1

#### DISCUSSION

Biosurfactant production conducted until the fourth day aims to ensure the viability condition of bacterial cells at the time of biosurfactant production test on the fourth day of incubation time. The growth curve of D1, D2, D3 and D4 bacteria on the fourth day (figure 1) in cultures of variation of the saccharide substrate mostly occurs at the end of the exponential phase to the stationary phase. The exponential phase occurring on the fourth day of incubation time is quite possible because the growth medium is still sufficient in culture so that bacteria can still utilize the medium well for its growth. This condition is in accordance with that of Ni'matuzahroh et al. (2010), the optimum biosurfactant activity with 2% molasses substrate occurs on the fourth day as the exponential phase ends or at the beginning of the stationary phase.

Figure 1 shows that when bacterial growth in the exponential phase, the culture pH decreases. The larger the number of bacterial cells that grow in a culture the lower the pH culture. The condition is indicated that bacteria D1, D2, D3 and D4 can utilize well the saccharide substrate as carbon source in metabolism process of its growth which is accompanied by decrease of pH culture. Hoog (2005) in his research also mentions that the bacteria can

9  
perform their metabolism by lowering the pH of the media environment due to the growth of bacteria in the growth medium. According to Pratiwi (2014), an increase in the number of bacteria along with a decrease in pH value due to the production of metabolites that can reduce the pH of the growth media environment.

Tests on biosurfactant products by indigenous oil sludge bacteria Dumai-Riau was first performed by hemolytic activity using blood. The positive test is characterized by the presence of a hemolysis zone formed. The inhibitory zone formed in the hemolytic activity test indicates the production of biosurfactant. This is consistent with that reported by Singh (2012) in his study, that the larger the diameter of lysis in the blood in order to result from the increasing concentration of biosurfactant. According to Zaragoza et al. (2010) the formation of a hemolytic zone (clear zone) in blood hemolysis tests is caused by two different mechanisms ie the dissolution of cell membranes that normally occur at high biosurfactant concentrations or due to increased membrane permeability to small soluble substances that normally occur when biosurfactant concentrations low, thus causing osmotic lysis.

After testing the hemolytic activity using blood agar, a stabilization test of surfactant droplets in an oil is coll drop and oil spreading test. According to Walter et al. (2010), drop collapse testing is associated with drip stability that depends on surfactant concentration and is related to surface and interface stresses. A supernatant containing biosurfactant will spread because of the interceptor force or tension between the supernatant and the reduced (hydrophobic) layer of the oil layer. The diameter of the clear zone produced by the oil spreading test indicates the presence of surfactant activity and there is a linear relationship between the surfactant and the resulting clear zone diameter.

Further biosurfactant detection is to use the principle of oil stabilization and water emulsion. In Figure 5, the value of D1, D2, D3 and D4 D1, D2, and D4 emulsification activity on D-glucose, sucrose, and molasses decreased from 1-hour emulsification activity (E1). This is due to the unstable emulsion properties due to the tendency of the emulsion particles to join the other particles, so the amount of emulsion produced tends to decrease at 24 hours. According to Karthik (2010), emulsification activity shows the stability of biosurfactant when incubated at room temperature.

A hydrophilic bacterial culture in contact with hydrophobic hydrocarbons (kerosene) is likely to have the ability to form emulsions that separate two distinct phases due to the high interface surface area. The spread and size of the emulsion grains may change over time. The result showed that the decrease of emulsification activity between the measurement time 1 hour and 24 hours. This is due to the unstable emulsion properties due to the tendency of the emulsion particles to join the other particles, so the amount of emulsion produced tends to decrease at 24 hours. The stability of the emulsion is proportional to the stability of the biosurfactant (emulgator) produced. As Karthik (2010) points out, the emulsification activity shows the stability of biosurfactant when incubated at room temperature.

The ability to stabilize emulsions indicates that bacteria can produce biosurfactants (Batista et al., 2006). Emulsification activity is also a mechanism used by bacteria to reach its growth substrate with the help of biosurfactants as emulsifiers produced by the bacteria itself. The value of different emulsification activities on different substrates indicates the effect of carbon sources on bacterial culture in generating emulsification. As reported by Hamed (2012), *S. aureus*, *Micrococcus spp.*, *Pseudomonas sp.*, *P. aeruginosa*, *Photobacter damsela* and *Chrysonnasluteola* strains are more efficient in emulsifying mineral oil than crude oil.

The average value of the surface tension of four bacterial isolates yields a value of more than 10 dyne / cm. This indicates that four bacterial isolates are capable of producing biosurfactant as a surface active agent when grown on a variety of saccharide substrate. As reported by Ni'matuzahroh et al. (2013) that the reduction of the surface value of the bacterial supernatant of  $\geq 10$  dyne / cm indicates that biosurfactant is potential as a surface active agent. The occurrence of surface tension decline is caused by the production of surface active compounds by microorganisms (Batista et al., 2006).

According to some biosurfactant detection results, the four indigenous oil sludge bacteria Dumai-Riau are known to produce biosurfactants grown on D-glucose, sucrose and molasses substrate. The best sequence of bacteria to produce biosurfactants based on the high average values in the biosurfactant production test on all variations of the saccharide substrate were bacteria D1, D2, D3 and D4. However, the bacteria that have the most potential ability to produce biosurfactant in this study is bacteria D1. The most potential growth substrate for producing biosurfactants is the molasses substrate, evidenced by the superior values in the growth response test and biosurfactant detection of several bacterial cultures compared to D-glucose and sucrose substrate.

Correlation test results show the relationship between methods of each substrate. This suggests that growth substrate may affect the biosurfactant product, so the type of biosurfactant produced may also influence which detection method is most suitable for use. For biosurfactant detection, it is recommended to use several methods in order to obtain accurate data results by linking some results from other methods used.

## CONCLUSION

The four indigenous oil sludge bacteria Dumai-Riau (D1, D2, D3 and D4) are able to produce biosurfactant on variations of saccharide substrate (molasses, sucrose, D-glucose) with clear zones of hemolytic activity, drop collapse value of 0.20-0.41 cm, the value of oil spreading ranges from 0.18-0.71 cm, the emulsification activity decrease value of 8.56% and the value of surface tension  $\geq 10$  dyne / cm. The average of the results of each biosurfactant production test on some bacteria, the molasses substrate has the best ability to produce biosurfactants. The most potent bacterial indigenous species produce bacterial biosurfactantA (4).

## ACKNOWLEDGMENTS

This research was conducted with funding from Kemenristek Dikti. Thanks for the support facilities and infrastructure of the Department of Biology and Department of Chemistry, Faculty of Science and Technology, Airlangga University and for the support delivery of oil sludge from BPPT, Tangerang.

## REFERENCES

- Banat, I. M., Samarah, N., Murad, M., Horne, R., dan Banerjee, S., 1991, Biosurfactant Production and Use In Oil Tank Clean-Up, *World Journal of Microbiology and Biotechnology* 7(1): 80-88.
- Batista, S. B., Mounteer, A. H., Amorim, F. R., dan Totola, M. R., 2006, Isolation and Characterization of Biosurfactant/Bioemulsifier-Producing Bacteria from Petroleum Contaminated Sites, *Journal of Bioresource Technology* 97(6): 868-875.
- Desai, J. D. dan Banat, I. M., 1997, Microbial Production of Surfactants and Their Commercial Potential, *Microbiology and Molecular Biology Reviews* 61(1): 47-64.
- Hogg, Stuart, 2005, *Essential Microbiology*, The University of Glamorgan, UK.
- Jain, D. K., Collins-Thomson, D. L., Lee, H., dan Trevors, J.T., 1991, A Drop-Collapsing Test for Screening Surfactant-Producing Microorganisms. *Journal of Microbiological Methods* 13(4): 271-279.
- Karthik, L., Kumar, G., dan Rao, K.V.B., 2010, Comparison of Methods and Screening of Biosurfactant Producing Marine Actinobacteria

- Katz, B. J., dan Dawson, W. C., 1997, A Structural and Tectonic Model of The Coastal Plain Block, Central Sumatra Basin, Indonesia, *Proceedings of Indonesian Petroleum Association*, 22th Annual Convention : 286-317p
- Kurniasari, L., 2005, Pengolahan Awal Lumpur Minyak dengan Metode Volatilisasi. *Momentum*, 1(2).
- Lin, S. C., 1996, Biosurfactants: Recent Advances. *Journal of Chemical Technology and Biotechnology* 66(2): 109-120.
- Liu, W., Luo, Y., Teng, Y., Li, Z., dan Ma, L. Q., 2010, Bioremediation of Oily Sludge-Contaminated Soil by Stimulating Indigenous Microbes, *Environmental geochemistry and health* 32(1) :23-29
- Morikawa, M., Hirata, Y., dan Imanaka, T., 2000, A Study on The Structure-Function Relationship of Lipopeptide Biosurfactants. *Biochimica et Biophysica Acta (BBA-Molecular and Cell Biology of Lipids)* 1488(3) : 211-218.
- Mulligan, C. N., 2005, Environmental Applications for Biosurfactants, *Environmental pollution* 133(2) : 183-198.
- Ni'matuzahroh, Alami, N.H., Khudlari, F., Fatimah, dan Tri, N., 2010, Studi Kinetika Produksi Biosurfaktan *Bacillus subtilis* 3KP pada Substrat Molase, *Berkala Penelitian Hayati* 16(33-38).
- Ni'matuzahroh, Fatimah, Tini, S., Sri, S., 2013, "Biocleanoil" Produk Berbasis Mikroba untuk Pengolahan Sludge Industri Minyak, *UPT Research Report Universitas Airlangga*. Surabaya.
- Ni'matuzahroh, Pratiwi, I.A., Irmayanti, I.S., Surtiningsih, T., Fatimah, Sumarsih, S., 2013, Combination of *Acinetobacter* sp. P2 (1) Biosurfactant and Crude Lipases on The Oil Sludge Solubility. *Proceedings of The "4<sup>th</sup> International Conference & Workshop on Basic and Applied Sciences*. Ibnu Sina Institute for Fundamental Science Studies Universiti Teknologi Malaysia 81310 UTM Johor Bahru. 172-177.671-674.
- Ni'matuzahroh, Puspitasari, A. O., Pratiwi, I.A., Fatimah, Sumarsih, S., Surtiningsih, T., Salamun., 2015. Oil Removal from Petroleum Sludge using Bacterial Culture with Molasses Substrate at Temperature Variation. *Proceedings 5th International Conference and Workshop on Basic and Applied Sciences (ICOWOBAS 2015) AIP Conf. Proc.* 1718, 030002-1–030002-8; doi: 10.1063/1.4943312 © 2016 AIP Publishing LLC 978-0-7354-1364-1
- Ni'matuzahroh, Trikurniadewi, N., Pramadita A. R. A. , Pratiwi I. A. Salamun, Fatimah, Sumarsih, S. 2017. Biodegradation of Naphthalene and Phenanthrene by *Bacillus subtilis* 3KP. *Proceedings 3rd International Biology Conference (IBOC) and 10 th KOREA-ASEAN Biomass Symposium: Biodiversity and biotechnology for human welfare*. AIP Conf. Proc.1854.020026. AIP Publishing ISBN : 978-0-7354-1528-7
- Pruthi, V. dan Cameotra, S. S., 1997, Rapid Identification of Biosurfactant-Producing Bacterial Strains Using a Cell Surface Hydrophobicity Technique, *Biotechnology Techniques*, 11(9):
- Shah, N., Nikam, R., Gaikwad, S., Sapre, V., dan Kaur, J, 2016, Biosurfactant: Types, Detection Methods, Importance and Applications. *Indian J Microbial Res* 2016 3(1):5-10.
- Singh, V., 2012, Biosurfactant - Isolation, Production, Purification and Significance, *Int J Sci Res Pub*, 2(7):1-4.
- Suastuti, N. G. A. M. A., 1998, Pemanfaatan Hasil Samping Industri Pertanian (Molase dan Limbah Cair Tahu) sebagai Sumber Karbon dan Nitrogen untuk Produksi Biosurfaktan oleh *Bacillus* sp. Galur Komersial dan Lokal, Institut Pertanian Bogor.

Suryatmana, P. E. Kardena, E. Ratmaningsih, dan Wisjnuaprpto, 2006, Karakteristik Biosurfaktan dari *Azotobacter chroococcum*. *Jurnal Microbiologi Indonesia*. 2(1): 30-34.

Walter, V., Syldatk, C. dan Hausmann, R., 2010, Screening Concepts for The Isolation of Biosurfactant Producing Microorganisms, *Biosurfactants* (pp. 1-13). Springer New York.

Youssef, N. H., Duncan, K. E., Nagle, D. P., Savage, K. N., Knapp, R. M., dan McInerney, M. J., 2004, Comparison of Methods to Detect Biosurfactant Production by Diverse Microorganisms, *Journal of Microbiological Methods* 56(3): 339-347.

Zahroh, Fatimatuz, 2011, Pengaruh konsentrasi Gula Cair dan Waktu Inkubasi Terhadap Produksi Biosurfaktan *Bacillus subtilis* 3KP, *Skripsi*, Universitas Airlangga, Surabaya.

Zaragoza, A., Fransisco J.A., Maria J.E., Jose A.T., Ana M., Angeles M. dan Antonio O., 2010, Hemolytic Activity of a Bacterial Trehalose Lipid Biosurfactant Produced by *Rhodococcus* sp.: Evidence for a Colloid-Osmotic Mechanism, *Langmuir* 26(11): 8567-8572.

# POTENCY OF OIL SLUDGE INDIGENOUS BACTERIA FROM DUMAI-RIAU IN PRODUCING BIOSURFACTANT ON VARIATION OF SACCHARIDE SUBSTRATES

## ORIGINALITY REPORT

17%

SIMILARITY INDEX

15%

INTERNET SOURCES

8%

PUBLICATIONS

8%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://repository.unair.ac.id">repository.unair.ac.id</a> Internet Source	6%
2	<a href="http://sinta3.ristekdikti.go.id">sinta3.ristekdikti.go.id</a> Internet Source	1%
3	<a href="http://www.omicsonline.org">www.omicsonline.org</a> Internet Source	1%
4	Fisseha Andualem Bezza, Mervyn Beukes, Evans M. Nkhalambayausi Chirwa. "Application of biosurfactant produced by <i>Ochrobactrum intermedium</i> CN3 for enhancing petroleum sludge bioremediation", <i>Process Biochemistry</i> , 2015 Publication	1%
5	Shima Afshar. "Comparative approach for detection of biosurfactant-producing bacteria isolated from Ahvaz petroleum excavation areas in south of Iran", <i>Annals of Microbiology</i> , 09/2008 Publication	<1%

6	Remon, S.. "Burlat cherry quality after long range transport: optimisation of packaging conditions", Innovative Food Science and Emerging Technologies, 200312 Publication	<1 %
7	link.springer.com Internet Source	<1 %
8	scienceandnature.org Internet Source	<1 %
9	worldwidescience.org Internet Source	<1 %
10	sevgiligiyim.com Internet Source	<1 %
11	www.adb.org Internet Source	<1 %
12	Ni'matuzahroh, S K Sari, N Trikurniadewi, A D Pufita, I P Ningrum, S N M M Ibrahim, T Nurhariyati, Fatimah, T Surtiningsih. " Utilization of Rice Straw Hydrolysis Product of sp. H9 as A Substrate of Biosurfactant Production by LII61 Hydrocarbonoclastic Bacteria ", IOP Conference Series: Earth and Environmental Science, 2019 Publication	<1 %
13	Noha H. Youssef, Kathleen E. Duncan, David P. Nagle, Kristen N. Savage, Roy M. Knapp,	<1 %

Michael J. McInerney. "Comparison of methods to detect biosurfactant production by diverse microorganisms", Journal of Microbiological Methods, 2004

Publication

14

Submitted to Al-Mashrek International School

Student Paper

<1 %

15

Submitted to University of Arizona

Student Paper

<1 %

16

Ebrahimi, Azizollah, and Najmeh Tashi. "Isolation of Biosurfactant Producing Bacteria From Poultry Breast Skin", Jundishapur Journal of Natural Pharmaceutical Products, 2012.

Publication

<1 %

17

[hagi-www.ics.es.osaka-u.ac.jp](http://hagi-www.ics.es.osaka-u.ac.jp)

Internet Source

<1 %

18

[www.investopedia.com](http://www.investopedia.com)

Internet Source

<1 %

19

Agus Triawan, Ni'matuzahroh, Agus Supriyanto. "Effects of the combination between bio-surfactant product types and washing times on the removal of crude oil in nonwoven fabric", AIP Publishing, 2017

Publication

<1 %

20

Ijeoma Vivian Nwaguma, Chioma Blaise Chikere, Gideon Chijioke Okpokwasili.

<1 %



"Isolation, characterization, and application of biosurfactant by Klebsiella pneumoniae strain IVN51 isolated from hydrocarbon-polluted soil in Ogoniland, Nigeria", Bioresources and Bioprocessing, 2016

Publication

---

21

Submitted to 76830

Student Paper

<1 %

---

22

Chirwa, Evans M. N., Tshepo Mampholo, and Oluwademilade Fayemiwo. "Biosurfactants as demulsifying agents for oil recovery from oily sludge – performance evaluation", Water Science & Technology, 2013.

Publication

---

23

I. M. Banat. "Characterization of biosurfactants and their use in pollution removal - State of the Art. (Review)", Acta Biotechnologica, 1995

Publication

---

24

Mohammed Haider Alley. "Four-quadrant assessment of gestational age-specific values of amniotic fluid volume in uncomplicated pregnancies", Acta Obstetricia Et Gynecologica Scandinavica, 3/1998

Publication

---

25

Pacwa-Płociniczak, Magdalena, Tomasz Płociniczak, Joanna Iwan, Monika Żarska, Mirosław Chorążewski, Marzena Dzida, and

<1 %

Zofia Piotrowska-Seget. "Isolation of hydrocarbon-degrading and biosurfactant-producing bacteria and assessment their plant growth-promoting traits", Journal of Environmental Management, 2016.

Publication

26

[aip.scitation.org](http://aip.scitation.org)

Internet Source

<1 %

27

[biocoreopen.org](http://biocoreopen.org)

Internet Source

<1 %

28

[repositorium.sdum.uminho.pt](http://repositorium.sdum.uminho.pt)

Internet Source

<1 %

29

[scholar.sun.ac.za](http://scholar.sun.ac.za)

Internet Source

<1 %

30

[www.frontiersin.org](http://www.frontiersin.org)

Internet Source

<1 %

31

[www.scribd.com](http://www.scribd.com)

Internet Source

<1 %

32

Vivek Kumar Gaur, Natesan Manickam.

"Microbial Biosurfactants: Production and Applications in Circular Bioeconomy", Elsevier BV, 2021

Publication

<1 %

33

Jing Li, Maocheng Deng, Yao Wang, Weixin Chen. "Production and characteristics of biosurfactant produced by Bacillus

<1 %

pseudomycoides BS6 utilizing soybean oil waste", International Biodeterioration & Biodegradation, 2016

Publication

---

34

Vanessa Walter, Christoph Syldatk, Rudolf Hausmann. "Chapter 1 Screening Concepts for the Isolation of Biosurfactant Producing Microorganisms", Springer Science and Business Media LLC, 2010

Publication

---

<1 %

---

Exclude quotes      Off

Exclude matches      Off

Exclude bibliography      On

# POTENCY OF OIL SLUDGE INDIGENOUS BACTERIA FROM DUMAI-RIAU IN PRODUCING BIOSURFACTANT ON VARIATION OF SACCHARIDE SUBSTRATES

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---

PAGE 9

---

PAGE 10

---

PAGE 11

---

PAGE 12

---