

Biodegradation of Naphthalene and Phenanthren by *Bacillus subtilis* 3KP

by Ni'matuzahroh Ni'matuzahroh

Submission date: 27-Apr-2021 10:53AM (UTC+0800)

Submission ID: 1570966974

File name: 4._Biodegradation_of_napthalene_-PI.pdf (511.63K)

Word count: 3183

Character count: 18338

Biodegradation of naphthalene and phenanthren by *Bacillus subtilis* 3KP

Cite as: AIP Conference Proceedings 1854, 020026 (2017); <https://doi.org/10.1063/1.4985417>
Published Online: 26 June 2017

Ni'matuzahroh, N. Trikurniadewi, A. R. A. Pramadita, I. A. Pratiwi, Salamun, Fatimah, and Sri Sumarsih



View Online



Export Citation

ARTICLES YOU MAY BE INTERESTED IN

11 [Bioremediation of oil sludge using a type of nitrogen source and the consortium of bacteria with composting method](#)

AIP Conference Proceedings 1854, 020011 (2017); <https://doi.org/10.1063/1.4985402>

1 [Effects of the combination between bio-surfactant product types and washing times on the removal of crude oil in nonwoven fabric](#)

AIP Conference Proceedings 1854, 020037 (2017); <https://doi.org/10.1063/1.4985428>

3 [Manganese \(Mn\) stress toward hyperaccumulators plants combination \(HPC\) using *Jatropha curcas* and lamtoro gung \(*L. leucocephala*\) in mycorrhizal addition on soybean \(*Glycine max*\) seedling stage](#)

AIP Conference Proceedings 1854, 020027 (2017); <https://doi.org/10.1063/1.4985418>

Lock-in Amplifiers
up to 600 MHz



12 [AIP Conference Proceedings 1854, 020026 \(2017\); <https://doi.org/10.1063/1.4985417>](#)

1854, 020026

© 2017 Author(s).

Biodegradation of Naphthalene and Phenanthrene by *Bacillus subtilis* 3KP

Ni'matuzahroh^{1,a)}, Trikurniadewi N.^{1,b)}, Pramadita A. R. A.¹, Pratiwi I. A.¹,
Salamun¹, Fatimah¹, Sumarsih Sri²

¹ Department of Biology, Faculty of Science and Technology, Universitas Airlangga, 60115 Surabaya, Indonesia.

² Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, 60115 Surabaya, Indonesia.

^{a)} Corresponding author: nimatuzahroh@fst.unair.ac.id

^{b)} nastiti.trikurniadewi-2015@fst.unair.ac.id

25

Abstract. The purposes of this research were to know growth response, degradation ability, and uptake mechanism of naphthalene and phenanthrene by *Bacillus subtilis* 3KP. *Bacillus subtilis* 3KP was grown on Mineral Synthetic (MS) medium with addition of 1% yeast extract and naphthalene and phenanthrene respectively 200 ppm in different cultures. *Bacillus subtilis* 3KP growth response was monitored by Total Plate Count (TPC) method, the degradation ability was monitored by UV-Vis spectrophotometer, and the uptake mechanism of hydrocarbon was monitored by emulsification activity, decrease of surface tension, and activity of Bacterial Adherence to Hydrocarbon (BATH). *Bacillus subtilis* 3KP was able to grow and show biphasic growth pattern on both of substrates. Naphthalene and phenanthrene were used as a carbon source for *Bacillus subtilis* 3KP growth that indicated by the reduction of substrate concomitant with the growth. At room temperature conditions ($\pm 30^{\circ}\text{C}$) and 90 rpm of agitation for 7 days, *Bacillus subtilis* 3KP could degrade naphthalene in the amount of 70.5% and phenanthrene in the amount of 24.8%. Based on the analysis of UV-Vis spectrophotometer, three metabolites, 1-hydroxy-2-naphthoic acid, salicylic acid, and pyrocatechol were found in both cultures. The metabolite identification became basis of propose degradation pathway of naphthalene and phenanthrene by *Bacillus subtilis* 3KP. The results of hydrocarbon uptake mechanism test show that *Bacillus subtilis* 3KP used all of the mechanism to degrade naphthalene and phenanthrene.

Keyword: *Bacillus subtilis* 3KP, Biodegradation, Biosurfactant, Phenanthrene, Naphthalene

INTRODUCTION

Naphthalene and phenanthrene are types of hydrocarbon that classified as polycyclic aromatic hydrocarbon (PAH). This kind of compound can cause damage on mammal tissue when it is exposure in high dose (Patnaik, 1994). The existence of PAH, as pollutant, is often yield from exploration and production in petroleum industry. Petroleum waste sludge (oil sludge) consists of hydrocarbon compounds such as aliphatic hydrocarbons and polycyclic aromatic hydrocarbons (PAH), water, metals and non-hydrocarbon compounds such as nitrogen, sulfur, oxygen, and asphalt (Connell and Miller, 1995). PAH almost reaches amount of 13.24% in petroleum (Yuliani, 2014).

A method called bioremediation has become an important method to solve the petroleum waste by using indigenous or exogenous microbes (Helmy et al., 2010). The progress on degradation study showed how difficult to degrade PAH because of its stability and recalcitrantly in soil. Less of solubility also limits bioavailability and efficiency of bioremediation process.

In biodegradation process of PAH, it is often mentioned about the role of biosurfactant on biodegradation. Biosurfactant could increase solubility and availability of substrate for bacteria (Li, 2009). Besides

7

Proceeding of International Biology Conference 2016
AIP Conf. Proc. 1854, 020026-1-020026-6; doi: 10.1063/1.4985417
Published by AIP Publishing. 978-0-7354-1528-7/\$30.00

020026-1

biosurfactant, some enzymes also have been linked with biodegradation process of hydrocarbon, especially PAH. The existence of enzyme enable bacterial cell degrade hydrocarbon polymer into simpler compound. So, it can be used as carbon source. Various types of enzyme also related with hydrocarbon uptake mechanism of bacteria.

Bacillus subtilis 3KP was bacteria which known as hydrocarbon degrading bacteria. These bacteria are also able to produce biosurfactant. According to Ni'matuzahroh et al. (2013), *Bacillus subtilis* 3KP was able to produce biosurfactant in the amount of 12.3 g/L using molasses substrate. *Bacillus subtilis* 3KP was potential to degrade PAH compound, but the research has not ever done yet. This research aims to observe the bacterial growth, biodegradation percentage of PAH, and mechanism of biodegradation that was probably developed by *Bacillus subtilis* 3KP.

19 MATERIALS AND METHODS

Chemicals

All chemicals were analytical grade or better. Naphthalene (99%), phenanthrene (99%), 1-hydroxy-naphtioic acid (99%), salicylic acid (99%), and pyrocatechoid acid (99%) were purchased from Sigma-Aldrich. Stock solution of 1-hydroxy-naphtioic acid, salicylic acid, and pyrocatechoid acid were prepared in ethyl acetate (MERCK). Naphthalene and phenanthrene were prepared in mineral salts medium (MS, see below) and sterilized by autoclaving before the bacteria were inoculated.

Microorganism and Growth Medium

The following bacteria used are *Bacillus subtilis* 3KP which isolated from petroleum-contaminated environment in Donan River, Cilacap, Indonesia. *Bacillus subtilis* 3KP was grown in Nutrient Broth and Nutrient Agar before inoculation.

All media used in these studies were prepared in distilled water. The MS medium from the method by Pruthi and Cameotra (1997) contained (L⁻¹) 3 g (NH₄)₂SO₄, 0.2 MgSO₄·7H₂O, 10 g NaCl, 0.01 CaCl₂, 0.001 g MnSO₄·H₂O, 0.001 g H₃BO₃, 0.001 g ZnSO₄·7H₂O, 0.001 g CuSO₄·5H₂O, 0.005 g CoCl₂·6H₂O, 0.001 g Na₂MoO₄·2H₂O, and 1% yeast extract. The buffer composition contained (L⁻¹) K₂HPO₄ and KH₂PO₄ for respectively 2.6207 g and 5 g in 50 ml. It was also added 0.0006 g FeSO₄·7H₂O which sterilized separately in 50 ml distilled water. The pH of the final medium was adjusted to 7.0. For preparing inoculation, bacterial cultures were suspended in 0.85% salt solution.

Determination of Residual Substrat

Biodegradation tests were performed in 250 ml flask and run in duplicate. *Bacillus subtilis* 3KP (5%) (v/v) was grown in 30 ml MS medium. Naphthalene and phenanthrene were added to the medium 200 ppm respectively. The flasks were shaken on a rotary shaker with 90 rpm at 30°C for 7 days. All cultures were started with cell density OD_{610 nm} of 0.5. Growth profile was observed by common serial dilution. The residual hydrocarbons and the bacterial growth were investigated at days 0, 1, 3, 5, and 7.

At the same time of interval, analysis of hydrocarbon residue in culture was observed by UV-vis spectrophotometer (Kumar et al., 2010). Before analysis, the culture was extracted by adding ethyl acetate to the separating funnel in ratio 1:1. Then it was shaken twice for 15 minute in each shaking. Separating of liquid phase, emulsion, and solvent phase was done by adding absolute ethanol. The data was shown as absorbance value at optimal wavelength 276 nm for naphthalene, 293 nm for phenanthrene, 339 nm for 1-hydroxy-2-naphthoic acid, 304 nm for salicylic acid, and 278 nm for pyrocatechol. Standard curve was used to determine concentration of naphthalene or phenanthrene in the culture. Hydrocarbons biodegradation percentage was calculated with the formula:

$$\text{Biodegradation (\%)} = \left(\frac{\text{initial concentration of hydrocarbon} - \text{final concentration of hydrocarbon}}{\text{initial concentration of hydrocarbon}} \right) \times 100\%$$

2 Cell surface Hydrophobicity

According to Rosenberg et al. (1980), the percentage of the cells adhering to an oily phase was investigated by using a modification method. The bacterial cells were harvested each 48 hours by centrifugation (3000 rpm, 15 minute). The cell pellets were washed twice and suspended in phosphate buffer (g L⁻¹; 7.3 KH₂PO₄ and 16.9 K₂HPO₄), then diluted to an optical density (OD) of 0.5 at 610 nm. Into the cell suspension (4 mL) in test tube,

100 μ L of hexadecane was added and vortex-shaken for 3 minute. After forming the suspension, it was allowed to separate for 60 minute, the absorbance OD₆₁₀ in the aqueous phase was measured directly. The clearing measurement of the aqueous phase indicated the capacity of adherence cell to the oily phase. The result is presented as percentage with formula:

$$H (\%) = \left(1 - \frac{A}{A_0}\right) \times 100\%$$

H was hydrophobicity; A was cell suspension absorbance after added by hexadecane; A₀ was initial cell suspension absorbance.

Surface Tension Reduction

All experiments were performed using two separately grown bacterial cultures. The cultures were centrifuged at 3000 rpm for 15 minute. Then, the surface tension of supernatant was measured by Du-Nouy tensiometer. The result is presented as mN m⁻¹ and reported as an average of triplicate each sample.

Emulsification Activity Assay

Emulsification activity was measured using the method by Cooper and Goldenberg (1987). The supernatant (1 mL) as yield of centrifugation was added to a test tube containing 1 ml kerosene. The mixture was homogenized by vortexing for 2 minute and the emulsion was allowed to stand for 60 minute to be measured. Emulsification index was determined as ratio of the emulsion height to total height of the mixture, then it was calculated their percentage.

RESULTS AND DISCUSSIONS

Biodegradation process of PAH is affected by several factors, those are microbial character, environmental factor that support the growth, and characteristic of substrate. Microbial ability in degradation of hydrocarbons depends on its adaptation, toxic resistance, and gene expression of the microbe (Abbaszhad et al., 2011). Microbial ability in degrade hydrocarbon could be known through the growth, the use of hydrocarbon as sole carbon, and the secondary metabolites that produced by microbe.

Bacillus subtilis 3KP grew up and showed positive responses to naphthalene and phenanthrene, 200 ppm respectively. *Bacillus subtilis* 3KP was able to adapt and use PAH as a sole carbon (Fig.1). This is proved by decreasing of concentration of substrate accompanied by increasing of the cell amount. The growth pattern of *Bacillus subtilis* 3KP on naphthalene and phenanthrene showed a biphasic pattern. In compare to control study that showed a normal phase, this kind of pattern indicated that *Bacillus subtilis* 3KP has adapted to the content of PAH in the medium.




FIGURE 1. (a) *Bacillus subtilis* 3KP growth response in the naphthalene and phenanthrene substrates, (b) Percentage of naphthalene and phenanthrene biodegradation at several days

Stationary phase occurs when cell growth was inhibited by cell density, accumulation of secondary metabolites that could be toxic and limited oxygen (Pelczar, 1986). In this study, accumulation of intermediate compound was found in the end of exponential phase, those are 1-hydroxy-2-naphthoic acid and salicylic acid. According to metabolic pathway of phenanthrene degradation by bacteria, 1-hydroxy-2-naphthoic acid would be transformed to 1-2-dihydroxynaphthalene before it became salicylic acid. Pumphrey and Madsen (2007) have explained that 1-2-dihydroxynaphthalene could be oxidized to a toxic 1-2-naphthaquinone which inhibited the growth of *Polaromonas naphthalenivorans*. The intermediate compounds of naphthalene and phenanthrene metabolism can be accumulated in culture because of their easy-soluble character and more toxic while they have increased continuously.

Some microorganisms are able to hold out from naphthalene and phenanthrene and to transform them to enter the TCA (tricarboxylic acid) cycle, *Bacillus subtilis* 3KP did it. At day 7, *Bacillus subtilis* 3KP has degraded 70.5% naphthalene and 24.8% phenanthrene. Intermediate compounds have been observed in the study, those are 1-hydroxy-2-naphthoic acid and salicylic acid. Pyrocatechol was used along with the use of main substrate. It showed that *Bacillus subtilis* 3KP has degraded naphthalene and phenanthrene in a same metabolic pathway. Data existence of intermediate was shown in table 1.

TABLE 1. The existence of intermediate compounds during the process of degradation of naphthalene and phenanthrene by *Bacillus subtilis* 3KP

Days	Existence of intermediate compounds					
	1-hydroxy-2-naphthoic acid		Salicylic acid		Pyrocatechol	
	Naphthalene	Phenanthrene	Naphthalene	Phenanthrene	Naphthalene	Phenanthrene
0	+	-	+	-	+	+
1	+	-	+	-	+	+
3	+	-	+	-	+	+
5	+	-	+	-	+	+
7	+	-	+	+	+	+

Biodegradation process of naphthalene and phenanthrene depends on the ability of microorganism to produce enzymes. It was assumed that there was an important role of dioxygenase, hydrogenase, and monooxygenase in biodegradation process of naphthalene and phenanthrene by *Bacillus subtilis* 3KP. Enzyme regulation in naphthalene and phenanthrene is affected by existence of gen. They were nah and phn which take the role in biodegradation of naphthalene and phenanthrene (Laurie and Jones, 1999). For this research, the existence of catabolic gen in *Bacillus subtilis* 3KP should be investigated as advanced research to affirm its ability of PAH degradation.

Moreover, hydrocarbon properties also affect biodegradation process. Complexity of structure, molecular weight, and solubility of the hydrocarbon become a limiting factor microbe's access to hydrocarbons. *Bacillus subtilis* 3KP was released biosurfactant to contact with the hydrocarbons. There was a similar growth pattern and uptake mechanism of substrate by *Bacillus subtilis* 3KP that indicated in Figure 2.




FIGURE 2. Mechanism of naphthalene (a) and phenanthrene (b) biodegradation by *Bacillus subtilis* 3KP

At the early stationary phase, emulsification activity and adherence activity increased, while surface tension reduced. Extracellular biosurfactant of *Bacillus subtilis* 3KP is able to reduce the surface tension of culture supernatant at the early stationary phase. It was in accordance to Cooper and Goldenberg (1987), the lysis of *Bacillus cereus* in sucrose was increasing the biosurfactant activity as a surface active agent because of its polysaccharide (emulsifier) and lipid reduced the surface tension. So that, at the early of stationary phase, emulsification activity increased together with the reducing of surface tension. It was assumed that *Bacillus subtilis* 3KP was secreted polypeptide-type biosurfactant.

Biosurfactant consists of surfactant monomer that would form stable micelle from about 10 – 200 molecules (Makkar and Rockne, 2003). Scippers et al. (2000) explained about the role of biosurfactant in biodegradation of PAH. The first role, micelle was used as a PAH porter to microbial cell wall surface, then PAH was entered diffusely the cell. The second role, micelle could increase PAH solubility in the medium. The last role, without micelle, surfactant still could facilitate the use of PAH by a cell-surfactant-PAH contact directly (bacterial adherence to hydrocarbons).

In pH 6 – 8, *Bacillus subtilis* 3KP are able to grow. Emulsification activity and surface tension reduction were not affected by increasing in pH. Therefore, at day 7 when the value of pH showed 8.0, emulsification activity was still rising. While in surface tension, there was not a significant difference.

Biosurfactant takes role in biodegradation process in accessing not soluble substrate. It was showed from the result that emulsification activity and surface tension reduction in culture with naphthalene was higher than with phenanthrene. However, the adherence activity was decreased after the end of exponential for culture with naphthalene. While in culture with phenanthrene, adherence activity has increased much higher at day 7. It was assumed that in culture with more complex structure, cells were adapted to transform its cell wall to be more hydrophobic. Prabhu and Phale (2003) have learned about uptake mechanism of aliphatic and aromatic hydrocarbons by *Pseudomonas* sp. strain PP2. Cell hydrophobicity raised after the decreased of emulsification activity in stationary phase. Level of cell hydrophobicity was higher in aromatic which is a lower soluble compound than aliphatic one.

At the same relatively environmental conditions and with the same type of microbe, phenanthrene degradation was longer than the naphthalene degradation. It proves that the type of substrate affects the biodegradation process. Although naphthalene and phenanthrene are classified as a some group of polycyclic aromatic hydrocarbons, naphthalene has much higher solubility than phenanthrene. Low substrate solubility in the medium can inhibit the bacterial access to the substrate. Limited of available substrate is the most important factor that involved in the slow degradation of PAHs (Makkar and Rockne, 2003).

15 ACKNOWLEDGMENTS

The authors thank to the Direktorat Riset dan Pengabdian Masyarakat (DRPM) KemenristekDikti, Indonesia for funding and to Universitas Airlangga for the support of the facility during the research.

REFERENCES

1. Abbasnezhad, H., Gray, M., Foght J. M., Influence of Adhesion on Aerobic Biodegradation and Bioremediation of Liquid Hydrocarbons. *Appl Microbiol Biotechnol* **92**, 653 – 675 (2011).
2. Connel, D. W., G. J. Miller, *Kimia dan Ekotoksikologi Pencemaran* (UI Press, Jakarta, 1995).
3. Cooper, David G., Goldenberg, Beena G., Surface-Active Agents from Two *Bacillus* Species. *American Society for Microbiology* **53**, 2 (1987).
4. Helmy, Q., E. Kardena, Z. Nurachman, Wisjnuprpto, Application of Biosurfactant Produced by *Azotobacter vinelandii* AV01 for Enhanced Oil Recovery and Biodegradation of Oil Sludge. *International Journal of Civil & Environmental Engineering IJCEE* **10**, 1 (2010)
5. Kumar, G., Singla R., Kumar R., Plasmid Associated Anthracene Degradation by *Pseudomonas* sp. Isolated from Filling Station Site. *Nature and Science* **8**, 89-94 (2010).
6. Laurie, A.D., Lloyd-Jones, G., The phn Genes of *Burkholderia* sp. strain RP007 Constitute a Divergent Gene Cluster for Polycyclic Aromatic Hydrocarbon Catabolism. *J Bacteriol* **181**, 2, 531 – 40 (1999).
7. Li, Liang-Jing, and Chen, Hung-Bing, Surfactant-Mediated Biodegradation of Polycyclic Aromatic Hydrocarbons. *Material* **2**, 76 – 94 (2009).
8. Makkar, S. Randdhir, Rockne, J. Karl., Comparizon of Syntetic Surfactant and Biosurfactant in Enhancing Biodegradation of Polycyclic Aromatic Hydrocarbons. *Environmental Toxycology and Chemistry* **22**, 10, 2280 – 2292 (2003).

9. Ni'matuzahroh, Fatimah, Sri Sumarsih, "Biocleanoil Produk Berbasis Mikroba untuk Pengolahan Sludge Industri Minyak". Laporan Akhir Penelitian Unggulan Perguruan Tinggi, Universitas Airlangga Surabaya, 2013.
10. Patnaik, Pradyot, *A Comprehensive Guide to The Hazardous Properties of Chemical Substances, Second Edition* (Canada, 1999).
11. Pelczar, J. Michel, *Dasar-Dasar Mikrobiologi* (Universitas Indonesia Press, Jakarta, 1986).
12. Prabhu Y., Phale S., Biodegradation of Phenanthrene by *Pseudomonas* sp. strain PP2: Novel Metabolic Pathway, Role of Biosurfactant and Cell Hydrophobicity in Hydrocarbon Assimilation. *Appl Microbiol Biotechnol* **61**, 342–351 (2003).
13. Pruthi, V. dan Cameotra, S.S., Rapid Identification of Biosurfactant Producing Bacterial Strains Using A Cell Surface Hydrophobicity Technique. *Biotechnology Techniques* **11**, 9, 671 – 674 (1997).
14. Pumphrey, Graham M., Madsen, Eugene L., Naphthalene metabolism and growth inhibition by naphthalene in *Polaromonas naphthalenivorans* strain CJ2. *Microbiology* **153**, 3730–3738 (2007).
15. Rosenberg, M., Gutnick, D., Rosenberg, E., Adherence of Bacteria to Hydrocarbons: A Simple Method for Measuring Cell-Surface Hydrophobicity. *FEMS Microbiol Lett* **9**, 29 –33 (1980).
16. Schippers, C., Gessner, K., Mueller, T., Scheper, T., Microbial Degradation of Phenanthrene by Addition of A Sphingolipid Mixture. *J Biotechnol* **83**, 189 – 198 (2000).
17. Yuliani, Hanif, "Biodegradasi Poliaromatik Hidrokarbon Pirena oleh Bakteri yang Diisolasi dari Laut Cilacap dan Laut Marina", Ringkasan Disertasi, Fakultas Teknik Universitas Indonesia Depok, 2014.

Biodegradation of Naphthalene and Phenanthren by Bacillus subtilis 3KP

ORIGINALITY REPORT

18%

SIMILARITY INDEX

12%

INTERNET SOURCES

15%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Almando Gernaldi, Ni'matuzahroh, Fatimah, Chang-Hao Cui, Thi Thuy Nguyen, Sun Chang Kim. "Enzymatic biotransformation of ginsenoside Rb1 by recombinant β -glucosidase of bacterial isolates from Indonesia", *Biocatalysis and Agricultural Biotechnology*, 2020
Publication 2%
 - 2 link.springer.com
Internet Source 1%
 - 3 fr.readkong.com
Internet Source 1%
 - 4 Mariathason Birdilla Selva Donio, Subbiahanadar Chelladurai Karthikeyan, Mariavincent Michaelbabu, Ganapathi Uma et al. " sp. JS1, a bioemulsifier producing halophilic bacterium isolated from Indian solar salt works ", *Journal of Basic Microbiology*, 2018
Publication 1%
-

5	Submitted to Universiti Brunei Darussalam Student Paper	1 %
6	setac.onlinelibrary.wiley.com Internet Source	1 %
7	Agus Triawan, Ni'matuzahroh, Agus Supriyanto. "Effects of the combination between bio-surfactant product types and washing times on the removal of crude oil in nonwoven fabric", AIP Publishing, 2017 Publication	1 %
8	Khusnul Ain, Akif Rahmatillah, Virgilius Rivan Seran. "Design of automatic water level measuring instrument IoT based on open pan evaporimeter", AIP Publishing, 2020 Publication	1 %
9	sevgiligiyim.com Internet Source	1 %
10	Xue-Qin Tao, Gui-Ning Lu, Zhi Dang, Xiao-Yun Yi, Chen Yang. "Isolation of phenanthrene-degrading bacteria and characterization of phenanthrene metabolites", World Journal of Microbiology and Biotechnology, 2006 Publication	1 %
11	ppjpi.unair.ac.id Internet Source	1 %
12	aip.scitation.org Internet Source	1 %

1 %

13

Submitted to Universitas Airlangga

Student Paper

1 %

14

ejbiotechnology.info

Internet Source

1 %

15

e-hir.org

Internet Source

<1 %

16

Asmaa M. M. Mawad, Abd El-Latif Hesham, Yasser M. Mostafa, Ahmed Shoriet. "Pyrene degrading *Achromobacter denitrificans* ASU-035: growth rate, enzymes activity, and cell surface properties", *Rendiconti Lincei*, 2016

Publication

<1 %

17

thesis.library.caltech.edu

Internet Source

<1 %

18

A. Hanani, S. Sitaresmi, Y. Yasman, A. Oetari. "Naphthalene biodegradation capability and characterization of bacteria isolate DRK 9.1 from volcanic mud in Renokenongo Village, Sidoarjo", AIP Publishing, 2020

Publication

<1 %

19

Xiaoyu Qi, Xiaoming Xu, Chuanqing Zhong, Tianyi Jiang, Wei Wei, Xin Song. "Removal of Cadmium and Lead from Contaminated Soils Using Sophorolipids from Fermentation

<1 %

Culture of *Starmarella bombicola* CGMCC
1576 Fermentation", International Journal of
Environmental Research and Public Health,
2018

Publication

20

arcabc.ca

Internet Source

<1 %

21

"Posters", FEMS Microbiology Letters, 2003

Publication

<1 %

22

Environmental Science and Engineering, 2012.

Publication

<1 %

23

Hanif Yuliani, Meka Saima Perdani, Imelda
Savitri, Meilani Manurung, Muhamad Sahlan,
Anondho Wijanarko, Heri Hermansyah.

"Antimicrobial activity of biosurfactant derived
from *Bacillus subtilis* C19", Energy Procedia,
2018

Publication

<1 %

24

academic.oup.com

Internet Source

<1 %

25

digilib.uinsgd.ac.id

Internet Source

<1 %

26

repositorio.unicamp.br

Internet Source

<1 %

27

Ankit Mishra, Aravind Krishnamoorthy, Pankaj
Rajak, Subodh Tiwari et al. " Free energy of

<1 %

hydration and heat capacity of calcium
dipicolinate in spore cores ", Applied Physics
Letters, 2018

Publication

28

Chongyang Wang, Yong Huang, Zuotao
Zhang, Hui Wang. "Salinity effect on the
metabolic pathway and microbial function in
phenanthrene degradation by a halophilic
consortium", AMB Express, 2018

Publication

<1 %

29

Deepak Sharma, Mohammad Javed Ansari,
Ahmad Al-Ghamdi, Nuru Adgaba, Khalid Ali
Khan, Vikas Pruthi, Noori Al-Waili.
"Biosurfactant production by Pseudomonas
aeruginosa DSVP20 isolated from petroleum
hydrocarbon-contaminated soil and its
physicochemical characterization",
Environmental Science and Pollution
Research, 2015

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Biodegradation of Naphthalene and Phenanthren by Bacillus subtilis 3KP

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
