


Volume 2002

 **Conference collection**

# Inventing Prosperous Future through Biological Research and Tropical Biodiversity Management Proceedings of the 5th International Conference on Biological Science



**Yogyakarta, Indonesia**

15-16 September 2017

**Editors**

Tuty Arisuryanti, Maryani, Zuliyati Rohmah, Lisna Hidayati and Ganies Riza Aristya

**AIP** | Conference Proceedings

[proceedings.aip.org](http://proceedings.aip.org)



AIP Conference Proceedings  
**The 18th International Conference  
 on Positron Annihilation**

ORDER PRINT EDITION

# AIP

Conference Proceedings

HOME

BROWSE

MORE ▼

To support global research during the COVID-19 pandemic, [AIP Publishing is making our content freely available](#) to scientists who register on Scitation.

To gain access, please [log in](#) or [create an account](#) and then [click here](#) to activate your free access. You must be logged in to Scitation to activate your free access.

## Table of Contents

### INVENTING PROSPEROUS FUTURE THROUGH BIOLOGICAL RESEARCH AND TROPICAL BIODIVERSITY MANAGEMENT: Proceedings of the 5th International Conference on Biological Science

< PREV NEXT >



Conference date: 15-16 September 2017

Location: Yogyakarta, Indonesia

ISBN: 978-0-7354-1718-2

Editors: Tuty Arisuryanti, Maryani, Zuliyati Rohmah, Lisna Hidayati and Ganies Riza Aristya


Volume number: 2002

Published: Aug 15, 2018

DISPLAY : 20 50 100 all

BROWSE VOLUMES

## ARTICLES

 Free . August 2018

### **Diversity and phenetic analysis of *tomboreso* (*Dioscorea pentaphylla* L.) in Yogyakarta based on morphological and leaf anatomical characters**


Purnomo, Malikah and Maryani

AIP Conference Proceedings **2002**, 020049 (2018); <https://doi.org/10.1063/1.5050145>

---

SHOW ABSTRACT



 Free . August 2018

### **Iron chelating activity of *Caesalpinia sappan* L. extract on iron status in iron overload rats (*Rattus norvegicus* L.)**


Ratu Safitri, Ani Melani Maskoen, Mas Rizky Anggun Adipurna Syamsunarno, Mohammad Chozali and Ramdan Panigoro

AIP Conference Proceedings **2002**, 020050 (2018); <https://doi.org/10.1063/1.5050146>

---

SHOW ABSTRACT



 Free . August 2018

### **Bioinformatics analysis of the expressed sequence tags from *Rhizophora stylosa* Griff. genomic library**


Ridha Wati, Mohammad Basyuni, Shigeyuki Baba and Hirotsuke Oku

AIP Conference Proceedings **2002**, 020051 (2018); <https://doi.org/10.1063/1.5050147>

---

SHOW ABSTRACT



 Free . August 2018

## **Detection of polymorphisms on strawberry (*Fragaria* spp.) based on cleavage amplified polymorphic sequence**


Rina Kasiamdari, Ganies Riza Aristya, Achmad Syahrian and Wakhidah Kurniawati

AIP Conference Proceedings **2002**, 020052 (2018); <https://doi.org/10.1063/1.5050148>

---

SHOW ABSTRACT



 Free . August 2018

## **Natural polyploidy in amaranths (*Amaranthus* spp.)**


Rita Andini, Muhammad Ikhsan Sulaiman and Ryo Ohsawa

AIP Conference Proceedings **2002**, 020053 (2018); <https://doi.org/10.1063/1.5050149>

---

SHOW ABSTRACT



 Free . August 2018

## **Potential of vascular plants as phytotourism products in Endau Rompin Johor National Park, Malaysia**


Salasiah Mohamad, Maryati Mohamed and Muhammad Shafiq Hamdin

AIP Conference Proceedings **2002**, 020054 (2018); <https://doi.org/10.1063/1.5050150>

---

SHOW ABSTRACT



 Free . August 2018


## **Ciprofloxacin down-regulates *Wnt5a* gene expression in chick embryo leg bud**

AIP Conference Proceedings 2002, 020055 (2018); <https://doi.org/10.1063/1.5050151>

---

SHOW ABSTRACT



 Free . August 2018

**The effect of supplemented probiotic on the laying performance and egg quality of Japanese quail (*Coturnix-coturnix japonica* Temminck and Schlegel, 1849)**


Slamet Widiyanto and Wahyu Indrawan

AIP Conference Proceedings 2002, 020056 (2018); <https://doi.org/10.1063/1.5050152>

---

SHOW ABSTRACT



 Free . August 2018

**Effect of coloring pH and mordant on fungal dyes quality using woolen yarn**


Suciatmih and Yuliar

AIP Conference Proceedings 2002, 020057 (2018); <https://doi.org/10.1063/1.5050153>

---

SHOW ABSTRACT



 Free . August 2018


**Breeding ecology of the rice field rat (*Rattus argentiventer* Rob & Kloss, 1916) in irrigated rice ecosystem in Indonesia**

Sudarmaji and Nur 'Aini Herawati

AIP Conference Proceedings 2002, 020058 (2018); <https://doi.org/10.1063/1.5050154>

SHOW ABSTRACT



 Free . August 2018

**Selection of potential probiotic lactic acid bacteria isolated from palm sap (*Borassus flabellifer* Linn.) origin Kupang, East Nusa Tenggara**


Sulistiani

AIP Conference Proceedings **2002**, 020059 (2018); <https://doi.org/10.1063/1.5050155>

---

SHOW ABSTRACT



 Free . August 2018

**Application biopreservatives produced by lactic acid bacteria (LAB) for preservation boiled-salted (*pindang*) tuna (*Euthynnus affinis* Cantor, 1849)**


Sulistiani and Rini Handayani

AIP Conference Proceedings **2002**, 020060 (2018); <https://doi.org/10.1063/1.5050156>

---

SHOW ABSTRACT



 Free . August 2018

**Pathological alteration of digestive tissues in blue mussel (*Mytilus galloprovincialis*) larvae induced by *Alexandrium fundyense* lysates**


Supono, Christopher Bolch and Graeme Knowles

AIP Conference Proceedings **2002**, 020061 (2018); <https://doi.org/10.1063/1.5050157>

---

SHOW ABSTRACT



 Free . August 2018

## **Effect of sequential extraction on total phenolic content (TPC) and antioxidant activities (AA) of *Luffa acutangula* Linnaeus dried pulps**


Supriatno and Lelifajri

AIP Conference Proceedings **2002**, 020062 (2018); <https://doi.org/10.1063/1.5050158>

---

SHOW ABSTRACT



 Free . August 2018

## **Study of the characteristic of neritidae: Shell and operculum**


Susintowati, Suwarno Hadisusanto, Nyoman Puniawati, Erny Poedjirahajoe and Niken Satuti Nur Handayani

AIP Conference Proceedings **2002**, 020063 (2018); <https://doi.org/10.1063/1.5050159>

---

SHOW ABSTRACT



 Free . August 2018


## **Screening and identifying of cellulolytic bacteria from Alas Purwo National Park**

Syahriar Nur Maulana Malik Ibrahim, Sibyandhita Erhaa Amalia, Bunga Agatha, Mustikayanti Yuni Lestari, Antonius Dwi Saputra, Nur Laily Fachira Ikmala, Fazareyna Hyuwan Argi Ryanto, Puspita Ayu Dewi, Mochamad Khoirul Anam, Farah Aisyah Nafidiastri, Nastiti Trikurniadewi, Ditta Putri Kumalasari, Erta Tri, Gendut Hariyanto, more...

AIP Conference Proceedings **2002**, 020064 (2018); <https://doi.org/10.1063/1.5050160>

SHOW ABSTRACT



 Free . August 2018

## **The contents of glucose and lactic acid in tube paste flour additional amyolytic *Lactobacillus bulgaricus* FNCC 004P and *Streptococcus thermophilus* FNCC 1.9.03**


Tatik Khusniati, Firda Fatimah Al Ansya and Erlinda Gangga

AIP Conference Proceedings **2002**, 020065 (2018); <https://doi.org/10.1063/1.5050161>

---

SHOW ABSTRACT



 Free . August 2018

## **Diversity of crabs in the intertidal zone at Sundak Beach, Gunungkidul, Indonesia**


Titis Sukmaningrum, Basith Kuncoro Adji, Eka Mulia Pratiwi, Bening Larasati, Puspa Restu Sayekti, Ikhsan Maulana and Rury Eprilurahman

AIP Conference Proceedings **2002**, 020066 (2018); <https://doi.org/10.1063/1.5050162>

---

SHOW ABSTRACT



 Free . August 2018

## **The antioxidant and cytotoxic activities of the chloroform extract of agarwood (*Gyrinops versteegii* (Gilg.) Domke) leaves on HeLa cell lines**


Tri Rini Nuringtyas, Rini Isromarina, Yovi Septia, Lisna Hidayati, Nastiti Wijayanti and Sukarti Moeljopawiro

AIP Conference Proceedings **2002**, 020067 (2018); <https://doi.org/10.1063/1.5050163>



SHOW ABSTRACT



 Free . August 2018

**Genetic identification of two mudskipper species (Pisces: Gobiidae) from Bogowonto Lagoon (Yogyakarta, Indonesia) using *COI* mitochondrial gene as a DNA barcoding marker**


Tuty Arisuryanti, Rika Lathif Hasan and Johan Putra Koentjana

AIP Conference Proceedings **2002**, 020068 (2018); <https://doi.org/10.1063/1.5050164>

---

SHOW ABSTRACT



 Free . August 2018

**Karyomorphological study on two local garlic cultivars (*Allium sativum* L.) from central Java, Indonesia**


Tuty Arisuryanti, Zulfa Kurniawati and Johan Putra Koentjana

AIP Conference Proceedings **2002**, 020069 (2018); <https://doi.org/10.1063/1.5050165>

---

SHOW ABSTRACT



 Free . August 2018

**Histologic observation, identification, and secondary metabolites analysis of endophytic fungi isolated from a medicinal plant, *hedychium acuminatum* Roscoe**

Utami Sri Hastuti, Putri Moortiyani Al Asna, and Dwi Rahmawati


AIP Conference Proceedings **2002**, 020070 (2018); <https://doi.org/10.1063/1.5050166>

---

SHOW ABSTRACT



BROWSE VOLUMES

 Free . August 2018

## **Length-Weight relationship, sex ratio and condition factor of blue swimming crab (*Portunus pelagicus* Linnaeus, 1758) from Java Sea Indonesia**


Vella Rohmayani, Muhammad Reza Pahlevi, Bambang Irawan and Agoes Soegianto

AIP Conference Proceedings **2002**, 020071 (2018); <https://doi.org/10.1063/1.5050167>

---

SHOW ABSTRACT



 Free . August 2018

## **Influence of bacterial consortium for copper biosorption and accumulation**

Wahyu Irawati, Nommensen Pangihutan Ompusunggu and Triwibowo Yuwono

AIP Conference Proceedings **2002**, 020072 (2018); <https://doi.org/10.1063/1.5050168>

---

SHOW ABSTRACT





## Academic paper editing

Get Professional academic paper editing by Subject Area Native English Editors Today!

[authorassists.com](http://authorassists.com)

OPEN

# AIP Conference Proceedings

**Country** [United States](#) - [SIR Ranking of United States](#)

**Subject Area and Category** [Physics and Astronomy](#)  
[Physics and Astronomy \(miscellaneous\)](#)

**Publisher** [American Institute of Physics](#)

**Publication type** Conferences and Proceedings

**ISSN** 0094243X, 15517616

**Coverage** 1974-1978, 1983-1984, 1993, 2000-2001, 2003-2020

**Scope** Today, AIP Conference Proceedings contain over 100,000 articles published in 1700+ proceedings and is growing by 100 volumes every year. This substantial body of scientific literature is testament to our 40-year history as a world-class publishing partner, recognized internationally and trusted by conference organizers worldwide. Whether you are planning a small specialist workshop or organizing the largest international conference, contact us, or read these testimonials, to find out why so many organizers publish with AIP Conference Proceedings.



[Homepage](#)

[How to publish in this journal](#)

[Contact](#)



[Join the conversation about this journal](#)

# 64








H Index

## Free Grammar Checker

Eliminate grammar errors instantly and enhance your writing with Grammarly

Grammarly

DOWNLOAD



**AIP Conference Proceedings** ← Show this widget in your own website

Not yet assigned quartile

**SJR 2019**  
**0.19**

powered by scimagojr.com

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com" style="border: 1px solid #ccc; padding: 2px 5px; display: inline-block;">https://www.scimagojr.com
```

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/327056010>

# Screening and identifying of cellulolytic bacteria from Alas Purwo National Park

Conference Paper in AIP Conference Proceedings · August 2018

DOI: 10.1063/1.5050160

CITATIONS

0

READS

76

17 authors, including:



**Syahrir Nur Maulana Malik Ibrahim**

Chulalongkorn University

4 PUBLICATIONS 3 CITATIONS

SEE PROFILE



**Sibyandhita Erhaa Amalia**

Airlangga University

1 PUBLICATION 0 CITATIONS

SEE PROFILE



**Antonius Dwi Saputra**

Universitas Jember

1 PUBLICATION 0 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Screening of Cellulolytic Bacteria From Alas Purwo National Park to Produce Cellulase [View project](#)



Utilization of Organic Waste for Biosurfactant Production Substrate by Hydrocarbonoclastic Bacteria [View project](#)

## Screening and identifying of cellulolytic bacteria from Alas Purwo National Park

Syahriar Nur Maulana Malik Ibrahim, Sibyandhita Erhaa Amalia, Bunga Agatha, Mustikayanti Yuni Lestari, Antonius Dwi Saputra, Nur Laily Fachira Ikmala, Fazareyna Hyuwan Argi Ryanto, Puspita Ayu Dewi, Mochamad Khoirul Anam, Farah Aisyah Nafidiasri, Nastiti Trikurniadewi, Ditta Putri Kumalasari, Erta Tri, Gendut Hariyanto, Suwarni, Agus Supriyanto, and Ni'matuzahroh

Citation: *AIP Conference Proceedings* **2002**, 020064 (2018); doi: 10.1063/1.5050160

View online: <https://doi.org/10.1063/1.5050160>

View Table of Contents: <http://aip.scitation.org/toc/apc/2002/1>

Published by the *American Institute of Physics*

---

---

**AIP** | Conference Proceedings

Get **30% off** all  
print proceedings!

Enter Promotion Code **PDF30** at checkout



# Screening and Identifying of Cellulolytic Bacteria from Alas Purwo National Park

Syahriar Nur Maulana Malik Ibrahim<sup>1, a)</sup>, Sibyandhita Erhaa Amalia<sup>1</sup>, Bunga Agatha<sup>1</sup>, Mustikayanti Yuni Lestari<sup>1</sup>, Antonius Dwi Saputra<sup>2</sup>, Nur Laily Fachira Ikmala<sup>1</sup>, Fazareyna Hyuwan Argi Ryanto<sup>1</sup>, Puspita Ayu Dewi<sup>1</sup>, Mochamad Khoirul Anam<sup>1</sup>, Farah Aisyah Nafidiastri<sup>1</sup>, Nastiti Trikurniadewi<sup>1</sup>, Ditta Putri Kumalasari<sup>1</sup>, Erta Tri<sup>1</sup>, Gendut Hariyanto<sup>3</sup>, Suwarni<sup>1</sup>, Agus Supriyanto<sup>1</sup>, and Ni'matuzahroh<sup>1</sup>

<sup>1</sup>Biology Department, Faculty of Science and Technology, Airlangga University, Surabaya, Indonesia.

<sup>2</sup>Faculty of Medicine, University of Jember, Jember, Indonesia.

<sup>3</sup>Alas Purwo National Park, Banyuwangi, Indonesia.

<sup>a)</sup>Corresponding author: r261196@gmail.com

**Abstract.** This research aimed to screen and identifies cellulolytic bacteria from Alas Purwo National Park. Soil samples were taken from five points in each location by indirect sampling method. Bacteria were isolated with CMC (Carboxymethylcellulose) agar. The result had successfully obtained 16 isolates which produced cellulase. The cellulase activity index assay of bacteria used CMC agar modified with adds 1 % YME (Yeast Malt Extract) and then was incubated during one day at room temperature. Bacterial colonies were soaked with Congo red 1 % and rinsed with NaOH 10 % to visualize the halo zone formed. Three isolates with highest cellulase activity index were identified by observing the colony morphology, cell morphology, and the biochemical characters. The results of this research showed that the three highest potential isolates producing cellulase were BPA-B, BSA-D, and BPA-A with cellulase activity indexes respectively 10.67, 9.86, and 8.16. BPA-B, BSA-D, and BPA-A each have the similarity with genera *Pimelobacter*, *Micrococcus*, and *Cellulomonas*.

**Keywords:** Alas Purwo National Park, cellulase activity index, cellulolytic bacteria

## INTRODUCTION

Alas Purwo National Park is geographically located at the eastern tip of Java Island South coast between 8.44°44'S and 114.22°60'E with an area of 44 037.30 ha. Based on the type of ecosystem, the area of Alas Purwo National Park is grouped into bamboo forests, coastal forests, mangrove forests, plantations, natural forests, and feeding ground [1]. The soil type in Alas Purwo National Park consists of four groups namely, red lithosol mediteran land complex, gray regosol soil, grumusol gray soil, and dominated by alluvial soil hidromorf. Leaves that fall on the ground cause high cellulose content [2]. Cellulose content on soil is used as a carbon source.

Almost half of the land on earth contains carbon components of cellulose. Cellulose is one of the carbon components of polysaccharides [3]. All carbon in the form of polysaccharides is mineralized with enzymes from microorganisms. One of the enzymes essential for the carbon cycle process is the polysaccharide hydrolysis enzyme. The enzyme that hydrolyzes the polysaccharide in the form of cellulose is cellulase. Cellulase can be found in soil microorganisms, one of them was cellulolytic bacteria. Cellulolytic bacteria are bacteria that have the ability to convert cellulose into oligosaccharides using cellulase enzyme [4]. This enzyme is used as a production material in the industry. Cellulase from bacteria is an extracellular enzyme consisting of the endo- $\beta$ -1,4-glucanase complex, exo- $\beta$ -

1,4-glucanase, and  $\beta$ -1,4-glucosidase [5–7]. It can have good prospects for fuel from lignocellulose by combining cellulase and lignocellulase systems. In addition to the prospect being used as a fuel, cellulase enzymes can increase the production of bioethanol or biofuel [9].

Screening and identifying cellulolytic bacteria need to be done because of the lack of indigenous cellulolytic bacteria data in Indonesia. This study aims to screen cellulolytic bacteria from Alas Purwo National Park that has the potential to produce cellulase. The result of this research is used to increase the isolate database of cellulase enzyme producer.

## MATERIALS AND METHODS

### Soil Sampling

The sampling process took place in Feb 2017. Soil sampling was conducted on four locations in Alas Purwo National Park. Selection was based on vegetation homogeneity and physicochemical parameter test. Air humidity was measured using sling psychrometer (Bacharach), ground temperature was measured using a thermometer (Yenaco), light intensity using lux meter (IEEE, LX1010B), pH, and soil moisture using soil tester (Takemura DM-13). Soil sampling was carried out randomly from center point by five samples in each location using handheld soil sampler (LaMotte 1055/EP). Soil sampler was inserted 10 cm deep from the ground surface of leaf litter or grass. Five soil sample of each location is composed and stored in sterile plastic. Then, the sample is stored in a closed container and taken to the laboratory for two days.

### Isolation and Screening of Cellulolytic Bacteria

A total of 25 g of composite soil from each location were homogenized with 225 mL of aquadest in Erlenmeyer (Pyrex). Then, the soil is agitated using a shaker (GFL MbH D-30938, Burgwedel) at a rate of 90 rpm for 15 min. After that, the supernatant was taken and macro dilution until  $10^{-7}$ . Furthermore, bacterial was isolated using pour plate technique with pure CMC media; Carboxymethylcellulose,  $10\,000\text{ mg} \cdot \text{L}^{-1}$  (PT Brataco) and agar powder  $20\,000\text{ g} \cdot \text{L}^{-1}$ . The culture was incubated for 24 h at  $27\text{ }^{\circ}\text{C}$  in incubator (Heraeus Instrument). Prior to the screening, the grow cultures were stocked in NA (Nutrient Agar (Oxoid)). Screening is done by soaking bacterial colonies and media with Congo red (XCWY) 1 % for 10 min. Then the immersion solution was replaced with 10 % NaOH (Oxoid) and allowed to stand for 15 min. After submersion, the solution is removed, and the halo zone will appear. The visible halo zone is measured and calculated by the formula [10]:

$$\text{Relative enzyme activity index (ICMC)} = \frac{\text{Halo zone diameter}}{\text{Colony zone diameter}}$$

### Identification of Bacteria

Bacterial isolates grown on CMC media were inoculated on NA medium at room temperature. The bacteria in the NA medium were used as insulating stock. Then, the bacteria growing on the NA medium were inoculated with a stab at the CMC medium for modification with the composition (Carboxymethylcellulose 1 % and Yeast Malt Extract 1 %). Three screening bacteria with potentially high cellulase production were inoculated on NA media. Identification of bacteria was based on macroscopic characters of colony, microscopic characters of cells, and also biochemical characterization. Macroscopic characters based on characteristic shape, size, color, texture, edges, and elevation of colonies. Microscopic characters were observed based on Gram characteristics with Gram staining, endospores staining with green Malachite dyes, and cell shape. Biochemical characterization of bacteria used ID Kit-12 A/B, catalase test, starch test, motility test, oxidative test, and  $37\text{ }^{\circ}\text{C}$  temperature. Yeast malt extract, Gram kit staining, green Malachite, and ID Kit-12 A/B were purchased from Oxoid. The results of identification are matched using Bergey's Determinative of Bacteria 9<sup>th</sup> edition.



## RESULTS AND DISCUSSIONS

Each location of Alas Purwo National Park has vegetation dominated by different plants. Savanna vegetation is dominated by *Cyperus rotundus*, Sadengan bamboo forest is dominated by *Bambusa jacobsii*, Pancur bamboo forest is dominated by *Bambusa arundinaceae*, and tropical rainforest is dominated by *Switenia macrophylla*. The physicochemical parameter data of location is shown in Table 1.

Vegetation dominance indicates the homogeneity of the soils [11]. The cellulose nutrients in the soil for cellulolytic bacteria from four locations are litter or dead plant. The soil moisture is important for bacterial growth. Different type of soil will make a different bacterial community [12]. Therefore, the physicochemical of soil and air measurement are needed.

Related to the physicochemical parameter from each location, the cellulolytic bacterial isolated was classified as mesophilic bacteria because of the temperatures from each location have rates range from 28.4 to 31.6 °C. The mesophilic bacteria temperature is below 50 °C [13].

**TABLE 1.** Vegetation dominant and physicochemical parameters of location in Alas Purwo National Park.

Locations (code)	Dominant Vegetation	Rate of Soils pH	Rate of Soils Humidity (RH %)	Rate of Soils Temperature (°C)	Rate of Airs Humidity (RH %)	Rate of Light Intensity (Lux)
Savanna (SAV)	<i>Cyperus rotundus</i>	6.3 ± 0.48	80	31.6 ± 1.94	70	3 000 ± 0
Sadengan bamboo forest (BSA)	<i>Bambusa jacobsii</i>	4.75 ± 0.46	80	29.2 ± 0.83	80	459 ± 104.84
Pancur bamboo forest (BPA)	<i>Bambusa arundinaceae</i>	5.45 ± 0.44	80	27.6 ± 1.34	91	868.4 ± 390.63
Tropical rain forest (HHT)	<i>Switenia macrophylla</i>	5,4 ± 0.45	80	28.4 ± 0.89	72	620 ± 76.48

The result of isolation of cellulolytic bacteria from the fourth soil samples of vegetation obtained sixteenth different isolates that potential to produce cellulase. The potency of cellulolytic bacteria is based on the result of the relative enzyme activity index ( $I_{cmc}$ ). From the sixteenth isolates, it was selected to be three isolates with the highest  $I_{cmc}$  value, shown in Table 2. The control used CMC media without a bacterial culture to indicate there is no enzyme activity. The halo zone forming because of the activity of cellulase to hydrolysis cellulose become glucose [14].

The three isolates selected were inoculated in NA media with incubation for one day. Visible form of macroscopic and microscopic characteristics of three bacteria is indicated in Table 3. Macroscopic characters observed are shape, size, color, texture, edges, and elevation. Macroscopic characters of bacterial colony are dominated by irregular forms. Obtained cellulolytic bacteria have two macroscopic colonies of large size, round, white bones, irregular edges and small, round, clear, irregular edges [15].

The microscopic characters of bacteria include Gram staining, endospore staining, and cell shape. The results of Gram staining are predominantly found Gram-positive with irregular rod-coccus form. The endospore stain has obtained in all isolate were not contain endospore (negative endospore).

Furthermore, the three isolates were performed to the physiological testing. Physiological test with ID 12A/B Kit shown in Table 4. Based on the test, three isolates can grow at 37 °C, catalase positive, motile, and oxidative hydrolysis. The BSA-D strain hydrolyzes ONPG, uses citrate, dilutes gelatin and inhibits malonate. The BPA-A strain performs fermentation of glucose, mannitol, sorbitol, rhamnose, sucrose, and arginine dihydrolase. The BPA-B strain can decarboxylate lysine, use citrate, dilute gelatin, and produce arginine dihydrolase. The results of macroscopic, microscopic, and physiological identification were matched on Bergey's Manual of Determinative Bacteriology 9<sup>th</sup> Edition.

Observed cellulolytic bacteria comprise several diverse physiological groups; first, fermentative anaerobes typically Gram-positive (*Clostridium*, *Ruminococcus*, and *Caldicellulosiruptor*) but containing a few Gram-negative

species (*Butryvibrio* and *Acetivibrio*); second, aerobic Gram-positive bacteria (*Cellulomonas* and *Thermobifida*); third, aerobic gliding bacteria (*Cytophaga* and *Sporocytophaga*) [9].

**TABLE 2.** Results of cellulase enzyme activity.

Locations	Strains	Relative Enzyme Activity Index ( $I_{cmc}$ )
SAV	SAV-A	$6.25 \pm 2.16$
	SAV-B	$5.33 \pm 1.44$
	SAV-C	$4.79 \pm 1.57$
	SAV-D	$4.86 \pm 1.32$
BSA	BSA-A	$6.66 \pm 4.61$
	BSA-B	$6.58 \pm 3.16$
	BSA-C	$7.88 \pm 4.47$
	BSA-D	$9.86 \pm 1.70$
BPA	BPA-A	$8.16 \pm 1.13$
	BPA-B	$10.60 \pm 1.57$
	BPA-C	$6.27 \pm 2.56$
	BPA-D	$6.67 \pm 2.51$
HHT	HHT-A	$2.00 \pm 0.46$
	HHT-B	$2.08 \pm 0.57$
	HHT-C	$1.86 \pm 0.12$
	HHT-D	$2.25 \pm 0.75$
Control		N/A

**TABLE 3.** The results of macroscopic and microscopic characteristics of bacteria.

Strains	Colony Morphology						Cell Morphology		
	Shape	Size	Color	Texture	Margin	Elevation	Gram	Endospore	Shape
BSA-D	Irregular	Medium	White	Smooth with granule	Flat	Flat	Positive	Negative	Coccus
BPA-A	Irregular	Medium	White	Smooth	Lobate	Flat	Positive	Negative	Irregular rod-coccus
BPA-B	Irregular	Small	White	Smooth	Lobate	Raised	Positive	Negative	Irregular rod-coccus

The results of colony morphology, cell morphology, and biochemical characterization were determined with bacteria determination book. Percentage similarity gets from divide of same amount characteristic between the results data and the total of identification characteristic from determination books. Table 5 shows the identification results obtained genus name [16].

BPA-B isolate was identified as *Pimelobacter* genus with similarity 85 %. *Pimelobacter* has characteristics cell Gram-positive, non-endosporic, and irregular-rod shape. *Pimelobacter* habitat is in soil with litter on the surface [16]. *Pimelobacter* has known can produce trehalose synthase. Trehalose synthase is an enzyme which converts maltose into trehalose. Trehalose ( $\alpha$ -D-glucopyranosyl  $\alpha$ -D-glucopyranoside) is a non-reducing disaccharide that is widespread. Conversion of maltose into trehalose reached 80 % [17]. Another side of *Pimelobacter* research to produce cellulase enzyme has not known yet before. BPA-B isolate can produce cellulase enzyme by the relative enzyme activity ( $I_{cmc}$ ) is  $10.67 \pm 1.57$ .

Identification result of BSA-D was genus *Micrococcus* with 80 % similarity. The characteristics colony were similar with *Micrococcus* is an irregular shape, white color, smooth texture, and flat while the same cell characteristics are Gram-positive and coccus shape. *Micrococcus* that produces cellulase has identified as *Micrococcus* sp. [18]. The bacteria can produce cellulase, xylanase, and carboxymethylcellulase but not showing the microscopic, macroscopic, and biochemical identifications of bacteria [19]. BSA-D has relative enzyme activities ( $I_{cmc}$ ) is  $9.86 \pm 1.70$ . Based on Bergey's Manual of Determinative Bacteriology 9<sup>th</sup> Edition, there are some biochemical tests same between two species, which are *M. sedentarius* and *M. lylae* [16].

The similarity of BPA-A isolate was closed with genus *Cellulomonas* [16]. The same cell characters of BPA-A with *Cellulomonas* were Gram-positive, irregular-rod shape, and non-endosporic. *Cellulomonas* also produces exoglucanase/xylanase, endoglucanase, cellulase, and  $\beta$ -glucosidase [20–22]. It can produce extracellular, intracellular, and cell-bound enzyme types. BPA-A just measured the cellulase activity index by halo zone of CMC medium and calculated the relative enzyme activities ( $I_{cmc}$ ). The relative enzyme activity of BPA-A is  $8.16 \pm 1.13$ . *Cellulomonas* sp. can use to electricity generations by the microbial fuel cell and photo-fermentation with consortium [23, 24].

**TABLE 4.** The results of biochemical characterization.

Physiological Tests	Strains		
	BSA-D	BPA-A	BPA-B
Lysin <sup>a</sup>	-	-	+
Ornithine	-	-	-
H <sub>2</sub> S <sup>b</sup>	-	-	-
Glucose <sup>c</sup>	-	+	-
Mannitol <sup>c</sup>	-	+	-
Xylose <sup>c</sup>	-	-	-
ONPG <sup>d</sup>	+	-	-
Indole from tryptophan <sup>b</sup>	-	+	-
Urease <sup>d</sup>	-	-	-
VP <sup>i</sup>	-	-	-
Citric <sup>e</sup>	+	-	+
TDA <sup>b</sup>	-	+	-
Gelatin <sup>f</sup>	+	+	+
Malonate <sup>g</sup>	+	-	-
Inositol <sup>c</sup>	-	-	-
Sorbitol <sup>c</sup>	-	+	-
Rhamnose <sup>c</sup>	-	+	-
Sucrose <sup>c</sup>	-	+	-
Lactose <sup>c</sup>	-	-	-
Arabinose <sup>c</sup>	-	-	-
Adonitol <sup>c</sup>	-	-	-
Raffinose <sup>c</sup>	-	-	-
Salicin <sup>c</sup>	-	-	-
Arginine <sup>h</sup>	+	+	+
Amylum <sup>d</sup>	-	-	-
Catalyses	+	+	+
Motility	+	+	+
Temperature 37 °C	+	+	+
Oxidative	+	+	+

<sup>a</sup> decarboxylation; <sup>b</sup> production; <sup>f</sup> fermentation; <sup>d</sup> hydrolysis; <sup>e</sup> used of; <sup>f</sup> dilution; <sup>g</sup> inhibition; <sup>h</sup> dihydrolase; <sup>i</sup> acetoin production.

**TABLE 5.** Results of three strains identification.

Strains	Locations	Similarity (%)	Strain of Closest Match	Identification results
BSA-D	Sadengan Bamboo Forest	80	<i>Micrococcus</i>	<i>Micrococcus</i>
BPA-A	Pancur Bamboo Forest	80	<i>Cellulomonas</i>	<i>Cellulomonas</i>
BPA-B	Forest	85	<i>Pimelobacter</i>	<i>Pimelobacter</i>

## CONCLUSIONS

It is concluded from this research that three of the highest potential productions of cellulase production were BPA-B, BSA-D, and BPA-A which has a cellulolytic index of 10.67, 9.86, and 8.16 respectively. Identification results BPA-B, BSA-D, and BPA-A each have in common with genus *Pimelobacter*, *Micrococcus*, and *Cellulomonas*. The identification can not conclude the species name because it needs a specific test as like as molecular sequencing and comparison with gene bank data.

## ACKNOWLEDGMENTS

The authors thank all staff of Alas Purwo National Park, all the P3L (Pelatihan Pengembangan dan Penelitian Lapangan) committee of Biology Universitas Airlangga 2017, alumni and senior of biology study program of Universitas Airlangga who has provided advice and assistance.

## REFERENCES

1. H. Syamsul, *Jurnal Biologi* **7**(1), 9–13 (2008).
2. P. P. Reanida, A. Supriyanto, and Salamun, “Eksplorasi bakteri selulolitik dari tanah mangrove Wonorejo Surabaya [Cellulolytic bacteria exploration from mangrove land Wonorejo Surabaya],” Bachelor thesis, Universitas Airlangga, 2012.
3. D. C. T. Betsy and J. Keogh, *Microbiology DeMYSTiFieD* (McGraw-Hill Companies, 2005), pp. 40–41.
4. A. S. S. Ibrahim and A. I. El-diwany, *Australian Journal of Basic and Applied Science* **1**(4), 473–478 (2007).
5. C. Zhang, X. Wang, W. Zhang, Y. Zhao, and X. Lu, *Appl. Microbiol. Biotechnol.* **101**(5), 1919–1926 (2017).
6. I. Haq and F. Akram, *Appl. Microbiol. Biotechnol.* **183**(1), 171–188 (2017).
7. V. Bhardwaj, G. Degrassi, and R. K. Bhardwaj, *International Research Journal of Engineering and Technology* **4**(1), 494–503 (2017).
8. M. Maki, K. T. Leung, and W. Qin, *Int. J. Biol. Sci.* **5**(5), 500–516 (2009).
9. M. J. Taherzadeh and K. Karimi, *Int. J. Mol. Sci.* **9**(9), 1621–1651 (2008).
10. M. Zyani, D. Mortabir, M. Mostakim, M. Iraqui, A. Habboud, and M. Ettayebi, *Ann. Microbiol.* **59**(4), 699–704 (2009).
11. H. Ruslan, *Vis Vitalis* **4**(1), 43–53 (2009).
12. B. F. T. Brockett, C. E. Prescott, and S. J. Grayston, *Soil Biol. Biochem.* **44**(1), 9–20 (2012).
13. R. H. Doi, A. Kosugi, K. Murashima, Y. Tamaru, and S. o. Han, *J. Bacteriol.* **185**(20), 5907–5914 (2003).
14. B. C. Behera, S. Parida, S. K. Dutta, and H. N. Thatoi, *American Journal of Microbiological Research* **2**(1), 41–46 (2014).
15. M. S. Alam, P. R. Sarjono, and A. L. N. Aminin, *Jurnal Sains & Matematika* **21**(2), 48–53 (2013).
16. D. H. Bergey and G. H. John, “Gram-Positive Cocci; Irregular, Nonsporing Gram-Positive Rods,” in *Bergey’s Manual of Determinative Bacteriology*, 9th edition, edited by P. H. Sneath and N. R. Krieg (Williams & Wilkins, Baltimore, 1994), pp. 530–579.
17. T. Nishimoto, M. Nakano, T. Nakada, H. Chaen, S. Fukuda, T. Sugimoto, *et al.*, *Biosci. Biotechnol. Biochem.* **60**(4), 37–41 (2014).
18. Z. M. Kaseke, K. Okaiyeto, U. U. Nwodo, I. V. Mabinya, and A. I. Okoh, *Sustainability* **8**(11), 1–15 (2016).
19. P. Nisha, A. Das, and S. Kv, *International Journal of Pharma Research and Health Sciences* **2**(3), 236–240 (2014).
20. K. O. Duedu and C. E. French, *Enzyme Microb. Technol.* **93–94**, 113–121 (2016).
21. A. K. N. Chan, A. K. L. Ng, K. K. Y. Ng, and W. K. R. Wong, *Gene* **642**, 367–375 (2018).
22. J. Gao and W. Wakarchuk, *J. Bacteriol.* **196**(23), 4103–4110 (2014).
23. Y. Takeuchi, W. Khawdas, Y. Aso, and O. H. Ohara, *J. Biosci. Bioeng.* **123**(3), 358–363 (2017).
24. Z. Y. Hitit, C. Z. Lazaro, and P. C. Hallenbeck, *Int. J. Hydrog. Energy* **42**(10), 1–11 (2016).