

Shopping Bag (Items)

[Home](#)[International Journals](#)[For Authors](#)[Books](#)[About Us](#)[Contact](#)

Ecology, Environment and Conservation Editorial Advisory Board

Chief Editor

Prof.(Dr.) R.K.Trivedy, Pune, India

Email id- rktrivedy@gmail.com

Tel: 91-20-46745119

Mobile No.-9975703363

EDITORIAL ADVISORY BOARD	
1. Dr. Prof. Moses Inbaraj, Professor, Department of Botany (Retired), Madras Christian College, Chennai, India	Indiainbarajmoses2004@yahoo.com
2. Dr. Prof. D.J. Lee, Professor, Department of Chemical Engineering, National Taiwan University, Taiwan	djlee@ntu.edu.tw
3. Dr. T. Bahorun, Professor, Department of Biosciences, Mauritius University, Mauritius	tbahorun@uom.ac.mu
4. Dr. Prof. A.R. Ghosh, Professor, Department of Environmental Sciences, University of Burdwan, Burdwan, India	apurbaghosh2010@gmail.com
5. Dr. Prof. M. Zaman, Professor Department of Botany, Dhaka University, Dhaka, Bangladesh	zashim@du.ac.bd
6. Dr. Marcantonio Bragadin, Professor Department of Chemistry, University of Venice, Venice, Italy	bragadin@unive.it
7. Dr. Z. Fuat Topark, Professor Dicle University, Diyarbakir, Turkey	toprakzey@itu.edu.tr

8. Dr. Wilson S. Tisera, Associate Professor Christian University, Kupang, Indonesia	wilson_tisera@yahoo.com
9. Dr. Prof. Philip C. Reid, Scientist Marine Biological Association, U.K	pcre@mba.ac.uk
10. Dr. Prof. Mohd. Yusuf, Professor Deptt. of Environmental Engineering, University Sans Malaysia, Malaysia	danielmy@iium.edu.my
11. Dr. Prof. Azni Idris, Deputy Vice-Chancellor, Research and Innov., Univ. of Putra Malaysia, Malaysia	azni@upm.edu.my
12. Dr. Vikas Sharma, Associate Professor, RARS of SKUAST, Jammu, J&K, India	vsagron@gmail.com
13. Dr. A.C. Pandey, Senior Scientist, Birsa Agricultural University, Ranchi, India	azni@upm.edu.my
14. Dr. Prof. Agoes Soegianto, Professor , Dept. Biology, Airlangga Univ. Kampus C, Surabaya, Indonesia	agoes_soegianto@fst.unair.ac.id
15. Dr. Ahmed L. Mahmoudi, Professor, Water Studies Centre, King Faisal University, Saudi Arabia	mahmoudi@kfu.edu.sa
16. Dr. Siroop Chaudhuri, Associate Professor Center for Environment, Sustainability and Human Development (CESH) O.P. Jindal Global University, Sonipat, India	schaudhuri@jgu.edu.in
17. Dr. Amin L. Setyo, Professor, Graduate School of International Crop, Hiroshima University, Hiroshima, Japan	amin28@ub.ac.id
18. Dr. Francis Gbogbo, Associate Professor, Department of Animal Biology, University of Ghana, Ghana	fgbogbo@ug.edu.gh
19. Dr. Prof. Deenbandhu Sahoo, Professor Department of Botany, university of Delhi, Delhi, India	dbasahoo@yahoo.com
20. Dr. Prof. J.K. Ladha, Professor Emeritus University of California, Davis, California, U.S.A.	jkladha@ucdavis.edu
21. Dr. Prof. M.Z.M. Nomani, Professor Faculty of Law, Aligarh Muslim University, Aligarh, India	zafarnomani@rediffmail.com
22. Dr. G.R. Pathade, Principal (Retired) H.V. Desai College, Pune, India	girishpathade@yahoo.co.in
23. Dr. Alfred Maroyi, Associate Professor Department of Botany, University of Fort Hare, Alice 5700, South Africa	amaroyi@ufh.ac.za
24. Dr. Prof. A.V.L.N.S.H. Hariharan, Professor Dept. of Chemistry, College of Science, GITAM University, Visakhapatnam, India	ahharan@gmail.com
25. Dr. Zeinab El-Sayed Zayed, Associate Professor Department of Biotechnology, The Central Lab of Date Palm Researches and Development Agriculture Research Center, Cairo, Egypt	zemz2005@yahoo.com
26. Dr. Raj Singh, Professor Department of Botany, MMD University, Mullana, Ambala, Haryana, India	dr.rajsingh09@gmail.com

27. Dr. Prof. Mohammed Latif Khan, Professor, Department of Botany Dr. Hari Singh Gour Vishwavidyalaya Sagar, India	khanml61@gmail.com
28. Dr. R. Dinesh Principal Scientist, ICAR-Indian Institute of Spices Research, Marikunnu, P.O. Kozhikode, India	rdinesh2005@gmail.com
29. Dr. Suresha Gududappa, Senior Environmental Scientist, Saudi Arabia	suresh@ebe.com.sa
30. Dr. Azma Hanim Ismail, School of Biological Sciences, Universiti Sains Malaysia, 11800 USM, Pulau Pinang, Malaysia.	
Back to EEC Journal Details	

[Home](#) | [International Journals](#) | [Books](#) | [About Us](#) | [Contact Us](#) | [Submit Paper](#) | [Search Journal Article](#) |

[Become a fan](#) on Facebook

[Follow us](#) on Twitter



© EM International 2012-2019 | **Developed by Eneblur Consulting**

The influence of ozone exposure on organoleptic and chlorophyll levels of curly lettuce (*Lactuca sativa* L.)

Suryani Dyah Astuti^{1,2,*}, Hery Purnobasuki³, Miratul Khasanah⁴, Siti Khoiriyatul¹,
Nurul Fitriyah^{1,2}, Deny Arifianto⁵ and Fadli Ama⁶

¹Departement of Physics, Faculty of Science and Technology, Universitas Airlangga, Surabaya, Indonesia 60115

²Biophysics and Medical Physics Research Group, Faculty of Sciences and Technology, Universitas Airlangga, Surabaya-Indonesia 60115

³Departement of Biology, Faculty of Science and Technology, Universitas Airlangga, Surabaya, Indonesia 60115

⁴Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Surabaya, Indonesia 60115

⁵Vocational Faculty, Universitas Airlangga, Surabaya, Indonesia 60115

⁶Biomedical Engineering, Faculty of Science and Technology, Universitas Airlangga, Surabaya, Indonesia 60115

(Received 22 February, 2020; Accepted 17 June, 2020)

ABSTRACT

Ozone has been utilized widely for sterilization application, for example to keep the freshness of both fruits and veggies yet keep them safe too for human consumption and chemical-free. This work aims to investigate the effect of ozone exposure through the air and water on organoleptic and chlorophyll level of curly lettuce (*Lactuca sativa* L.). The given exposure has been done with various ozone concentrations corresponding with increasing exposure time, namely 0.0147 mg/L (120 second exposure time), 0.0294 mg/L (240 second exposure time), 0.0371 mg/L (360 second exposure time) and 0.0479 mg/L (480 second exposure time). The curly lettuce (*Lactuca sativa* L.) was originally taken from veggies farmland located in Sumberojo village, Batu sub-district, Batu city East Java, Indonesia. This project was carried out by using Completed Random Contrivance (CRC) technique. The observation was focused on organoleptic and mass loss of curly lettuce (*Lactuca sativa* L.) and it was carried out on every 8 hours for two days straight non-stop. The chlorophyll level measurement was employed by using UV-Vis spectrophotometer after two days from the last observation. Organoleptic showed that ozone exposure technique through water mediator with ozone concentration 0.0147 mg/L gave the best results to detain organoleptic damages of curly lettuce (*Lactuca sativa* L.). The chlorophyll level was analyzed by employing *Statistical Package for The Social Sciences* (SPSS) with factorial technique (Two Way ANOVA). Therefore, all the analysis result convinced that this ozone exposure technique with water mediator gave the best results compared to air mediator, in term of chlorophyll level preserving of *Lactuca sativa* L. This technique is very effective to maintain the quality and curly lettuce (*Lactuca sativa* L.) freshness.

Key words: Ozon, Dried lettuce (*Lactuca sativa* L.), Organoleptic, Chlorophyll level

Introduction

Geographically, Indonesia is located between two continents, Asia and Australia, and two oceans, the Pacific Ocean and the Indian Ocean. While astronomically, Indonesia is located at 6° NL – 11° SL and 95° EL - 141° EL. Since its location Indonesia becomes one of the 14 countries crossed by the equator. Due to this astronomical location, Indonesia has a tropical climate that has plentiful rainfall and fertile soil structure. As this soil fertility, Indonesia has abundant flora wealth. One of them is vegetables (Mattos *et al.*, 2014).

Curly lettuce is one type of horticulture plant with high economic value (Nübling *et al.*, 2019). This is represented from the high number of markets demands to meet market needs, especially in hospitality, large restaurants to foreign countries as export commodities (Adimiharja *et al.*, 2013). But lettuce is one type of vegetable with a low resistance, thus it cannot last long. In spite of lettuce leaves have interesting appearance, they are often used as decorations in serving food and have crispy tastes that are very suitable for fresh vegetables, yet they are easy to rot (Adimihardja *et al.*, 2013). Hence the high economic value of lettuce is not commensurate with its cell durability.

Cell damage is induced by water content that begins to decrease after being picked as well as the alleged existence of microorganisms in the form of bacteria that grow on lettuce. In addition, cell damage is also caused by enzymes, temperature, air and light (Wulfkuehler *et al.*, 2014). As a result, lettuce does not last long, thereby reducing its economic value. The economic value is the comparison of costs incurred with the costs received. If lettuce is easily damaged, the economic value will also be reduced. This will harm the lettuce farmer. Besides the reduced value of exports will reduce the value of state income.

Storing fruits and vegetables to keep them fresh have been carried out with several ways, such as gamma-ray exposure, conventional chlorine or chlorine disinfectant (Das *et al.*, 2012). However, gamma ray radiation is expensive, the scent of chlorine is irritating and cause side effects in the form of carcinogenic trihalomethane compounds (Astuti *et al.*, 2020). Compared with chlorine, ozone speeds are 3250 times faster in killing organisms and 150% stronger in oxidative power. In addition, storing fruits and vegetables can be in the refrigerator, yet

the bacteria in vegetables are still active (Baur *et al.*, 2004). Chemicals are also used to storing vegetables, but the usage of chemicals produces is harmful since it contains toxins. Another way to storing fruits and vegetables is using immersion in a solution of acetic acid (C₂H₄O₂) or often called vinegar water. This solution is food safe, yet the taste of food will change to be acidic. Hence, currently the ozone method is being developed which are considered to be more eco-friendly and safe (Karaca and Velioglu, 2014).

Ozone is the smallest part of the Earth's atmosphere that is unstable and quickly decomposes at ambient temperatures (Take *et al.*, 2014). Ozone is naturally formed from ultraviolet (UV) radiation from sunlight (Bocci *et al.*, 2006). Various research stated that ozone can be utilized in photodynamic therapy (Astuty *et al.*, 2018; Sunarko *et al.*, 2017; Puspita *et al.*, 2020) to inactivate the beneficial bacteria in order to prevent the population of pathogenic bacteria on the skin and food. Ozone in the mechanism of photosensitization acts as oxygen which absorbs light energy to form radical oxygen species (ROS) that will kill both planktonic bacteria (Astuti *et al.*, 2017), biofilms (Astuti *et al.*, 2017; Shie *et al.*, 2019) and fungi (Astuti *et al.*, 2019). According to Asgar *et al.*, (2017) ozone method can also shed the remnants of pesticides and heavy metals attached to the fruit or leaves that it is food safe and ozone treatment can increase the life-extending of tomatoes. Ozone can inactivate microorganisms without leaving chemical residues and will re-create into oxygen, therefore it is very eco-friendly (Syafarudin *et al.*, 2013).

The most important mechanism of ozone is the sulfhydryl oxidation from the enzyme. This layer is the first subject to be attacked by ozone. Ozone invasion of the cell wall leads to changes in the permeability of cells and causes cell lysis (Asgar *et al.*, 2017). There are several studies indicated that ozone can be used as a natural preservative. As ozone contains antibacterial which can inactivate microorganisms thus extending the storage period (Zorlugenç *et al.*, 2008). Ozone as the most powerful oxidizer after hydroxide radicals (OH^{*}), can oxidize heavy metals, degrades organic compounds including organo-chlorides and aromatics, eliminate color, odor, taste. Justification of parameters and the best treatment for flower cabbage, namely at 1.5 mg/L ozone concentrations with leaves in cold storage due to the characteristic of ozone which acts as a

sterilizer (ozone can inactivate microbes such as pathogenic bacteria, fungi, and viruses), deodorization (removes smells owing to microbes and organic compounds), decolorization (removes organic dyes) and degradation (decomposes organic compounds and oxidizes heavy metals) (Borelli and Bocci, 2018).

Ozone exposure techniques applied in this study are exposure through air mediator (gasification) and through water mediator. Ozone exposure to red chili can increase its life-extension. Hence in this case ozone is engaged for sterilization (Calvosa *et al.*, 1991). Besides cell damage is caused by breakage to enzymes due to oxidation by ozone that initiate through the stomata, thus respiration becomes slow and water levels are not easily lost as a result of the vegetables will remain fresh (Asgar *et al.*, 2011).

Based on the introduction and existing research, this study aims to investigate the concentration and radiation techniques that are most effective in inhibiting cell damage in curly lettuce (*Lactuca sativa* L).

Materials and Methods

This study was conducted at the Analytical Chemistry Laboratory, Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga for characterization of ozone concentration and Biophysical Laboratory, Department of Physics, Faculty of Science and Technology, Universitas Airlangga for ozone exposure on curly lettuce (*Lactuca sativa* L) and testing chlorophyll level of curly lettuce (*Lactuca sativa* L).

Sample Treatment

The sample used in this study was 81 curly lettuce (*Lactuca sativa* L) obtained from the Vegetable Garden in Sumberejo Village, Batu sub-district, Batu City. The study was conducted with a Completed Random Contrivance (CRC). The sample consisted of group C: control without treatment, G1 : ozone exposure through air mediator with concentration of 0.0147 mg/L (120 s exposure time), G2 : ozone exposure in the air concentration of 0.0294 mg/L (240 s exposure time), G3 : ozone exposure through air mediator with concentration of 0.0371 mg/L (360 s exposure time), and G4 ozone exposure through air mediator with concentration of 0.0479 mg/L (480 s exposure time), W1 : ozone exposure through water mediator with concentration of 0.0147 mg/L (120 s exposure time), W2 : ozone exposure through water with concentration 0.0294 mg / L (240 s exposure

time), W3 : ozone exposure through water with concentration 0.0371 mg/L (360 s exposure time), and W4 : ozone exposure through water with concentration of 0.0479 mg/L (480 s exposure time).

The study was conducted with 3 repetitions. Exposure through air mediator was held by placing the sample in an isolated container and then ozone was exposed according to the time needed to reach the desired concentration. Ozone exposure through water mediator was done by exposure the ozone in 3 liters of water according to the time needed to reach the desired concentration. The sample then soaked in ozone for 6 minutes. Washing and soaking vegetables and fruit in water gives the best results between 5-10 minutes.

Organoleptic and chlorophyll observation

Organoleptic observation of curly lettuce was carried out every 4 hours for two days. Chlorophyll levels were tested after 2 days using a UV-Vis spectrophotometer (Milenkovie *et al.*, 2012 ; Torres *et al.*, 2016).

Statistical Analysis

Quantitative data analysis was performed using Statistical Package for Social Science (SPSS) analysis in the form of factorial tests (Two Way Anova). The data tested statistically are chlorophyll level. This test aims to determine whether there are differences in chlorophyll level in curly lettuce (*Lactuca sativa* L) after treatment.

Results and Discussion

The results of statistical testing of curly lettuce chlorophyll (*Lactuca sativa* L) levels indicate that there is an impact of exposure techniques on chlorophyll level with p value = 0.02 (p < 0.05), there is no effect of variations in ozone concentration on chlorophyll content with p value = 0,65 (p > 0.05) and there is no influence of the interaction of the two factors on chlorophyll content with p value = 0.58 (p > 0.05). The results of testing the exposure technique with the Independent sample T-Test showed the technique of exposure to ozone through water mediator produced the highest average value of 2.18.

The Effect of Ozon Flow Time on Ozone Concentration

The value of ozone concentration (O₃) for each flow time is illustrated in Figure 1. Based on the graph, it




Fig. 1. The graph of ozon concentration againts ozon flow time

represents that the greater the ozone concentration time, the greater the flow.

Organoleptic

Based on observations, it stated that organoleptic curly lettuce (*Lactuca sativa* L) which was not exposed to ozone or control samples began to show appearance of wilting leaves at intervals of 16 hours and increased with time. Curly lettuce (*Lactuca sativa* L) exposed to ozone through the air mediator can be seen in Figure 2.

Observations indicated that curly lettuce (*Lactuca sativa* L) exposed to ozone through the air mediator was more easily damaged than the control group without ozone exposure. There was a brownish

Time	Control	0,0147 mg/L	0,0294 mg/L	0,0371 mg/L	0,0479 mg/L
8 hours					
16 hours					
24 hours					
32 hours					
48 hours					

Fig. 2. The organoleptic of curly lettuce (*Lactuca sativa* L) control group with ozon exposure through air mediator

color to the leaves such as the effects of burning, wilting and drying. These effects were convenient with increasing time and concentration used. Hence the exposure to ozone through the air mediator accelerated organoleptic damage of curly lettuce (*Lactuca sativa* L). Curly lettuce (*Lactuca sativa* L) with treatment of ozone exposure through water mediator is illustrated in Figure 3.

The results of ozone exposure treatment in curly lettuce through water mediator (*Lactuca sativa* L) showed the effect of damage after 40 hours of storage in the form of leaf texture that started wilting and yellowing increased with time. Ozone exposure to curly lettuce (*Lactuca sativa* L) through water mediator are able to maintain its freshness longer than the control samples or samples that are exposed to


























Time	Control	0,0147 mg/L	0,0294 mg/L	0,0371 mg/L	0,0479 mg/L
8 hours					
16 hours					
24 hours					
32 hours					
40 hours					

Fig. 3. The control group organoleptic and ozon exposure through water mediator to wash curly lettuce (*Lactuca sativa* L)

ozone through the air mediator. The most durable curly lettuce (*Lactuca sativa* L) was indicated by a concentration of 0.0147 mg / L.

Mass Loss of Curly Lettuce (*Lactuca sativa* L)

Graph of mass loss curly lettuce (*Lactuca sativa* L) with ozone treatment through air and water mediator is showed in Figure 4 and 5. Both graphs in Figure 4 and 5 represent that the mass loss of curly lettuce (*Lactuca sativa* L) on ozone exposure through air mediator and water mediator tends to be the same for all concentrations. Thus, in this study the weight loss cannot be applied as a reference in investigating the effective treatment. This is suspected due to the concentration used is too small hence it does not provide a significant difference.

While mass loss of curly lettuce (*Lactuca sativa* L) after ozone exposure through water mediator is illustrated in Figure 6.




Fig. 4. Mass loss graph of curly lettuce (*Lactuca sativa* L) after ozone exposure treatment through air mediator against time interval




Fig. 5. Mass loss graph of curly lettuce (*Lactuca sativa* L) after ozone exposure treatment through water mediator against time interval




Fig. 6. The graph of chlorophylls level on ozone exposure sample for both applied mediator (water and air) versus ozone concentration

Chlorophyll Level of Curly Lettuce (*Lactuca sativa* L)

Graph of chlorophyll level of curly lettuce on both samples after ozone exposure trough air mediator and water mediator are illustrated in Figure 6.

Based on the graph, it represents that curly lettuce (*Lactuca sativa* L) exposed to ozone through water mediator gives a higher value of chlorophyll level than through air mediator. The highest levels of chlorophyll are found in curly lettuce (*Lactuca sativa* L) which is exposed to ozone through water mediator with a concentration of 0.0147 mg/L which is equal to 2,559 mg/L. While the lowest chlorophyll level is found in curly lettuce (*Lactuca sativa* L) which is exposed to ozone through the air mediator with a concentration of 0.0479 mg / L which is equal to 1,228 mg/L. This is in accordance with research by Zorlugenc *et al.*, (2008) which stated that ozone exposure through water mediator was more effective than through air mediator. The mechanism of ozone exposure to curly lettuce (*Lactuca sativa* L) is displayed in Figure 7.

The interaction of ozone when added to water is by oxidizing metal ions contained in water. This leads the electron transfer process to produce ozone radical anions. Ozone radical anions will initiate the formation of ozone decomposition chains. Ozone which reacts with carbon in water would produce hydrogen peroxide and carbonyl (Calvosa *et al.*, 1991). According to Hong (2010) the time of washing with ozone water greatly determined the efficiency.

Ozone maintains freshness by damaging the en-

zymes contained in cells. Ozone will infiltrate the tissue of curly lettuce (*Lactuca sativa* L) through the stomata valves in epidermis when washing with ozone. Furthermore, ozone will oxidize cell walls into lysis or rupture. This induces the enzymes in the membrane and cell nucleus in the chloroplast to be inactive, hence the process of respiration will be inhibited (Asgar *et al.*, 2011). Respiration is a metabolic process in the form of combustion of complex compounds by oxygen (O_2) which produces simple compounds such as carbon dioxide (CO_2), water and energy for synthesis reactions by cells. The inhibition of respiration will lead the vegetable water content to be maintained thereby inhibiting cell damage. Thus, it will increase the freshness of vegetables (Okpala *et al.*, 2015).

In addition, the new compounds contained in the water will interact with microorganisms that may be carried in curly lettuce plants (*Lactuca sativa* L). In line with Nurjanah (2002), plants after being picked are still doing respiration and producing a ripening hormone, namely ethylene hormone. Microorganisms stimulate ethylene production and induce the vegetables to quickly rot.

Ozone will infiltrate microbial cells, change the permeability of microbes and lead microbes into lysis or rupture (Asgar *et al.*, 2017). Some researchers have shown that only with low ozone concentration (less than 0.5 mg/L) ozone can inactivate microorganisms in water, moreover ozone is able to

sterilize water (Asgar *et al.*, 2011). The sensitivity of microorganisms with ozone is affected by several factors including microorganism strains, cell physiological state, growth rate, pH of the medium, temperature, humidity (Zorlugenc *et al.*, 2008).

Ozone treatment on curly lettuce (*Lactuca sativa* L) can maintain the freshness up to 5 days compare with controls that only last for 2 days in the form of wilted leaves and soft, runny texture. Observation of curly lettuce (*Lactuca sativa* L) was carried out in every 8 hours in term of mass loss.

Mass loss is mainly due to transpiration and infrequently by respiration (Asgar *et al.*, 2011). Shrinkage of weight indicates reduced water content. Decreased water content during storage can induce vegetables wilt and decrease their fertility. In this study the weight loss does not provide a real difference allegedly owing to the concentration used is too small. Thus it does not provide a significant difference.

Curly lettuce (*Lactuca sativa* L) gives changes to the organoleptic. Organoleptic testing was carried out based on the sensing process. In accordance with this study, it is able to be discovered that there are changes in the curly lettuce such as the texture is softer and changes in the color of the leaves. Asgar *et al.*, (2011) stated that the softening of fruits or vegetables was owing to changes that occur in cell walls and other pectin substance compounds, namely the solubility of the depolymeration of pectin sub-




Fig. 7. Ozon exposure mechanism on curly lettuce (*Lactuca sativa* L), (a) given exposure was divided in two kind of exposure technique, (b) ozone exposure through air mediator, (c) through water mediator that inhibit cell respiration thus it can maintain its durability

stances. Insoluble pectin was transformed by various enzymes to water-soluble pectin. According to Asgar *et al.* (2011) on flower cabbage, excess oxygen from the decomposition of the remaining ozone actually became the main menu for the growth of the aloive microbes, these microbes that induced the texture of plants to be soft.

The transformation of green leaves to yellow on curly lettuce (*Lactuca sativa* L) is due to the degraded chlorophyll pigment structure which was followed by the formation or appearance of yellow pigment (Milenkoviæ *et al.*, 2012). In consonance influential factors in the formation of yellow color were abscisic and ethylene hormones. Both of these hormones triggered the aging of leaves which induced the loss of chlorophyll, RNA, protein, and stimulated the formation of carotenoids. In line with Asgar *et al.* (2011) discoloration in vegetables caused by the majority of chlorophyll pigments undergo changes during storage due to maturity, besides the loss of Mg + from the tetrapyrrole structure to mold feotin in chlorophyll induced the brown discoloration.

Chlorophyll level in curly lettuce (*Lactuca sativa* L) were tested after two days using a UV-Vis spectrophotometer. UV-Vis spectrophotometer tested the chlorophyll level by detecting the concentration of the sample solution (Gokmen *et al.*, 2002). Light would absorb the color of the sample solution at certain wavelengths. The research applied a Genesys 30 type UV-Vis spectrophotometer that could detect wavelengths in the range 325-1000 nm. The level of curly lettuce chlorophyll (*Lactuca sativa* L) was calculated using the formula Wintermans and Mots (Pratama and Ainun, 2015) which used 2 wavelengths in their analysis, namely at 665 nm and 649 nm. In this study the chlorophyll level produced from various treatments had almost the same value. This was presumably due to the difference in concentration used was too small.

Conclusion

Ozone exposure through the air mediator can accelerate organoleptic damage of curly lettuce (*Lactuca sativa* L) while ozone exposure through water mediator can inhibit organoleptic damage of curly lettuce (*Lactuca sativa* L). Nevertheless, ozone exposure through air mediator and through water mediator do not indicate an impact on the chlorophyll level of curly lettuce as it gives nearly same value. This is presumably due to the concentration range of ozone

exposure was too small. The most effective treatment in preventing damage to curly lettuce (*Lactuca sativa* L) is ozone exposure through water mediator at a concentration of 0.0147 mg/L. Other parameters freshness of some specific fruits and various veggies using this technique need further investigation.

References

- Adimiharja, S.A., Hamid, G. and Rosa, E. 2013. The effect of giving a combination of cow copos and fertimix on the growth and production of two lettuce cultivars (*Lactuca sativa* L) in the floating raft hydroponic system. *Journal of Agriculture*. 4 : 6-20
- Asgar, Ali, Darkam Musaddad, and Rahmat Sutarya, 2017. The effect of ozone and packaging in storage for decreasing in pesticide residue and keeping the freshness of red chili characteristic. *Jurnal Holtikultura*. 27: 241-252.
- Asgar, Ali, A.T. Sugiarto, Sumartini and Ariani, D. 2011. Study of ozone exposure to the characteristics of fresh cabbage flower (*Brassica oleracea* var. Botrytis) during storage at cold temperatures. *Jurnal Holtikultura*. 10: 787-795.
- Astuti, S.D., Fina, W.F., Darmanto, W., Purnobasuki, H., Fitriyah, N. and Ama, F. 2020. Effects of Co 60 gamma ray Ionizing radiation exposure on the variability of *Adenium obesum* growth, *Eco. Env. & Cons.* 26 (June Suppl. Issue): S128-S134.
- Astuti, S.D., Drantantiyas, N.D.G., Putra, A.P., Puspita, P.S., Syahrom, A. and Suhariningsih, 2019. Photodynamic effectiveness of laser diode combined with ozone to reduce *Staphylococcus aureus* biofilm with exogenous chlorophyll of *Dracaena angustifolia* leaves. *Biomedical Photonic*. 8(2): 4-13.
- Astuti, SD., Wibowo, A., Abdurachman and Triyana, K. 2017. Antimicrobial Photodynamic Effects of Polychromatic Light Activated by Magnetic Fields to Bacterial Viability. *J. Int. Dent. Med. Res.* 10 (1) : 111-117.
- Astuti, S.D., Arifianto, D., Drantantiyas, N.D.G., Aulia M.T.N. and Abdurachman, 2017. Efficacy of CNC-diode laser combine with chlorophylls to eliminate *Staphylococcus aureus* biofilm, Publishing in *International Seminar Sensors, Instrumentation, Measurement, and Metrology (ISSIMM)*, IEEE Xplore 5 Januaryry INSPEC Accession Number: 16563895
- Astuty, S.D., Suhariningsih, Baktir, A. and Astuti, S.D. 2019. The Efficacy of Photodynamic Inactivation of the Diode Laser in Inactivation of the *Candida albicans* Biofilms with Exogenous Photosensitizer of Papaya Leaf Chlorophyll. *Journal of Lasers in Medical Sciences*. 10(3) : 215-224.
- Baur, S., Klaiber, R., Hammes, W.P. and Carle, R. 2004. Sen-

- sory and Microbiological Quality of Shredded, Packaged Iceberg Lettuce as Affected by Pre-Washing Procedures with Chlorinated and Ozonated Water. *Innovative Food Science & Emerging Technologies*. 5(1): 45-55.
- Bocci, V., Borrelli, E., Travagli, V. and Zanardi, I. 2006. The Ozone/Paradox: Ozone Is a Strong Oxidant as Well as a Medical Drug. Wiley Interscience. 2006.
- Borrelli, E. and Bocci, V. 2018. Basic Biological and Therapeutic Effects of Ozone Therapy in Human Medicine. in *Ozone Science and Technology*, [Ed. Rein Munter], In: *Encyclopedia of Life Support Systems (EOLSS)*, Developed under the Auspices of the UNESCO, Eolss Publishers, Oxford, UK, 2010, [http://www.eolss.net] [Retrieved June 22, 2018]
- Calvosa, L., Monteverdi, A., Rindone, B. and Riva, G. 1991. Ozone oxidation of compounds resistant to biological degradation. *Journal of Technology*. 25 : 985-993.
- Das, A., Utpal Raychaudhuri, U. and Chakraborty, R. 2012. Effect of freeze drying and oven drying on antioxidant properties of fresh wheatgrass. *International Journal of Food Sciences and Nutrition*. 63 (6) : 718-721.
- Gokmen, V., Bahçeci, S. and Acar, J. 2002. Liquid Chromatographic method for the Determination of Chlorophylls, Carotenoids, and their Derivatives in fresh and Processed Vegetables. *Journal of Liquid Chromatography & Related Technologies* 25 (8): 1201-1213.
- Hong, S.G., Park, H.M. and Kwang, H.C. 2010. Development of washing and sterilization system for leafy vegetables. *Engineering in Agriculture, Environment and Food*. 3 : 87-92.
- Karaca, H. and Velioglu, Y.S. 2014. Effects of ozone treatments on microbial quality and some chemical properties of lettuce, spinach, and parsley. *Postharvest Biology and Technology*. 88 : 46-53.
- Leisner, C. 2020. Review: Climate change impacts on food security- focus on perennial cropping systems and nutritional value. *Plant Science*. 293 (2020) 110412.
- Mattos, L.M., Moretti, C.L., Jan, S., Sargent, S.A., Lima, C.E.P. and Fontenelle, M.R. 2014. Climate Changes and Potential Impacts on Quality of Fruit and Vegetable Crops. *Emerging Technologies and Management of Crop Stress Tolerance*. 467-486.
- Milenkovic, S.M., Zvezdanovic, J.B., Andelkovic, T.D. 2012. The Identification of Chlorophyll and its Derivatives in the Pigment Mixtures : Hplc-Chromatography. *Visible and Mass Spectroscopy Studies*. 1(1): 16-24.
- Nübling, S., Hägele, F., Schweiggert, R.M. Carle R. Schmidt, H. and Weiss, A. 2019. Effect of Different Wash Water Additives and Deep-Frozen Storage on the Quality of Curly Parsley (*Petroselinum crispum* var. *crispum*). *Food Bioprocess Technol.* 12 : 158-165. <https://doi.org/10.1007/s11947-018-2201-7>
- Nurjanah, S. 2002. Kajian laju respirasi dan produksi etilen sebagai dasar penentuan waktu simpan sayuran dan buah-buahan. *Jurnal Bionatura*. 4 : 148-156
- Okpala, Odilichukwu, C.R., Gioacchino, B, Abdurahim, Abdulkadir, and Chukwuka U. Madumelu, 2015. Ozone (O3) Process Technology (OPT): an exploratory brief of minimal ozone discharge applied to shrimp product. *Journal of Energy*. 75 : 2427-2435.
- Pratama, A.J. and Ainun, N.L. 2015. Analisis kandungan klorofil gandasuli (*hedychium gardnerianum* shephard ex ker-gawl) pada tiga daerah perkembangan daun yang berbeda. *Jurnal Pendidikan Biologi*. 1 : 216-219.
- Puspita, P.S., Astuti, S.D., Nasution, A.M.T., Pradhana, A.A.S. and Mawaddah, A. 2020. Photodynamic Therapy with Ozone aids to *Staphylococcus aureus* Biofilm Reduction. *Indian Vet. Journal*. 97 (02) : 24-26.
- Shie, M.Y., Shen, Y.F., Astuti, S.D., Xing Lee, A.K., Lin, S.H., Dwijaksana, L.N.B. and Chen, Y.W. 2019. Review of Polymeric Materials in 4D Printing Biomedical Applications. *Polymers*. 11 (1864) : 1-17
- Syafarudin., Angki, and Novia. 2013. Produksi ozon dengan bahan baku oksigen menggunakan alat ozon generator. *Jurnal Teknik Kimia*. 19: 1-9.
- Sunarko, S.A., Ekasari, W. and Astuti S.D. 2017. Antimicrobial effect of pleomeleangustifolia pheophytin A activation with diode laser to *Streptococcus mutans*. *Journal of Physics: Conference Series*. 85(2017)12038
- Take, Sosiawati, Nur M. and Winarni, T.A. 2014. Produksi ozon dalam reaktor dielektrik barrier discharge plasma (DBDP) terkait panjang reaktor dan laju alir udara serta pemanfaatannya untuk menjaga kualitas asam amino ikan. *Jurnal Fisika Teori, Eksperimen dan Fisika Aplikasi*. 17 : 25-32.
- Torres, P.B., Chow, F., Furlan, C.M. and Mandelli, F. 2014. Standardization of A Protocol to Extract and Analyze Chlorophyll A and Carotenoids. *Brazilian J. Oceanogr.* 62 (1) : 57-63.
- Wulfkuehler, S., Stark, S., Dietz, J., Schmidt, H., Weiss, A. and Carle, R. 2014. Effect of Water Jet Cutting and Moderate Heat Treatment on Quality of Fresh-Cut Red Oak Leaf Lettuce (*Lactuca sativa* L. var. *crispa*). *Food Bioprocess Technol.* 7 : 3478-3492 (2014). <https://doi.org/10.1007/s11947-014-1360-4>
- Zorlugenç, Bülent., Feyza Kýrog, çlu Zorlugenç, Serdar Öztekin, I. and Bülend Evliya. 2008. The influence of gaseous ozone and ozonated water on microbial flora and degradation of aflatoxin B1 in dried figs. *Food and Chemical Toxicology*. 46 : 3593-3597.



SJR

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)

Ecology, Environment and Conservation

Discontinued in Scopus as of 2021

COUNTRY

India



Universities and research institutions in India



Media Ranking in India

SUBJECT AREA AND CATEGORY

Agricultural and Biological Sciences
Ecology, Evolution, Behavior and Systematics

Environmental Science
Ecology
Nature and Landscape Conservation

PUBLISHER

EM International

H-INDEX

17

PUBLICATION TYPE

Journals

ISSN

0971765X

COVERAGE

1997-2021

INFORMATION

[Homepage](#)

[How to publish in this journal](#)

r_trivedy@bharatmail.co.in

SCOPE

Published Quarterly Since 1995. Ecology, Environment and Conservation is published in March, June, September and December every year. ECOLOGY, ENVIRONMENT AND CONSERVATION is one of the leading International environmental journal. It is widely subscribed in India and abroad by Institutions and Individuals in education and research as well as by Industries, Govt. Departments and Research Institutes.

Join the conversation about this journal

Quartiles

FIND SIMILAR JOURNALS ?

options

1 Applied Ecology and Environmental Research HUN 38% similarity	2 Nature Environment and Pollution Technology IND 36% similarity	3 DYNA (Colombia) COL 35% similarity	4 EnvironmentAsia THA 33% similarity	5 Jurnal Teknologi (Sciences and Engineering) MYS 33% similarity
---	--	--	--	--

SJR








Total Documents



Citations per document





Ecology, Environment and Conservation

Not yet assigned quartile

SJR 2021
0

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com" style="display: inline-block; border: 1px solid black; padding: 2px 5px;">https://www.scimagojr.com
```



Explore, visually communicate and make sense of data with our **new data visualization tool.**



Metrics based on Scopus® data as of April 2022