

# 20. Differences in interleukin-6 and interleukin-17 expression

*by* Etty Hary Kusumastuti

---

**Submission date:** 09-Jan-2023 09:10AM (UTC+0800)

**Submission ID:** 1989935269

**File name:** .Differences\_in\_interleukin-6\_and\_interleukin-17\_expression.pdf (446.34K)

**Word count:** 5131

**Character count:** 27943

# Differences in interleukin-6 and interleukin-17 expression in covid-19 post-mortem lung tissue biopsy compared with non-covid-19

Etty Hary Kusumastuti<sup>1,\*</sup>, Priangga Adi Wiratama<sup>1</sup>, Grace Ariani<sup>1</sup>, Stephanie Natasha Djuanda<sup>1</sup>, Alphania Rahniayu<sup>1</sup>, Nila Kurniasari<sup>1</sup>, Dyah Fauziah<sup>1</sup>, Anny Setijo Rahaju<sup>1</sup>, Isnin Anang Marhana<sup>2</sup>, Alfian Nur Rosyid<sup>2</sup>, Dwi Wahyu<sup>2</sup>, Gilang Muhammad Setyo Nugroho<sup>2</sup>, Adhatri Anggoro<sup>2</sup>, I Komang Rusgi Yandi<sup>2</sup>, Bambang Pujo Semedi<sup>3</sup>, Jilientasia Godrace Lilihata<sup>3</sup>, Umami Maimunah<sup>4</sup>, Supriadi<sup>4</sup>, Achmad Lefi<sup>5</sup>, Lalu Galih Pratama Rinjani<sup>5</sup>, Edi Suyanto<sup>6</sup>, Ricardo Ardian Nugraha<sup>6</sup>

<sup>1</sup>Department of Anatomical Pathology, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Surabaya, INDONESIA.  
<sup>2</sup>Department of Pulmonology and Respiratory Medicine, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Surabaya, INDONESIA.  
<sup>3</sup>Department of Anesthesiology and Reanimation, Faculty of Medicine, Universitas Airlangga University – Dr. Soetomo General Academic Hospital, Surabaya, INDONESIA.  
<sup>4</sup>Department of Internal Medicine, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Surabaya, INDONESIA.  
<sup>5</sup>Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Surabaya, INDONESIA.  
<sup>6</sup>Department of Forensics and Medicolegal Medicine, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Surabaya, INDONESIA.

## Correspondence

**Etty Hary Kusumastuti**  
Department of Anatomical Pathology, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Surabaya, INDONESIA.  
Email: ettyhary@fk.unair.ac.id

## History

- Submission Date: 08-10-2022;
- Review completed: 28-11-2022;
- Accepted Date: 05-12-2022.

DOI : 10.5530/pj.2022.14.184

Article Available online

<http://www.phcogj.com/v14/i6>

## Copyright

© 2022 Phcogj.Com. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.



## ABSTRACT

**Background:** COVID-19 has spread rapidly around the world. It is necessary to study lung tissue of post-mortem COVID-19 patients to determine the molecular alteration particularly the role of IL-6 and IL-17 in causing fatality. **Objective:** This study aims to determine the differences in the expressions of IL-6 and IL-17 in lung tissue of post-mortem COVID-19 patients compared to non-COVID-19 patients. This study also aimed to analyze the correlation between the expressions of IL-6 and IL-17 in lung tissue of post-mortem COVID-19 patients. **Methods:** This research is an observational analytic study with cross-sectional approach. The samples were 15 paraffin blocks of post-mortem lung tissue biopsy of COVID-19 patients, and 15 paraffin blocks of inflammatory lung tissue biopsy or surgery of non-COVID-19 patients. IL-6 and IL-17 expressions were evaluated by immunohistochemical procedure. **Result:** There was a significant difference in the expression of IL-6 in the COVID-19 group and the non-COVID-19 group with a p-value = 0.001 (p < 0.05). There was a significant difference in the expression of IL-17 in the COVID-19 group and the non-COVID-19 group with p-value = 0.001 (p < 0.05). There was a significant correlation between the expressions of IL-6 and IL-17 in the COVID-19 group, with the Spearman coefficient value (rs) of 0.548 with p = 0.034 (p < 0.05). **Conclusion:** There are differences in the expression of IL-6 and IL-17 between COVID-19 and non-COVID-19 lung tissue. There is a significant correlation between the expressions of IL-6 and IL-17 in post-mortem lung tissue of COVID-19 patients.

**Key words:** Biopsy, COVID-19, IL-6, IL-17, Post mortem lung tissue.

## INTRODUCTION

Since December 2019, Corona Virus Disease (COVID-19) caused by Severe Acute Respiratory Syndrome Corona Virus 2 (SARS CoV-2) has spread rapidly around the world. SARS-CoV2 is an RNA (ribonucleic acid) virus with a new variant belonging to the Coronaviridae family which was first described in February 2020.<sup>1</sup> The World Health Organization stated that until March 2021, the number of confirmed cases was 125 million with a death toll of 2.8 million. COVID-19 has also had a negative impact on the global economy and health system. Lack of knowledge regarding the characteristics and pathogenesis of the SARS-CoV-2 virus causes difficulties in controlling the course of the COVID-19 disease. It is necessary to conduct research on lung tissue of post-mortem COVID-19 patients to determine the molecular alteration that plays a role in causing fatality.

The pathogenesis of COVID-19 can generally be categorized into three successive and overlapping phases, namely the pulmonary phase, the proinflammatory phase and the prothrombotic phase.<sup>2</sup> In the proinflammatory phase, SARS-CoV-2 causes the release of proinflammatory cytokines in very large quantities that cause interstitial pneumonia and acute respiratory distress syndrome (ARDS). In a more advanced

state will develop sepsis and cytokine storm (cytokine storm) with hypercoagulability and multiorgan dysfunction.<sup>2,3</sup> Several types of cytokines that play a role in the pathogenesis of COVID-19 are IL-1 $\beta$ , IL-2, IL-4, IL-6, IL-8, IL12, IL-17, G-CSF, GM-CSF, IFN $\gamma$ , MCP-1 and TNF $\alpha$ . Cytokines IL-6 and IL-17 are known to play an important role and influence disease fatality, which levels of both are known to be elevated in blood.<sup>1,4-6</sup>

Interleukin-6 (IL-6) is one of the cytokines that is activated in the presence of infection or injury. IL-6 is secreted by various cell types and plays a role in the regulation of various physiological processes. Elevated IL-6 serum levels are associated with an inflammatory response, respiratory failure and affect mortality in COVID-19 patients.<sup>7-9</sup> Interleukin-17 (IL-17) is a cytokine synthesized by Th17 lymphocytes and is increased in inflammatory processes and autoimmune diseases. IL-17 is a proinflammatory cytokine that plays a role in tissue damage, physiological stress, and infection.<sup>5,10</sup>

A number of studies have been carried out through blood serum analysis to study the important role of IL-6 and IL-17 in the pathogenesis of COVID-19, but lack studies to analyze the role of IL-6 and IL-17 in lung tissue of patients with COVID-19. This study aims to determine the differences in the expression of IL-6 and IL-17 in lung tissue of post-mortem

**Cite this article:** Kusumastuti EH, Wiratama PA, Ariani G, Djuanda SN, Rahniayu A, Kurniasari N, et al. Differences in interleukin-6 and interleukin-17 expression in covid-19 post-mortem lung tissue biopsy compared with non-covid-19. Pharmacogn J. 2022;14(6)Suppl: 887-892.

44

COVID-19 patients compared to non-COVID-19 patients and to analyze the correlation between the expression of IL-6 and IL-17 in lung tissue of post-mortem COVID-19 patients.

## MATERIAL AND METHODS

The design of this research is an observational analytic with cross-sectional approach. This research is a branch of research titled 'Multi-organ pathological finding in COVID-19 infection through post-mortem core biopsy' and has received ethical approval from the Health Research Ethics Committee of Dr. Soetomo general hospital number 0022/KEPK/VII/2020.

The number of research samples is 30, consisting of 15 paraffin blocks of post-mortem pulmonary tissue biopsy of patients with COVID-19 and 15 paraffin blocks of pulmonary tissue biopsy or surgery of non-COVID-19 pneumonia patients. IL-6 expression was evaluated by an immunohistochemical procedure using IL-6 antibody GeneTex Antibodies Rabbit Polyclonal antibody, unconjugated (GTX17623). Antigen retrieval using pH 6, with a dilution of 1:100. IL-17 expression was also evaluated by an immunohistochemical procedure using anti-GeneTex Antibodies Rabbit Recombinant-Human IL-17 (IL-17A) polyclonal antibody, unconjugated (GTX31174) antibody, with a concentration of 0.25 g/ml. The expression of IL-6 was positive for brown color in the cytoplasm of lymphocytes and macrophages, while the expression of IL-17 was positive for brown color in the cytoplasm of lymphocyte cells. The assessment is carried out in five fields of view with high power fields. The number of positive cells is calculated as a percentage. The assessment was carried out by two pathologists in a double-blind manner and then an average of the two was made. Analysis of expression differences was done with independent T-test. Analysis of correlation was done with Spearman correlation test. Statistic result is significant if  $p < 0.05$ .

## RESULT

The data obtained in this study showed the sex distribution in the COVID-19 group, the number of subjects being male was 8 people (53%) and women were 7 people (47%). While in the non-COVID-19 group, the number of male subjects was 11 people (73%) and 4 women (27%).

Age data showed that in the COVID-19 group, the highest number was from the 41-50-year-old and 51-60-year age group, each of which amounted to 5 people (33.3%). Meanwhile, from the non-COVID-19 group, the highest number was in the age range of 61-70 years, namely 5 people (33.3%).

The results of this study showed the mean expression of IL-6 in the COVID-19 group was  $80.67\% \pm 7.03$ , while in the non-COVID-19 group it was  $21.50 \pm 11.60$ . The difference between the two was then assessed using the Independent T test. The results of statistical analysis showed a significant difference between the expression of IL-6 in the COVID-19 group and the non-COVID-19 group with  $p$  value = 0.001 ( $p < 0.05$ ).

The results of IL-6 expression in the COVID-19 and non-COVID-19 groups are presented in table 3. The average percentage of IL-6 expression in the COVID-19 group is higher than the non-COVID-19 group.

The results of this study showed the mean expression of IL-17 in the COVID-19 group was  $75.33\% \pm 14.93$ , while in the non-COVID-19 group it was  $45.67\% \pm 17.31$ . The difference between the two was then assessed using the Independent T test. The results of statistical analysis showed a significant difference between the expression of IL-17 in the COVID-19 group and the non-COVID-19 group with  $p$  value = 0.001 ( $p < 0.05$ ).

**Table 1: Distribution COVID-19 and non COVID-19 groups by gender.**

	COVID-19 (n)	Non COVID-19 (n)
Male	8	11
Female	7	4

**Table 2: Distribution COVID-19 and non-COVID-19 groups by age.**

	COVID-19		Non COVID-19	
	Number	Percentage	Number	Percentage
< 10 years	0	0%	1	6.67%
11-20 years	0	0%	1	6.67%
21-30 years	1	6.67%	0	0%
31-40 years	3	20%	1	6.67%
41-50 years	5	33.3%	3	20%
51-60 years	5	33.3%	3	20%
61-70 years	1	6.67%	5	33.3%
> 70 years	0	0%	1	6.67%
Total	15	100%	15	100%

**Table 3: Table of results of the assessment of IL-6 expression in the COVID-19 and non-COVID-19 groups.**

Groups	n	Average (%)	SD	Min	Max	p
COVID-19	15	80.67	7.03	70	90	0.001
Non COVID-19	15	21.50	11.60	7.5	55	

**Table 4: Table of results of the assessment of IL-17 expression in the COVID-19 and non-COVID-19 groups.**

Groups	n	Average (%)	SD	Min	Max	P
COVID-19	15	75.33	14.93	30	90	0.001
Non COVID-19	15	45.67	17.31	20	75	

**Table 5: Correlation of IL-6 and IL-17 expression in all samples.**

		IL-17 expression
IL-6 expression	$r_s$	0.779
	p	0.001
	n	30


**Table 6: Correlation of IL-6 and IL-17 expression in the COVID-19 patient group.**

		IL-17 expression
IL-6 expression	$r_s$	0.548
	p	0.034
	n	15


The results of IL-17 expression in the COVID-19 and non-COVID-19 groups are presented in table 4. The average percentage of IL-17 expression in the COVID-19 group is higher than the non-COVID-19 group. Figure 2 shows an overview of IL-17 expression in the COVID-19 and non-COVID-19 groups.

The correlation between the expression of IL-6 and IL-17 in lung tissue of COVID-19 and non-COVID-19 patients was statistically tested using the Spearman correlation test. The results of the analysis showed a significant correlation between the expression of IL-6 and IL-17 in all samples (COVID-19 and non-COVID-19 groups) with a Spearman correlation coefficient ( $r_s$ ) of 0.779 with  $p = 0.001$  ( $p < 0.05$ ), which means that the higher the expression of IL-6, the higher the expression of IL-17.

In the group of COVID-19 patients, a significant correlation was found between the expression of IL-6 and IL-17. The results of statistical analysis using the Spearman correlation test showed the Spearman coefficient ( $r_s$ ) of 0.548 with  $p = 0.034$  ( $p < 0.05$ ) which means that the



**Figure 1:** IL-6 expression in the COVID-19 and non-COVID-19 groups, magnification 400x. (a) IL-6 is positively expressed in lung tissue of COVID-19 patients (white arrow), (b) IL-6 is negatively expressed in lung tissue of non-COVID-19 patients (black arrow).



**Figure 2:** IL-17 expression in the COVID-19 and non-COVID-19 groups, magnification 400x. (a) IL-17 is positively expressed in lung tissue of COVID-19 patients (white arrow), (b) IL-17 is negatively expressed in lung tissue of non-COVID-19 patients (black arrow).

high expression of IL-6 in the tissue will be directly proportional to the expression of IL-17.

## DISCUSSION

The data obtained in this study shows the sex distribution of COVID-19 cases, where the incidence of COVID-19 in men (53%) and women (47%) is almost the same. The incidence of disease and fatality in males was found to be slightly higher than that of females. This is in line with WHO data where the percentage of male sufferers is more (58%) in the UK 57% and 51.1% in China.<sup>10-12</sup>

Almost the same figure was also found in the incidence of COVID 19 in Indonesia. Several studies have shown higher morbidity and mortality rates in men, namely 58.94%,<sup>13</sup> 56.5% with a fatality rate of 63.8%.<sup>14</sup> Several studies have linked this event to several factors including sex hormones, the X chromosome (playing a role in high innate and adaptive immunity) and high expression of ACE 2 receptors, and lifestyle. Men smoke and consume alcohol more often.<sup>15,16</sup>

The age range of the sample is between 5 and 88 years, with the mean age of the entire sample being 48.93 years. From these data, it was found that in the COVID-19 group, the highest number was from the 41-50-year-old and 51-60-year age group, each of which amounted to

5 people (33.3%). This is slightly different from several other studies where the highest fatality rate was found in the age range > 60 years, followed by the 40-59-year age group. While the smallest mortality rate was obtained from the age group < 10 years.<sup>11,14</sup> The high mortality rate in old age and the number of comorbid diseases that aggravate the course of COVID-19. In this study, the highest number was from the age group of 41-50 years and 51-60 years. This is different from other studies because this study did not assess the proportion of fatalities in each age group.

In addition to old age and male gender, several other risk factors that play a role in increasing mortality rates in COVID patients are the presence of co-morbidities such as diabetes, cardiovascular, cerebrovascular and kidney diseases, and obesity.<sup>15,17</sup> Several socioeconomic conditions can also play a role, for example lack of socioeconomic conditions, high levels of air pollution and diurnal temperature variations.<sup>17</sup>

The data obtained from this study showed that the mean expression of IL-6 in the COVID-19 group was significantly higher than in the non-COVID-19 group. Previous studies have also shown high expression of IL-6 in lung tissue of COVID-19 patients, and is a predictor of fatality and disease progression.<sup>7,9,18,19</sup>

Interleukin-6 is a pleiotropic cytokine that is activated in the presence of infection or injury, one of which is a viral infection. Interleukin-6 is secreted by various cell types (eg: T lymphocytes, macrophages, and fibroblasts) and plays a role in the regulation of various physiological processes.<sup>9</sup> The role of IL-6 in the inflammatory process including COVID infection is like a double-edged sword, because it has two effects, namely pro- and anti-inflammatory. There are two pathways in the IL-6 signaling process, namely the classical pathway and the IL-6 signal trans pathway. The classical pathway is considered the anti-inflammatory pathway whereas the signal trans pathway is the pro-inflammatory pathway. Therefore, in various inflammatory conditions and autoimmune diseases, the role of the IL-6 signaling trans pathway is that inhibiting this process is needed to relieve inflammation and damage that occurs.<sup>20</sup> One of the roles of IL-6 as part of the pro-inflammatory function is in the activation and differentiation of Th17, and induces the expression of granzyme B and perforin on CD8 T lymphocytes that can eliminate the virus. Activation of these cytotoxic T cells will stimulate the release of TNF- $\alpha$  and IFN- $\gamma$  which causes neutrophil migration. In addition, IL-6 also plays a role in thermoregulation (as an endogenous pyrogenic cytokine) and amplifies the immune response to viral infections. The IL-6 signal is known to induce the production of IL-27 by monocytes and macrophages in the respiratory tract, thereby stimulating T cell maturation in the lungs.<sup>20</sup> Interleukin 6 has also been shown to be associated with inflammatory responses in the acute phase, including CRP and LDH.<sup>21</sup> On the other hand, the anti-inflammatory function of IL-6 plays a role in the progression of viral infection, thereby causing viral persistence in the host. Interleukin-6 stimulates the Th2 response while inhibiting Th1 proliferation through two mechanisms. First, by stimulating CD4 T cells to secrete IL-4 which will stimulate Th2 and suppress the production of IFN- $\gamma$  by CD4 T cells and cause Th1 polarization (antiviral response). Interleukin-6 will cause Th17 differentiation which will then produce IL-17 which will increase the expression of anti-apoptotic molecules so that the virus is stronger and can survive.<sup>19</sup> A study proved that increased expression of IL-6 in lung tissue indicates the severity of COVID-19 sufferers.<sup>22</sup>

Interleukin-17 plays a role in the pathogenesis of ARDS by increasing the infiltration of neutrophils into the lung. Increased signaling of IL-17A and Th17 or regulatory T cells (Treg) is known to be positively correlated with disease severity in MERS infection, where the cause is a type of virus that belongs to the same family as MERS. The overexpression of Th17 cells and the excessive cytotoxic effect of CD8+ T cells will cause tissue damage in COVID-19 patients with pneumonia. Like IL-6, high expression of IL-17A in lung tissue is also directly proportional to lung inflammation and poor outcomes (ARDS) and multi-organ dysfunction.<sup>23,24</sup>

Various studies have shown the important role of IL-17A as a regulator of PMN infiltration. IL-17 is known to play a role in neutrophil maturation and differentiation by increasing the release of granulocyte-colony stimulating factor (G-CSF). This situation will cause the differentiation of hematopoietic progenitor cells CD34 into neutrophils. In addition, IL-17 can stimulate granulopoiesis markers and other chemokines, such as Growth-regulated Oncogene- $\alpha$  (GRO- $\alpha$ ) which will regulate neutrophil penetration into tissues. Interleukin-17 also plays a role in the release of cytokines and other inflammatory chemokines such as IL-1, IL-6, TNF- $\alpha$ , macrophage inflammatory protein-2 (MIP-2), IL-8, Interferon-inducible protein-10 (IP-10). All these molecules will be used by neutrophils in the process of chemotaxis.<sup>25</sup>

According to Hou *et al.*, IL-17 can also increase the expression of antiapoptotic molecules that increase the survival of virus-infected cells and block the process of destruction by cytotoxic T cells.<sup>26</sup> Therefore, the role of IL-17 in COVID inflammation is important in amplification of the inflammatory process.<sup>25</sup>

The results of this study showed that the mean expression of IL-17 in the COVID-19 group was significantly higher than the non-COVID-19 group. Several previous studies showed high IL-17 expression in lung tissue of COVID-19 patients and was a predictor of fatality and disease progression.<sup>23-27</sup> A similar study conducted by Avezedo *et al.*, which assessed the expression of IL-17 and IL-8 in post-mortem lung tissue of patients with COVID-19 also showed results that were in line with this study.<sup>28</sup>

The results of the analysis of this study showed that there was a positive and significant correlation between the expression of IL-6 and IL-17 in all samples (COVID-19 and non-COVID-19 groups) as well as when analyzed only in the COVID-19 group. The COVID-19 group showed a higher correlation. This phenomenon can be seen in general that the pathogenesis of COVID-19 may be different from other infectious processes. In COVID-19, there is an exaggerated immune response that is characteristic of this disease. The state of hypercytokinemia or what is called a cytokine storm is a collection of symptoms caused by the activation of T lymphocytes, macrophages and the release of various cytokines which then activate other immune cells. In the laboratory, cytokine storm is characterized by an increase in various proinflammatory cytokines such as IL-1, IL-6, IL-12, IFN- $\gamma$ . Measurement of these levels, especially IL-6 is considered useful as a predictor of fatality rates and the need for respiratory equipment, and can be used as a potential therapeutic target.<sup>7,9,19</sup>

The SARS-CoV2 virus which is the etiology of COVID-19 has a surface protein, namely the surface spike (S) protein, which will bind to the Angiotensin Converting Enzyme-2 (ACE-2) receptor found in pulmonary alveolar epithelial cells to enter the body. The ACE-2 receptor is very important in viral virulence, because it has been shown that cells that do not have this receptor are resistant to viral infection. Angiotensin Converting Enzyme-2 is not only expressed in the respiratory tract, but also in the small intestine, pancreas, kidney, heart, esophagus, bladder and brain. This has led to multi-organ involvement in COVID-19. Under normal conditions, ACE-2 functions to cause the conversion of angiotensin II to angiotensin 1-7 in the renin-angiotensin-aldosterone system (RAA system). Various studies have reported increased levels of Angiotensin II in COVID-19 patients compared to healthy individuals, which is associated with viral load and lung damage. This situation is caused by the virus binding to ACE-2, so that Angiotensin II cannot be converted, accumulates and causes an imbalance in the RAA system. Angiotensin II buildup will cause a change in equilibrium towards a proinflammatory state.<sup>19</sup> In the absence of ACE-2, angiotensin II will bind to the angiotensin I receptor which will cause an oxidative stress cascade, resulting in the formation of Reactive Oxygen Species (ROS) and an increase in IL-6 expression. On the other hand, the increased expression of IL-6 will also increase the expression of angiotensin I receptors and attach to endothelial cell walls and cause cellular and vascular inflammation. Interleukin-6 and angiotensin II will cause an increase in the expression of each other continuously so that over time it will aggravate the state of oxidative stress.<sup>19</sup>

The role of IL-6 in inflammatory processes including COVID infection is like a double-edged sword, because it has two effects, namely pro- and anti-inflammatory.<sup>9,19</sup> In pro-inflammatory, one of the functions of IL-6 is in the activation and differentiation of Th17 and the induction of granzyme B and perforin in CD8 T lymphocyte cells. Interleukin-6 causes the production of IL-17 which will further increase the expression of antiapoptosis molecules leading to increasing the survival of virus-infected cells and blocking the process of digestion by cytotoxic T cells.<sup>26</sup> Interleukin-17 mainly causes neutrophil migration to the lungs thus causing tissue damage during the inflammatory process.<sup>19</sup> Interleukin-17 also plays a role in the release of cytokines and other inflammatory chemokines such as IL-1, IL-6, TNF- $\alpha$ , macrophage

inflammatory protein-2 (MIP-2), IL-8, Interferon-inducible protein-10 (IP-10).<sup>19</sup>

The synergistic interaction between IL-17A and IL-6 plays an important role in the formation of pulmonary fibrosis and imbalances of the respiration system. IL-6 not only plays a role in the differentiation of T helper (Th) cells into Th17 cells through ROR $\gamma$ t which is the JAK-STAT3 pathway but also in the process of pulmonary fibrosis (causing collagen deposition) due to the function of epithelial cells and fibroblasts that are aberrant (works synergistically with IL-8).<sup>6,29</sup>

## CONCLUSION

There are significant differences between IL-6 and IL-17, which are higher expressed in the lung tissue of covid-19 patients than the lung tissue of the non-COVID-19 group. There was also a strong correlation between IL-6 and IL-17 in the lung tissue of COVID-19 patients, as well as in the entire sample. Although not a single factor, IL-6 and IL-17 play a role in the pathogenesis of COVID-19.

## ACKNOWLEDGEMENT

The authors are grateful for support from the director and research and development unit Dr. Soetomo General Academic Hospital.

## ETHICS AND CONSENT

This research is a branch of research titled 'Multi-organ pathological finding in COVID-19 infection through post-mortem core biopsy' and has received ethical approval from the Health Research Ethics Committee of Dr. Soetomo general hospital number 0022/KEPK/VII/2020.


## CONFLICTS OF INTEREST/PLAGIARISM REPORT

There are no conflicts of interest in this research.

## REFERENCES

- Zhou P, Yang XL, Wang XG. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*. 2020;579(7798):270-3.
- Lee C, Choi WJ. Overview of COVID-19 inflammatory pathogenesis from the therapeutic perspective. *Arch Pharm Res*. 2021;44(11):99-116.
- Maiese A, Manetti AC, La Russa R. Autopsy findings in COVID-19-related deaths: a literature review. *Forensic Sci Med Pathol*. 2021;17(2):279-96.
- Mendoza VM. Interleukin-17: A potential therapeutic target in COVID-19. *J Infection*. 2020;81(2):136-8.
- Costela-Ruiz VJ, Illescas-Montes R, Puerta-Puerta JM, Ruiz C, Melguizo-Rodríguez L. SARS-CoV-2 infection: The role of cytokines in COVID-19 disease. *Cytokine Growth Factor Rev*. 2020;54:62-75.
- Shibabaw T. Inflammatory Cytokine: IL-17A Signaling Pathway in Patients Present with COVID-19 and Current Treatment Strategy. *J Inflamm Res*. 2020;13(3):673-80.
- Herold T, Jurinovic V, Amreich C. Elevated levels of IL-6 and CRP predict the need for mechanical ventilation in COVID-19. *J Allergy Clin Immunol*. 2020;146(1):128-36.
- Zhang ZL, Hou YL, Li DT, Li FZ. Laboratory findings of COVID-19: a systematic review and meta-analysis. *Scand J Clin Lab Invest*. 2020;80(6):441-7.
- Santa Cruz A, Mendes-Frias A, Oliveira AI. Interleukin-6 Is a Biomarker for the Development of Fatal Severe Acute Respiratory Syndrome Coronavirus 2 Pneumonia. *Front Immunol*. 2021;12(1):613422.
- Fadlallah S, Sham Eddin MS, Rahal EA. IL-17A in COVID-19 Cases: a meta-analysis. *J Infect Dev Ctries*. 2021;15(11):1630-9.
- Brown AE, Heinsbroek E, Kall MM. Epidemiology of Confirmed COVID-19 Deaths in Adults, England, March-December 2020. *Emerg Infect Dis*. 2021;27(5):1468-71.
- Park SE. Epidemiology, virology, and clinical features of severe acute respiratory syndrome -coronavirus-2 (SARS-CoV-2; Coronavirus Disease-19). *Clin Exp Pediatr*. 2020;63(4):119-24.
- Sutaryono, Andasari SD, Kasjono HS. Diagnosis and epidemiology of Coronavirus (COVID-19) outbreak in Indonesia. *J Teknologi Laboratorium*. 2020;9(1):49-57.
- Hikmawati I, Setiyabudi R. Epidemiology of COVID-19 in Indonesia: common source and propagated source as a cause for outbreaks. *J Infect Dev Ctries*. 2021;15(5):646-52.
- Noor AU, Maqbool F, Bhatti ZA, Khan AU. Epidemiology of COVID-19 Pandemic: Recovery and mortality ratio around the globe. *Pak J Med Sci*. 2020;36(COVID19-S4):S79-84.
- Bwire GM. Coronavirus: why men are more vulnerable to Covid-19 than women? *SN Compr Clin Med*. 2020;2(7):874-6.
- Cao Y, Hiyoshi A, Montgomery S. COVID-19 case-fatality rate and demographic and socioeconomic influencers: worldwide spatial regression analysis based on country-level data. *BMJ Open*. 2020;10(11):e043560.
- Smetana K, Brabek J. Role of interleukin-6 in lung complications in patients with COVID-19: therapeutic implications. *In Vivo*. 2020;34(3 Suppl):1589-92.
- Shekhawat J, Gauba K, Gupta S. Interleukin-6 Perpetrator of the COVID-19 Cytokine Storm. *Indian J Clin Biochem*. 2021;36(4):440-50.
- Su H, Lei CT, Zhang C. Interleukin-6 Signaling Pathway and Its Role in Kidney Disease: An Update. *Front Immunol*. 2017;8:405.
- Adiatmaja CO, Nugraha J, Utari A. Correlation between Interleukin-6, CRP, and LDH in COVID-19 patients of Dr. Soetomo Teaching Hospital, Surabaya, Indonesia. *Res J Pharm Technol*. 2022;15(10):4753-7.
- Nugroho GMS, Marhana IA, Kusumastuti EH, Semedi BP, Maimunah U, Lefi A, *et al.* Interleukin-6 (IL-6) expression of lung tissue in COVID-19 patient severity through core biopsy post mortem. *Ann Med Surg*. 2022;82(2022):104648.
- Kang YW, Lee SC, Jeon SM, Jo EK. Roles of interleukin-17 and Th17 responses in COVID-19. *J Bacteriol Virol*. 2021;51(3):89-102.
- Feng CM, Wang XM, Li MD. Serum interleukin-17 predicts severity and prognosis in patients with community acquired pneumonia: a prospective cohort study. *BMC Pulm Med*. 2021;21(1):393.
- Maione F, Casillo GM, Raucci F. Interleukin-17A (IL-17A): A silent amplifier of COVID-19. *Biomed Pharmacother*. 2021;142:111980.
- Hou W, Kang HS, Kim BS. Th17 cells enhance viral persistence and inhibit T cell cytotoxicity in a model of chronic virus infection. *J Exp Med*. 2009;206(2):313-28.
- Pourgholaminejad A, Pahlavanneshan S, Basiri M. COVID-19 immunopathology with emphasis on Th17 response and cell-based immunomodulation therapy: Potential targets and challenges. *Scand J Immunol*. 2022;95(2):e13131.
- Azevedo MLV, Zanchettin AC, Vaz de Paula CB. Lung Neutrophilic Recruitment and IL-8/IL-17A Tissue Expression in COVID-19. *Front Immunol*. 2021;12:656350.
- Darif D, Hammi I, Kihel A, El Idrissi Saik I, Guessous F, Akarid K. The pro-inflammatory cytokines in COVID-19 pathogenesis: What goes wrong? *Microb Pathog*. 2021;153:104799.

## GRAPHICAL ABSTRACT



## ABOUT AUTHORS



Etty Hary Kusumastuti is a Pathologist at Dr. Soetomo General Academic Hospital and as a lecturer in Department of Anatomical Pathology, Faculty of Medicine, Universitas Airlangga, Indonesia. She completed her residency in Anatomical Pathology at the Faculty of Medicine of Universitas Airlangga, Indonesia. She graduated doctoral degree in clinical medicine at the Faculty of Medicine of Universitas Airlangga, Indonesia. She is one of the most expert in cytopathology of respiratory tract. She has participated in many conferences, workshops, and courses as a speaker and participant. She writes a number of scientific papers on various subjects of pathology.



Isnin Anang Marhana is an interventional pulmonologist at Dr. Soetomo General Academic Hospital, as well as a lecturer in Pulmonology and Respiratory Medicine, Faculty of Medicine, Universitas Airlangga, Indonesia. After graduating as a resident of pulmonology at Faculty of Medicine, Universitas Airlangga, he then underwent interventional bronchoscopy training in St. Marianna University, Kawasaki, Japan. He is a researcher and the author of a number of studies in the field of lungs and especially COVID19.

**Cite this article:** Kusumastuti EH, Wiratama PA, Ariani G, Djuanda SN, Rahniayu A, Kurniasari N, et al. Differences in interleukin-6 and interleukin-17 expression in covid-19 post-mortem lung tissue biopsy compared with non-covid-19. *Pharmacogn J.* 2022;14(6) Suppl: 887-892.

## 20. Differences in interleukin-6 and interleukin-17 expression

### ORIGINALITY REPORT

15%

SIMILARITY INDEX

12%

INTERNET SOURCES

11%

PUBLICATIONS

0%

STUDENT PAPERS

### PRIMARY SOURCES

1	<a href="http://rebus.us.edu.pl">rebus.us.edu.pl</a> Internet Source	<1 %
2	Mehmet Hursitoglu, Nilgun Isiksacan, Betul Erismis, Faruk Karandere, Alev Kural, A.Baki Kumbasar, Kadriye Kart Yasar. "In-vitro cytokine production and nasopharyngeal microbiota composition in the early stage of COVID-19 infection", Cytokine, 2021 Publication	<1 %
3	<a href="http://www.gastrores.org">www.gastrores.org</a> Internet Source	<1 %
4	<a href="http://www.isindexing.com">www.isindexing.com</a> Internet Source	<1 %
5	<a href="http://academic.oup.com">academic.oup.com</a> Internet Source	<1 %
6	<a href="http://jnsbm.com">jnsbm.com</a> Internet Source	<1 %
7	Henrry Diaz Londres, Jorge Jiménez Armada, Aray Hernández Martínez, Anselmo A Abdo Cuza et al. "Blocking EGFR with nimotuzumab:	<1 %



a novel strategy for COVID-19 treatment",  
Immunotherapy, 2022

Publication

8

[digitallibrary.usc.edu](https://digitallibrary.usc.edu)

Internet Source

<1 %

9

[search.bvsalud.org](https://search.bvsalud.org)

Internet Source

<1 %

10

Jyoti Kanwar Shekhawat, Mithu Banerjee.  
"Role of breath biopsy in COVID-19", The  
Journal of Applied Laboratory Medicine, 2022

Publication

<1 %

11

[rjoas.com](https://rjoas.com)

Internet Source

<1 %

12

Arunkumar Krishnan, James P. Hamilton,  
Saleh A. Alqahtani, Tinsay A.Woreta. "A  
narrative review of coronavirus disease 2019  
(COVID-19): clinical, epidemiological  
characteristics, and systemic manifestations",  
Internal and Emergency Medicine, 2021

Publication

<1 %

13

Zoia R. Korobova, Natalia A. Arsentieva,  
Natalia E. Liubimova, Oleg K. Batsunov et al.  
"Cytokine Profiling in Different SARS-CoV-2  
Genetic Variants", International Journal of  
Molecular Sciences, 2022

Publication

<1 %

14 Razi Ullah, Jadoon Khan, Nosheen Basharat, Danqun Huo, Ahmad Ud Din, Guixue Wang. "Evaluation of Cardiac Biomarkers and Expression Analysis of IL-1, IL-6, IL-10, IL-17, and IL-25 among COVID-19 Patients from Pakistan", *Viruses*, 2022  
Publication

---

15 [humaniora.journal.ugm.ac.id](http://humaniora.journal.ugm.ac.id)  
Internet Source

---

16 [repositori.usu.ac.id](http://repositori.usu.ac.id)  
Internet Source

---

17 [www.coursehero.com](http://www.coursehero.com)  
Internet Source

---

18 Abraham P Bautista. "Impact of alcohol on the ability of Kupffer cells to produce chemokines and its role in alcoholic liver disease", *Journal of Gastroenterology and Hepatology*, 2001  
Publication

---

19 [covid19-data.nist.gov](http://covid19-data.nist.gov)  
Internet Source

---

20 [e-journal.unair.ac.id](http://e-journal.unair.ac.id)  
Internet Source

---

21 [www.medrxiv.org](http://www.medrxiv.org)  
Internet Source

---

22 [go.gale.com](http://go.gale.com)  
Internet Source

---

23	<a href="http://japsonline.com">japsonline.com</a> Internet Source	<1 %
24	<a href="http://jurnalrespirologi.org">jurnalrespirologi.org</a> Internet Source	<1 %
25	<a href="http://www.longdom.org">www.longdom.org</a> Internet Source	<1 %
26	<a href="http://www2.mdpi.com">www2.mdpi.com</a> Internet Source	<1 %
27	"Wednesday July 5, 2006 9:30?11:30 Hall 5A Main Session Brain maturation in epilepsy evolution", <i>Epilepsia</i> , 12/2006 Publication	<1 %
28	<a href="http://cdn.publisher.gn1.link">cdn.publisher.gn1.link</a> Internet Source	<1 %
29	<a href="http://link.springer.com">link.springer.com</a> Internet Source	<1 %
30	<a href="http://publication.babcock.edu.ng">publication.babcock.edu.ng</a> Internet Source	<1 %
31	<a href="http://www.ijcmas.com">www.ijcmas.com</a> Internet Source	<1 %
32	<a href="http://www.thieme-connect.com">www.thieme-connect.com</a> Internet Source	<1 %
33	Kun-Chang Lin, Chun-Chieh Wang, Wei-Chun Huang and Juey-Jen Hwang. "Considerations	<1 %

When Managing Heart Failure during the COVID-19 Pandemic-Consensus from the Taiwan Society of Cardiology", Acta Cardiologica Sinica, 2021

Publication

34

MH Sohn, SY Noh, W Chang, KM Shin, DS Kim. "Circulating interleukin 17 is increased in the acute stage of Kawasaki disease", Scandinavian Journal of Rheumatology, 2009

Publication

<1 %

35

[congress-ph.ru](http://congress-ph.ru)

Internet Source

<1 %

36

[hmj.hums.ac.ir](http://hmj.hums.ac.ir)

Internet Source

<1 %

37

[immunetwork.org](http://immunetwork.org)

Internet Source

<1 %

38

[mdpi-res.com](http://mdpi-res.com)

Internet Source

<1 %

39

[myemail.constantcontact.com](http://myemail.constantcontact.com)

Internet Source

<1 %

40

[online-journal.unja.ac.id](http://online-journal.unja.ac.id)

Internet Source

<1 %

41

[staging-mhealth.jmir.org](http://staging-mhealth.jmir.org)

Internet Source

<1 %

42

[www.dovepress.com](http://www.dovepress.com)

Internet Source

<1 %

---

43

[www.europeanproceedings.com](http://www.europeanproceedings.com)

Internet Source

<1 %

---

44

[www.jelsciences.com](http://www.jelsciences.com)

Internet Source

<1 %

---

45

[www.srpiskiarhiv.rs](http://www.srpiskiarhiv.rs)

Internet Source

<1 %

---

46

[www.wjgnet.com](http://www.wjgnet.com)

Internet Source

<1 %

---

47

Amita Jain, Nidhi Pandey, R. K. Garg, Rashmi Kumar. "IL-17 Level in Patients with Dengue Virus Infection & its Association with Severity of Illness", Journal of Clinical Immunology, 2012

Publication

<1 %

---

48

Aziz Rodan Sarohan, Murat Kızıl, Ahmet Çağkan İnkaya, Shokhan Mahmud, Muhammad Akram, Osman Cen. "A novel hypothesis for COVID-19 pathogenesis: Retinol depletion and retinoid signaling disorder", Cellular Signalling, 2021

Publication

<1 %

---

49

Heather Amsden, Olena Kourko, Madison Roth, Katrina Gee. "Antiviral Activities of Interleukin-27: A Partner for Interferons?", Frontiers in Immunology, 2022

Publication

<1 %

---

50	<p>Manal M. El-Desoky, Samar Tharwat, Nora Mostafa, Asem A. Hewidy et al. "Association of Interleukin-17F Polymorphism and Mortality Predictors with the Risk of COVID-19", International Journal of Clinical Practice, 2022</p> <p>Publication</p>	<1 %
51	<p>Sylvain Perruche, Pin Zhang, Takashi Maruyama, Jeffrey A. Bluestone, Philippe Saas, WanJun Chen. "Lethal Effect of CD3-Specific Antibody in Mice Deficient in TGF-<math>\beta</math>1 by Uncontrolled Flu-Like Syndrome", The Journal of Immunology, 2009</p> <p>Publication</p>	<1 %
52	<p><a href="http://bc.wydawnictwo-tygiel.pl">bc.wydawnictwo-tygiel.pl</a></p> <p>Internet Source</p>	<1 %
53	<p><a href="http://plus.mcmaster.ca">plus.mcmaster.ca</a></p> <p>Internet Source</p>	<1 %
54	<p><a href="http://wjgnet.com">wjgnet.com</a></p> <p>Internet Source</p>	<1 %
55	<p><a href="http://www.cyprusjmedsci.com">www.cyprusjmedsci.com</a></p> <p>Internet Source</p>	<1 %
56	<p><a href="http://www.daten-quadrat.de">www.daten-quadrat.de</a></p> <p>Internet Source</p>	<1 %
57	<p><a href="http://www.e-cep.org">www.e-cep.org</a></p> <p>Internet Source</p>	<1 %

58

[www.frontiersin.org](http://www.frontiersin.org)

Internet Source

<1 %

---

59

[www.iberreamjmed.com](http://www.iberreamjmed.com)

Internet Source

<1 %

---

60

[www.medsci.org](http://www.medsci.org)

Internet Source

<1 %

---

61

[www.researchsquare.com](http://www.researchsquare.com)

Internet Source

<1 %

---

62

Brigitta U. Mueller, Phillip A. Pizzo. "Chapter 8 Cytokines and biological response modifiers in the treatment of infection", Springer Science and Business Media LLC, 1998

Publication

<1 %

---

63

Radhakrishnan Vishnubalaji, Hibah Shaath, Nehad M. Alajez. "Protein Coding and Long Noncoding RNA (lncRNA) Transcriptional Landscape in SARS-CoV-2 Infected Bronchial Epithelial Cells Highlight a Role for Interferon and Inflammatory Response", Genes, 2020

Publication

<1 %

---

64

Samir Awasthi, Tyler Wagner, Aj Venkatakrishnan, Arjun Puranik et al. "Plasma IL-6 Levels following Corticosteroid Therapy as an Indicator of ICU Length of Stay in Critically ill COVID-19 Patients", Cold Spring Harbor Laboratory, 2020

Publication

<1 %

---

65

Elena Campione, Terenzio Cosio, Luigi Rosa, Caterina Lanna, Stefano Di Girolamo, Roberta Gaziano, Piera Valenti, Luca Bianchi.

"Lactoferrin as Protective Natural Barrier of Respiratory and Intestinal Mucosa against Coronavirus Infection and Inflammation", International Journal of Molecular Sciences, 2020

Publication

<1 %

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On



# 20. Differences in interleukin-6 and interleukin-17 expression

---

GRADEMARK REPORT

---

FINAL GRADE

**/100**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---