

Uterine conservative–resective surgery for selected placenta

by Rozi Adhitya Aryananda

Submission date: 27-Apr-2023 09:12PM (UTC+0800)

Submission ID: 2077138619

File name: Uterine_conservative_resective_surgery_for_selected_placenta.pdf (974.92K)

Word count: 6158

Character count: 32947

Uterine conservative–resective surgery for selected placenta accreta spectrum cases: Surgical–vascular control methods

Rozi Aditya Aryananda¹ | Aditiawarman Aditiawarman¹ | Khanisyah Erza Gumilar¹ | Manggala Pasca Wardhana¹ | M. Ilham Aldika Akbar¹ | Nareswari Cininta¹ | Ernawati Ernawati¹ | Budi Wicaksono¹ | Hermanto Tri Joewono¹ | Erry Gumilar Dachlan¹ | Citra Aulia Bachtiar¹ | Devita Kurniawati¹ | Dian Puspita Virdayanti¹ | Grace Ariani² | Gustaaf Albert Dekker^{1,3} | Agus Sulistyono¹

¹Maternal–Fetal Medicine Division, Obstetrics & Gynecology Department, Dr. Soetomo Academic General Hospital, Universitas Airlangga, Surabaya, Indonesia

²Anatomical Pathology Department, Dr. Soetomo General Hospital, Universitas Airlangga, Surabaya, Indonesia

³Women & Children's Division, Department of Obstetrics and Gynecology, Lyell McEwin Hospital, University of Adelaide, Adelaide, Australia

Correspondence

Rozi Aditya Aryananda, Maternal–Fetal Medicine, Department of Obstetrics & Gynecology, Dr Soetomo Academic General Hospital, Universitas Airlangga, Surabaya, Indonesia.
Email: rozi.odi@gmail.com

Abstract

Introduction: The incidence of placenta accreta spectrum (PAS) has increased, but the optimal management and the optimal way to achieve vascular control are still controversial. This study aims to compare maternal outcomes between different methods of vascular control in surgical PAS management.

Material and methods: A retrospective cohort study on consecutive cases diagnosed with PAS between 2013 and 2020 in single tertiary hospital. The final diagnosis of PAS was made following preoperative ultrasound and confirmation during surgery. Management of PAS using cesarean hysterectomy with internal iliac artery ligation (IIAL) was compared with two types of vascular control in uterine conservative–resective surgery (IIAL vs identification–ligation of the upper vesical, upper vaginal, and uterine arteries).

Results: Over an 8-year period, 234 pregnant women were diagnosed with PAS meeting the inclusion criteria. Uterine conservative–resective surgery (200 cases) was associated with lower mean blood loss compared with cesarean hysterectomy with IIAL (34 cases) in all PAS cases (1379 ± 769 mL vs 3168 ± 1916 mL; $p < 0.001$). In sub-analysis of the two uterine conservative–resective surgery subgroups, the group with identification–ligation of the upper vesical, upper vaginal, and uterine arteries had a significantly lower blood loss compared with uterine conservative–resective surgery with IIAL (1307 ± 743 mL vs 1701 ± 813 mL; $p = 0.005$). Women in the hysterectomy with IIAL group had more massive transfusion (35.3% vs 2.5%; $p < 0.001$; odds ratio [OR] 21.3, 95% confidence interval [CI] 6.9–66), major blood loss (>1500 mL) (70.6% vs 34%, $p < 0.001$; OR 4.7; 95% CI 2.1–10.3), catastrophic blood loss (>2500 mL) (64.7% vs 12.5%; $p < 0.001$; OR 12.8, 95% CI 5.7–29.1), other complications (32% vs

Abbreviations: IIAL, internal iliac artery ligation; PAS, Placenta Accreta Spectrum.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2022 The Authors. *Acta Obstetrica et Gynecologica Scandinavica* published by John Wiley & Sons Ltd on behalf of Nordic Federation of Societies of Obstetrics and Gynecology (NFOG).

12.4%; $p = 0.007$; OR 3.4, 95% CI 1.5–7.7), and intensive care unit admission (32.4% vs 1.5%; $p < 0.001$; OR 31.4, 95% CI 8.2–120.7) compared with the uterine conservative–resective surgery groups. The identification–ligation of the upper vesical, upper vaginal and uterine arteries had a significant lower risk for major blood loss (30.5% vs 50%; $p = 0.041$; OR 0.44, 95% CI = 0.2–0.9) compared with IIAL for vascular control of uterine conservative–resective surgery.

Conclusions: Cesarean hysterectomy is not the default treatment for PAS, PAS with invasion above the vesical trigone are suitable for uterine conservative–resective surgery with upper vesical, upper vaginal and uterine artery vascular control.

24 KEYWORDS

placenta accreta spectrum, uterine conservative–resective surgery, vascular control

19

1 | INTRODUCTION

Placenta accreta spectrum (PAS) disorders are rapidly becoming one of most common serious potentially life-threatening maternal perinatal problems. The incidence of PAS has increased significantly over the last decades and it is now estimated to occur in one in 533 deliveries;¹ this increase has followed the increase of cesarean scar pregnancy as the precursor of PAS from 1:2216 to 1:1800.² PAS may lead to massive intractable hemorrhage, disseminated intravascular coagulopathy, adult respiratory distress syndrome, massive blood transfusion, electrolyte imbalance, and renal failure.³ The three main types of PAS management discussed in the literature include total hysterectomy, leaving the placenta in situ, and uterine conservative–resective surgery, but the optimal surgical management of PAS is still controversial.^{4–6}

The American College of Obstetricians and Gynecologists recommends cesarean hysterectomy as the main management for PAS.⁷ Cesarean hysterectomy is a life-saving procedure, associated with major risks for complications like injury to ureters, bladder, bowel, or neurovascular structures, and permanent loss of fertility with its associated psychological insult. PAS accounts for 38% of pregnant women requiring peripartum hysterectomy.⁸ Maternal mortality may occur despite adequate planning, multidisciplinary management, and appropriate transfusion management.⁹ Fertility preservation is desirable in many cases because of the relatively young age of the PAS pregnant women. The first described method to preserve the uterus was leaving the placenta in situ. This method requires the woman to stay in the vicinity of the hospital for several months of monitoring due to the high risk of sudden severe hemorrhage.¹⁰ Some experts tried to develop several conservative–resective approaches in managing PAS for fertility preservation.¹¹ Palacios-Jaraquemada et al reported the first study about uterine conservative–resective surgery in 68 cases, 10 women had a subsequent pregnancy.⁴ Palacios-Jaraquemada et al described that pregnant women with placental invasion above the bladder trigone are particularly suitable for uterine conservative–resective surgery.^{4,12} Chandrharan et al reported a successful approach for avoiding peripartum hysterectomy in four women with PAS using the triple P procedure (Perioperative

Key message

Cesarean hysterectomy is not the default treatment for PAS. Internal iliac artery ligation is not the best option for vascular control in PAS management. Identification and ligation of the upper vesical, upper vaginal and uterine arteries are the key procedures reducing the blood loss during PAS management.

ultrasound—Pelvic de-vascularization using internal iliac artery balloon catheter—Placental non-separation with myometrial excision),⁵ but this method is difficult to apply in many hospitals, particularly in many low-income countries.

As part of the uterine conservative–resective surgical procedure, two methods of vascular control have been used: internal iliac artery ligation (IIAL) and the other focusing on vascular control of the vesical–vaginal–uterine arteries.

Dr. Soetomo Hospital in Surabaya is the main referral hospital for East Java (population about 38 million). Over the past decade we have witnessed a great increase in pregnant women presenting with PAS.

This has allowed us to build on the uterine conservative–resective surgical management initially championed by Palacios-Jaraquemada et al.⁴ The aim of this large retrospective cohort study was to compare the maternal outcomes between two methods of vascular control used during the current uterine conservative–resective surgical approach of women presenting with PAS with the initial approach consisting of hysterectomy plus IIAL.

11

2 | MATERIAL AND METHODS

This is a retrospective cohort study, using medical, surgical, and pathological reports and ultrasound images on consecutive pregnant

women diagnosed with PAS from 2013 to 2020 in the main tertiary referral hospital in East Java. All women were followed for at least 6 month after surgery.

Inclusion criteria were based on the final preoperative diagnosis following preoperative ultrasound imaging consistent with PAS and on intraoperative confirmation during surgery.¹³⁻¹⁵ All pregnant women were selected on the basis of specific preoperative ultrasound imaging and intraoperative staging^{13,15-17} in our hospital, no woman had massive bleeding immediately before surgery, and all surgery was performed as an elective procedure in our hospital. Antepartum hemorrhage in PAS can occur and cause an emergency situation where preoperative diagnosis cannot be made. This situation is not included because the diagnosis was made intraoperatively with different preparation, such as aortic compression to reduce the blood loss during surgery and this can bias the maternal outcome.

Intraoperative diagnosis was based on guidelines published by Collins et al, which have now been updated by The International Federation of Gynecology and Obstetrics (FIGO).^{13,14} PAS with invasion below the bladder trigone were excluded for uterine conservative-resective surgery because of the high risk for failure. For the purpose of this study, pregnant women with invasion below the bladder trigone were also excluded from the hysterectomy group in order to only include comparable outcomes.^{4,12} Specimens were microscopically examined by the pathologist to confirm the definitive diagnosis and determine the area of placental invasion.¹⁸

Other exclusion criteria included uterine scar dehiscence (bluish smooth appearance of lower uterine segment due to thin uterine wall without any newly formed vessel, where the placenta can mostly separate with gentle traction),¹⁹ massive preoperative bleeding, diffuse placental invasion identified during intraoperative staging, advanced PAS grading with lower bladder-cervical invasion and/or lower parametrial invasion.²⁰

Invasion below the bladder trigone can be distinguished by experienced sonographers using preoperative ultrasound identifying the increase in ultrasound abnormality signs of PAS in the lower uterine segment to the cervix.^{21,22} Intraoperative staging can be carried out by opening the anterior leaf of the broad ligament and evaluating the vesico-uterine space.²³

2.1 | Cesarean hysterectomy

In the early part of our experience, cesarean hysterectomy in women with PAS was performed after delivery of the baby via a fundal incision and IIAL, if there was persistent active bleeding from the lower uterine segment following partial manual placental removal (blunt placental removal). In these cases, we placed a modified Cho suture (brace suture) in the anterior lower uterine segment to reduce the blood loss before cesarean hysterectomy. Planned cesarean hysterectomy was also performed in cases with preoperative diagnosis of placenta increta or percreta and/or intraoperative confirmation of

a bluish appearance with neovascularization (placenta increta) or placental tissue visible in the uterine serous layer with neovascularization (placenta percreta); in these cases, hysterectomy was performed after IIAL procedure without placental removal.

2.2 | Uterine conservative-resective surgery with IIAL

Over the study period, two types of uterine conservative-resective surgery evolved. The first procedure was the Surabaya Modified Procedure for Uterine Conservation (SuMPUC) surgery. With the initial SuMPUC method the baby is delivered via a fundal midline incision followed by IIAL^{5,24,25} and placement of a tourniquet below the lower uterine segment to achieve vascular control of this region.²⁶ After obtaining initial hemostasis, a bladder flap is carefully dissected with ligation of all new vessels from the bladder while stepwise lowering of the tourniquet into the para-cervical area, dissection of the abnormal placentation, and removal of the placenta along with the pathologically invaded area.⁴ A modified Cho suture (brace suture) below the placental dissection area²⁷ was placed if bleeding occurred after the uterine tourniquet was released. Uterine closure of both incisions is completed after complete hemostasis is achieved. In most cases (32 out of 36 cases) this was followed by bilateral salpingectomy.

2.3 | Uterine conservative-resective surgery with identification-ligation of the upper vesical, upper vaginal, and uterine arteries

A midline or transversal fundal incision is performed to deliver the baby, followed by careful development of the bladder flap using the Pelosi maneuver (Figure 1),²⁸ and identification-ligation of vesical and vaginal arteries up to the uterine arteries.

The new concept of uterine conservative-resective surgery consisted of replacing the tourniquet with cervical area sutures in anterior and lateral cervix to control bleeding from the vascular supply coming from the lower uterine segment.²⁹ Just lateral of the abnormal placental invasion site, on both sites, hemostatic sutures are placed and the abnormal placental invasion is dissected, with removal of the whole placenta and closure of the anterior and fundal uterine wall. Bilateral salpingectomy was performed in most cases to prevent the next pregnancy, but not in women opting for another pregnancy.

The most recent development of this technique consisted of a one-step conservative surgical approach; instead of fundal incision, the first uterine incision is performed above the abnormal placental invasion to allow delivery of the baby.

In all cases of uterine conservative surgery, intravenous oxytocin 10 IU is administered after surgical hemostasis of the lower uterine segment has been achieved.

After surgery, PAS was confirmed by pathological examination (macroscopic and microscopic) in all placental specimens. Pathology results were compared with ultrasound findings, and




FIGURE 1 Bladder flap using Pelosi maneuver starting from lateral bladder to medial posterior bladder

the intraoperative diagnosis was compared with the final diagnosis based on agreement between surgeon and pathologist.^{13,14,18}

Uterine conservative surgery was considered to have failed if bleeding did not stop after placental removal/dissection and the aforementioned hemostatic sutures of the uterine, vesical, and vaginal arteries were in place, or in case of recurrent bleeding after the abdominal wall was closed.⁴

In addition to the blood loss and laboratory results, other maternal outcomes were recorded such as maternal complications, major blood loss (>1500 mL),^{30,31} catastrophic blood loss (>2500 mL),³² massive blood transfusion (>4 units of packed red blood cells in 1 hour, >10 units of packed red blood cells in 24 hours, or replacement of 50% of total blood volume within 3 hours),³³ disseminated intravascular coagulation (platelets <100000/ μ L, prolonged prothrombin time-activated partial thromboplastin time, and fibrinogen level <3 g/L)³⁴ and intensive care unit admission.

Independent t test, Mann-Whitney U test, and chi-squared test were performed (SPSS 23, IBM) for the various maternal outcome variables where appropriate.

2.4 | Ethical approval

Ethical approval was obtained from the Ethics Committee in Health Research Dr. Soetomo Academic General Hospital number 1711/KEPK/XII/2019 on December 17, 2019.

3 | RESULTS

Out of 396 surgery-confirmed PAS cases, 234 pregnant women met the inclusion criteria (Figure 2).

Cesarean hysterectomy and IIAL were performed in all PAS cases before November 2016. Uterine conservative surgery started in November 2016 after the sharp increase in PAS cases and the massive blood loss associated with total hysterectomy. The uterine conservative-resective surgery with IIAL was performed as default from November 2016 up to November 2017, the purpose of IIAL was to reduce the blood loss in particular from the very vascular lower uterine segment.^{5,6,24,25}

Delivery of the baby and final surgical procedures were performed at 34–36 weeks of gestation except if there was vaginal bleeding or there were fetal indications, and in some pregnant women as a result of late referral (Table 1). Heavy menstrual bleeding occurred in eight women 3 months after uterine conservative-resective surgery, these women had a thick endometrial line on ultrasound examination, and after treatment with tranexamic acid had further normal periods (Table 2).

Two women in the uterine conservative-resective surgery with IIAL group have been pregnant again, both delivering at 37 weeks of gestation without any complication (Table 3).

Attempts to complete conservative surgery failed in eight women (these were the eight excluded cases) because of anterior-parametrial placental invasion (two cases with blood loss 5000 mL and 4500 mL,




FIGURE 2 Enrollment data based on ultrasound reports, combined medical records, surgical reports, and pathology reports

TABLE 1 Maternal characteristics comparison of cesarean hysterectomy with internal iliac artery ligation vs uterine conservative–resective surgery

	Uterine conservative–resective surgery (n = 200)	Cesarean hysterectomy with IIAL (n = 34)	p
Characteristics			
Age (y), median (range) ^a	33 (19–44)	35 (26–41)	0.013
GA at diagnosis (wk), median (range) ^a	34 (21–40)	34 (28–39)	0.270
GA at surgery (wk), median (range) ^a	35 (28–40)	36 (30–39)	0.385
Number of CS, n (%) ^b			
1 CS	147 (73.5%)	20 (58.8%)	0.120
2 CS	42 (21%)	13 (38.2%)	
>2 CS	11 (5.5%)	1 (2.9%)	
Major implantation, n (%) ^b			
Accreta	60 (30%)	3 (8.8%)	0.031
Increta	93 (46.5%)	19 (55.9%)	
Percreta	47 (23.5%)	12 (35.3%)	

Abbreviations: CS, cesarean section; GA, gestational age; IIAL, internal iliac artery ligation.

^aMann–Whitney U test.

^bChi-squared test.

posterior placental invasion (one case with blood loss 3500 mL), cervical invasion (two cases with blood loss 4000 mL and 3500 mL), and bladder invasion below the bladder trigone (three cases with blood loss from 4000 to 5000 mL). The decision to perform a hysterectomy was taken immediately after massive bleeding occurred during surgery after resection of the placenta and failed attempts to achieve vascular control in the lower segment of the uterus.

Thirty-four of the women underwent surgery before 34 weeks of pregnancy because of recurrent minor vaginal bleeding and major vaginal bleeding while still being hemodynamically stable.

The results demonstrate that cesarean hysterectomy with IIAL is associated with more blood loss and complications compared with both uterine resective–reconstructive surgery methods. Pregnant women managed using uterine conservative–resective surgery

TABLE 2 Maternal outcome comparison of cesarean hysterectomy with internal iliac artery ligation vs uterine conservative–resective surgery

	Uterine conservative–resective surgery (n = 200)	Cesarean hysterectomy with IIAL (n = 34)	p	OR (95% CI)
Outcomes				
Mean blood loss (mL), mean ± SD ^a	1379 ± 769	3168 ± 1916	<0.001	
Hb presurgery (g/dL), mean ± SD ^a	10.5 ± 1.3	9.9 ± 1.5	0.018	
Hb post-surgery (g/dL), mean ± SD ^c	9.6 ± 1.5	8.6 ± 1.7	0.002	
Plt presurgery (/ μ L), mean ± SD ^a	278615 ± 147976	276059 ± 88890	0.968	
Plt post-surgery (/ μ L), mean ± SD ^a	248415 ± 222828	165382 ± 71261	<0.001	
PRBC units transfused, median (range) ^a	1 (0–6)	3 (0–10)	<0.001	
Major blood loss (>1500 mL), n (%) ^b	68 (34%)	24 (70.6%)	<0.001	4.7 (2.1–10.3)
Catastrophic blood loss (>2500 mL), n (%) ^b	25 (12.5%)	22 (64.7%)	<0.001	12.8 (5.7–29.1)
Massive transfusion, n (%) ^b	5 (2.5%)	12 (35.3%)	<0.001	21.3 (6.9–66)
ICU admission, n (%) ^b	3 (1.5%)	11 (32.4%)	<0.001	31.4 (8.2–120.86)
DIC, n (%) ^b	1 (0.5%)	2 (5.9%)	0.056	–
Other maternal complications, n (%) ^b	25 (12.4%)	11 (32.4%)	0.007	3.5 (1.5–8.1)
Re-laparotomy for recurrent bleeding	2 (1%)	2 (5.9%)	0.423	–
Bladder injury	9 (4.5%)	7 (20.6%)	0.002	5.2 (1.9–16)
Vascular injury	1 (0.5%)	2 (5.9%)	0.056	–
Uterine atony	21 (10.5%)	0	0.143	–
Maternal death	0	0	–	–
Long-term follow up (6 mo)				
Fistula	1 (uterocutaneous fistula)	2 (vesicovaginal fistula)		
Heavy menstrual bleeding	8	0		
Percreta invasion				
	n = 47	n = 12		
GA at surgery (wk), median (range) ^a	36 (28–39)	36 (32–38)	0.797	
Mean blood loss for percreta invasion (mL), mean ± SD ^c	1926 ± 792	4875 ± 1680	<0.001	
Hb presurgery (g/dL) ^c	10.5 ± 1.5	9.1 ± 1.9	0.012	
Hb post-surgery (g/dL) ^c	9.5 ± 1.5	8.4 ± 1.3	0.024	
Plt presurgery (/ μ L), mean ± SD ^c	276787 ± 85033	273833 ± 105629	0.919	
Plt post-surgery (/ μ L), mean ± SD ^c	211638 ± 67705	134833 ± 63941	0.001	
PRBC units transfused, median (range) ^a	2 (0–6)	7 (3–10)	<0.001	
Major blood loss for percreta invasion, n (%) ^b	27 (57.4%)	12 (100%)	0.005	18.6 (1–333.3)
Catastrophic blood loss for percreta invasion (>2500 mL), n (%) ^b	13 (27.7%)	12 (100%)	<0.001	63.9 (3.5–1156.4)
Massive transfusion for percreta, n (%) ^b	4 (8.5%)	9 (75%)	<0.001	32.3 (6.1–169.7)
Admission to ICU, n (%) ^b	2 (4.3%)	8 (66.7%)	<0.001	45 (7–288)
DIC, n (%) ^b	0	2 (16.7%)	0.039	22.6 (1–506.7)

Abbreviations: Hb, hemoglobin; ICU, intensive care unit; IIAL, internal iliac artery ligation; PAS, placenta accrete spectrum; Plt, platelets; PRBC, packed red blood cells; DIC, disseminated intravascular coagulation.

aMann–Whitney U test.

bChi-squared test.

cIndependent t test.

TABLE 3 Maternal outcome comparison of uterine conservative–resective surgery with internal iliac artery vs identification–ligation of the upper vesical, upper vaginal and uterine arteries

	Uterine conservative–resective surgery with IIAL (n = 36)	Identification–ligation of the upper vesical, upper vaginal, and uterine arteries (n = 164)	p	OR (95% CI)
Characteristics				
GA at surgery (wk), median (range) ^a	35 (28–40)	35 (28–39)	0.483	
Major implantation, n (%) ^b				
Accreta	8 (22.2%)	52 (31.7%)	0.522	
Increta	19 (52.8%)	74 (45.1%)		
Percreta	9 (25%)	38 (23.2%)		
Outcomes				
Mean blood loss (mL), mean ± SD ^a	1701 ± 813	1307 ± 743	0.005	
Hb presurgery (g/dL), mean ± SD	10.3 ± 1.5	10.5 ± 3	0.255	
Hb post-surgery (g/dL), mean ± SD	9.4 ± 1.5	9.5 ± 1.5	0.376	
Plt presurgery (/ μ L), mean ± SD	267 666 ± 77802	281018 ± 159377	0.789	
Plt post-surgery (/ μ L), mean ± SD	218 896 ± 71 676	254915 ± 243 473	0.295	
PRBC units transfused, median (range)	2 (0–6)	1 (0–6)	0.052	
Major blood loss (>1500 mL), n (%) ^b	18 (50%)	50 (30.5%)	0.041	0.44 (0.21–0.91)
Catastrophic blood loss (>2500 mL), n (%) ^b	7 (19.4%)	18 (11%)	0.170	–
Massive transfusion, n (%)	1 (2.8%)	4 (2.4%)	1.000	–
Admission to intensive care unit, n (%) ^b	0	3 (1.8%)	1.000	–
DIC, n (%)	0	1 (0.6%)	1.000	–
Other maternal complication, n (%) ^b	6 (16.7%)	19 (11.6%)	0.408	
Re-laparotomy for recurrent bleeding	0	2 (1.2%)	0.940	–
Bladder injury	4 (11.8%)	5 (3%)	0.048	0.25 (0.06–0.99)
Vascular injury	1 (2.8%)	0	0.109	–
Uterine atony	2 (5.6%)	19 (11.6%)	0.297	–
Percreta invasion				
	n = 9	n = 38		
GA at surgery (wk), median (range) ^a	34 (28–37)	36 (29–39)	0.081	
Mean blood loss for percreta invasion (mL), mean ± SD ^a	2288 ± 509	1674 ± 855	0.020	
Hb presurgery (g/dL), mean ± SD ^c	10.2 ± 2.3	10.6 ± 1.3	0.447	
Hb post-surgery (g/dL), mean ± SD ^c	9.3 ± 1.6	9.6 ± 1.5	0.678	
Plt presurgery (/ μ L), mean ± SD ^c	230 777 ± 67 952	287 684 ± 85 757	0.071	
Plt post-surgery (/ μ L), mean ± SD ^c	163 222 ± 42 693	223 105 ± 67 822	0.015	
PRBC units transfused, median (range) ^a	3 (2–6)	2 (0–5)	0.086	
Major blood loss for percreta invasion (>1500 mL), n (%) ^b	8 (88.9%)	19 (50%)	0.059	–
Catastrophic blood loss for percreta invasion (>2500 mL), n (%) ^b	4 (44.4%)	9 (23.7%)	0.237	–
Massive transfusion for percreta, n (%) ^b	1 (11.1%)	3 (7.9%)	1.000	–
Admission to ICU, n (%) ^b	0	2 (5.3%)	1.000	–
DIC, n (%) ^b	0	0		

Abbreviations: DIC, disseminated intravascular coagulation; Hb, hemoglobin; ICU, intensive care unit; IIAL, internal iliac artery ligation; PAS, placenta accreta spectrum; Plt, platelets; PRBC, packed red blood cells.

^aMann–Whitney *U* test.

^bChi-squared test.

^cIndependent *t* test.

with identification–ligation of the upper vesical, upper vaginal, and uterine arteries have lower blood loss compared with the uterine conservative–resective surgery with IIAL.

4 | DISCUSSION

The findings based on this large series of pregnant women with selective PAS managed in one tertiary referral hospital in East Java demonstrate that uterine conservative–resective surgery is associated with a significantly lower rate of major maternal complications when compared with cesarean hysterectomy and IIAL.

The steep increase in the number of PAS managed in our center primarily reflects the increased recognition that these pregnant women need their care provided by recognized tertiary centers with a dedicated “accreta team” in addition to the well-known increased incidence of cesarean sections in Indonesia (from 9.8% in 2013 to 17.6% in 2018).³⁵

The principal of surgical strategies depends on the placental topography and on detailed knowledge of the placental–uterine vascularity.^{4,29} Research by Palacios-Jaraquemada et al on uterine conservative resective surgery has demonstrated the high success rate of uterine conservative surgery for placental invasion above the bladder trigone.¹²

Mean blood loss, rate of maternal complications, and the need for intensive care unit admission for catastrophic (>2500 mL) blood loss was significantly higher in the original hysterectomy plus IIAL group compared with the uterine conservative–resective surgery group, even in placenta percreta cases. With this procedure, the possibility of bleeding from the extrauterine vascular anastomosis–damaged when clamping the lower uterine segment—can lead to massive bleeding.²⁹

The first uterine conservative–resective surgery with IIAL uses a similar concept to the triple P procedure;⁵ both methods focus on reducing blood loss from the uterine arteries as the major branches of IIA. Additional sutures were often needed in the lower uterine segment to control the bleeding, especially from the placental bed after placental resection and removal. However, the difference is the use of the lower uterine tourniquet approach to reduce the blood loss.

Uterine conservative–resective surgery with IIAL was associated with a higher risk of major blood loss compared with the modified uterine conservative–resective surgery with identification–ligation of the upper vesical, upper vaginal, and uterine arteries ($p = 0.041$): mean blood loss was, respectively, 1701 mL vs 1307 mL ($p = 0.005$) for all PAS grade and 2288 mL vs 1674 mL ($p = 0.020$) for percreta invasion. Kutuk et al also found higher blood loss in a hysterectomy group compared with uterine conservative–resective surgery and leaving the placental in situ for all grades of PAS, but described higher blood loss associated with uterine conservative–resective surgery in women with placenta increta and percreta while using IIAL, uterine artery, and utero-ovarian artery ligation.³⁶

The higher risk of major blood loss associated with uterine conservative–resective surgery with IIAL is probably because this particular technique only focuses on the uterine artery as the branch of the internal iliac artery, while “ignoring” the vaginal arteries.

Other studies using a similar approach of vascular control in cases with invasion above the bladder trigone also found blood loss of approximately 1500 mL;^{12,37} this level of 1500 mL can be used as a reasonable benchmark in uterine conservative–resective surgery.

Strategies required for developing the bladder flap are pivotal to control vascularity from the bladder to placental invasion. The Pelosi maneuver is an important step to open the vesico–uterine space and identify the location of the placental invasion during surgery.²⁸ This maneuver will be difficult if the placenta invades to below the lower trigonal bladder¹² and may risk damaging the new vessel and, as a result, massive bleeding.

We no longer recommend IIAL in conservative–resective uterine surgery for PAS because of the associated higher blood loss, surgical complexities, and the associated 1.8% risk of vascular injury (internal iliac artery injury).^{38,39}

The International Federation of Gynecology and Obstetrics (FIGO) have reached consensus on how to standardize clinical grading of PAS; the placental topography (accreta–increta–percreta) and abnormal placental location are the critical factors determining the success rate of surgical conservative approaches. Treatment of abnormal placental invasion above the lower uterine segment has higher success rates compared with placental invasion in the lower uterine segment (lower bladder, cervix, parametrium, and posterior placental invasion) in PAS with placenta previa. Detailed preoperative placental mapping using MRI or ultrasound (abdominal/transvaginal gray-scale, abdominal/transvaginal Doppler, three-dimensional Doppler, three-dimensional volume-rendering ultrasound) have a critical role in predetermining the success of uterine conservative surgery.⁴⁰ In emergency situations like preoperative massive vaginal bleeding with hemodynamically compromised pregnant women, we do not recommend uterine conservative surgery.

The limitation of this retrospective cohort study is mainly that over the years, our experience and expertise have improved significantly and the current high success rates are closely linked to the presence of a dedicated team of experienced surgeons with the backup of a large tertiary facility. These procedures are better not done in smaller units lacking the experience.

In case an obstetrician encounters a major PAS during a cesarean section in a small peripheral hospital, the best strategy is to deliver the baby, clamp the cord, close the uterus and transfer the pregnant women to the regional tertiary facility.

5 | CONCLUSION

Cesarean hysterectomy is not the default treatment for all PAS, the pregnant woman with PAS with invasion above the vesical trigone should have uterine conservative–resective surgery because it is much safer than cesarean hysterectomy.

13

ACKNOWLEDGMENTS

These new uterine conservative–resective surgery concepts were developed in close collaboration with Prof. José Miguel Palacios-Jaraquemada.

47

CONFLICT OF INTEREST

None.

AUTHOR CONTRIBUTIONS

The study was design by RAA, AA, KEG, MPW, MIA, NIC, EE, BW, HTJ, EGD, CAB, DK, DPV, GA, GAD, and AS. It was planned by RAA, AA, KEG, MPW, MIA, NIC, EE, EGD, GAD and AS and conducted by RAA, MPW, NIC, EGD, GA, DK, DPV, and AS.

54

Data were analyzed by RAA, CAB, GA, GAD, and AS. All authors contributed to writing the manuscript.

22

ORCID

Rozi Aditya Aryananda  <https://orcid.org/0000-0001-6674-7682>

REFERENCES

- Wu S, Kocherginsky M, Hibbard JU. Abnormal placentation: twenty-year analysis. *Am J Obstet Gynecol.* 2005;192:1458-1461.
- Grechukhina O, Deshmukh U, Fan L, et al. Cesarean scar pregnancy, incidence, and recurrence: five-year experience at a single tertiary care referral center. *Obstet Gynecol.* 2018;132:1285-1295.
- Likis FE, Sathe NA, Morgans AK, et al. *Management of Postpartum Hemorrhage.* Agency for Healthcare Research and Quality (US); 2015.
- Palacios Jaraquemada JM, Pesaresi M, Nassif JC, Hermosid S. Anterior placenta percreta: surgical approach, hemostasis and uterine repair. *Acta Obstet Gynecol Scand.* 2004;83:738-744.
- Chandrarahan E, Rao S, Belli A-M, Arulkumaran S. The triple-P procedure as a conservative surgical alternative to peripartum hysterectomy for placenta percreta. *Int J Gynaecol Obstet.* 2012;117:191-194.
- Sentilhes L, Goffinet F, Kayem G. Management of placenta accreta. *Acta Obstet Gynecol Scand.* 2013;92:1125-1134.
- American College of Obstetricians and Gynecologists; Society for Maternal-Fetal Medicine. Obstetric Care Consensus No. 7: Placenta Accreta Spectrum. *Obstet Gynecol.* 2018;132:e259-e275.
- Committee on Obstetric Practice. Committee opinion no. 529: placenta accreta. *Obstet Gynecol.* 2012;120:207-211.
- Jolley JA, Nageotte MP, Wing DA, Shrivastava VK. Management of placenta accreta: a survey of maternal-fetal medicine practitioners. *J Matern Fetal Neonatal Med.* 2012;25:756-760.
- Sentilhes L, Ambroselli C, Kayem G, et al. Maternal outcome after conservative treatment of placenta accreta. *Obstet Gynecol.* 2010;115:526-534.
- Provansal M, Courbiere B, Agostini A, D'Ercole C, Boubli L, Bretelle F. Fertility and obstetric outcome after conservative management of placenta accreta. *Int J Gynaecol Obstet.* 2010;109:147-150.
- Palacios-Jaraquemada JM, Fiorillo A, Hamer J, Martinez M, Bruno C. Placenta accreta spectrum: a hysterectomy can be prevented in almost 80% of cases using a resective-reconstructive technique. *J Matern Fetal Neonatal Med.* 2022;35:275-282.
- Collins SL, Stevenson GN, Al-Khan A, et al. Three-dimensional power doppler ultrasonography for diagnosing abnormally invasive placenta and quantifying the risk. *Obstet Gynecol.* 2015;126:645-653.
- Jauniaux E, Bhide A, Kennedy A, et al. FIGO consensus guidelines on placenta accreta spectrum disorders: prenatal diagnosis and screening. *Int J Gynecol Obstet.* 2018;140:274-280.
- Rac MWF, Dashe JS, Wells CE, Moschos E, McIntire DD, Twickler DM. Ultrasound predictors of placental invasion: the Placenta Accreta Index. *Am J Obstet Gynecol.* 2015;212:343.e1-343.e7.
- Alfirevic Z, Tang AW, Collins SL, Robson SC, Palacios-Jaraquemada J, Ad-hoc International AIP Expert Group. Pro forma for ultrasound reporting in suspected abnormally invasive placenta (AIP): an international consensus. *Ultrasound Obstet Gynecol.* 2016;47:276-278.
- Nieto-Calvache AJ, Palacios-Jaraquemada JM, Aryananda RA. Factors to consider when seeking better results in placenta accreta spectrum. *Acta Obstet Gynecol Scand.* 2021;100:1932-1933.
- Dannheim K, Shaiker SA, Hecht JL. Hysterectomy for placenta accreta; methods for gross and microscopic pathology examination. *Arch Gynecol Obstet.* 2016;293:951-958.
- Jauniaux E, Collins S, Burton GJ. Placenta accreta spectrum: pathophysiology and evidence-based anatomy for prenatal ultrasound imaging. *Am J Obstet Gynecol.* 2018;218:75-87.
- Palacios-Jaraquemada JM, D'Antonio F, Buca D, Fiorillo A, Larrazza P. Systematic review on near miss cases of placenta accreta spectrum disorders: correlation with invasion topography, prenatal imaging, and surgical outcome. *J Matern Fetal Neonatal Med.* 2020;33:3377-3384.
- Cali G, Forlani F, Lees C, et al. Prenatal ultrasound staging system for placenta accreta spectrum disorders. *Ultrasound Obstet Gynecol.* 2019;53:752-760.
- di Pasquo E, Ghi T, Cali G, et al. Intracervical lakes as sonographic marker of placenta accreta spectrum disorder in patients with placenta previa or low-lying placenta. *Ultrasound Obstet Gynecol.* 2020;55:460-466.
- Nieto-Calvache AJ, Palacios-Jaraquemada JM, Aryananda RA, et al. How to identify patients who require aortic vascular control in placenta accreta spectrum disorders? *Am J Obstet Gynecol MFM.* 2022;4:100498.
- Tan CH, Tay KH, Sheah K, et al. Perioperative endovascular internal iliac artery occlusion balloon placement in Management of Placenta Accreta. *Am J Roentgenol.* 2007;189:1158-1163.
- Shehata A. Uterine sparing techniques in placenta Accreta. *Obstet Gynecol Int J.* 2016;5:258-262.
- Yan J, Shi C-Y, Yu L, Yang H-X. Folding sutures following tourniquet binding as a conservative surgical approach for placenta previa combined with morbidly adherent placenta. *Chin Med J (Engl).* 2015;128:2818-2820.
- Cho JY, Kim SJ, Cha KY, Kay CW, Kim MI, Cha KS. Interrupted circular suture: bleeding control during cesarean delivery in placenta previa accreta. *Obstet Gynecol.* 1991;78:876-879.
- Pelosi MA 3rd, Pelosi MA. Modified cesarean hysterectomy for placenta previa percreta with bladder invasion: retrovesical lower uterine segment bypass. *Obstet Gynecol.* 1999;93(5 Pt 2):830-833.
- Palacios Jaraquemada JM, Garcia Monaco R, Barbosa NE, Ferle L, Iriarte H, Conesa HA. Lower uterine blood supply: extrauterine anastomotic system and its application in surgical devascularization techniques. *Acta Obstet Gynecol Scand.* 2007;86:228-234.
- Prevention and Management of Postpartum Haemorrhage: Green-top Guideline No. 52. *BJOG.* 2017;124:e106-e149.
- Jadon A, Bagai R. Blood transfusion practices in obstetric anaesthesia. *Indian J Anaesth.* 2014;58:629-636.
- Wise A, Clark V. Strategies to manage major obstetric haemorrhage. *Curr Opin Anaesthesiol.* 2008;21:281-287.
- Patil V, Shetmahajan M. Massive transfusion and massive transfusion protocol. *Indian J Anaesth.* 2014;58:590-595.
- Levi M, Toh CH, Thachil J, Watson HG. Guidelines for the diagnosis and management of disseminated intravascular coagulation. *Br J Haematol.* 2009;145:24-33.
- Laporan Hasil Riset Kesehatan Dasar (Rikesdas). Available from: <https://www.litbang.kemkes.go.id/laporan-riset-kesehatan-dasar-rikesdas/>
- Kutuk MS, Ak M, Ozgun MT. Leaving the placenta in situ vs conservative and radical surgery in the treatment of placenta accreta spectrum disorders. *Int J Gynaecol Obstet.* 2018;140:338-344.

37. Shih J-C, Liu K-L, Kang J, Yang J-H, Lin M-W, Yu C-U. 'Nausicaa' compression suture: a simple and effective alternative to hysterectomy in placenta accreta spectrum and other causes of severe postpartum haemorrhage. *BJOG*. 2019;126:412-417.
38. Iwata A, Murayama Y, Itakura A, Baba K, Seki H, Takeda S. Limitations of internal iliac artery ligation for the reduction of intraoperative hemorrhage during cesarean hysterectomy in cases of placenta previa accreta. *J Obstet Gynaecol Res*. 2010;36:254-259.
39. Kaya B, Damarer Z, Guralp O. Is there still a role of internal iliac artery ligation in obstetric hemorrhage with the current rise in popularity of other uterus sparing techniques? *Eur J Obstet Gynecol Reprod Biol*. 2016;206:e57-e58.
40. Polat M, Kahramanoglu I, Senol T, Ozkaya E, Karateke A. Shorter the cervix, more difficult the placenta percreta operations. *J Mat Fetal Neonat Med*. 2016;29:2327-2331.

How to cite this article: Aryananda RA, Aditiawarman A, Gumilar KE, et al. Uterine conservative-resective surgery for selected placenta accreta spectrum cases: Surgical-vascular control methods. *Acta Obstet Gynecol Scand*. 2022;00:1-10. doi: [10.1111/aogs.14348](https://doi.org/10.1111/aogs.14348)

Uterine conservative–resective surgery for selected placenta

ORIGINALITY REPORT

16%

SIMILARITY INDEX

12%

INTERNET SOURCES

13%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Flore - Anne Pain, Anthony Dohan, Gilles Grange, Louis Marcellin et al. " Percreta score to differentiate between placenta accreta and placenta percreta with ultrasound and imaging ", Acta Obstetricia et Gynecologica Scandinavica, 2022
Publication 2%
- 2** [medcraveonline.com](https://www.medcraveonline.com)
Internet Source 1%
- 3** repository.unair.ac.id
Internet Source 1%
- 4** Albaro José Nieto-Calvache, José Miguel Palacios-Jaraquemada, Rozi Aditya Aryananda, Fernando Rodriguez et al. "How to identify patients who require aortic vascular control in placenta accreta spectrum disorders?", American Journal of Obstetrics & Gynecology MFM, 2022
Publication 1%
- 5** www.balimedicaljournal.org
Internet Source 1%

6	arcjournals.org Internet Source	<1 %
7	Rozi Aditya Aryananda, Grace Ariani. "Chapter 5 Classification of Placenta Accreta Spectrum", Springer Science and Business Media LLC, 2023 Publication	<1 %
8	www.scribd.com Internet Source	<1 %
9	Silvia Gianola, Greta Castellini, Annalisa Biffi, Gloria Porcu et al. "Accuracy of risk tools to predict critical bleeding in major trauma", Journal of Trauma and Acute Care Surgery, 2021 Publication	<1 %
10	www.arcjournals.org Internet Source	<1 %
11	pure.eur.nl Internet Source	<1 %
12	"Main Program", Australian and New Zealand Journal of Obstetrics and Gynaecology, 2018 Publication	<1 %
13	edoc.pub Internet Source	<1 %
14	women.texaschildrens.org Internet Source	<1 %

15

"Moderated Poster Sessions", Journal of Endourology, 09/15/2003

Publication

<1 %

16

Perez-Delboy, A, and JD Wright. "Surgical management of placenta accreta: to leave or remove the placenta?", BJOG An International Journal of Obstetrics & Gynaecology, 2014.

Publication

<1 %

17

Yüksel, Hasan. "A novel approach to primary lower uterine segment atony", Taiwanese Journal of Obstetrics and Gynecology, 2015.

Publication

<1 %

18

curis.ku.dk

Internet Source

<1 %

19

Margarida Cal, Diogo Ayres-de-Campos, Eric Jauniaux. "International survey of practices used in the diagnosis and management of placenta accreta spectrum disorders", International Journal of Gynecology & Obstetrics, 2018

Publication

<1 %

20

www.mdpi.com

Internet Source

<1 %

21

Edwin Chandraharan, Richard Hartopp, Baskaran Thilaganathan, Conrado Milani Coutinho. "How to set up a regional specialist referral service for Placenta Accreta Spectrum

<1 %

(PAS) disorders?", Best Practice & Research
Clinical Obstetrics & Gynaecology, 2020

Publication

22

Rozi Aditya Aryananda, Aldika Akbar,
Manggala Pasca Wardhana, Khanisyah Erza
Gumilar et al. "New three-dimensional/four-
dimensional volume rendering imaging
software for detecting the abnormally
invasive placenta", Journal of Clinical
Ultrasound, 2019

Publication

23

tel.archives-ouvertes.fr

Internet Source

24

J. C. Shih. "Role of three-dimensional power
Doppler in the antenatal diagnosis of placenta
accreta: comparison with gray-scale and color
Doppler techniques", Ultrasound in Obstetrics
and Gynecology, 02/2009

Publication

25

MINNA TIKKANEN. "Antenatal diagnosis of
placenta accreta leads to reduced blood loss :
Placenta accreta", Acta Obstetricia Et
Gynecologica Scandinavica, 10/2011

Publication

26

S K H Wong. "Prediction of therapeutic failure
after adrenaline injection plus heater probe
treatment in patients with bleeding peptic
ulcer", Gut, 2002

<1 %

<1 %

<1 %

<1 %

<1 %

27

openarchive.ki.se

Internet Source

<1 %

28

researchspace.ukzn.ac.za

Internet Source

<1 %

29

www.science.gov

Internet Source

<1 %

30

www.slideshare.net

Internet Source

<1 %

31

"Maternal Fetal Health", Journal of Obstetrics and Gynaecology Research, 2020

Publication

<1 %

32

"Maternal Fetal Medicine", Journal of Obstetrics and Gynaecology Research, 2023

Publication

<1 %

33

Ding Huang, Sha Chen, Yanqing Cai, Liye Shi, Yiru Shi, Min Zeng, Weiwei Cheng, Jian Zhang, Hongjie Pan. "Adjunctive azithromycin prophylaxis protects women from uterine cesarean scar defect: A randomized controlled trial", Acta Obstetrica et Gynecologica Scandinavica, 2022

Publication

<1 %

34

daneshyari.com

Internet Source

<1 %

medi-guide.meditool.cn

35

Internet Source

<1 %

36

repub.eur.nl

Internet Source

<1 %

37

www.acenet.edu

Internet Source

<1 %

38

www.minervamedica.it

Internet Source

<1 %

39

www.researchgate.net

Internet Source

<1 %

40

Andreia Fonseca, Diogo Ayres de Campos.
"Maternal morbidity and mortality due to
placenta accreta spectrum disorders", Best
Practice & Research Clinical Obstetrics &
Gynaecology, 2020

Publication

<1 %

41

Kenneth G. Saag, Michael A. Becker, Andrew
Whelton, Barbara Hunt, Majin Castillo,
Krisztina Kisfalvi, Lhanoo Gunawardhana.
"Efficacy and Safety of Febuxostat Extended
and Immediate Release in Patients with Gout
and Renal Impairment: Phase III Placebo-
Controlled Study", Arthritis & Rheumatology,
2018

Publication

<1 %

42

Oncology, 2006.

Publication

<1 %

43

assets.researchsquare.com

Internet Source

<1 %

44

ejhs.ju.edu.et

Internet Source

<1 %

45

qdoc.tips

Internet Source

<1 %

46

www.frontiersin.org

Internet Source

<1 %

47

www.ijrcog.org

Internet Source

<1 %

48

ELIZA M. BERKLEY, ALFRED ABUHAMAD.
"Imaging of Placenta Accreta Spectrum",
Clinical Obstetrics and Gynecology, 2018

Publication

<1 %

49

Lior Friedrich, Nizan Mor, Alina Weissmann -
Brenner, Eran Kassif et al. "Risk factors for
bladder injury during placenta accreta
spectrum surgery", International Journal of
Gynecology & Obstetrics, 2022

Publication

<1 %

50

W. Merz. "Diagnosis, Treatment and
Application of Color Doppler in Conservative
Management of Abnormally Adherent

<1 %

Placenta", Ultraschall in der Medizin -
European Journal of Ultrasound, 12/2009

Publication

51

F. Finazzo, F. D'antonio, G. Masselli, F. Forlani et al. "Interobserver agreement in MRI assessment of severity of placenta accreta spectrum disorders", Ultrasound in Obstetrics & Gynecology, 2020

Publication

<1 %

52

Mehmet Kutuk, Aysegul Kilic, Mehmet Ak, Mahmut Ozgun. " Infectious complications in morbidly adherent placenta treated with leaving placenta : A cohort series and suggested approach ", The Journal of Maternal-Fetal & Neonatal Medicine, 2018

Publication

<1 %

53

Principles of Critical Care in Obstetrics, 2016.

Publication

<1 %

54

Saad El Gelany, Mohammed H. Mosbeh, Emad M. Ibrahim, Mo'men Mohammed et al. "Placenta Accreta Spectrum (PAS) disorders: incidence, risk factors and outcomes of different management strategies in a tertiary referral hospital in Minia, Egypt: a prospective study", BMC Pregnancy and Childbirth, 2019

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Uterine conservative–resective surgery for selected placenta

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10