

Comparison of short-term

by Aditiawarman Aditiawarman

Submission date: 27-Apr-2023 09:37PM (UTC+0800)

Submission ID: 2077157408

File name: 15_Comparison_of_short-term.pdf (313.46K)

Word count: 3696

Character count: 18797



Comparison of short-term outcome and serum magnesium levels in premature infants of mothers after magnesium sulfate therapy

Aditiawarman ¹, Jacob Trisusilo Salean ², Mahendra Tri Arif Sampurna ³, Ernawati ¹, Hermanto Tri Joewono ¹

¹ Division of Maternal-Fetal Medicine, Department of Obstetrics and Gynecology, Faculty of Medicine-Dr. Soetomo Teaching Hospital, Universitas Airlangga, Surabaya 60131, INDONESIA

² Department of Obstetrics and Gynecology, Faculty of Medicine-Dr. Soetomo Teaching Hospital, Universitas Airlangga, Surabaya 60131, INDONESIA

³ Department of Pediatrics, Faculty of Medicine-Dr. Soetomo Teaching Hospital, Universitas Airlangga, Surabaya 60131, INDONESIA

*Corresponding author: aditiawarman@fk.unair.ac.id

Abstract

Background: Magnesium sulfate (MgSO₄) is proven to be a neuroprotectant which functions to reduce the risk of cerebral palsy in premature infants, but the effects on the short-term outcome of newborns are still contradictory. This study aimed to analyze differences in short-term outcomes and serum magnesium levels in newborns after given MgSO₄ therapy.

Methods: This was a prospective cohort study. Subjects were divided into two groups involved preterm infants from mothers who received MgSO₄ and pregnant women who did not received MgSO₄ as a control.

Results: Among sixty subjects, there were significant differences in outcomes for gestational age infants and respiratory distress syndrome (RDS) between two groups ($p = 0.011$ and $p = 0.013$, respectively). Whereas, the significance results were found in birth weight ($p = 0.306$), Apgar scores 1 ($p = 0.816$) and 5 minutes ($p = 0.924$), the use of ventilators ($p = 0.335$), sepsis ($p = 0.276$), mortality ($p = 0.754$), and umbilical artery pH ($p = 0.123$). The magnesium serum levels of mothers who received MgSO₄ and control were 3.45 ± 1.65 mg/dl and 1.78 ± 0.23 mg/dl, respectively ($p = 0.001$), and infants who received MgSO₄ and control were 3.10 ± 1.35 mg/dl and 1.74 ± 0.19 mg/dl, respectively ($p = 0.001$).

Conclusion: There were significant differences in infant outcomes only at gestational age and RDS. The magnesium serum levels of newborns in therapy group were higher than control.

Keywords: MgSO₄, short-term outcome, premature infant

Aditiawarman, Salean JT, Sampurna MTA, Ernawati, Joewono HT (2020) Comparison of short-term outcome and serum magnesium levels in premature infants of mothers after magnesium sulfate therapy. Eurasia J Biosci 14: 1863-1867.

© 2020 Aditiawarman et al.

This is an open-access article distributed under the terms of the Creative Commons Attribution License.

INTRODUCTION

In the late 1990s, studies of infants born to mothers given magnesium sulfate (MgSO₄) to prevent eclampsia seizures or to tocolism showed decreased levels of Periventricular Cystic Leukomalacia (PVL) and cerebral palsy (CP). The meta-analysis stated that the administration of MgSO₄ as a neuroprotection is proven to prevent CP. Cochrane in 2009 also revealed that patients given MgSO₄ could reduce the incidence of CP by 1.6% compared to those not given antenatal MgSO₄. For moderate to severe CP levels, MgSO₄ shows the ability to reduce the risk of CP by a statistically significant difference (Salfi et al. 2019, Zeng et al. 2016).

The use of MgSO₄ as a neuroprotection begins with studies in preeclampsia patients where there is a decrease in CP in patients receiving MgSO₄ therapy.

MgSO₄ can reduce blood pressure thereby reducing the risk of eclampsia and maternal death in which no adverse effects can be seen on the mother or baby in the short term. Thus, Magnesium sulphate for Prevention of Eclampsia trial (MAGPIE trial) concluded that the short-term effects on the administration of MgSO₄ therapy are safe (Group 2007, Pasaribu et al. 2016). Preterm birth is the leading cause of neonatal death (Sukatendel et al. 2018). The meta-analysis concluded that there is a reduced risk of developing CP in fetuses exposed to MgSO₄ to mothers at risk of giving

Received: May 2019

Accepted: April 2020

Printed: June 2020



birth to premature babies without increasing the risk of fetal or newborn death (Costantine et al. 2009).

However, if seen from the study conducted by Dwight in 2008, fetal exposure to MgSO₄ prior to the anticipation of early preterm labor does not reduce the combined risk of CP or death. Side effects have also been reported in several studies of neonates exposed to antenatal MgSO₄. The risk of respiratory distress, the need for resuscitation, neonatal seizures, and Apgar scores less than 7 at 5 minutes will increase with increasing MgSO₄ concentrations in the baby's body (Rouse et al. 2008).

The use of MgSO₄ for neuroprotection is not yet a fixed procedure in Indonesia. Short-term side effects are a concern because the infant mortality rate at less than 32 weeks' gestation in Indonesia is quite high. Premature infants with high exposure to MgSO₄ (≥ 4.5 Meq/L) can increase perinatal mortality due to lack of birth weight and gestational age (Basu et al. 2011). However, there are a number of studies of MgSO₄ for neuroprotection in the short term that show insufficient but not significant outcomes (Basu et al. 2011, Costantine et al. 2009). Thus, we aimed to analyze the differences in short-term outcomes and serum magnesium levels in newborns and mothers given MgSO₄ therapy and not treated with MgSO₄ (Kuchits S.S et al., 2019).

MATERIALS AND METHODS

This was an observational cohort study conducted in a group of newborns with a diagnosis of premature pregnant women. The study sample was selected from a population of infants with preterm pregnant women who received MgSO₄ taken from severe preeclampsia patients at 28-34 weeks gestational age who received MgSO₄, and the control sample was premature pregnant women at 28-34 weeks gestational age born at Dr. Soetomo Hospital in the period July 2018 to December 2018. This research had received ethical feasibility from the ethics commission for research at Dr Soetomo Hospital, Surabaya with the number 0403/KEPK/VII/2018.

Inclusion criteria included premature pregnant women who gave birth at a gestational age of more than 28 to 34 weeks, giving birth by vaginal delivery or by fetal delivery with a single pregnancy, no intrauterine asphyxia (Non Stress test (NST) category 1). Whereas, the exclusion criteria included major congenital abnormalities which are circulatory system malformations including congenital heart block, transposition of large arteries, left ventricular hypoplasia, pulmonary atresia and tetralogy of fallot. Abnormalities in the digestive system include abdominal defects, gastroschisis, central nervous system, and chromosomal abnormalities, mothers with cardiac abnormalities, smokers, consumption of opiates or alcohol during pregnancy, and autoimmune diseases.

A sample of 60 subjects was divided into groups who received MgSO₄ therapy and controls who did not receive MgSO₄ therapy. The group that received MgSO₄ therapy was given MgSO₄ therapy at a dose of 4 grams intravenously then continued with maintenance of 1 gram/hour of syringe pump up to 24 hours followed up until delivery was performed. The control group was not given MgSO₄ therapy but was followed up until delivery. In both groups, maternal and neonatal serum magnesium levels, postpartum infant blood gas analysis and short-term outcome assessments were performed, including first and fifth minute Apgar scores, respiratory distress syndrome (RDS), sepsis, type of oxygen support and infant mortality for 7 days of treatment. Data were analyzed using SPSS statistics 21 software. Chi-square analysis, Fisher's exact test, independent-samples test, and one-way Anova were used to analyze differences between variables.

RESULTS

There were 98 premature deliveries, with 9 babies born dead and excluded, 11 babies with congenital abnormalities, 18 babies with gestational age <28 weeks, then 60 samples that met the inclusion criteria were taken. Characteristics of mothers giving birth to premature babies during the study period are presented in Table 1. Table 1 shows 30 patients with body mass index (BMI) above 30 (obesity). Of the 30 patients, 20 patients (83.3%) were diagnosed with severe preeclampsia and received MgSO₄ therapy, and an analysis using chi-square on BMI with MgSO₄ therapy showed a significance value of $p = 0.001$. Because the p value < 0.05 , there was a difference between BMI with MgSO₄ therapy. Of the 20 patients who received MgSO₄ therapy, 4 patients obtained BMI values above 40 (grade 3 obesity). Of the 19 infants with RDS outcomes, 5 were obtained in infants born to mothers who received MgSO₄ therapy.

Table 2 shows the comparison of infant outcomes between the two groups analyzed using chi-square. There were significant differences in outcomes for gestational age infants and respiratory distress syndrome ($p = 0.011$ and $p = 0.013$, respectively). Whereas, the significance value of $p > 0.05$ was found in birth weight, Apgar scores 1 and 2 minutes, the use of ventilators, sepsis, mortality, and umbilical artery pH.

Table 3 shows the mean serum magnesium levels of mothers in those receiving MgSO₄ therapy of 3.45 ± 1.65 mg/dl with a significance value of $p = 0.001$. Because the p value < 0.05 , the effect of MgSO₄ administration on maternal serum magnesium levels was obtained. The mean magnesium level of infants in those receiving MgSO₄ therapy was 3.10 ± 1.35 mg/dl with a significance value of $p = 0.001$. Because the p value < 0.05 , the effect of MgSO₄ administration on infant serum magnesium levels was obtained.

Table 1. Characteristics of mothers and infants with or without MgSO₄ therapy

Variables	Therapy		Control		P-Value
	N	%	n	%	
Characteristics of Mothers					
Mothers' Age					
16-35 years	21	50	21	50	1.000
>35 years	9	50	9	50	
Age of Pregnancy					
28 - <32 weeks	6	40.0	10	62.5	0.375
32 - ≤34 weeks	24	54.5	20	45.5	
Parity					
Multiparous	22	51.2	21	48.8	1.000
Primiparous	8	47.1	9	52.9	
Body Mass Index (BMI)					
BMI <30	10	27.8	26	72.2	0.001
BMI ≥30	20	83.3	4	16.7	
Lung Maturation					
Yes	22	51.2	22	51.2	1.000
No	8	47.1	8	47.1	
Labor Method					
Abdominal	28	50.9	27	49.1	1.000
Vaginal	2	40	3	60	
Characteristics of Infants					
Breathing					
Spontaneous	17	58.6	12	41.4	0.389
CPAP	11	44	14	56	
Ventilator	2	33.3	4	66.7	
Mortality					
No	29	50	29	50	1.000
Yes	1	50	1	50	
Respiratory distress syndrome					
Yes	5	26.3	14	73.7	0.013
No	25	61	16	39	
Sepsis					
Yes	9	60	6	40	0.552
No	21	46.7	24	53.3	

Table 2. Outcomes of infants born prematurely with or without MgSO₄ therapy

Variables	Therapy (n = 30)	Control (n = 30)	P-Value
Birth Weight (grams)	1743.33±334.19	1880.40±642.86	0.306
Pregnancy Age (weeks)	33.43±1.25	32.2±2.23	0.011
AS in first minute	5.73±1.83	5.83±1.46	0.816
AS in fifth minute	7.10±1.51	7.13±1.16	0.924
CPAP	11 (36.67)	14 (46.67)	0.432
Ventilator	2 (6.67)	4 (13.33)	0.335
Respiratory distress syndrome	5 (16.67)	14 (46.66)	0.013
Sepsis	9 (30)	6 (20)	0.276
Mortality	1 (3.33)	1 (3.33)	0.754
Umbilical arterial pH	7.19±0.09	7.23±0.08	0.123

Data presented as Mean±SD and number (%)

Table 3. Maternal and infant magnesium levels of each group

Variables	Therapy (n = 30)	Control (n = 30)	P-Value
Maternal Magnesium Levels (mg/dl)	3.45±1.65	1.78±0.23	0.001
Infant Magnesium Serum Levels (mg/dl)	3.10±1.35	1.74±0.19	0.001

Table 4. Outcome of infants in groups based on magnesium levels

Groups based on magnesium levels (n = 60)	Group A (<2.5 mg/dl)	Group B (>2.5 - <3.5 mg/dl)	Group C (>3.5 mg/dl)	P-Value
	n = 41	n = 10	n = 9	
AS in first minute	5.8±1.61	6.30±1.05	5.11±2.20	0.499
AS in fifth minute	7.10±1.28	7.60±0.69	6.67±2.00	0.608
CPAP	18 (43.9)	3 (30)	4 (44.44)	0.714
Ventilator	5 (12.19)	0 (0)	1 (11.11)	0.619
Respiratory distress syndrome	16 (39.02)	0 (0)	3 (33.33)	0.013
Sepsis	9 (21.95)	4 (40)	2 (22.22)	0.487
Mortality	2 (4.87)	0 (0)	0 (0)	0.619
Umbilical arterial pH	7.23±0.07	7.22±0.10	7.15±0.11	0.106

Data presented as Mean±SD and number (%)

Table 4 shows the outcomes of respiratory distress syndrome in Group A (<2.5 mg/dl) of 16 patients (39.02%) and Group C (≥3.5 mg/dl) of 3 patients (33.33%). Analysis using chi-square with a significance

value of p = 0.013 was performed. Because the p value <0.05, differences in respiratory distress syndrome outcomes with groups of serum magnesium levels in infants were obtained.

DISCUSSION

In this study, there were significant differences in infant outcomes only in gestational age and RDS between the two groups. For serum magnesium levels, there was a statistically significant difference between serum magnesium levels both in the group receiving MgSO₄ therapy and those not receiving MgSO₄ therapy.

Previous research concluded that the biggest risk factor for RDS was premature birth; from 86.5% of preterm infants, the incidence rate of RDS ranged from 86% at 24 weeks gestation compared to 39 weeks gestation which was only 1% (Qari et al. 2018). Obese mothers have long been considered a risk factor for neonatal morbidity and mortality. Neonatal morbidity is defined as an adverse outcome in the first four weeks of a baby's life. The assessed outcomes can be injuries at birth, low Apgar scores, RDS, sepsis, hypoglycemia, neonatal seizures, treatment at the NICU and even death (Burn et al. 2018). RDS is also a complication of sepsis (Dewi et al. 2016). However, previous study stated antenatal corticosteroid therapy might be a protective factor that can reduce the incidence of RDS in prematurity (Zenita et al. 2018).

A study measured blood magnesium levels. This found that umbilical cord blood magnesium levels were found to increase significantly with an increase in maternal serum magnesium levels (Das et al. 2015). In addition to the dosage regimen, several other factors such as gestational age, birth weight, and maternal serum magnesium levels, can affect higher serum levels of fetal magnesium. Another factor that may contribute to high fetal serum on the umbilical cord is the ability of the placenta to actively transport magnesium to the fetus. Furthermore, magnesium renal clearance in the early days of life is reduced in neonates and especially in preterm neonates, causing magnesium levels to remain high for a long time (Basu et al. 2011).

In this study, no statistically significant differences were found in the umbilical artery pH between groups with or without MgSO₄ and groups with various levels of MgSO₄ levels. Cord blood is the most objective way to assess the metabolic condition of the fetus at birth. Association has been investigated between umbilical artery pH and important clinical outcomes in the prenatal period. Although there is debate about definite pH levels to determine the significance of acidemia, in this study we used pH <7.14 as suggested as a pathological marker of premature infants with acidemia (Simhan 2018).

A study comparing umbilical cord blood gas analysis in premature neonates born to healthy mothers versus preeclampsia mothers found lower pH compared to those in healthy mothers and higher BE in the

preeclampsia group. In addition, metabolic acidosis was more detectable in preterm neonates born to preeclampsia women compared to those born to healthy women (22.2% versus 8.4%, p=0.001) (Sheikh et al. 2016).

In the first and second minute Apgar scores, the use of ventilators, sepsis, mortality and umbilical arterial pH did not differ significantly between groups with or without MgSO₄ and groups with various levels of MgSO₄ levels. This is consistent with WHO recommendation that there is no statistically significant difference between the group that receives MgSO₄ and the group that does not receive active treatment. One measure of infant morbidity reported: RDS, neonatal infection, severe IVH or periventricular leukomalacia, NEC, and use of mechanical ventilation did not have a significant difference in those who used MgSO₄ therapy and those who did not receive therapy (World Health Organization 2015).

The main results from other studies were neonatal morbidity, which is defined as a 5 minute Apgar score of less than 7, hospitalization for more than 5 days, being treated at the NICU, and diagnosed with hypoglycemia, RDS, sepsis, and infant mortality. Results showed a statistically significant increase (P <0.001) in the overall incidence of neonatal morbidity with increasing BMI (Burn et al. 2018). Similar with other studies that infant experienced neonatal mortality occurred with APGAR score ≤ 6 which indicates asphyxia (Putri et al. 2018, Yudrika et al. 2018).

In another study, there was no significant difference between the first and fifth minute score scores and the need for resuscitation, both in groups with and without administration of MgSO₄ or in groups with different levels of MgSO₄ (Basu et al. 2011). A similar review was found in a systematic review that MgSO₄ given specifically to the fetus as neuroprotection did not affect the incidence of Apgar score of 5 minutes <7 (RR 1.03; 95% CI 0.90 in 4387 infants) and hypotonia (RR 1.02 95% CI 0.77 in 2444 infants). Similar to these findings, our study also found no significant differences in the Apgar score (Magee et al. 2011).

This research has various limitations that might be anticipated when conducting further research or other similar research. The limitations include the use of populations with preeclampsia mothers so that the outcomes of MgSO₄ could not be fully evaluated.

CONCLUSION

Significant differences in infant outcomes were only found in gestational age and RDS. Magnesium levels in the blood of newborns with mothers given MgSO₄ therapy were higher than those not treated with MgSO₄.

REFERENCES

- Basu SK, Chickajajur V, Lopez V, Bhutada A, Pagala M, Rastogi S, (2011) Immediate clinical outcomes in preterm neonates receiving antenatal magnesium for neuroprotection. *Journal of perinatal medicine* 40(2): 185-189. <https://doi.org/10.1515/JPM.2011.094>
- Burn SC, Burn MS, Burn P, (2018) Maternal Obesity and its Impact on Neonatal Morbidity.
- Costantine MM, Weiner SJ, (2009) Effects of antenatal exposure to magnesium sulfate on neuroprotection and mortality in preterm infants: a meta-analysis. *Obstetrics and gynecology* 114(2 Pt 1): 354-364. <https://doi.org/10.1097/AOG.0b013e3181ae98c2>
- Das M, Chaudhuri PR, Mondal BC, Mitra S, Bandyopadhyay D, Pramanik S, (2015) Assessment of serum magnesium levels and its outcome in neonates of eclamptic mothers treated with low-dose magnesium sulfate regimen. *Indian journal of pharmacology* 47(5): 502-508. <https://doi.org/10.4103/0253-7613.165183>
- Dewi R, Supriyatno B, Madjid AS, Gunanti, Lubis M, (2016) The effects of colloids or crystalloids on acute respiratory distress syndrome in swine (*Sus scrofa*) models with severe sepsis: Analysis on extravascular lung water, IL-8, and VCAM-1. *Medical Journal of Indonesia* 25(1): 33-38. <https://doi.org/10.13181/mji.v25i1.1204>
- Group MTF-USC, (2007) The Magpie Trial: a randomised trial comparing magnesium sulphate with placebo for pre-eclampsia. Outcome for women at 2 years. *BJOG : an international journal of obstetrics and gynaecology* 114(3): 300-309. <https://doi.org/10.1111/j.1471-0528.2006.01166.x>
- Magee L, Sawchuck D, Synnes A, von Dadelszen P, Basso M, Crane JM, Doyle L, Ehman W, Gagnon R, Grobman W, (2011) Magnesium sulphate for fetal neuroprotection. *Journal of Obstetrics and Gynaecology Canada* 33(5): 516-529.
- Pasaribu HP, Hariman H, Roeshadi RH, Koh SCL, (2016) Soluble vascular cell adhesion molecule-1 and magnesium sulfate with nifedipine treatment in Indonesian women with severe pre-eclampsia. *Interventional Medicine and Applied Science* 8(3): 97-102. <https://doi.org/10.1556/1646.8.2016.3.4>
- Putri AR, Ansar J, Arsyad DS, (2018) Descriptive epidemiology of neonatal mortality in Gowa district 2015. In: *International Conference on Medical and Health Informatics*. Association for Computing Machinery <https://doi.org/10.1145/3242789.3242827>
- Qari SA, Alsufyani AA, Muathin SH, El Margoushy NM, (2018) Prevalence of respiratory distress syndrome in neonates. *The Egyptian journal of hospital medicine* 70(2): 257-264.
- Rouse DJ, Hirtz DG, Thom E, Varner MW, Spong CY, Mercer BM, Iams JD, Wapner RJ, Sorokin Y, Alexander JM, (2008) A randomized, controlled trial of magnesium sulfate for the prevention of cerebral palsy. *New England Journal of Medicine* 359(9): 895-905.
- Salfi QN, Saharso D, Atika A, (2019) Profile of Cerebral Palsy Patients in Dr. Soetomo General Hospital Surabaya, Indonesia. *Biomolecular and Health Science Journal* 2(1): 13-16.
- Sheikh M, Zoham MH, Hantoushzadeh S, Shariat M, Dalili H, Amini E, (2016) Umbilical blood gas analysis in preeclamptic versus healthy pregnancies with preterm birth. *The Journal of Maternal-Fetal & Neonatal Medicine* 29(15): 2548-2553.
- Simhan HN, (2018) Umbilical cord blood acid-base analysis at delivery.
- Sukatendel K, Hasibuan CL, Pasaribu HP, Sihite H, Ardyansah E, Situmorang MF, (2018) Risk factor for preterm labor in Haji Adam Malik General Hospital, Pirmgadi General hospital and satellite hospitals in Medan from January 2014 to December 2016. In: *IOP Conference Series Earth and Environmental Science* Vol. 125. Institute of Physics Publishing <https://doi.org/10.1088/1755-1315/125/1/012192>
- World Health Organization, (2015) WHO recommendations on interventions to improve preterm birth outcomes: evidence base. World Health Organization
- Yudrika A, Zulkiffi A, Muis M, (2018) Risk factors of neonatal mortality incidence in Muna regency in 2014. In: *International Conference on Medical and Health Informatics*. <https://doi.org/10.1145/3239438.3239489>
- Zeng X, Xue Y, Tian Q, Sun R, An R, (2016) Effects and safety of magnesium sulfate on neuroprotection: a meta-analysis based on PRISMA guidelines. *Medicine* 95(1):.
- Zenita DP, Utomo MT, Darmawan E, (2018) The Effect of Antenatal Corticosteroid Therapy to Respiratory Distress Syndrome Event on Preterm Infants in Surabaya. *Biomolecular and Health Science Journal* 1(1): 40-42.

Comparison of short-term

ORIGINALITY REPORT

18%

SIMILARITY INDEX

14%

INTERNET SOURCES

12%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Mahdi Sheikh, Mojdeh Habibi Zoham, Sedigheh Hantoushzadeh, Mamak Shariat, Hosein Dalili, Elaheh Amini. "Umbilical blood gas analysis in preeclamptic versus healthy pregnancies with preterm birth", The Journal of Maternal-Fetal & Neonatal Medicine, 2015
Publication 2%
 - 2 www.medcraveonline.com
Internet Source 1%
 - 3 Sudepta Kumar Basu, Vijay Chickajajur, Vivian Lopez, Alok Bhutada, Murali Pagala, Shantanu Rastogi. "Immediate clinical outcomes in preterm neonates receiving antenatal magnesium for neuroprotection", Journal of Perinatal Medicine, 2012
Publication 1%
 - 4 "Oral Abstracts", Journal of Paediatrics and Child Health, 4/2007
Publication 1%
 - 5 thieme-connect.com
Internet Source 1%
-

6	paper.researchbib.com Internet Source	1 %
7	docplayer.net Internet Source	1 %
8	www.frontiersin.org Internet Source	1 %
9	www.ghrnet.org Internet Source	1 %
10	www.openaccess.hacettepe.edu.tr:8080 Internet Source	1 %
11	escholarship.org Internet Source	1 %
12	Submitted to iGroup Student Paper	1 %
13	repository-tnmgrmu.ac.in Internet Source	<1 %
14	L. García Alonso, M. Pumarada Prieto, E. González Colmenero, A. Concheiro Guisán et al. "Prenatal Therapy with Magnesium Sulfate and Its Correlation with Neonatal Serum Magnesium Concentration", American Journal of Perinatology, 2017 Publication	<1 %
15	www.ijp-online.com Internet Source	<1 %

- 16 Dalla Doohan, Muhammad Miftahussurur, Yuichi Matsuo, Yasutoshi Kido et al. "Characterization of a novel Helicobacter pylori East Asian-type CagA ELISA for detecting patients infected with various cagA genotypes", Medical Microbiology and Immunology, 2019
Publication <1 %
-
- 17 repository.binausadabali.ac.id
Internet Source <1 %
-
- 18 pubmed.ncbi.nlm.nih.gov
Internet Source <1 %
-
- 19 www.tjoddergisi.org
Internet Source <1 %
-
- 20 Adra Yudrika, Andi Zulkifli, Masyita Muis. "Risk Factors of Neonatal Mortality Incidence in Muna Regency in 2014", Proceedings of the 2nd International Conference on Medical and Health Informatics - ICMHI '18, 2018
Publication <1 %
-
- 21 Etsuko Shimada, Masaki Ogawa, Yoshio Matsuda, Minoru Mitani, Hideo Matsui. "Umbilical artery pH may be a possible confounder for neonatal adverse outcomes in preterm infants exposed to antenatal magnesium", The Journal of Maternal-Fetal & Neonatal Medicine, 2012 <1 %

22 www.coursehero.com <1 %
Internet Source

23 archive.ircmj.com <1 %
Internet Source

24 bmcpediatr.biomedcentral.com <1 %
Internet Source

25 onlinelibrary.wiley.com <1 %
Internet Source

26 www.researchsquare.com <1 %
Internet Source

27 www.slideshare.net <1 %
Internet Source

28 www.thieme-connect.de <1 %
Internet Source

29 Alicia K. Yee, Leon S. Siriwardhana, Gillian M. Nixon, Lisa M. Walter, Flora Y. Wong, Rosemary S.C. Horne. "Duration and Consequences of Periodic Breathing in Infants Born Preterm Before and After Hospital Discharge", The Journal of Pediatrics, 2023
Publication

30 Nirmal Chandra Sahu, Pravat Kumar Thatoi, Sarada Priyadarsini Suna, Sai Swaroop, Mohanty C B K. "SERUM MAGNESIUM IN <1 %

CRITICALLY ILL PATIENTS AND ITS
CORRELATION WITH OUTCOME", Journal of
Evolution of Medical and Dental Sciences,
2018

Publication

31

Ainhoa Gochi Valdovinos, María Arriaga-Redondo, Ester Dejuan Bitriá, Isabel Pérez Rodríguez et al. "Prenatal therapy with magnesium sulphate and intestinal obstruction due to meconium in preterm newborns", Anales de Pediatría (English Edition), 2022

Publication

<1 %

32

Falkner, D.. "Serum magnesium levels in asthmatic patients during acute exacerbations of asthma", American Journal of Emergency Medicine, 199201

Publication

<1 %

33

Robert Galinsky, Justin M. Dean, Ingran Lingam, Nicola J. Robertson, Carina Mallard, Laura Bennet, Alistair J. Gunn. "A Systematic Review of Magnesium Sulfate for Perinatal Neuroprotection: What Have We Learnt From the Past Decade?", Frontiers in Neurology, 2020

Publication

<1 %

34

Soroush Aalipour, Sedigheh Hantoushzadeh, Mamak Shariat, Sadaf Sahraian, Mahdi

<1 %

Sheikh. "Umbilical Cord Blood Acidosis in Term Pregnancies With Gestational Diabetes Mellitus and Its Relations to Maternal Factors and Neonatal Outcomes", Iranian Red Crescent Medical Journal, 2018

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Comparison of short-term

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
