

Low molecular weight heparin is associated with better outcomes than unfractionated heparin

by Mochamad Yusuf Alsagaff

Submission date: 03-Mar-2023 10:59AM (UTC+0800)

Submission ID: 2027562604

File name: associated_with_better_outcomes_than_unfractionated_heparin.pdf (807.26K)

Word count: 5825

Character count: 32559

Low molecular weight heparin is associated with better outcomes than unfractionated heparin for thromboprophylaxis in hospitalized COVID-19 patients: a meta-analysis

Mochamad Yusuf Alsagaff^{1,*}, Eka Prasetya Budi Mulia¹, Irma Maghfirah¹, Yusuf Azmi¹, Dita Aulia Rachmi¹, Alqi Yutha¹, Luqman Hakim Andira¹ and Bambang Pujo Semedi²

¹Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga—Dr Soetomo General Hospital, Mayjen Prof. Dr Moestopo Street No. 47, Surabaya 60132, Indonesia; and ²Department of Anesthesiology and Reanimation, Faculty of Medicine, Universitas Airlangga—Dr Soetomo General Hospital, Surabaya 60132, Indonesia

Received 16 May 2022; revised 21 July 2022; accepted 29 July 2022; online publish-ahead-of-print 3 August 2022

Aims

This study aimed to compare the outcomes of the administration of LMWH and UFH in hospitalized COVID-19 patients.

Methods and results

We systematically searched several databases and included observational studies or clinical trials that compared the outcomes of the administration of LMWH and UFH in hospitalized COVID-19 patients. A total of nine studies comprising 9637 patients were included. Meta-analysis showed that LMWH administration was associated with a lower in-hospital mortality and 28/30-day mortality compared with UFH administration {[relative risk (RR) 0.44; 95% confidence interval (95% CI) 0.32–0.61; I^2 : 87.9%] and (RR 0.45; 95% CI 0.24–0.86; I^2 : 78.4%), respectively}. Patient with LMWH had shorter duration of hospital and ICU length of stay compared with UFH {[weighted mean difference (WMD) –2.20; 95% CI –3.01 to –1.40; I^2 : 0%] and (WMD –1.41; 95% CI –2.20 to –0.63; I^2 : 0%), respectively}. The risk of ICU admission or mechanical ventilation was lower in patients who received LMWH than in those who received UFH (RR 0.67; 95% CI 0.55–0.81; I^2 : 67.3%). However, there was no difference in the incidence of bleeding with LMWH compared with UFH (RR 0.27; 95% CI 0.07–1.01; I^2 : 64.6%).

Conclusion


Our meta-analysis showed that administration of LMWH was associated with better outcomes compared with UFH in hospitalized COVID-19 patients. Prospective cohorts and RCTs are urgently needed to explore the definitive effect of LMWH to provide direct high-certainty evidence.

PROSPERO registration number: CRD42021271977

* Corresponding author. Tel: +6231-5501601, Fax: +6231-5031752, Email: yusuf_505@fk.unair.ac.id

© The Author(s) 2022. Published by Oxford University Press on behalf of the European Society of Cardiology. All rights reserved. For permissions, please e-mail: journals.permissions@oup.com

Graphical Abstract



Keywords

COVID-19 • Low molecular weight heparin • Unfractionated heparin • Thromboprophylaxis • Mortality

Introduction

Coronavirus Disease 2019 (COVID-19) has spread rapidly around the world, causing high morbidity and mortality. Although respiratory symptoms dominate the clinical manifestations of COVID-19, some patients have an increased risk of thromboembolism due to coagulopathy complications.^{1,2} Coagulopathy is one of the severe complications with a high incidence in COVID-19 patients.^{3,4} Elevated coagulation parameters, including D-dimer, prothrombin time, and fibrinogen, are significantly associated with poor prognosis in patients with COVID-19.^{5–7} Activation of the coagulation pathway during the immunologic response to infection may also lead to overproduction of pro-inflammatory cytokines resulting in multiorgan damage.⁸

Early reports demonstrated that anticoagulant thromboprophylaxis was associated with better outcomes in hospitalized COVID-19 patients.^{9,10} Currently, several recommendations suggest venous thromboembolism (VTE) risk stratification and the use of prophylactic anticoagulants in all hospitalized COVID-19 patients. In contrast, evidence is insufficient to support thromboprophylaxis with therapeutic doses of anticoagulants.^{11–13} A meta-analysis showed that anticoagulant regimens in therapeutic and prophylactic doses decreased in-hospital mortality in COVID-19 patients. However, prophylactic doses may be preferred due to the increased risk of bleeding in therapeutic doses of anticoagulants.¹⁴ Another study also demonstrated a similar result that intermediate-to-therapeutic doses of anticoagulant increased mortality and major bleeding compared with prophylactic doses.¹⁵ These findings demonstrate a favourable profile of the efficacy and safety of prophylactic anticoagulation in hospitalized COVID-19 patients.

Although the choice of prophylactic dosing regimen for anticoagulation becomes more evident, the type of anticoagulant itself remains a question. A spectrum of anticoagulants such as low molecular weight heparin (LMWH), unfractionated heparin (UFH), fondaparinux, and direct oral anticoagulant (DOAC) are used in the management of COVID-19 patients, with UFH and LMWH as the most commonly prescribed.¹⁶ A recent study in matched cohorts of COVID-19 patients showed that enoxaparin was associated with lower 28-day mortality compared with UFH.¹⁷ The plausible superiority of LMWH over UFH was also demonstrated in previous studies with different settings.^{18–21} Therefore, the objective of our systematic review and meta-analysis was to compare the outcomes of administration of the LMWH and UFH in hospitalized COVID-19 patients.

Methods

Eligibility criteria

Our analysis included studies that assessed the outcomes of LMWH and UFH administration in hospitalized COVID-19 adult patients. We included articles published in English and studies with extractable data

for analysis. Other publications such as case reports, case series, review articles, editorials, and *in vitro* or animal studies were excluded. Studies on special populations such as pregnant women and children were also excluded.

Search strategy and study selection

We conducted a systematic literature search based on the search strategy presented in Supplementary material online, Table S1 and finalized the search on 21 June 2022. We searched several databases, including PubMed, ScienceDirect, Scopus, Cochrane Library, and Europe PMC. After eliminating duplication from the literature search, three authors (E.P.B.M., I.M., and Y.A.) independently screened the title and abstract. Eligibility criteria were used to assess the full article. The differences in article assessment were solved by a discussion with the senior authors (M.Y.A. and B.P.S.). The research protocol was registered in the International Prospective Register of Systematic Reviews (PROSPERO) database (registration number CRD42021271977). This research followed the recommendations described in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement.²²

Data collection process


Piloted data extraction forms were used independently by three authors (D.A.R., A.Y., and L.H.A.) to extract data that consisted of the author, publication date, design of the study, population studied, intervention administered, comparison or control, and outcomes. The comparative intervention assessed in this study was the administration of LMWH and UFH in hospitalized COVID-19 patients. The primary outcome was in-hospital mortality, while the additional outcomes were 28- or 30-day mortality, intensive care unit (ICU) admission or mechanical ventilation, hospital and ICU length of stay, and bleeding complications. We presented categorical and continuous variables as mean \pm standard deviation (SD) and frequency (percentage), respectively.

Quality assessment

All included observational studies were assessed by three authors (E.P.B.M., I.M., and Y.A.) independently using the Newcastle–Ottawa score (NOS), which consisted of three domains that included sample selection, comparability of cohorts, and outcomes assessment.²³ The risk of bias for the included randomized controlled trial (RCT) was assessed by the modified Cochrane Collaboration tool.²⁴ Discussions were held to resolve the discrepancies between the authors.

Data analysis

Data used for each included study was following univariate analysis in the original study. Pooled effect estimates of the outcomes were reported as weighted mean difference (WMD) and relative risk (RR) for the continuous and dichotomous variables, respectively. Data with low heterogeneity



(I^2 statistic < 50% or P -value > 0.1) were pooled with a fixed-effects model, and data with high heterogeneity (I^2 statistic > 50% or P -value < 0.1) were pooled with a random-effects model. P -value < 0.05 was determined as statistical significance. We used funnel-plot analysis to assess the publication bias qualitatively. Egger's test and Harbord's test were used to further assess publication bias on continuous and dichotomous variables, respectively. Furthermore, univariate meta-regression analysis was performed to determine which factors contributed to in-hospital mortality in hospitalized COVID-19 patients receiving two different anticoagulants of LMWH and UFH. Meta-analysis was performed on Stata software V.14.0 (College Station).

Results

Study characteristics

We identified 6243 articles from the initial database search and 2 articles through additional search and then removed the duplicates. The remaining 4701 articles were screened for titles and abstracts, which resulted in 4406 articles being excluded. Using the eligibility criteria, 295 potential full-text articles were assessed. Finally, seven observational studies^{10,17,25-29} and two RCTs^{30,31} consisting of 9637 patients were included for qualitative analysis and meta-analysis (Figure 1 and Table 1).

Table 1 Characteristics of included studies

No	Author, year	Study design	Population	Setting	Time period	Prophylactic doses	Therapeutic doses	Outcome extracted	Bleeding definition	N	Age, year	Male, %	HTN	DM	CKD	Cancer	Quality
1	Voltes, 2022 ²⁹	Retrospective single centre	Critically ill adult patients	Stony Brook University Hospital	7 February–17 May 2020	Enoxaparin: 40 mg daily (D-dimer < 1000 ng/mL) or 40 mg twice a day (D-dimer ≥ 1000 ng/mL but < 3000 ng/mL)	Enoxaparin: 1 mg/kg twice a day or UFH: IV starting rate of 18 units/kg/h to achieve a goal PTT of 60–90	Mortality	NA	240 (LMWH ther: 70; UFH ther: 65; proph: 11; ther: 72)	≥ 18	NA	NA	NA	NA	NA	7
2	Olynyk, 2021 ³¹	Randomized control trial single centre	Severe hospitalized adult patients	Kyiv City Clinical Hospital No. 4	1 July 2020–1 March 2021	Enoxaparin: 50 anti-Xa IU/kg SC QD	Enoxaparin: 100 anti-Xa IU/kg SC BID; UFH: IV initial 80 U/kg/h, followed by 18 U/kg/h with APTT of 40–70 s	28-day mortality, mechanical ventilation	NA	126 (LMWH ther: 42; UFH ther: 42; UFH ther: 42)	70.3 ± 2.8	60.3	NA	NA	NA	NA	Low overall risk of bias
3	GiL, 2021 ²⁵	Retrospective single centre	Hospitalized adult patients	Tertiary care centre	20–31 March 2020	Enoxaparin: 40 mg SC once a day (BMI < 40; GFR = 30) or 30 mg SC twice a day (BMI = 40); UFH: NA	Enoxaparin: 1–5 mg/kg/day or 1 mg/kg twice a day; UFH 80 units/kg IV bolus followed by continuous IV infusion	Mortality	NA	225 (LMWH proph: 127; UFH proph: 4; UFH proph: 20; UFH ther: 10)	62.1 ± 14.4	57.8	64.9	38.7	19.1	8.4	8
4	Kirkup, 2021 ²⁶	Retrospective multicentre	Hospitalized adult patients	192 hospitals in 20 countries	Start time NA–4 January 2021	NA	NA	Mortality, 28-day mortality, ICU admission, length of stay	NA	3196 (LMWH: 1939; UFH: 1012)	63.6 ± 28.2	57.1	54.0	34.9	12.8	6.7	9
5	Lopes, 2021 ³⁰	Randomized control trial multicentre	Hospitalized adult patients and elevated D-dimer concentration	31 hospitals in Brazil	24 June 2020–26 February 2021	Enoxaparin: 40 mg QD SC (GFR ≥ 30, BMI < 40), 60 mg QD SC or 40 mg BID SC (GFR ≥ 30, BMI < 40); UFH: 5000 units TID/BID SC (BMI < 40, age ≥ 75), 1 mg/kg BID SC (BMI ≥ 40)	Enoxaparin: 1 mg/kg BID SC or 1.5 QD SC mg/kg (GFR ≥ 30, BMI < 40); 0.75 mg/kg BID SC (GFR ≥ 30, BMI < 40, age ≥ 75), 1 mg/kg BID SC (GFR ≥ 30, BMI ≥ 40); 1 mg/kg QD SC (GFR < 30, BMI < 40); UFH: 60 unit/kg IV bolus, then 12 units/kg/h	Mortality, bleeding	Major bleeding by ISTH criteria	615 (LMWH: 256; UFH: 47)	56.6 ± 14.3	59.8	49.1	24.4	NA	0	Low overall risk of bias
6	Nadarni, 2020 ⁷	Retrospective multicentre	Hospitalized adult patients	5 New York City hospitals	1 March–3 April 2020	Enoxaparin: once daily; UFH: SC	Enoxaparin: 1 mg/kg twice daily or 1.5 mg/kg daily; UFH: continuous IV infusions	Mortality, mechanical ventilation, major bleeding	Major bleeding by CD 10th Revision codes	2859 (LMWH: 445; UFH: 941)	66.5 ± 16.2	57.9	37.4	24.7	12.1	8.3	7

Table 1 Continued

Author, year	Study design	Population	Setting	Time period	Prophylactic doses	Therapeutic doses	Outcome extracted	Bleeding definition	N	Age, year	Male, %	HTN	DM	CKD	Cancer	Quality
7 Pawlowski, 2020 ¹⁷	Retrospective multicentre	Patients admitted to hospital	19 Mayo Clinic sites in Arizona, Florida, Minnesota, and Wisconsin	4 April–31 August 2020	Enoxaparin: ≤ 40 mg/day SC; UFH: periodic administration	Enoxaparin: > 40 mg/day SC; UFH: continuous administration	Mortality: 28-day mortality; ICU admission, length of stay, bleeding	Bleeding complications from the physician notes	557 (LMWH: 441; UFH: 166)	58.4 ± 17.9	54.5	3.1	180	112	5.9	8
8 Piazza, 2020 ²⁸	Retrospective multicentre	Hospitalized adult patients	The Mass General Brigham integrated health network	13 March–3 April 2020	NA	NA	ICU admission	NA	399 (LMWH: 277; UFH: 102)	61.1 ± 17.1	42.6	56.1	336	NA	6.5	7
9 Rentsch, 2020 ¹⁰	Retrospective multicentre	Patients admitted to hospital	> 1200 points of care nationwide of US Department of Veterans Affairs	1 March–31 July 2020	Enoxaparin: 40 mg QD or 30 mg BID SC; UFH: 5000 units BID or TID SC	Enoxaparin: > 40 mg QD SC; UFH: IV dose adjusted on PTT	Mortality: 30-day mortality	NA	3627 (LMWH: 2506; UFH: 1094)	67.0 ± 12.3	93.6	66.6	434	19.1	13.6	9

N, number of samples; UFH, unfractionated heparin; HTN, hypertension; DM, diabetes mellitus; CKD, chronic kidney disease; BMI, body mass index; GFR, glomerular filtration rate; NA, not available; QD, quaque die means once a day; BID, bis in die means twice a day; TID, ter in die means three times a day; IV, intravenous; SC, subcutaneous; ICU, intensive care unit; ISTH, International Society on Thrombosis and Haemostasis; proph, prophylactic; ther, therapeutic.




Panel A: in-hospital mortality**Panel B: 28/30-day mortality**

Figure 2 (A) forest plot for the association of two different types of anticoagulant (LMWH vs. UFH) with in-hospital mortality in hospitalized COVID-19 patients. (B) forest plot for the association of two different types of anticoagulant (LMWH vs. UFH) with 28- or 30-day mortality in COVID-19 patients.



Outcome comparison between low molecular weight heparin and unfractionated heparin

The meta-analysis showed that LMWH administration was associated with a lower risk of in-hospital mortality compared with UFH administration [(RR 0.44; 95% CI 0.32–0.61; I^2 : 87.9%) and (RR 0.45; 95% CI 0.24–0.86; I^2 : 78.4%), respectively]. These effects were comparable to the pooled risk estimates comprising both doses of therapeutic and prophylactic (RR 0.28; 95% CI 0.17–0.47; P = 0.001; I^2 : 86.3%) and solely prophylactic doses (RR 0.63; 95% CI 0.46–0.87; P = 0.005; I^2 : 65.6%) (Figure 2). Similar results were obtained at 1-month follow-up that LMWH was associated with a lower 28- or 30-day mortality compared with UFH (RR 0.45; 95% CI 0.24–0.86; I^2 : 78.4%) (Figure 2).

Patients who received therapeutic and prophylactic doses of LMWH had a shorter duration of hospital and ICU length of stay than those who received therapeutic and prophylactic doses of UFH [(WMD -2.20; 95% CI -3.01 to -1.40; P < 0.001; I^2 : 0%) and (WMD -1.41; 95% CI -2.20 to -0.63; P < 0.001; I^2 : 0%), respectively] (Figure 3). The risk of ICU admission or mechanical ventilation was also lower in patients who received LMWH than in those who received UFH (RR 0.67; 95% CI 0.55–0.81; P = 0.016; I^2 : 67.3%) (Figure 4). However, there was no significant difference in the incidence of bleeding with LMWH compared with UFH (RR 0.27; 95% CI 0.07–1.01; P = 0.052; I^2 : 64.6%) (Supplementary material online, Figure S1).

Meta-regression

Univariate meta-regression analysis revealed that the association of two different anticoagulants of LMWH and UFH with in-hospital mortality in COVID-19 patients was not significantly affected by age (P = 0.463), gender (P = 0.954), hypertension (P = 0.571), diabetes (P = 0.912), chronic kidney disease (P = 0.597), cancer (P = 0.802), chronic lung disease (P = 0.604), heart failure (P = 0.812), coronary artery disease (P = 0.901), asthma (P = 0.742), chronic liver disease (P = 0.358), and stroke and cerebrovascular disease (P = 0.398) (Supplementary material online, Table S2).

Quality assessment and publication bias

Seven studies were of high quality based on the assessment using NOS (Supplementary material online, Table S3), and two RCTs showed a low overall risk of bias based on the assessment using the modified Cochrane Collaboration tool (Supplementary material online, Table S4). The asymmetric shape was obtained by visually qualitative assessment of the funnel plot for analysis on the variables of anticoagulant selection and mortality, indicating possible publication bias (Supplementary material online, Figure S2). However, quantitative analysis using the regression-based Harbord's test showed no small-study effects (P = 0.876) on the same variable.




Figure 4 Forest plot for the association of two different types of anticoagulant (LMWH vs. UFH) with ICU admission or mechanical ventilation in all hospitalized COVID-19 patients.

GRADE assessment

Grading of Recommendations Assessment, Development, and Evaluation (GRADE) assessment indicated a very low certainty of the evidence for the effect of LMWH and UFH on in-hospital mortality, 28- or 30-day mortality, hospital and ICU length of stay, ICU admission or mechanical ventilation, and bleeding complications. The certainty of the evidence was very low for both subgroups of in-hospital mortality outcomes (Supplementary material online, Table S5).

Discussion

This systematic review and meta-analysis highlighted the differences in outcomes associated with the administration of LMWH and UFH in hospitalized COVID-19 patients. We found that LMWH was associated with lower in-hospital mortality, 28- or 30-day mortality, ICU admission or mechanical ventilation, and bleeding complications compared with UFH. Furthermore, we found that patients treated with LMWH had a shorter hospital and ICU length of stay compared with those treated with UFH.

A spectrum of anticoagulants, including LMWH, UFH, fondaparinux, and DOAC, are used in managing COVID-19 patients, both as prophylactic and therapeutic approaches.³² Heparin is one of the anticoagulant drugs that is classified according to its molecular weight and is used for the prophylaxis and treatment of venous and arterial thrombosis.³³ Only UFH comprising at least 18 saccharide sequences can bind to and potentiate the activity of antithrombin. However, UFH of any length comprising a unique pentasaccharide sequence can inhibit the action of factor Xa.³⁴ This pentasaccharide sequence is used as the basis for developing LMWH. In contrast to UFH, whose main action is mediated by thrombin inhibition, LMWH inhibits prothrombin activation more robustly through its action on factor Xa.³⁵

The proposed mechanisms of using heparin in COVID-19 management are to block uncontrolled blood clotting and prevent VTE and other thrombotic events.³⁶ However, heparin may also offer anti-inflammatory, anti-complement activity, anti-viral, and immunomodulatory effects, which may benefit beyond the anticoagulation in COVID-19.³⁷ Recent studies have also reported that soluble heparin inhibited viral entry by competing with heparan sulfate proteoglycan for binding to the SARS-CoV-2 spike protein, thereby inhibiting the attachment of SARS-CoV-2 to the surface of the host cell.³⁸

The superiority of LMWH over UFH has been demonstrated by previous studies in several different settings. A meta-analysis of RCTs comparing the administration of anticoagulant thromboprophylaxis in intensive care patients showed that LMWH was associated with a better net clinical benefit and a lower risk of DVT compared with UFH.²¹ Another meta-analysis of patients with VTE showed that compared with UFH, initial treatment with LMWH was more effective in preventing recurrent VTE, reduced the risk of bleeding, and was associated with a lower mortality rate at follow-up.¹⁹ In addition, the incidence of pulmonary embolism and proximal DVT was lower in patients undergoing orthopaedic surgery who received LMWH than UFH.²⁰

There are several reasons that explain the plausible superiority of LMWH over UFH. Heparin-induced thrombocytopenia (HIT) is less common with LMWH than with UFH. A meta-analysis evaluating the incidence of HIT in patients at risk for VTE treated with prophylactic doses of anticoagulant showed that the incidence of HIT was lower with LMWH (0.2%) than with UFH (2.6%).³⁹ The peak anti-Xa activity (C_{max}) and area under the curve (AUC) are higher with enoxaparin and dalteparin compared with UFH.⁴⁰ In addition, administration of heparin increases the release of TPI antigen, an inhibitor of tissue factor that plays a role in the antithrombotic effect of heparin. However, unlike UFH, the LMWH administration can

increase TPF that lasts longer in circulation. This finding may explain the different antithrombotic efficacy of LMWH and UFH in clinical settings.⁴¹ LMWH is also better than UFH at suppressing inflammatory tissue factor expression and contributes to endothelial haemostatic properties in the microvasculature, where endothelial dysfunction is associated with poor outcomes in COVID-19 patients.^{42,43} Finally, LMWH is preferred over UFH because of its good predictability, dose-dependent plasma levels, and longer plasma half-life, allowing it to be administered subcutaneously once or twice daily and reducing healthcare worker exposure.³³

Clinical implication

Our meta-analysis suggests that LMWH may provide better outcomes in COVID-19 patients. These results may provide evidence for the current recommendation that LMWH is preferred over UFH as thromboprophylaxis for COVID-19.^{12,16,44,45} However, this new evidence does not justify changing existing guidelines and is insufficient to support or oppose using UFH as thromboprophylaxis in hospitalized COVID-19 patients. Currently, the selection of anticoagulant agents may still be based on the availability of anticoagulants, the experience of the physicians, treatment objectives, and other patient-specific factors.¹¹ Further evidence from RCTs is urgently needed to determine the most effective anticoagulant agents for thromboprophylaxis in COVID-19 patients.

Limitations

There are several limitations to this meta-analysis. Possible publication bias was noted in several outcomes, as well as substantial heterogeneity. Most of the included studies were retrospective observational, which were not matched or adjusted for confounders, so the strength of the association could not be measured accurately. The dose definition of the prophylactic or therapeutic LMWH or UFH, including the route of administration, also varied slightly across studies. Another issue is that the presence of both prophylactic and therapeutic dosing in the studies included outcomes, especially in non-critically ill COVID-19 patients, as shown in recent studies.^{46,47} Some studies did not mention the specific type of LMWH, and not all studies provided details regarding the selection criteria for LMWH or UFH (the reason for using LMWH or UFH at an individual level cannot be identified), such as the use of UFH, which is the preferred choice for patients with renal dysfunction or disseminated intravascular coagulation. These problems may translate into uncertain effect estimates from these individual studies. Most studies also did not mention pre-existing conditions that have been anticoagulated before admission. The definition of bleeding also varied across studies. Lastly, one study by Lopes *et al.*³⁰ only included hospitalized adult patients with elevated D-dimer levels.

Conclusion

Our meta-analysis highlighted current evidence that administration of LMWH was associated with better outcomes compared with UFH in hospitalized COVID-19 patients. Nevertheless, this meta-analysis does not provide guidance for therapeutic decision-making or change the existing guidelines for COVID-19 thromboprophylaxis. Until direct high-certainty evidence comparing these two types of anticoagulants is available, clinical inference from this analysis should be drawn with caution. Prospective cohorts and RCTs are urgently needed to explore the definitive effects of LMWH and UFH to provide direct high-certainty evidence.

Supplementary material

Supplementary material is available at *European Heart Journal—Quality of Care and Clinical Outcomes* online.

Acknowledgement

None.

Funding

None.

Conflict of interest: The authors declared no conflict of interest.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Data availability

All data generated or analysed during this study are included in this published article. The corresponding author (A) can be contacted for more information.

Credit authorship contribution statement

M.Y.A. and B.P.S. contributed to the study's conception and design. E.P.B.M., I.M., and Y.A. performed article screening and quality assessment of the studies. D.A.R., A.Y., and L.H.A. performed literature retrieval and data extraction. E.P.B.M. and Y.A. wrote the first draft of the manuscript. All authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

References

- Clerkin KJ, Fried JA, Raikhelkar J, Sayer G, Griffin JM, Masoumi A *et al.* COVID-19 and cardiovascular disease. *Circulation* 2020;**141**:1648–1655.
- Zheng Y-Y, Ma Yi-T, Zhang J-Y, Xie X. COVID-19 and the cardiovascular system. *Nat Rev Cardiol* 2020;**17**:259–260.
- Klok FA, Kruip MJHA, Van Der Meer NJM, Arbous MS, Gommers DAMPJ, Kant KM *et al.* Incidence of thrombotic complications in critically ill ICU patients with COVID-19. *Thromb Res* 2020;**191**:145–147.
- Helms J, Tacquard C, Severac F, Leonard-Lorant I, Ohana M, Delabranche X *et al.* High risk of thrombosis in patients with severe SARS-CoV-2 infection: a multicenter prospective cohort study. *Intensive Care Med* 2020;**46**:1089–1098.
- Polimeni A, Leo I, Spaccarotella C, Mongiardo A, Sorrentino S, Sabatino J *et al.* Differences in coagulopathy indices in patients with severe versus non-severe COVID-19: a meta-analysis of 35 studies and 6427 patients. *Sci Rep* 2021;**11**:10464.
- Nugroho J, Wardhana A, Mulia EP, Maghfirah I, Rachmi DA, Ayun MQ *et al.* Elevated fibrinogen and fibrin degradation product are associated with poor outcome in COVID-19 patients: a meta-analysis. *Clin Hemorheol Microcirc* 2021;**77**:221–231.
- Nugroho J, Wardhana A, Maghfirah I, Mulia EP, Rachmi DA, Ayun MQ *et al.* Relationship of D-dimer with severity and mortality in SARS-CoV-2 patients: a meta-analysis. *Int J Lab Hematol* 2021;**43**:110–115.
- Jose RJ, Manuel A. COVID-19 cytokine storm: the interplay between inflammation and coagulation. *Lancet Respir Med* 2020;**8**:e46–e47.
- Tang N, Bai H, Chen X, Gong J, Li D, Sun Z. Anticoagulant treatment is associated with decreased mortality in severe coronavirus disease 2019 patients with coagulopathy. *J Thromb Haemost* 2020;**18**:1094–1099.
- Rentsch CT, Beckman JA, Tomlinson L *et al.* Early initiation of prophylactic anticoagulation for prevention of coronavirus disease 2019 mortality in patients admitted to hospital in the United States: Cohort study. *BMJ* 2021;**372**:n311.
- Cuker A, Tseng EK, Nieuwaat R, Anchaisuksiri P, Blair C, Dane K *et al.* American Society of Hematology 2021 guidelines on the use of anticoagulation for thromboprophylaxis in patients with COVID-19. *Blood Adv* 2021;**5**:872–888.

12. Thachil J, Tang N, Gando S, Falanga A, Cattaneo M, Levi M et al. ISTH interim guidance on recognition and management of coagulopathy in COVID-19. *J Thromb Haemost* 2020;**18**:1023–1026.
13. Marietta M, Ageno W, Artoni A, De Candia E, Greslele P, Marchetti M et al. COVID-19 and haemostasis: a position paper from Italian Society on Thrombosis and Haemostasis (SISET). *Blood Transfus* 2020;**18**:167–169.
14. Parisi R, Costanzo S, Di Castelnuovo A, De Gaetano G, Donati MB, Iacoviello L. Different anticoagulant regimens, mortality, and bleeding in hospitalized patients with COVID-19: a systematic review and an updated meta-analysis. *Semin Thromb Hemost* 2021;**47**:372–391.
15. Moonla C, Sosothikul D, Chiasakul T, Rojnuckarin P, Uaprasert N. Anticoagulation and in-hospital mortality from coronavirus disease 2019: a systematic review and meta-analysis. *Clin Appl Thromb Hemost* 2021;**27**. doi:10.1177/10760296211008999.
16. Chandra A, Chakraborty U, Ghosh S, Dasgupta S. Anticoagulation in COVID-19: current concepts and controversies. *Postgrad Med J* 2022;**98**:395–402.
17. Pawlowski C, Venkatakrishnan Aj, Kirkup C, Berner G, Puranik A, O'horo JC et al. Enoxaparin is associated with lower rates of mortality than unfractionated Heparin in hospitalized COVID-19 patients. *EClinicalMedicine* 2021;**33**:100774.
18. Jacobs BN, Cain-Nielsen AH, Jakubus JL, Mikhail JN, Fath JJ, Regenbogen SE et al. Unfractionated heparin versus low-molecular-weight heparin for venous thromboembolism prophylaxis in trauma. *J Trauma Acute Care Surg* 2017;**83**:151–158.
19. Erkens PMG, Prins MH. Fixed dose subcutaneous low molecular weight heparins versus adjusted dose unfractionated heparin for venous thromboembolism. *Cochrane Database Syst Rev* 2010;**2**:CD001100.
20. Koch A, Ziegler S, Breitschwerdt H, Victor N. Low molecular weight heparin and unfractionated heparin in thrombosis prophylaxis: meta-analysis based on original patient data. *Thromb Res* 2001;**102**:295–309.
21. Beitland S, Sandven I, Kjærøvik L-K, Sandset PM, Sunde K, Eken T. Thromboprophylaxis with low molecular weight heparin versus unfractionated heparin in intensive care patients: a systematic review with meta-analysis and trial sequential analysis. *Intensive Care Med* 2015;**41**:1209–1219.
22. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;**6**:e1000097.
23. Stang A. Critical evaluation of the Newcastle–Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol* 2010;**25**:603–605.
24. Higgins JPT, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD et al. The cochrane collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011;**343**:d5928.
25. Gil MR, Gonzalez-Lugo JD, Rahman S, Barouqa M, Szymanski J, Ikemura K et al. Correlation of coagulation parameters with clinical outcomes during the Coronavirus-19 surge in New York: observational cohort. *Front Physiol* 2021;**12**:618929.
26. Kirkup C, Pawlowski C, Puranik A, Conrad I, O'horo JC, Gomaa D et al. Healthcare disparities among anticoagulation therapies for severe COVID-19 patients in the multi-site VIRUS registry. *J Med Virol* 2021;**93**:4303–4318.
27. Nadkarni GN, Lala A, Bagiella E, Chang HL, Moreno PR, Pujadas E et al. Anticoagulation, bleeding, mortality, and pathology in hospitalized patients with COVID-19. *J Am Coll Cardiol* 2020;**76**:1815–1826.
28. Piazza G, Campia U, Hurwitz S, Snyder JE, Rizzo SM, Pfeferman MB et al. Registry of arterial and venous thromboembolic complications in patients with COVID-19. *J Am Coll Cardiol* 2020;**76**:2060–2072.
29. Volteas P, Drakos P, Alkadea LN, Cleri NA, Asencio AA, Oganov A et al. Low-molecular-weight heparin compared to unfractionated heparin in critically ill COVID-19 patients. *J Vasc Surg Venous Lymphat Disord*; doi:10.1016/j.jvs.2022.04.019. Published online ahead of print 15 June 2022.
30. Lopes RD, De Barros E Silva PGM, Furtado RHM, Macedo AVS, Bronhara B, Damiani LP et al. Therapeutic versus prophylactic anticoagulation for patients admitted to hospital with COVID-19 and elevated D-dimer concentration (ACTION): an open-label, multicentre, randomised, controlled trial. *Lancet (London, England)* 2021;**397**:2253–2263.
31. Oliynyk O, Barg W, Slifirczyk A, Oliynyk Y, Dubrov S, Gurianov V et al. Comparison of the effect of unfractionated heparin and enoxaparin sodium at different doses on the course of COVID-19-associated coagulopathy. *Life* 2021;**11**:1032.
32. Connors JM, Levy JH. COVID-19 and its implications for thrombosis and anticoagulation. *Blood* 2020;**135**:2033–2040.
33. Hirsh J, Anand SS, Halperin JL, Fuster V. Guide to anticoagulant therapy: heparin. *Circulation* 2001;**103**:2994–3018.
34. Costanzo L, Palumbo FP, Ardità G, Antignani PL, Arosio E, Failla G. Coagulopathy, thromboembolic complications, and the use of heparin in COVID-19 pneumonia. *J Vasc surgery Venous Lymphat Disord* 2020;**8**:711–716.
35. Newall F. *Anti-Factor Xa (Anti-Xa) Assay BT - Haemostasis: Methods and Protocols*, Monagle P, ed. Totowa, NJ: Humana Press; 2013. pp. 265–272.
36. Tiwari NR, Khatib KI, Dixit SB, Rathore PK, Melinker S, Ganapule A et al. Anticoagulation in COVID-19: an update. *J Crit Care Med (Universitae Med si Farm din Targu-Mures)* 2020;**6**:217–223.
37. Gozzo L, Viale P, Longo L, Vitale DC, Drago F. The potential role of heparin in patients with COVID-19: beyond the anticoagulant effect. A review. *Front Pharmacol* 2020;**11**:307.
38. Shi C, Tingting W, Li J-P, Sullivan MA, Wang C, Wang H et al. Comprehensive landscape of heparin therapy for COVID-19. *Carbohydr Polym* 2020;**254**:117232.
39. Martel N, Wells PS. A meta-analysis to determine the risk of heparin induced thrombocytopenia (HIT) and isolated thrombocytopenia in prophylaxis studies comparing unfractionated heparin (UFH) and low molecular weight heparin (LMWH). *Blood* 2004;**104**:2587.
40. Eriksson BI, Söderberg K, Widlund L, Wandeli B, Tengborn L, Risberg B. A comparative study of three low-molecular weight heparins (LMWH) and unfractionated heparin (UH) in healthy volunteers. *Thromb Haemost* 1995;**73**:398–401.
41. Hansen JB, Sandset PM, Huseby KR, Huseby NE, Bendz B, Ostergaard P et al. Differential effect of unfractionated heparin and low molecular weight heparin on intravascular tissue factor pathway inhibitor: evidence for a difference in antithrombotic action. *Br J Haematol* 1998;**101**:638–646.
42. Vignoli A, Marchetti M, Balducci D, Barbui T, Falanga A. Differential effect of the low-molecular-weight heparin, dalteparin, and unfractionated heparin on microvascular endothelial cell hemostatic properties. *Haematologica* 2006;**91**:207–214.
43. Andrianto, Al-Farabi MJ, Nugraha RA, Marsudi BA, Azmi Y. Biomarkers of endothelial dysfunction and outcomes in coronavirus disease 2019 (COVID-19) patients: a systematic review and meta-analysis. *Microvasc Res* 2021;**138**:104224.
44. Moores LK, Tritschler T, Brosnahan S, Carrier M, Collen JF, Doerschug K et al. Prevention, diagnosis, and treatment of VTE in patients with coronavirus disease 2019: CHEST Guideline and Expert Panel Report. *Chest* 2020;**158**:1143–1163.
45. Spyropoulos AC, Levy JH, Ageno W, Connors JM, Hunt BJ, Iba T et al. Scientific and standardization committee communication: clinical guidance on the diagnosis, prevention, and treatment of venous thromboembolism in hospitalized patients with COVID-19. *J Thromb Haemost* 2020;**18**:1859–1865.
46. REMAP-CAP Investigators; ACTIV-4a Investigators; ATTACC Investigators; Goligher EC, Bradbury CA, McVerry BJ et al. Therapeutic anticoagulation with heparin in noncritically ill patients with Covid-19. *N Engl J Med* 2021;**385**:790–802.
47. Sholzberg M, Tang GH, Rahhal H, AlHamzah M, Kreuziger LB, Ni Ainle F et al. Effectiveness of therapeutic heparin versus prophylactic heparin on death, mechanical ventilation, or intensive care unit admission in moderately ill patients with covid-19 admitted to hospital: RAPID randomised clinical trial. *BMJ* 2021;**375**:n2400.

Low molecular weight heparin is associated with better outcomes than unfractionated heparin

ORIGINALITY REPORT

18%

SIMILARITY INDEX

12%

INTERNET SOURCES

15%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1

repository.tno.nl

Internet Source

<1 %

2

tingting bai, Ling Ling, Tamaki Sono, Fumie Hisanaga, Jinlian Zhang. "Integrating meta-analysis and visual analysis to evaluate the validity of different therapies for patients with schizophrenia", Research Square Platform LLC, 2022

Publication

<1 %

3

www.medicaid.nv.gov

Internet Source

<1 %

4

Akl, Elie A, Srinivasa Rao Vasireddi, Sameer Gunukula, Maddalena Barba, Francesca Sperati, Irene Terrenato, Paola Muti, Holger Schünemann, and Elie A Akl. "Anticoagulation for the initial treatment of venous thromboembolism in patients with cancer", Cochrane Database of Systematic Reviews Protocols, 2011.

Publication

<1 %

5

Purnami Nyilo, Manyakori Serafika Permoni Putri. "The Association of Reactive Oxygen Species Levels on Noise Induced Hearing Loss of High Risk Workers in Dr. Soetomo General Hospital Surabaya, Indonesia", Indian Journal of Otolaryngology and Head & Neck Surgery, 2018

Publication

<1 %

6

bmcophthalmol.biomedcentral.com

Internet Source

<1 %

7

canada.explore.openaire.eu

Internet Source

<1 %

8

C C Lerro. "A systematic review and meta-analysis of the relationship between body size and testicular cancer", British Journal of Cancer, 10/26/2010

Publication

<1 %

9

Satoshi Ikeda, Yuki Ueno, Koji Maemura, Sen Yachi et al. "Association Between the Development of Thrombosis and Worsening of Disease Severity in Patients With Moderate COVID-19 on Admission — From the CLOT-COVID Study —", Circulation Journal, 2023

Publication

<1 %

10

bmjopenquality.bmj.com

Internet Source

<1 %

obrieniph.ucalgary.ca

11	Internet Source	<1 %
12	openaccess.biruni.edu.tr Internet Source	<1 %
13	pdfs.semanticscholar.org Internet Source	<1 %
14	rbmb.net Internet Source	<1 %
15	www.ahcmedia.com Internet Source	<1 %
16	www.casereportsinternational.com Internet Source	<1 %
17	www.nature.com Internet Source	<1 %
18	www.termidia.pl Internet Source	<1 %
19	www.thelancet.com Internet Source	<1 %
20	Fahmeeda Murtaza, Rachel Goud, Setti Belhouari, Kenneth T. Eng et al. "Prognostic Features of Preoperative OCT in Retinal Detachments: A Systematic Review and Meta-analysis", Ophthalmology Retina, 2022 Publication	<1 %

bmcinfectdis.biomedcentral.com

21

Internet Source

<1 %

22

[bmcpregnancychildbirth.biomedcentral.com](https://www.bmcpregnancychildbirth.biomedcentral.com)

Internet Source

<1 %

23

[f1000research.com](https://www.f1000research.com)

Internet Source

<1 %

24

[globalhealthmedicine.com](https://www.globalhealthmedicine.com)

Internet Source

<1 %

25

[themodernsolution.com](https://www.themodernsolution.com)

Internet Source

<1 %

26

www.nice.org.uk

Internet Source

<1 %

27

Corrado Lodigiani, Giacomo Iapichino, Luca Carenzo, Maurizio Cecconi et al. "Venous and arterial thromboembolic complications in COVID-19 patients admitted to an academic hospital in Milan, Italy", Thrombosis Research, 2020

Publication

<1 %

28

Mieke Deschodt, Johan Flamaing, Patrick Haentjens, Steven Boonen, Koen Milisen. "Impact of geriatric consultation teams on clinical outcome in acute hospitals: a systematic review and meta-analysis", BMC Medicine, 2013

Publication

<1 %

- | | | |
|----|--|------|
| 29 | worldwidescience.org
Internet Source | <1 % |
| 30 | www.lincei.it
Internet Source | <1 % |
| 31 | www.repository.cam.ac.uk
Internet Source | <1 % |
| 32 | D.F. O'Shaughnessy. "Current perspectives on the treatment of venous thromboembolism: need for effective, safe and convenient new antithrombotic drugs", International Journal of Clinical Practice, 2004
Publication | <1 % |
| 33 | Martin Husovec. "The DSA as a creator's charter?", Journal of Intellectual Property Law & Practice, 2023
Publication | <1 % |
| 34 | Mary McGrae McDermott, Joe Feinglass, Rael Slavensky, William H. Pearce. "The ankle—brachial index as a predictor of survival in patients with peripheral vascular disease", Journal of General Internal Medicine, 1994
Publication | <1 % |
| 35 | Mishra, Shiraz I, Roberta W Scherer, Paula M Geigle, Debra R Berlanstein, Ozlem Topaloglu, Carolyn C Gotay, Claire Snyder, and Shiraz I Mishra. "Exercise interventions on health-related quality of life for cancer survivors", | <1 % |

Cochrane Database of Systematic Reviews Reviews, 2012.

Publication

36

Nasibeh Zerangian, Gisou Erabi, Mohadeseh Poudineh, Kosar Monajjem et al. "Venous thromboembolism in viral diseases: A comprehensive literature review", Health Science Reports, 2023

Publication

<1 %

37

Orhan Kursat Poyrazoglu, Ayhan Dogukan, Mehmet Yalnız, Dilara Seckin, Ali Ihsan Gunal. "Acute Effect of Standard Heparin Versus Low Molecular Weight Heparin on Oxidative Stress and Inflammation in Hemodialysis Patients", Renal Failure, 2009

Publication

<1 %

38

Wanting Ma, Zhaolei Zou, Lisa Yang, Dongjia Lin, Junyi Guo, Zhongyan Shan, Qiannan Hu, Zhi Wang, Bin Li, Juan Fang. "Exploring the bi-directional relationship between periodontitis and dyslipidemia A comprehensive systematic review and meta-analysis", Research Square Platform LLC, 2022

Publication

<1 %

39

Zeyang Zhang, Ziheng Zhang, Cao Zou, Zhongcai Pei, Zheyuan Yang, Jing Wu, Shikun Sun, Fei Gu. "ECGNet: An Efficient Network for Detecting Premature Ventricular Complexes

<1 %

Based on ECG Images", IEEE Transactions on Biomedical Engineering, 2023

Publication

40	ashpublications.org Internet Source	<1 %
41	epdf.pub Internet Source	<1 %
42	journal.publications.chestnet.org Internet Source	<1 %
43	pureadmin.qub.ac.uk Internet Source	<1 %
44	repositoriosalud.es Internet Source	<1 %
45	research-repository.griffith.edu.au Internet Source	<1 %
46	thejh.org Internet Source	<1 %
47	www.chdr.nl Internet Source	<1 %
48	www.ncbi.nlm.nih.gov Internet Source	<1 %
49	www.unirepository.svkri.uniri.hr Internet Source	<1 %
50	www.wjgnet.com Internet Source	<1 %

51

www2.mdpi.com

Internet Source

<1 %

52

Bharat Narasimhan, Marta Lorente-Ros, Jose S. Aguilar-Gallardo, Christopher Perez Lizardo et al. "Anticoagulation in COVID-19: a review of current literature and guidelines", Hospital Practice, 2021

Publication

<1 %

53

Eman Mohisn Mansory, Mohammed Abu Farhaneh, Alla Iansavitchene, Alejandro Lazo-Langner. "Venous and arterial thrombosis in ambulatory and discharged COVID-19 Patients: a systematic review and meta-analysis", TH Open, 2022

Publication

<1 %

54

Kyoung Ae Kong, Sodam Jung, Mina Yu, Junbeom Park, In Sook Kang. "Association Between Cardiovascular Risk Factors and the Severity of Coronavirus Disease 2019: Nationwide Epidemiological Study in Korea", Frontiers in Cardiovascular Medicine, 2021

Publication

<1 %

55

Lisa K. Moores, Tobias Tritschler, Shari Brosnahan, Marc Carrier et al. "Thromboprophylaxis in Patients with COVID-19. A Brief Update to the CHEST Guideline and Expert Panel Report", Chest, 2022

Publication

<1 %

56

Marco Marietta, Valeria Coluccio, Mario Luppi.
"COVID-19, coagulopathy and venous
thromboembolism: more questions than
answers", Internal and Emergency Medicine,
2020

Publication

<1 %

57

Ping Sun, Yanlong Li, Lijun Chang, Xudong
Tian. "Prognostic and clinicopathological
significance of Gamma-Glutamyltransferase in
patients with hepatocellular carcinoma",
Medicine, 2019

Publication

<1 %

58

Radek Cihlar, Vladimir Sramek, Adriana
Papiez, Miroslav Penka, Pavel Suk.
"Pharmacokinetic Comparison of
Subcutaneous and Intravenous Nadroparin
Administration for Thromboprophylaxis in
Critically Ill Patients on Vasopressors",
Pharmacology, 2020

Publication

<1 %

59

afef.asso.fr

Internet Source

<1 %

60

doczz.net

Internet Source

<1 %

61

jamanetwork.com

Internet Source

<1 %

www.covid19treatmentguidelines.nih.gov

62

Internet Source

<1 %

63

www.cureus.com

Internet Source

<1 %

64

www.degruyter.com

Internet Source

<1 %

65

www.dovepress.com

Internet Source

<1 %

66

www.frontiersin.org

Internet Source

<1 %

67

www.mayoclinic.org

Internet Source

<1 %

68

www.mdpi.com

Internet Source

<1 %

69

www.minervamedica.it

Internet Source

<1 %

70

Alex C. Spyropoulos, Jerrold H. Levy, Walter Ageno, Jean Marie Connors et al. "Scientific and Standardization Committee Communication: Clinical Guidance on the Diagnosis, Prevention and Treatment of Venous Thromboembolism in Hospitalized Patients with COVID - 19", Journal of Thrombosis and Haemostasis, 2020

Publication

<1 %

71

Ayush Batra, Jeffrey R. Clark, Anthony K. Kang, Sareen Ali et al. "Persistent viral RNA shedding of SARS-CoV-2 is associated with delirium incidence and six-month mortality in hospitalized COVID-19 patients", GeroScience, 2022

Publication

<1 %

72

Konstantinos Kyriakoulis, Evangelos Dimakakos, Ioannis Kyriakoulis, Mariella Catalano et al. "Practical Recommendations for Optimal Thromboprophylaxis in Patients with COVID-19: A Consensus Statement Based on Available Clinical Trials", Journal of Clinical Medicine, 2022

Publication

<1 %

73

Kun Yang, Matthew Holt, Min Fan, Victor Lam, Yong Yang, Tuanzhu Ha, David L. Williams, Chuanfu Li, Xiaohui Wang. "Cardiovascular Dysfunction in COVID-19: Association Between Endothelial Cell Injury and Lactate", Frontiers in Immunology, 2022

Publication

<1 %

74

Kuo-Chuan Hung, Yen-Ta Huang, Ying-Jen Chang, Chia-Hung Yu et al. "Association between Fibrinogen-to-Albumin Ratio and Prognosis of Hospitalized Patients with COVID-19: A Systematic Review and Meta-Analysis", Diagnostics, 2022

<1 %

75

Litao Zhang, Xinsheng Yan, Qingkun Fan, Haiyan Liu, Xintian Liu, Zejin Liu, Zhenlu Zhang. "D - dimer levels on admission to predict in - hospital mortality in patients with Covid - 19", Journal of Thrombosis and Haemostasis, 2020

Publication

<1 %

76

Romil Singh, Sawai Singh Rathore, Hira Khan, Abhishek Bhurwal et al. "Mortality and Severity in COVID-19 Patients on ACEIs and ARBs—A Systematic Review, Meta-Analysis, and Meta-Regression Analysis", Frontiers in Medicine, 2022

Publication

<1 %

77

www.pure.ed.ac.uk
Internet Source

<1 %

78

Davide Salmi, Ingrid Andreea Ilie, Nadia Pagnoncelli, Greta Gandolfi et al. "Chapter 20 Pre-hospital Management of COVID-19: Looking for a Future Perspective", Springer Science and Business Media LLC, 2022

Publication

<1 %

79

Emmanuel J. Favaloro, Giuseppe Lippi. "Maintaining Hemostasis and Preventing Thrombosis in Coronavirus Disease 2019 (COVID-19): Part II", Seminars in Thrombosis and Hemostasis, 2021

<1 %

80

Heng Zhang, Qifang Lao, Jue Zhang, Jieqing Zhu. "Coagulopathy in COVID-19 and anticoagulation clinical trials", Best Practice & Research Clinical Haematology, 2022

Publication

<1 %

81

Hong-Lin Chen, Xiao-Dong Wang. "Heparin for venous thromboembolism prophylaxis in patients with acute spinal cord injury: a systematic review and meta-analysis", Spinal Cord, 2013

Publication

<1 %

82

Jeffrey A. Lafranca, Sander M. Hagen, Leonienke F.C. Dols, Lidia R. Arends, Willem Weimar, Jan N.M. IJzermans, Frank J.M.F Dor. "Systematic review and meta-analysis of the relation between body mass index and short-term donor outcome of laparoscopic donor nephrectomy", Kidney International, 2013

Publication

<1 %

83

Liang Yu, Xiaojuan Shen, He Liu. "The efficacy of pericapsular nerve group block for postoperative analgesia in patients undergoing hip surgery: A systematic review and meta-analysis of randomized controlled trials", Frontiers in Medicine, 2023

Publication

<1 %

84

Lilas Dagher, Hanyuan Shi, Yan Zhao, Andrew Wetherbie et al. "New - Onset Atrial Arrhythmias Associated With Mortality in Black and White Patients Hospitalized with COVID - 19", Pacing and Clinical Electrophysiology, 2021

Publication

<1 %

85

Vanesa Bellou, Ioanna Tzoulaki, Maarten van Smeden, Karel G.M. Moons, Evangelos Evangelou, Lazaros Belbasis. "Prognostic factors for adverse outcomes in patients with COVID-19: a field-wide systematic review and meta-analysis", European Respiratory Journal, 2022

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Low molecular weight heparin is associated with better outcomes than unfractionated heparin

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10
