

February 2021

Pakistan Journal of Medical and Health Sciences

Vol. 15, Issue 2, February 2021

Website: www.pjmhsonline.com

Domestic Violence and Pregnancy Outcome Naila Javaid, Sadia Kanwal, Jaweria Faisal, Nazneen Akhtar, Zobia Jawad, Nasreen Akhtar	391
Early Outcome of Patients with Poor Hemodynamics Undergoing Open Chest Management Muhammad Sher-i-Murtaza, Muhammad Hamid Chaudhary, Mirza Ahmad Raza Baig, M. M. Khan	395
Frequency of Different Gynecological Cancers in Suspected Cases of Gynecological Cancers Hafiza Urooj Shoukat, Ujala Aslam, Fiza Jamil, Muhammad Usman Aslam, Faisal Javed, Noman Sadiq	399
Does Gabapentin make any difference in Post-Operative Pain in Modified Radical Mastectomy patients? Anum Arif, Minahil Iram, Syed Hashim Ali Inam,, Sabih Nofil, Saman Tanveer, Farhat Jaleel	402
Effectiveness of Dexmedetomidine and Fentanyl with Intrathecal Levobupivacaine in C-Section Tariq Iqbal, Noman Tariq, Asaad Rizwan Rana, Tayyaba Rasheed, Syeda Gulrukh Saba Shah	405
Examine the Prevalence of Albuminuria in Diabetic Patients Presented With Stroke Abdul Ghafoor Magsi, Muhammad Imran Khan, Mushtaq Ahmed Shahid, Abrar Shaikh	408
Frequency of Atrial Fibrillation within 48 Hours of Acute Myocardial Infarction Abdul Latif, Atif-Ur-Rahman, Farhat Shireen, Sher Bahadar Khan	411
Ultrasound Features of Ovarian Cysts in married females Zahid Nazir, Samina Raza, Shamaila Hussain, Muhammad Azam, Syed Waqas Ahmed	415
Course of Covid 19 and Fetomaternal Outcome at a tertiary care hospital Robina Furrukh, Sumera Kanwal, Qurat Ul Ain, Farah Kalsoom, Farrukh Kamal	418
Frequency of Skin Disorders Among Paediatric Population of Charity Hospital of Lahore Mariam Sheikh, Rabail Majeed, Rabia Anjum, Amna Shahid	421
Gross Anatomical Placental Anomalies Observed at the Time of Delivery Areeba Riasat Nahra, Riasat Ali Nehra, Arif Gulzar, Kishwar Naheed, Hafsa Riasat, Ali Zafar	424
Prevalence and Risk Factors of Violence in Schizophrenic Inpatients of Psychiatry, SIMS, Lahore Hafiz M I Afzal, A Bashir, N Mazhar, M A Awab Sarwar, M Ishtiaq, S Afzal, T S A M Ali, S Younas, K Junaid	428

How Low Socioeconomic Status During Pregnancy Causes Orofacial Clefts In Newborns	Riasat Ali Nehra, Areeba Riasat Nahra, Arif Gulzar, Noor Ul Mobeen, Hafsa Riasat, Ali Zafar	432
HRCT chest finding and clinical course of asymptomatic cases with COVID-19 pneumonia	Saman Ch K R Yousaf, S Saeed, H Mehmood, A A Chatha, V Ashraf, M. Z Asgher, H Riaz, S Amjad, A Aslam	434
Impact of Sociodemographic Factors on the Access to Oral Healthcare Facilities among Adults	Daud Anthoney, F Ali Syed, Mehwish Khan, Asadullah Rathore, Erum Zahid, A Ayub Shah, I Hameed Khaliq	438
Ocular, Medical and Psychological Effects in Medical Students who use Smart Phones	Arif Hussain, Hijab Farrukh, Arif Gulzar, Kishwar Naheed, Hafiza Fatima Mishal, Zoya Arif	443
Prevalence of Complicated Appendectomy in Patients with Acute Appendicitis: Comparison the Outcomes between Laparoscopic and Open Approach	Aqeel Ahmad, Ali Gohar Bozdar, Dileep Kumar, Khumair Asif	446
Hypoalbuminemia in Patients of Systolic Heart Failure	Salma Kadir, Bashir Ullah, Mushtaq Ahmed Shahid	449
Gastric Duplication Cyst Mimicking a Liver Cyst Case Report and Review of the Literature	Hussam S. Hassan, Hany A. Elhady, Mohamed Shawky Elfaragy	451
The Effects of Hydrogen Peroxide Solution on Various Properties of CAD/CAM based Polymethylmethacrylate (PMMA)	Mustafa Asaad Mohammed, Wasmaa Sadik Mahmood	455
Frequency of Complications of Acute Appendicitis	Abdul Hafeez Arain, Shahnawaz Khatti, Riaz Ahmed Memon, Abdul Salam Memon, Shaheen Gul, Shahida Khatoun	460
Cultural Elements in the Rituals of Traditional Sports among Turkic Peoples	Mehmet Türkmen	464
Effect of Learning Styles on Self-Esteem and Stress Levels of University Students	Sadaf Aijaz Abdul Rahman, Hafeezullah Wazir Ali, Ghazala Rasool, Irum Siddique	469
Entamoeba histolytica, Identification In Asymptomatic Infection	Fadia Abd Al-Muhsin Al-Khayat, Hanadi J. Al-Zubaidi, Zainab A.Aldhaher	473
Comparing undergraduate Nursing student academic engagement and achievement in traditional versus Blended Learning Models	Enas Hassan Saad, Magda Abd Alhamid Abd Al Fattah, Ahmed Mahmoud Fakhry, And Mohammed Ali Pessa	478
Association of Self-Esteem with Gender and Medical Year: A Cross Sectional Study	Sadaf Aijaz Abdul Rahman, Ghazala Rasool, Hafeezullah Wazir Ali, Irum Siddique	483
Protective effects of Lactobacillus casei on Lead-induced biochemical deteriorations in Rats	Sundus Hamed Ahmed, Amin Suleiman Badawi, Abdullah Khamis Abdullah	486
Antidiabetic effect of Aqueous Extract of Medicago Sativa with Enhanced Histopathology of Pancreas in Alloxan Induced Diabetic Rats	Salim J. Khalaf, Gadeer Hatem Aljader, Entedhar R. Sarhat, Thuraia Rifaat Sarhat, Kasim Sakran Abass	492



Impact of Serum Adropin and Irisin in Iraqi patients with Congestive Heart Failure Mahde Salih Hamad, Entedhar Rifaat Sarhat, Thuraia Rifaat Sarhat, Kasim Sakran Abass	497
Endothelial Function and Heart Rate Variability Parameters before and after Body Correction Program Kostenchak Svystak OE, Klymuk AR, Nemesh MI, Palamarchuk OS, Feketa VP	500
Lesson from Indonesia: Covid-19 Testing Strategy In Obstetric Emergency Cases At Low-Resource Health Care Setting Manggala Pasca Wardhana, Aditiawarman, Nareswari Cininta Maniora, Rozi Aditya, Khanisyah Erza Gumilar, Budi Wicaksono, Muhammad Ilham Aldika Akbar, Ernawati, Agus Sulistyono, Hermanto Tri Juwono, Erry Gumilar Dachlan	508
Stigma, Discrimination, Treatment Effectiveness And Policy: Public Views About Drug Addiction In Malaysia Dharshini Navanethan, Erna Faryza Mohd Poot, Nathan Vytialingam, Mohammad Nazmul Hasan Maziz	514
General Pedagogy of Traditional Wrestling: The example of Turkish people Siyamik Arstanbekov*, Ali Bayazit	520
National Wrestling of Tuva Turks: Huresh (Hüres) Inga Mendot*, Mehmet Turkmen	524
Archaic Tracks of Kyrgyz Traditional Sports Games Mehmet Türkmen	530
Accuracy of RT-PCR Test in diagnosing COVID-19: A retrospective Study Razieh Zahedi, Haniye Khorshidsavar, Vahid Rahmanian, Nader Sharifi	534
Effects of Hip and Hip Core Muscles Strengthening Versus Knee Muscle Strengthening for the Management of Anterior Knee Pain Luqman Ali, Sana Hafeez, Zarwa Hafeez, Saima Riaz, Rehan Ramzan Khan, Hayatullah Khalid	539
Basophils are seeking our attention in Multan region Zertaj Kashif, Sehar Shamshad Ali, Faiza Shafqat, Babar Mumtaz, Noman Khalid, Sonia Zafar Warraich	542
Effects of Motor Control Training of Hip Muscles on Pain and Physical Function After Total Hip Arthroplasty Bilal Fayyaz, Saima Riaz, Rehan Ramzan Khan, Muhammad Asif Javed, Muhammad Sulman, Hayatullah Khalid	546
Assessment and Management of Unilateral Spatial Neglect by Using Mobile Application "Visual Attention Lite" in Acute Stroke Patients Nadia Siddique, Zeest Hashmi, Rehan Ramzan Khan, Saima Riaz, Sana Hafeez, Bilal Fayyaz	549
Parity with Risk and Increased Incidence of Postpartum Hemorrhage Nuli Nuryanti Zulala, Sunarti	552
Characteristics, Source of Information and Knowledge of Housewives about Transmission of HIV and Aids Herlin Fitriani Kurniawati, Herlin Fitriana Kurniawati	555
Nutritional Status of Children under Five in Public Health Care Girmulyo II Evi Wahyuntari, Feby Ramadhani Dewi	559
The Role of Peers on Adolescent Sexual Behavior in Indonesia Fitriyani Bahriyah, Yekti Satriyandari	563



The Correlation between Prenatal Gentle Yoga and the Success of Normal Childbirth in Bidan Kita Clinic Klaten In 2017 Dwi Rahmawati, Anjarwati	567
The Association between Maternal Age and History of Cesarean Section with Antepartum Hemorrhage in Indonesia Anjeli Ratih Syamlingga Putri, Herlin Fitriana	571
The Correlation between Parity and Husband's Support with the Choice of Intra-Uterine Device Contraception at Work Area of Sleman Health Center Mita Meilani, Dhesi Ari Astuti	575
Medical History of Pregnant Women with Preeclampsia Fayakun Nur Rohmah, Yesi Ardila	579
Antenatal Depressive Symptoms in Asian Women: A Review of the Articles Perwitasari, Umu Hani Edi N	582
The Image of Age and Occupation Toward Exclusive Breastfeeding Success in the Working Area of Turi Primary Health Center, Sleman District, Yogyakarta Fitriana Yuni Permana Sari, Dhesi Ari Astuti	587

[|<<1234>>|](#)

Showing 51 to 100 of 173 (4 Pages)





[Home](#) / [Editorial Team](#)

Editorial Team

EDITOR IN CHIEF

Prof. Abdul Majeed Chaudhry

Asadullah Malik

EDITORS

Prof. Yaseen Rafi

Khalid Irshad (UK)

ASSOCIATE EDITORS

Prof. Syed Asghar Naq

Prof. M. Nadeem Aslam

Prof. Abrar Ashraf Ali

ASSISTANT EDITORS

Prof. Wasim Amir

Prof. Hammad Naeem Rana

BIOSTATICIAN

Minahil Irum

ADVISORY BOARD

NATIONAL

Farid Ahmad Khan

Muhammad Javaid Athar

Mudassira Saqib

Shahzad Shams Goraya

Syed Irfan Hussain

Masood Rashid

Nighat Nadeem

INTERNATIONAL

Afaq Zaman Khan (USA)

Naeem Akhtar (UK)

Munir Ahmad Rathore (UK)

Muhammad Ahmad (UK)

Donald B Reid (UK)

Amer Farooq Majeed (UK)

PJMHS is a peer-reviewed,
open access, monthly journal

ISSN

Online: 2957-899X

Print: 1996-7195

INDEXED WITH

[Elsevier in EMBASE Scopus](#)

[EMR Index Medicus \(IMEMR\) of WHO](#)

[International Scientific Indexing \(ISI\)](#)

[SCImago Journal & Country Rank](#)

[Google Scholar](#)

[Digital Object Identifier \(DOI\)](#)

[Crossref](#)

RECOGNIZED BY

1. Higher Education Commission
2. Pakistan Medical Commission

ABBREVIATION

PJMHS

Platform &
workflow by
OJS / PKP

Lesson from Indonesia: Covid-19 Testing Strategy In Obstetric Emergency Cases At Low-Resource Health Care Setting

MANGGALA PASCA WARDHANA¹, ADITIAWARMAN¹, NARESWARI CININTA MANIORA¹, ROZI ADITYA¹, KHANISYAH ERZA GUMILAR¹, BUDI WICAKSONO¹, MUHAMMAD ILHAM ALDIKA AKBAR¹, ERNAWATI¹, AGUS SULISTYONO¹, HERMANTO TRI JUWONO¹, ERRY GUMILAR DACHLAN¹

¹Department of Obstetrics and Gynaecology, Faculty of Medicine, Universitas Airlangga/ Dr. Soetomo General Hospital, Surabaya Indonesia

²Faculty of Medicine, Universitas Airlangga, Surabaya Indonesia

Correspondence to Aditiawarman, Email: aditiawarman@fk.unair.ac.id

ABSTRACT

Background: COVID-19 identification in obstetric emergency cases with limited resources is problematic, particularly in asymptomatic cases.

Aim: To examine the screening strategies of COVID-19 obstetric emergency cases in low-resource health care settings.

Method: A retrospective cohort design was carried out on patients with COVID-19 positive screening results. It was assessed based on symptoms, contact history, lymphocytopenia, chest X-rays, and rapid antibody tests compared to RT-PCR results SARS-COV-2.

Result: Out of the 190 cases that came to the delivery room, the staff suspected 69 COVID-19 cases (36.3%) through the first screening protocol. Positive SARS-COV-2 RT-PCR was found in 23 cases with a majority asymptomatic (52.2%). The percentages of sensitivity and specificity from the parameters as follow: 48% and 74% in COVID-19 symptoms (febris or respiratory symptoms); 9% and 100% in contact history; 22% and 83% in lymphocytopenia; 52% and 48% in chest x-ray; 78% and 30% in rapid antibody test. Rapid antibody tests have the highest sensitivity to increase the identification of 12 asymptomatic cases.

Conclusion: Other screening beyond symptoms and contact history such as lymphocytopenia, chest x-ray, and rapid antibody test can improve the identification, especially for asymptomatic cases in areas with the limited testing ability and high Covid-19 transmission

Keywords: Covid-19, Screening, Obstetric Emergency, Low-resource health care

INTRODUCTION

The condition of the COVID-19 pandemic in Indonesia remains to increase exponentially. Surabaya, the second-largest city in Indonesia, is one of the COVID-19 centers for local transmission, and it was handling almost 10% of total COVID-19 cases in Indonesia¹. Cases finding, detecting, and tracing are the main approaches to mitigate COVID-19 transmission in communities and hospitals. Identifying COVID-19 in obstetric emergencies has an essential role since it will affect managing appropriate patient care, allocating isolation rooms, implementing neonatal care management, and preventing transmission to other patients and health workers.

A study from Vintzileos (2020) showed that many COVID-19 patients were asymptomatic in the delivery room². Another study by Sutton (2020) also found almost 88% of pregnant women with COVID-19 who admitted to the hospital were asymptomatic³. According to these reports, universal COVID-19 testing in pregnant women is recommended to detect an asymptomatic patient population. However, this condition will cause other obstacles, mainly in areas with limited testing capacity.

Generally, Indonesia's testing capacity is reported to be 2,378 tests/1 million population. While the testing capacity in developed countries, such as the US, is up to 37 times with a nearly comparable population⁴. Therefore, it is crucial to modify the COVID-19 screening strategy to identify symptomatic cases and be used to identify patients who come to the hospital for obstetric emergencies. Based on the above evidence, it is vital to find a COVID-19 testing

strategy for obstetric emergency cases, primarily in large local transmissions areas with inadequate testing capacity.

METHOD

The study applied a retrospective cohort design using COVID-19 screening protocols at the Department of Obstetrics and Gynecology, Soetomo General Hospital in Surabaya. Since 20th April 2020, we have developed the testing criteria. From testing only based on COVID-19 symptomatic and contact history, we examined several other parameters. The COVID-19 screening protocol includes evaluating the signs and symptoms, COVID-19 history, fever ($\geq 37.8^{\circ}\text{C}$), and respiratory problems (cough and shortness of breath) within 14 days. The presence of any or all of the symptoms was inserted as symptomatic cases. Evaluation of patient contact with the suspect or confirmed COVID-19 patients, COVID-19 laboratory results, and lymphocytopenia (lymphocyte count $< 1 \times 10^9/\text{L}$). Based on Huang's study⁵, we also took a radiological examination through the chest X-Ray and assessed the results as COVID-19 suspected or not based on Jacobi's pictorial review⁶. Subsequently, we arranged a rapid test of COVID-19 antibodies (Wondfo One Step COVID-19; Severe Acute Respiratory Syndrome, Coronavirus 2 / SARS-CoV-2 Antibody Test) based on immunochromatographic assay with lateral flow method in detecting qualitative SARS-CoV-2 IgG/IgM antibody from serum samples.

If any of the first screening tests were positive, the patient would proceed with a diagnostic examination using

Real-Time Reverse Transcriptase Polymerase Chain Reaction (RT-PCR) SARS-COV-2 for confirmation. The SARS-COV-2 diagnostic test was performed by taking a swab in the nasopharyngeal and putting it in a viral transport medium. The Abbott m2000 with Abbott Real-time SARS-COV-2 assay was used for the qualitative detection of nucleic acids from SARS-COV-2. Results were reported as positive if RdRp or N-gene examined positive. All processes were carried out in the hospital laboratory.


The study was conducted from 20th April 2020 to 10th June 2020, and the COVID-19 risk screening was carried out in all patients coming to the delivery room. If a positive screening result is received, the patient will be included in the study, and a swab test will be performed as confirmation and analyzed. This research has been approved by the ethics committee of Soetomo General Hospital Surabaya.

Categorical variables were shown as numbers (percentage), and continuous variables were represented as means (Standard Deviation / SD) or medians (interquartile ranges /IQRs). SPSS version 24.0 software for Windows (IBM Corp., Armonk. NY, USA) was used for statistical analysis. Chi-Square and Fisher Exact Test as an alternative was applied to compare the difference in the categorical variables, Independent T-Test and Mann-Whitney Test as an alternative was utilized to compare the continuous variables between groups. The value of $p < 0.05$ was considered significant. The accuracy of screening is performed by calculating the sensitivity, specificity, Positive Predictive Value (PPV), and Negative Predictive Value (NPV).

RESULT

During the study, a total of 190 patients came to the delivery room. Sixty-nine patients (36.3%) were involved in the study, with suspected COVID-19 based on the COVID-19 screening protocol, and RT-PCR diagnostic tests were performed. Twenty-three patients (33%) with positive initial screening (12.1% of all 190 patients came to the delivery room) had positive RT-PCR results. This research framework can be observed in Figure 1.

Figure 1. Research Framework



Sixty-nine patient characteristics with a positive screening test were divided into positive and negative RT-PCR tests (table 1). The majority of COVID-19 positive patients were found to have asymptomatic cases (52.2%). Significant contact history of COVID-19 was only detected in 2 cases. Lymphocytopenia was observed in 21.7% of cases and abnormal chest X-rays in 52.2%. The majority of RT-PCR COVID-19 (+) cases had positive rapid antibody tests (78.3%). There were no significant differences in the characteristics and examination parameters between positive and negative RT-PCR results.

Based on the result, the calculation of several parameters' screening ability is shown in the following table (Table 2). The highest sensitivity was found in the antibody test, while the highest specificity and PPV were obtained from COVID-19 contact history.

The specific findings of the complete examination in the positive RT-PCR case are presented in figure 2. It will comprehend what tests were positive from the screening protocol in each case.

Table 1. Distribution of Patient Characteristics

Variable	COVID-19 PCR (+)	COVID-19 PCR (-)	p
	cases (%) N=23	cases (%) N=46	
Maternal Age (mean±SD)	29±5.34	30.85±5.99	0.216
< 20 y.o	0 (0%)	1 (2.2%)	0.634
20-35 y.o	18 (78.3%)	32 (69.6%)	
> 35 y.o	5 (21.7%)	13 (28.3%)	
Gestational Age			0.447
< 20 weeks	2 (8.7%)	2 (4.3%)	
20 - 37 weeks	5 (21.7%)	12 (26.1%)	
> 37 weeks postpartum	15 (62.5%)	32 (69.6%)	
Pregnancy Planning			0.282
Conservative Delivery / Pregnancy Termination	9 (39.1%) 14 (60.9%)	12 (26.1%) 34 (73.9%)	
Parity			0.267
Primiparity Multiparity	9 (39.1%) 14 (60.9%)	12 (26.1%) 34 (73.9%)	
Positive Symptoms	11 (47.8%)	12 (26.1%)	0.071
Febris	7 (30.4%)	6 (13%)	0.107
Cough	9 (39.1%)	8 (17.4%)	0.058
Dyspnea	5 (21.7%)	6 (13%)	0.352
Contact History	2 (8.7%)	0	0.108
Obstetric Complication			
Obesity	4 (17.4%)	14 (30.4%)	0.245
Diabetes	2 (8.7%)	1 (2.2%)	0.256
Chronic Hypertension	2 (8.7%)	2 (4.3%)	0.596
Preeclampsia	3 (13%)	11 (23.9%)	0.29
White Blood Count x10⁹/L (median[IQR])	10.19 (5.54)	11.49 (4.69)	0.093
Lymphocytopenia	5 (21.7%)	8 (17.4%)	0.663
Abnormal Chest X-Ray	12 (52.2%)	24 (52.2%)	1
Positive Rapid Antibody Test	18 (78.3%)	32 (69.6%)	0.446

Table 2. COVID-19 Screening Test Performance

Variable	COVID-19 PCR (+)	COVID-19 PCR (-)	Sensitivity (%)	Specificity (%)	PPV (%)	NPV (%)
	cases (%) N=23	cases (%) N=46	(95% CI)	(95% CI)	(95% CI)	(95% CI)
Positive Symptom	11 (47.8%)	12 (26.1%)	47.83% (26.82%-69.41%)	73.91% (58.97%-85.73%)	47.83% (32.43%-63.65%)	73.91% (64.89%-81.29%)
Febris	7 (30.4%)	6 (13%)	30.43% (13.21%-52.92%)	86.96% (73.74%-95.06%)	53.85% (30.69%-75.45%)	71.43% (65.11%-77.01%)
Cough	9 (39.1%)	8 (17.4%)	39.13% (19.71%-61.46%)	82.61% (68.58%-92.18%)	52.94% (33.35%-71.67%)	73.08% (65.59%-79.45%)
Dyspnea	5 (21.7%)	6 (13%)	21.74% (7.46%-43.70%)	86.96% (73.74%-95.06%)	45.45% (22.12%-70.97%)	68.97% (63.55%-73.91%)
Contact History	2 (8.7%)	0 (0%)	8.7% (1.07%-28.04%)	100% -	100% -	68.66% (65.88%-71.30%)
Lymphocytopenia	5 (21.7%)	8 (17.4%)	21.74% (7.46%-43.70%)	82.61% (68.58%-92.18%)	38.46% (18.71%-62.92%)	67.86% (62.11%-73.11%)
Chest X-Ray	12 (52.2%)	24 (52.2%)	52.17% (30.59%-73.18%)	47.83% (32.89%-63.05%)	33.33% (23.64%-44.67%)	66.67% (54.25%-77.13%)
Rapid Antibody Test	18 (78.3%)	32 (69.6%)	78.26% (56.30%-92.54%)	30.43% (17.74%-45.75%)	36% (29.66%-42.86%)	73.68% (53.48-87.21%)

Figure 2. Screening Results in SARS-COV-2 Positive by RT-PCR

Symptomatic Cases												
Parameters	Case 1	Case 2	Case 3	Case 4	Case 5	Case 6	Case 7	Case 8	Case 9	Case 10	Case 11	
Symptoms	+	+	+	+	+	+	+	+	+	+	+	
Contact (+)	+	-	-	-	-	-	-	-	-	-	-	
Lymphocytopenia	-	+	-	+	-	-	-	+	-	+	-	
Chest X-Ray	-	+	+	-	+	-	+	+	-	+	+	
Rapid Antibody Test	-	-	+	+	+	+	+	+	-	+	-	
Asymptomatic Cases												
Parameters	Case 12	Case 13	Case 14	Case 15	Case 16	Case 17	Case 18	Case 19	Case 20	Case 21	Case 22	Case 23
Symptoms	-	-	-	-	-	-	-	-	-	-	-	-
Contact (+)	-	+	-	-	-	-	-	-	-	-	-	-
Lymphocytopenia	-	-	-	-	-	-	-	+	-	-	-	-
Chest X-Ray	-	-	+	-	+	-	+	-	+	+	-	-
Rapid Antibody Test	+	+	+	+	+	+	+	+	+	-	+	+

DISCUSSION

Currently, the world is suffering from a global crisis in fronting the COVID-19 pandemic. It sets enormous pressure on the health system around the world. Nearly all health service centers in entire countries are overwhelmed. Excellent anticipation with appropriate and adaptive strategies to reduce transmission is wholly needed and considered to increase exponentially. Even though some specific health services have been postponed, maternal health services must remain to be performed and cannot be discontinued at any time. Recent evidence from areas with a high transmission prevalence will also present many pregnant women with COVID-19³.

Out of 190 patients admitted to the delivery room, it found 23 COVID-19 positive cases (12.1%). Related studies from the American Journal of Obstetrics and Gynecology (AJOG) revealed a higher proportion of maternal COVID-19 cases (19.9%)². Sutton's investigation also reported a similar result, with a proportion of 15.3% from 33 positive cases in 215 pregnant women³. The different numbers of overall COVID-19 cases between the US and Indonesia can differentiate this study's results. Furthermore, the two comparative studies conducted universal RT-PCR. In contrast, this study conducted RT-PCR testing based on the initial screening.

The mean COVID-19 positive maternal age was 29 years old, and the majority was in the range of 20 to – 35 years old. Most patients came for delivery or pregnancy termination (60.9%). Then most subjects at term gestational age (62.5%). This study is compatible with a systematic review study, where 108 COVID-19 pregnancy women were obtained with the average age from 29 to 32 years and mostly came to the delivery room in the 3rd trimester. In this review, labor also occurred in 80% of cases⁷. The characteristics of obstetrics patients were different compared to the general population. Guan's research explained a median age of 47 years old⁸. The pregnant patient's population always has a younger age range. Moreover, maternal COVID-19 patients' identification is not due to the COVID-19 problem but correlates with specific obstetric problems in the delivery process. Therefore, the screening and identification of COVID-19 in pregnant women who came to the hospital have a vital role.

The most obstetric complication related to COVID-19 comorbidities in the studies was obesity (4 cases, 17.4%). Other studies have also confirmed that obesity is the most commonly found in diabetic mellitus and hypertension during pregnancy^{9,10}. Breslin's investigation showed that patients who need health services at the Intensive Care Unit (ICU) are at a Body Mass Index (BMI) > 35, however further studies are needed to reconfirm the correlated morbidity findings¹¹.

The majority of patients with COVID-19 positive were asymptomatic (12 cases, 52.2%). Coughing is the most often complained (39.1% from all cases and 81.8% from symptomatic cases). Insignificantly different from Zaigham's study, fever was found to be the most frequent symptom⁷, while Breslin's observation explained that cough was the most prevalent ailment¹⁰. Breslin's study is more in line with this study because the samples were taken by universal screening. Typical screening protocols will also give different characteristics. Significant COVID-19 contact history was only found in 2 cases (8.7%). In contrast, a different study showed contact history in a higher proportion of about 34.5% of cases¹⁰. Yan's study revealed 38 out of 116 cases with a positive contact history⁹. In contrast, a different study showed contact history in a higher proportion of about 34.5% of cases. Yan's study revealed 38 out of 116 cases with a positive contact history⁹. Evaluating contacts is quite challenging, notably in a developing country due to patients' negative stigma, fear of isolation care, and risk of separating newborns from their mothers. This affects the contact history covered.

Lymphocytopenia obtained only 21.7%, while other studies from Zaigham reported lymphocytopenia in 59%⁷. This study also examined a chest X-ray and revealed that half of COVID-19 positive showed abnormal results (52%). A retrospective study from Yan on 116 COVID-19 pregnant women informed abnormal radiologic findings in 96.3% of cases and a higher proportion of symptomatic cases (76.7%)⁹.

The Centers for Disease Control and Prevention (CDC) does not recommend that chest radiography to diagnose COVID-19. The findings on chest imaging are not specific and overlap with other infections. Therefore, this modality should not be accepted as a first-line test to diagnose COVID-19. However, in the low-resource testing areas, this option can be considered. The most obstetric

complication related to COVID-19 comorbidities in the studies was obesity (4 cases, 17.4%). Other studies have also confirmed that obesity is the most commonly found in diabetic mellitus and hypertension during pregnancy⁹. Breslin's investigation showed that patients who need health services at the Intensive Care Unit (ICU) are at a Body Mass Index (BMI) > 35, however further studies are needed to reconfirm the correlated morbidity findings.

The majority of patients with COVID-19 positive were asymptomatic (12 cases, 52.2%). Coughing is the most often complained (39.1% from all cases and 81.8% from symptomatic cases). Insignificantly different from Zaigham's study, fever was found to be the most frequent symptom⁷, while Breslin's observation explained that cough was the most prevalent ailment¹². Breslin's study is more in line with this study because the samples were taken by universal screening. Typical screening protocols will also give different characteristics. Significant COVID-19 contact history was only found in 2 cases (8.7%).

In contrast, a different study showed contact history in a higher proportion of about 34.5% of cases. Yan's study revealed 38 out of 116 cases with a positive contact history⁹. In contrast, a different study showed contact history in a higher proportion of about 34.5% of cases¹². Yan's study revealed 38 out of 116 cases with a positive contact history⁹. Evaluating contacts is quite challenging, notably in a developing country due to patients' negative stigma, fear of isolation care, and risk of separating newborns from their mothers. This affects the contact history covered.

Lymphocytopenia obtained only 21.7%, while other studies from Zaigham reported lymphocytopenia in 59%⁷. This study also examined a chest X-ray and revealed that half of COVID-19 positive showed abnormal results (52%). A retrospective study from Yan on 116 COVID-19 pregnant women informed abnormal radiologic findings in 96.3% of cases and a higher proportion of symptomatic cases (76.7%)⁹. The Centers for Disease Control and Prevention (CDC) does not recommend that chest radiography to diagnose COVID-19. The findings on chest imaging are not specific and overlap with other infections. Therefore, this modality should not be accepted as a first-line test to diagnose COVID-19¹³. However, in the low-resource testing areas, this option can be considered.

The protocol in this study involved a rapid antibody test due to our limited testing capability. This revealed that the majority (78.3%) of maternal COVID-19 cases had a positive antibody test. Almost all patients with COVID-19 will test positive for antibodies within 10–20 days after symptoms, but the antibody test's clinical significance cannot be explained in either the pregnant population or obstetric cases. There were also various methods for antibody testing for SARS-COV-2 with different sensitivity and specificity¹⁴. In general, if the results of rapid antibody tests are positive, we will continue with swab RT-PCR to distinguish whether this patient is still infected¹⁵.

It should be known that triage or COVID-19 cases sorting must be performed at the hospital¹⁶. Unlike many gynecological visits that can be postponed, obstetric patients' arrival, especially during labor, frequently cannot be planned. It is fundamental to develop strategies to secure the safety of patients and health workers.

Therefore, the implementation of universal testing must be strongly considered in all obstetric cases¹⁷. The safety of women, babies, partners, and health staff remains an absolute priority. Offering testing to women receiving maternity care can reduce nosocomial transmission. However, universal testing can only be arranged in high-resource hospitals with adequate laboratories. Bowling's recommendations were to take universal testing if the hospital's capacity could adhere to the high number of asymptomatic COVID-19¹⁸. Therefore, other approaches need to be considered.

This study added more parameters (lymphocytopenia, abnormal chest x-ray, and rapid antibody test) in COVID-19 screening to increase the testing capacity. With this screening protocol, we can detect more COVID-19 patients. If only symptoms are used, this strategy can only detect 11 out of 190 patients (5.8%), while if we expanded the screening strategy also by attending at lymphocytopenia, abnormal chest X-ray, and rapid antibody tests, the detected cases increased more than 2-fold to 23 cases per 190 patients (12.1%). Tests based on symptoms will not be effective and endanger the maternal delivery ward of new COVID-19 transmissions.

The symptoms only have a sensitivity and specificity of 48% and 74%. There is still a few research that looks at COVID-19 screening capabilities. A study of Vintzileos revealed lower sensitivity (34.4%) but higher specificity (96.1%) in maternal symptoms. The highest specificity was found in the history of COVID-19 contact, but as explained earlier, evaluating contacts especially in extensive local transmission conditions is not easy, patients sometimes not know whether she was exposed or try to avoid COVID-19 testing by hiding past symptoms and contact history².

No other studies have observed the ability to select other parameters (lymphocytopenia, chest X-rays, and rapid antibody tests). As revealed earlier, by supplementing these parameters, the COVID-19 detection raised almost twice. The antibody test had the highest sensitivity (78.3%), while chest X-rays and lymphocytopenia had a lower sensitivity (52% and 22%). A good screening test has several characteristics, including being used for critical health problems, diagnosis and therapy facilities are available, not expensive, and has good sensitivity¹⁹. COVID-19 can be detected indirectly by measuring the immune response; therefore, serological diagnosis is vital for mild problems¹⁴. Although the highest antibody detection is obtained after the first week, Guo's analysis confirmed that the fastest detection even starts from the first day symptoms appear²⁰.

This study focuses on finding ways to detect asymptomatic cases of COVID-19 in RT-PCR limited testing situations. Asymptomatic cases are important because the potential for transmission remains reported by asymptomatic transmission from Bai²¹ and several Lancet publications that showed asymptomatic pediatric patients in family clusters with abnormal radiographs. A different study has shown that the detectable viral load in asymptomatic cases is almost the same as in symptomatic shows the potential transmission ability from asymptomatic case²². Some studies have shown that most obstetric cases with COVID-19 are asymptomatic; hence universal testing is extremely recommended²³. Although this cannot be

generalized in low infection states, COVID-19 status is essential to determine isolation in hospitals, bed allocation, CDC's newborn protocol, and management of Personal Protective Equipment (PPE) procedure^{24,25}. This study presents an excellent alternative to the COVID-19 screening protocol in maternal health services. This protocol can increase the detection of up to 12 asymptomatic patients. Indeed, if universal testing is implemented as in developed countries, we need 190 swab tests, whereas, with this protocol, we only perform RT-PCR swab tests only on 69 patients at risk, which can reduce costs and keep the test for other patients who also need it.

A limitation of this study was that we did not perform RT-PCR swabs on all patients who had negative results on initial screening. So that the possibility of an undetected COVID-19 case was not ruled out. Further studies to compare our screening protocols and universal testing will allow us to see the screening protocol's capabilities more precisely. Our screening protocol can detect more asymptomatic patients and saving medical personnel who have essential inpatient care roles. The conditions in Italy that reported the death of several health workers with 3000 or more affected by COVID-19 indicate the importance of identifying cases, especially in maternal care that can never be delayed and always requires direct contact services for delivery¹⁸.

CONCLUSION

More asymptomatic cases of COVID-19 can be detected using our hospital screening protocol (Dr. Soetomo General Hospital) than the RT-PCT swab test in selected symptomatic cases. Rapid antibody testing has high sensitivity and a beneficial role in screening for COVID-19, particularly in areas with high COVID-19 rates with limited testing capacity. This study reveals the importance of identifying COVID-19, notably in asymptomatic cases, to prevent infection transmission to newborns, other patients, patients' families, and health workers urgently required in this pandemic.

Conflict of interest: The authors declared there were no competing interests in the study.

Ethical approval: This research was declared an ethical pass test by the Ethics Committee of Dr. Soetomo General Hospital, Surabaya East Java Indonesia.

Funding: This study was funded by the authors.

REFERENCES

- Setiati S, Azwar MK. COVID-19 and Indonesia. *Acta Med Indones.* 2020;52(1):84–9.
- Vintzileos WS, Muscat J, Hoffmann E, John NS, Vertichio R, Vintzileos AM, et al. Screening all pregnant women admitted to labor and delivery for the virus responsible for coronavirus disease 2019. *Am J Obstet Gynecol.* 2020;
- Sutton D, Fuchs K, D'alton M, Goffman D. Universal screening for SARS-CoV-2 in women admitted for delivery. *N Engl J Med.* 2020;382(22):2163–4.
- Covid W. Coronavirus Pandemic. Accessed May. 19AD;16:2020.
- Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet.* 2020;395(10223):497–506.
- Jacobi A, Chung M, Bernheim A, Eber C. Portable chest X-ray in coronavirus disease-19 (COVID-19): A pictorial review. *Clin Imaging.* 2020;
- Zaigham M, Andersson O. Maternal and perinatal outcomes with COVID-19: a systematic review of 108 pregnancies. *Acta Obstet Gynecol Scand.* 2020;99(7):823–9.
- Guan W, Ni Z, Hu Y, Liang W, Ou C, He J, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med.* 2020;382(18):1708–20.
- Yan J, Guo J, Fan C, Juan J, Yu X, Li J, et al. Coronavirus disease 2019 in pregnant women: a report based on 116 cases. *Am J Obstet Gynecol.* 2020;223(1):111–e1.
- Breslin N, Baptiste C, Gyamfi-Bannerman C, Miller R, Martinez R, Bernstein K, et al. COVID-19 infection among asymptomatic and symptomatic pregnant women: Two weeks of confirmed presentations to an affiliated pair of New York City hospitals. *Am J Obstet Gynecol MFM.* 2020;100118.
- Breslin N, Baptiste C, Miller R, Fuchs K, Goffman D, Gyamfi-Bannerman C, et al. COVID-19 in pregnancy: early lessons. *American Journal of Obstetrics and Gynecology MFM.* 2020;
- Noelle B, Caitlin B, Cynthia G-B, Russell M, Rebecca M, Kyra B, et al. COVID-19 infection among asymptomatic and symptomatic pregnant women: two weeks of confirmed presentations to an affiliated pair of New York City hospitals. *Am J Obs Gynecol MFM.* 2020;
- Henkel M, Weikert T, Marston K, Schwab N, Sommer G, Haslbauer J, et al. Lethal COVID-19: Radiological-Pathological Correlation of the Lungs. *Radiol Cardiothorac Imaging.* 2020;2(6):e200406.
- Sethuraman N, Jeremiah SS, Ryo A. Interpreting diagnostic tests for SARS-CoV-2. *Jama.* 2020;323(22):2249–51.
- Zullo F, Di Mascio D, Saccone G. Coronavirus disease 2019 antibody testing in pregnancy. *Am J Obstet Gynecol MFM.* 2020;2(3):100142.
- Capanna F, Haydar A, McCarey C, Bernini Carri E, Bartha Rasero J, Tsibizova V, et al. Preparing an obstetric unit in the heart of the epidemic strike of COVID-19: quick reorganization tips. *J Matern Neonatal Med.* 2020;1–7.
- Werner EF, Louis JM, Hughes B, Han CS, Norton ME, Srinivas SK. Community obstetrical units less likely than academic units to have universal COVID-19 testing. *Am J Perinatol.* 2020;37(10):1074–6.
- Boelig RC, Manuck T, Oliver EA, Di Mascio D, Saccone G, Bellussi F, et al. Labor and delivery guidance for COVID-19. *Am J Obstet Gynecol MFM.* 2020;2(2):100110.
- Stoll BJ, Hansen NI, Higgins RD, Fanaroff AA, Duara S, Goldberg R, et al. Very low birth weight preterm infants with early onset neonatal sepsis: the predominance of gram-negative infections continues in the National Institute of Child Health and Human Development Neonatal Research Network, 2002–2003. *Pediatr Infect Dis J.* 2005;24(7):635–9.
- Guo L, Ren L, Yang S, Xiao M, Chang D, Yang F, et al. Profiling early humoral response to diagnose novel coronavirus disease (COVID-19). *Clin Infect Dis.* 2020;71(15):778–85.
- Bai Y, Yao L, Wei T, Tian F, Jin D-Y, Chen L, et al. Presumed asymptomatic carrier transmission of COVID-19. *Jama.* 2020;323(14):1406–7.
- Zou L, Ruan F, Huang M, Liang L, Huang H, Hong Z, et al. SARS-CoV-2 viral load in upper respiratory specimens of infected patients. *N Engl J Med.* 2020;382(12):1177–9.
- Ferrazzi EM, Frigerio L, Cetin I, Vergani P, Spinillo A, Prefumo F, et al. COVID-19 Obstetrics Task Force, Lombardy, Italy: executive management summary and short report of outcome. *Int J Gynecol Obstet.* 2020;149(3):377–8.
- Mayor S. Covid-19: Nine in 10 pregnant women with infection when admitted for delivery are asymptomatic, small study finds. *British Medical Journal Publishing Group;* 2020.
- Poon LC, Yang H, Lee JCS, Copel JA, Leung TY, Zhang Y, et al. ISUOG Interim Guidance on 2019 novel coronavirus infection during pregnancy and puerperium: information for healthcare professionals. *Ultrasound Obstet Gynecol.* 2020;55(5):700–8