

the risk persistent hypertension

by Ernawati Ernawati

Submission date: 27-Apr-2023 10:12PM (UTC+0800)

Submission ID: 2077189554

File name: 27_the_risk_persistent_hypertension.pdf (431.48K)

Word count: 5279

Character count: 27385

The risk of persistent hypertension and chronic kidney disease in early and late-onset preeclampsia, a report from a low middle income country

Emawati Emawati (✉ emawati@fk.unair.ac.id)

Universitas Airlangga – Dr. Soetomo Academic General Hospital

Agus Sulistyono

Universitas Airlangga – Dr. Soetomo Academic General Hospital

Aditiawardana Aditiawardana

Universitas Airlangga

Kamalia Kamalia

Universitas Airlangga – Dr. Soetomo Academic General Hospital

Salsabila Nabilah Rifdah

Universitas Airlangga – Dr. Soetomo Academic General Hospital

M. Ilham Aldika Akbar

Universitas Airlangga – Dr. Soetomo Academic General Hospital

Erry Gumilar

Universitas Airlangga – Dr. Soetomo Academic General Hospital

Aditiawarman Aditiawarman


Universitas Airlangga – Dr. Soetomo Academic General Hospital

Research Article

Keywords: ¹¹ Early-onset of preeclampsia (EOP), ¹¹ late-onset of preeclampsia (LOP), chronic kidney disease, persisten hypertension, long-term effects of preeclampsia

Posted Date: January 12th, 2023

²⁷
DOI: <https://doi.org/10.21203/rs.3.rs-2460877/v1>

²¹
License:  This work is licensed under a ²¹ Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Abstract

Background

⁵⁴ Early-onset preeclampsia (EOP) and late-onset preeclampsia (LOP) are associated with different maternal and fetal outcomes, biochemical markers, and clinical characteristics. Nevertheless, only a few data were presented about its long-term effects on kidney function ³⁹ later in life.

Objective

This study aimed to explore the risk of persistent hypertension and kidney failure in EOP and LOP in five years after delivery. **Study Design:** This retrospective cohort study included ¹³ women with prior history of severe preeclampsia or normotensive pregnancy ¹³ admitted to one of tertiary hospitals in Indonesia. The ¹³ blood pressure, body mass index, urea and creatinine serum, and protein urine were analyzed, and risk of chronic kidney disease (CKD) was performed using KDIGO classification.

Results

A total of 27 EOP, 35 LOP and 30 normotensive cases were included. Mean blood pressure after five years recorded respectively 115.6 ± 14.25 mmHg in normotensive group, 131.82 ± 19.34 mmHg in LOP group, and 154.96 ± 23.48 mmHg in EOP group. Percentage of women with positive protein urine varied from 13.3–31.4% and 66.7% in normotensive, LOP, and EOP, respectively. According to KDIGO classification, normotensive group had 90% of low-risk CKD, whereas the severe preeclampsia group had 41.9% of high-risk CKD. In the severe preeclampsia group, the risk of CKD was 20.94 times higher compared to normotensive women (OR 20.94; 95% CI [2.67-163.72], $p = 0.004$). Then risk of CKD in EOP group was 6.75 times higher than LOP group (OR 6.75; 95% CI [2.19–20.76], $p = 0.001$), whereas persistent hypertension in EOP group was 5.78 times higher than LOP group (OR 5.78; 95% CI [1.91-17.395], $p = 0.002$).

Conclusions

⁴ Preeclampsia women have a higher risk of CKD than normotensive women. Women with a history of EOP are more likely to develop persistent hypertension and CKD than women with prior LOP history.

³⁵ Introduction

Pre-eclampsia (PE), a multisystem disorder characterized by hypertension and frequently coinciding with new-onset proteinuria, is associated with a number of problems that place women and their fetuses at a disproportionate risk for further complications, as well as life-long sequelae.¹ PE has been

disproportionately prevalent in developing countries and constitutes a leading cause of maternal mortality in low-income countries.² Women with PE in developing countries have a higher risk of mortality than those in developed countries, and hypertension is one of the leading causes of maternal mortality in PE.^{3,4} Currently, PE is generally classified into EOP and LOP, which exhibit different clinical manifestations and pathogenesis. EOP is associated with placental insufficiency and defective vascular remodelling, whereas LOP is most likely caused by maternal factors, particularly vascular maladaptation.^{5,6,7}

Prior research has linked PE to permanent kidney damage, including an increased risk of albuminuria⁸, chronic kidney disease (CKD)⁹, and end-stage kidney disease (ESKD).¹⁰ Some studies report that kidney dysfunction can resolve in most women with a history of PE^{11,12}, however, some women with PE may experience persistently decreased GFR and/or proteinuria and/or an increased risk of CKD.^{13,14} Only few data were presented about its long-term effects on kidney function later in life, mostly in early-onset preeclampsia (EOP) and late-onset preeclampsia (LOP). Since EOP and LOP are associated with different maternal and fetal outcomes, biochemical markers, and clinical characteristics.¹⁵ It is thought that EOP poses a substantial risk to both mother and fetus¹⁶, whereas LOP may manifest with less severe clinical symptoms.¹⁷ It has been suggested that EOP and LOP have different risks of developing renal impairment in women with preeclampsia. Unfortunately, to our knowledge there is no data or international publication on the risk of CKD following preeclampsia from developing country. Thus, this study intends to explore the risk of renal failure in EOP and LOP five years after preeclampsia in Indonesia as one of low middle income country in South East Asia.

Methods

This was a retrospective cohort study of women who had previously been diagnosed with severe PE or eclampsia and delivered in Dr. Soetomo Academic Hospital, the largest tertiary referral hospitals in East Indonesia between January 2013 and June 2014. Using the medical records, we identified all patients who met our criteria and included them in the exposed group, whereas women who recorded having uncomplicated pregnancy constituted as control group. All exposed cases who lived in Surabaya and willing to engage in this study were then enrolled. However, women with pre-existing comorbidities such as chronic hypertension, kidney disease, autoimmune disease, cardiac disease at the time of their pregnancy, women who had twins or multiple fetuses, and patients who died over the course of this study were excluded. All samples were contacted and/or visited at their residences before being invited to the hospital for a medical examination and blood sampling test.

PE or eclampsia were defined according to revised ISSHP criteria: the presence of hypertension (systolic blood pressure ≥ 140 mmHg and/or diastolic blood pressure ≥ 90 mmHg), which developed after 20 weeks of pregnancy and the coexistence of one or more of new onset conditions: proteinuria (spot urine protein/creatinine ratio ≥ 430 mg/mmol or $\geq 4,300$ mg protein/24 h or at least "2+" on dipstick testing), and/or other maternal organ dysfunction and/or suspected of intrauterine growth restriction (IUGR). Depending on time, EOP was defined as PE that develops before 34 weeks of gestation, whereas LOP was defined as

PE that develops at or after 34 weeks of gestation.¹⁸ An uncomplicated pregnancy history was defined as women who giving birth between 37 and 42 weeks of gestation and having a normal blood pressure and with absence of IUGR.

The primary outcome of the study was risk of chronic kidney disease defined according to Kidney Disease Improvement Global Outcomes (KDIGO) 2012 definitions for the classification of chronic kidney disease based on renal function measured by GFR. Renal function was measured using serum creatinine values determined by the Jaffe method and calibrated using isotope dilution mass spectrometry (IDMS) method. Estimated GFR (eGFR) was calculated using the modified Cockcroft and Gault: $GFR = \frac{((140 - \text{age (year)}) \times \text{weight (kg)})}{(72 \times \text{serum creatinin (mg/dL)})} \times 0.85$. Proteinuria was measured using urinalysis examination. According to these criteria, risk of chronic kidney disease classified into low risk if the eGFR ≥ 60 mL/min/1.73m² and proteinuria < 30 mg/g, whereas high risk if the eGFR 30–44 mL/min/1.73m² and proteinuria < 30mg/g or the eGFR ≥ 60 mL/min/1.73m² and proteinuria > 300 mg/g.¹⁹

Basic characteristics data were presented for women with priorly diagnosed EOP or LOP compared to those healthy pregnant woman. Categorical data was presented as frequencies (percentage), while continuous variables was presented either as mean (standard deviation/SD) or median (interquartile range/IQR). Differences were analyzed using Kruskal Wallis and the Fisher exact test was applied as an alternative test. Differences in values between groups considered statistically significant if the P-value was < 0.05. Odds ratios for the primary outcomes were calculated using logistic regression with 95% confidence intervals (CI). All statistical analysis were performed using SPSS 21 software (IBM Corp, Armonk, NY). The Human Research and Ethics Committee for Basic Science and Clinical Research of Dr. Soetomo General Academic Hospital approved the research protocol (Ref. No: 0842/KEPK/XII/2018). Informed consent was obtained from all participants before the initiation of the study.

Results

During periods of study, 673 women who were previously diagnosed with PE between January 2013 and June 2014 were listed from medical record. At the beginning, all patients were assessed for eligibility criteria and of whom 235 who met inclusion and exclusion criteria were enrolled to this study. After excluding some patients due to some reasons, finally we included 62 exposed women with prior PE history, consisting of 27 with EOP and 35 with LOP. We also recruited 30 healthy pregnant women to participate in this study as control group. Details regarding the study selection are documented in Fig. 1.

Basic characteristics and laboratory outcomes of included participants when first diagnosed with PE were presented on Table 1. Our study obtained that women with prior PE tend to have a higher mean of maternal age compared to control group with normotensive blood pressure at the time of delivery. The LOP group was dominated by multiparity, whereas in the EOP group was dominated by nulliparity. Our results also obtained that women with EOP had significantly higher mean of body mass index (BMI) compared to women with LOP and control group ($p = 0.019$) and were more likely to have chronic

hypertension, kidney disease and diabetes. However, those with chronic hypertension or CKD at baseline were excluded from further analysis. As expected, the mean systolic blood pressure of women with prior EOP and LOP were significantly higher compared to control group ($p = 0.001$), as well as diastolic blood pressure ($p = 0.001$). Our study also obtained that gestational age at delivery were significantly differed between groups ($p = 0.001$). Moreover, there was no statistically significant difference of eGFR between EOP, LOP and control group ($p = 0.577$).

Table 1
Baseline characteristics of the sample at delivery

Characteristics	Control (n = 30)	Pre-eclampsia		P-value		
		LOP (n = 35)	EOP (n = 27)	C LOP	C EOP	LOP EOP
Age, mean (SD), years	30.1 ± 6.06	34.46 ± 7.38	33.07 ± 7.01	0.058		
Parity, n (%)						
Nullipara	11 (36.6)	14 (40)	20 (74.1)	0.785	0.005*	0.006*
Multipara	19 (63.4)	21 (60)	7 (25.9)			
BMI, mean (SD), (kg/m ²)	28.27 ± 4.07	27.07 ± 4.51	30.23 ± 4.94	0.019*		
Gestational age at delivery, mean (SD), (weeks)	38.89 ± 1.38	37.20 ± 2.86	33.3 ± 3.01	0.001*		
Blood pressure at admission mean (SD), (mmHg)						
Systolic	111.66 ± 12.05	160.34 ± 18.22	168.88 ± 15.63	0.001*		
Diastolic	70.66 ± 9.07	102.11 ± 8.20	90.00 ± 16.40	0.001*		
Renal function test at admission						
BUN, n (%)	0 (0)	2 (5.7)	5 (18.5)	0.187	0.014*	1.117
BUN > 21 mg/dL						
Serum creatinine, n (%)	1 (3.3)	0 (0)	2 (7.4)	0.280	0.495	0.104
Cr > 1.1 mg/dL						
eGFR, mean (SD) (ml/min/1.73m ²)	194.93 ± 33.03	171.37 ± 50.86	163.79 ± 57.78	0.577		
Proteinuria (+), n (%)	0 (0)	35 (100)	27 (100)	0.001*	0.001*	

*P-value from Kruskal-Wallis test for three groups, while fisher exact test for each two groups; C, control group; EOP, early-onset severe preeclampsia; LOP, late-onset severe preeclampsia; SD, standard deviation; BMI, body mass index; BUN, blood urea nitrogen; Cr, serum creatinine; eGFR, estimated glomerular filtration rate.

After five years from being first diagnosed with severe PE, the mean ⁴⁵ systolic and diastolic blood pressure showed significantly higher result in EOP group compared to all groups ($p = 0.001$). We obtained that women with EOP also had higher mean of BMI ($30.23 \pm 4.94 \text{ kg/m}^2$) compared to LOP and control group ($27.07 \pm 4.51 \text{ kg/m}^2$ and $28.27 \pm 4.07 \text{ kg/m}^2$, respectively), although failed to show significant difference ($p = 0.084$). All parameters in renal function showed statistically significant difference between groups, except for the number of positive proteinuria and abnormal protein-to-creatinine ratio between LOP compared to control group ($p > 0.05$). Additionally, our analysis obtained that women with prior EOP history showed a significant decrease of eGFR compared to all groups ($p = 0.001$), indicating that the group with prior history of EOP may pose worse renal outcome compared to other groups. Detailed information regarding other characteristics is documented in Table 2.

Table 2
Characteristics of the sample five years after delivery

Characteristics	Control (n = 30)	Pre-eclampsia		P-value		
		LOP (n = 35)	EOP (n = 27)	C LOP	C EOP	LOP EOP
BMI, mean (SD), (kg/m ²)	28.27 ± 4.07	27.07 ± 4.51	30.23 ± 4.94	0.084		
Blood pressure at admission mean (SD), (mmHg)						
Systolic	115.6 ± 14.25	131.82 ± 19.34	154.96 ± 23.48	0.001*		
Diastolic	66.53 ± 11.41	81.74 ± 14.49	96.00 ± 16.16	0.001*		
Persistent hypertension, n (%)	0 (0)	9 (25.7)	18 (66.7)	0.003*	0.001*	0.001*
Renal function test, mean (SD)						
BUN, n (%)	0 (0)	0 (0)	10 (37.1)		0.001*	0.001*
BUN > 21 mg/dL						
Serum creatinine, n (%)	0 (0)	0 (0)	10 (37.1)		0.001*	0.001*
Cr > 1.1 mg/dL						
eGFR, mean (SD) (ml/min/1.73m ²)	143.67 ± 33.77	120.80 ± 42.81	97.22 ± 28.71	0.001*		
Proteinuria (+), n (%)	4 (13.3)	11 (31.4)	18 (66.7)	0.087	0.001*	0.006*
Abnormal albumine-to- cratinine ratio, n (%)	4 (13.3)	13 (37.1)	19 (70.4)	0.031*	0.001*	0.010*
Abnormal protein-to- creatinine ratio, n (%)	6 (20)	9 (25.7)	18 (66.7)	0.589	0.001*	0.001*
Persistent proteinuria, n (%)	0 (0)	11 (31.4)	18 (66.7)	0.001*	0.001*	0.006*

*P-value from Kruskal-Wallis test for three groups, while fisher exact test for each two groups; C, control group; EOP, early-onset severe preeclampsia; LOP, late-onset severe preeclampsia; SD, standard deviation; BMI, body mass index; BUN, blood urea nitrogen; Cr, serum creatinine; eGFR, estimated glomerular filtration rate.

Further analysis regarding the incidence of persistent hypertension among women with prior history of PE were varied from 66.7% and 25.7% in EOP and LOP groups respectively. According to the logistic regression analysis, RR of developing persistent hypertension is significantly higher among women with

prior history of EOP (RR 5.778; P-value = 0.002; 95% CI 1.919–17.395), as well as the risk of developing CKD (RR 6.75; P-value = 0.001; 95% CI 2.194–20.764) compared to women with prior history of LOP (Table 3). Likewise, according to KDIQO 2012 classification, women with prior severe PE history had significantly higher risk of further developing CKD (RR 20.94; P-value = 0.004; 95% CI 2.679–163.723) compared to normotensive control group.

Table 3
The association between the type of PE with the incidence of persistent hypertension and the risk of developing CKD 5 years after diagnosis.

Variables	Type of PE		Total	P-value	RR (95%CI)
	LOP	EOP			
Persistent Hypertension					
Yes	9 (25.7%)	18 (66.7%)	27	0.002*	5.778 (1.919–17.395)
No	26 (74.3%)	9 (33.3%)	35		
Risk of CKD					
High	8 (22.9%)	18 (66.7%)	26	0.001*	6.75 (2.194–20.764)
Low	27(77.1%)	9 (33.3%)	36		

*P-value from Kruskal-Wallis test for three groups; PE, pre-eclampsia; EOP, early-onset pre-eclampsia; LOP, late-onset pre-eclampsia; CKD, chronic kidney disease; RR, relative risk; CI, confidence interval.

Table 4
Association between severe PE and risk of developing CKD five years after diagnosis.

Variable	CKD risk		Total	P-value	RR(95%CI)
	High	Low			
Severe PE					
Yes	26 (28.3%)	36 (39.1%)	62	0.004*	20.94 (2.679–163.723)
No	3 (3.2%)	27 (29.3%)	30		

*P-value from Kruskal-Wallis test between groups; PE, pre-eclampsia; CKD, chronic kidney disease; RR, relative risk; CI, confidence interval.

Discussion

This study revealed that 5 years after delivery, women with a history of preeclampsia were at risk for persistent hypertension. EOP and LOP cases had higher blood pressure than normal pregnant women. Result from prior study indicated that roughly 20% and 8% of women with a history of PE still had

hypertension and proteinuria six months postpartum, respectively.²⁰ The study by Berks et al., showed 39% and 14% of women with a prior history of PE remained had high blood pressure and proteinuria three months postpartum, while 18% and 8% remained had hypertension and proteinuria two years afterwards.²¹ Lykke et al., found that women with a history of severe PE had the higher risk of chronic or persistent hypertension compared to those with PE (RR 6.07 vs. 3.61).²²

Our study found that women with EOP have a higher blood pressure than LOP. Women with EOP have a risk of developing persistent hypertension 5.7 times higher than LOP. Study by Veerbeek et al. observed that over nearly half of women with a history of EOP developed persistent postpartum hypertension. Moreover, the blood pressure of women with an EOP history and pregnancy-induced hypertension was considerably greater than that of women with a LOP history.⁶ Comparing EOP and LOP, the maternal vascular response and remodelling pattern revealed distinct vascular adaptations. Therefore, increased vascular resistance can contribute to systolic and diastolic dysfunction and could be a driving force behind the development of chronic hypertension in women with EOP.⁷ Consequently, our findings support the idea that cardiovascular risk during pregnancy is predictive of cardiovascular risk later in life, particularly the risk of persistent hypertension.^{23,24}

Our study indicated that women with a history of severe PE had a greater risk of CKD than the normotensive group. The risk of developing CKD at 5 years after delivery in severe PE patients is 20 times higher than in normal pregnancies. Patients with a previous history of preeclampsia presented lower eGFR and had more cases of persistent protein urine than normal pregnant women. Moreover, eGFR was the lowest in the EOP group. Previous report including a large cohort study found hypertensive disorders of pregnancy are associated with an increased risk of subsequent CKD. Renal impairment was also found earlier in women with GH and PE than in normotensive women.¹³ Another study also reported closely association between PE and gestational Hypertension with risk of renal disorder in the the future.^{9,25-27}

Risk of CKD in EOP and LOP differed significantly in this study. EOP had higher risk to develop CKD than LOP. The large cohort study by Vikse et al., with a sample of 570,433 women, showed that preeclampsia was a risk factor for the development of end-stage renal disease (ESRD). The risk is higher in preeclampsia patients who give birth to premature babies or children with low birth weight, which indicates EOP cases.²⁵ Irreversible vascular damage due to more severe endothelial damage and inflammatory stress in EOP than in LOP cannot be disregarded.²⁸ Renal histology of postpartum biopsies on PE patients showed glomerular endotheliosis and vascular injury as classic pathologies features²⁹ support this finding. Preeclampsia is suggested to develop kidney disease by causing acute renal impairment, endothelial damage, and podocyte loss.³⁰

Endothelial dysfunction induced by PE persists after preeclampsia in many patients.³¹ The remaining endothelial dysfunction is unknown; It could be assumed that endothelial cell disturbance enhances by a high level of sFlt1 in women with a history of preeclampsia and also due to epigenetic changes induced by preeclampsia.³²

Increased levels of sFlt1 have been found in formerly preeclamptic women.^{33,34} This persistence of elevated levels of sFlt1 in women with a history of preeclampsia is expected due to an extra-placental source such as endothelial cells and monocytes. This increased sFlt1 may lead to changes in the vascular endothelium, increasing the risk of renovascular diseases in later life. Interestingly, increasing sFlt1 was also found in CKD patients without a history of preeclampsia, which positively correlates with proteinuria.³⁵

This fact follows EOP. The combination of insufficient immune tolerance to the fetus and poor placentation resulted in the elevation of serum sFlt-1 and decreasing of PIGF level, thus causing vascular endothelial dysfunction, which led to Preeclampsia manifestation by 34 weeks gestation.³⁶ It may explain that the risk of developing CKD was higher in EOP than in LOP.

The strengths of this study are mostly related to our hospital (Dr. Soetomo General Academic Hospital), a level 3 and top referral centre hospital in eastern Indonesia. At level 3, we have many cases of preeclampsia, and almost all cases of early-onset preeclampsia and all conservative management are referred to our hospital. Therefore, we have a large number of EOP and LOP cases.

As the limitation of the study, we have to consider that this was a retrospective cohort study. Thus, some information may be missed, like no assessment of renal anatomic abnormalities before pregnancy or when the patient was diagnosed with severe PE, as it is not a standard procedure for initial examination before pregnancy or at the time of diagnosis in our hospital. The high mobility of the population makes this research even more challenging. Since most of the patients in this study were seasonal residents who moved frequently, it was difficult to ascertain their whereabouts, reducing the number of participating patients. However, these limitations do not invalidate our conclusion that EOP is associated with a higher risk of CKD than LOP.

Conclusion

In women with a history of PE, the outcome of persistent hypertension and/or proteinuria may have renal repercussions. The systolic and diastolic blood pressures were found to be substantially associated with prior history of PE, which influences future renal consequences in both the EOP and LOP groups. In a five-year follow-up, women with severe PE had a higher risk of developing CKD than normotensive women. In addition, women with a history of EOP are more likely to develop persistent hypertension and CKD than women with prior LOP history. Further comprehensive prospective studies with well-designed and longer follow-ups are required to confirm the findings of this study.

Abbreviations

EOP
Early-onset preeclampsia
LOP

late-onset preeclampsia

PE

Pre-eclampsia

GH

gestational Hypertension

CKD

chronic kidney disease

ESKD

end-stage kidney disease

ESRD

end-stage renal disease

KDIGO

Kidney Disease Improvement Global Outcome

GFR

glomerular filtration rate

eGFR

estimated glomerular filtration rate

ISSHP

International Society for the Study of Hypertension in Pregnancy

IDMS

isotope dilution mass spectrometry

IUGR

intrauterine growth restriction

SD

standard deviation

IQR

interquartile range

CI

confidence intervals

BMI

body mass index.

Declarations

Ethics approval and consent to participate

The study was approved by the Human Research and Ethics Committee for Basic Science and Clinical Research of Dr. Soetomo General Academic Hospital approved the research protocol (Ref. No: 0842/KEPK/XII/2018). Informed consent was obtained from all participants before the initiation of the study.

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

⁵ This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Authors' contributions

ERN, ADM: conceived research idea, supervised, designed questionnaire, data collection, manuscript draft and review. MIA, EGD: co-supervised, manuscript review. ERN, KML, ASL: designed questionnaire, data collection, manuscript draft. KML, SNR: data analysis, manuscript draft and review. ERN, KML,ADW, ADM: design of research, data collection, data analysis. All authors read and approved the manuscript.

Acknowledgements

Not applicable.

References

1. Sibai BM, Dekker G, Kupferminc M. Pre-eclampsia. *The Lancet*. 2005;365(9461):785. doi: 10.1016/S0140-6736(05)17987-2
2. Akbar MIA, Wicaksono B, Dachlan E. Maternal mortality and its mainly possible cause pre-eclampsia/eclampsia in developing country (Surabaya-Indonesia as the model). *Pregnancy Hypertension: An International Journal of Women's Cardiovascular Health*. 2012;2(3):184. doi: 10.1016/j.preghy.2012.04.019
3. Lo JO, Mission JF, Caughey AB. Hypertensive disease of pregnancy and maternal mortality. *Curr Opin Obstet Gynecol*. 2013;25(2):124-32. doi: 10.1097/GCO.0b013e32835e0ef5.
4. Wagner LK. Diagnosis and management of preeclampsia. *Am Fam Physician*. 2004;70(12): 2317-24.
5. Stergiotou I, Crispi F, Valenzuela-Alcaraz B, Bijmens B, Gratacos E. Patterns of maternal vascular remodeling and responsiveness in early-versus late-onset preeclampsia. *Am J Obstet Gynecol*. 2013;209:558.e1-14. <http://dx.doi.org/10.1016/j.ajog.2013.07.030>

6. Veerbeek JHW, Hermes W, Breimer AY, van Rijn BB, Koenen SV, Mol BW, et al. Cardiovascular Disease Risk Factors After Early-Onset Preeclampsia, Late-Onset Preeclampsia, and Pregnancy-Induced Hypertension. *Hypertension*. 2015;65:600-606. DOI: 10.1161/HYPERTENSIONAHA.114.04850.
7. Valensise H, Vasapollo B, Gagliardi G, Novelli GP. Early and late pre-eclampsia: two different maternal hemodynamic states in the latent phase of the disease. *Hypertension*. 2008;52:873–880. doi: 10.1161/HYPERTENSIONAHA.108.117358
8. Kattah AG, Asad R, Scantlebury DC, Bailey KR, Wiste HJ, Hunt SC, et al. Hypertension in pregnancy is a risk factor for microalbuminuria later in life. *J Clin Hypertens (Greenwich)*. 2013;15(9):617-623. doi:10.1111/jch.12116
9. Kristensen JH, Basit S, Wohlfahrt J, Damholt MB, Boyd HA. Pre-eclampsia and risk of later kidney disease: nationwide cohort study. *BMJ*. 2019;365:l1516. doi:10.1136/bmj.l1516
10. Covella B, Vinturache AE, Cabiddu G, Attini R, Gesualdo L, Versino E, et al. A systematic review and meta-analysis indicates long-term risk of chronic and end-stage kidney disease after preeclampsia. *Kidney Int*. 2019;96(3):711-727. doi:10.1016/j.kint.2019.03.033
11. Paauw ND, van der Graaf AM, Bozoglan R, van der Ham DP, Navis G, Gansevoort RT, et al. Kidney Function After a Hypertensive Disorder of Pregnancy: A Longitudinal Study. *Am J Kidney Dis*. 2018;71(5):619-626. doi:10.1053/j.ajkd.2017.10.014
12. Lopes van Balen VA, Spaan JJ, Comelis T, Spaanderman MEA. Prevalence of chronic kidney disease after preeclampsia. *J Nephrol*. 2017;30(3):403-409. doi:10.1007/s40620-016-0342-1
13. Ayansina D, Black C, Hall SJ, Marks A, Millar C, Prescott GJ, et al. Long term effects of gestational hypertension and pre-eclampsia on kidney function: Record linkage study. *Pregnancy Hypertens*. 2016;6(4):344-349. doi:10.1016/j.preghy.2016.08.231
14. McDonald SD, Han Z, Walsh MW, Gerstein HC, Devereaux PJ. Kidney disease after preeclampsia: a systematic review and meta-analysis. *Am J Kidney Dis*. 2010;55(6):1026-1039. doi:10.1053/j.ajkd.2009.12.036
15. Pettit F, Mangos G, Davis G, Henry A, Brown MA. Pre-eclampsia causes adverse maternal outcomes across the gestational spectrum. *Pregnancy Hypertens*. 2015;5(2):198-204. doi:10.1016/j.preghy.2015.02.002
16. Fox R, Kitt J, Leeson P, Aye CYL, Lewandowski AJ. Preeclampsia: Risk Factors, Diagnosis, Management, and the Cardiovascular Impact on the Offspring. *J Clin Med*. 2019;8(10):1625. doi:10.3390/jcm8101625
17. Wójtowicz A, Zembala-Szczerba M, Babczyk D, Kołodziejczyk-Pietruszka M, Lewaczyńska O, Huras H. Early- and Late-Onset Preeclampsia: A Comprehensive Cohort Study of Laboratory and Clinical Findings according to the New ISHHP Criteria. *Int J Hypertens*. 2019;2019:4108271. doi:10.1155/2019/4108271
18. Tranquilli AL. Early and late-onset pre-eclampsia. *Pregnancy Hypertens*. 2014;4(3):241. doi:10.1016/j.preghy.2014.04.007

19. KDIGO CKD Work Group. KDIGO 2012 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. *Kidney Int Suppl.* 2013;3(1):1-150
20. Girsberger M, Muff C, Hösli I, Jan Dickenmann M. Short term sequelae of preeclampsia: a single center cohort study. *BMC Pregnancy and Childbirth.* 2018;18:177. <https://doi.org/10.1186/s12884-018-1796-z>.
21. Berks D, Steegers EAP, Molas M, Visser W. Resolution of hypertension and proteinuria after preeclampsia. *Obstet Gynecol.* 2009;114:1307–14. <https://doi.org/10.1097/AOG.0b013e3181c14e3e>.
22. Lykke JA, Langhoff-Roos J, Sibai BM, Funai EF, Triche EW, Paidas MJ. Hypertensive pregnancy disorders and subsequent cardiovascular morbidity and type 2 diabetes mellitus in the mother. *Hypertension.* 2009;53:944–951. doi: 10.1161/HYPERTENSIONAHA.109.130765.
23. Magee LA, Von Dadelszen P. Pre-eclampsia and increased cardiovascular risk. *Br. Med. J.* 2007;335(7627):945–946.
24. Amaral LM, Cunningham MW, Cornelius DC, LaMarca B. Preeclampsia: Long-term Consequences for Vascular Health. *Vascular Health and Risk Management* 2015;11:403-415. <http://dx.doi.org/10.2147/VHRM.S64798>
25. Vikse BE, Irgens LM, Leivestad T, Skjærven R, Iversen BM. Preeclampsia and the risk of end-stage renal disease. *N. Engl. J. Med.* 2008;359(8):800–809.
26. Nisell H, Lintu H, Lunell NO, Möllerström G, Pettersson E. Blood pressure and renal function seven years after pregnancy complicated by hypertension. *Br J Obstet Gynaecol.* 1995;102(11):876-881. doi:10.1111/j.1471-0528.1995.tb10874.x
27. Shammash AG, Maayah JF. Hypertension and its relation to renal function 10 years after pregnancy complicated by pre-eclampsia and pregnancy induced hypertension. *Saudi Med J.* 2000;21(2):190-192
28. Staff AC, Dechend R, Redman CW. Review: Preeclampsia, acute atherosclerosis of the spiral arteries and future cardiovascular disease: two new hypotheses. *Placenta.* 2013;34 Suppl:S73-S78. doi:10.1016/j.placenta.2012.11.022
29. Fisher KA, Luger A, Spargo BH, Lindheimer MD. Hypertension in pregnancy: clinical-pathological correlations and remote prognosis. *Medicine (Baltimore).* 1981;60(4):267–76.
30. Kattah A. Preeclampsia and Kidney Disease: Deciphering Cause and Effect. *Curr Hypertens Rep.* 2020;22(11):91. doi:10.1007/s11906-020-01099-1
31. Blaauw J, Graaff R, van Pampus MG, van Doormaal JJ, Smit AJ, Rakhorst G, et al. Abnormal endothelium-dependent microvascular reactivity in recently preeclamptic women. *Obstet Gynecol.* 2005;105(3):626-632. doi:10.1097/01.AOG.0000153490.41973.e0
32. van der Graaf AM, Toering TJ, Faas MM, Lely AT. From preeclampsia to renal disease: a role of angiogenic factors and the renin-angiotensin aldosterone system?. *Nephrol Dial Transplant.* 2012;27 Suppl 3:iii51-iii57. doi:10.1093/ndt/gfs278

33. Wolf M, Hubel CA, Lam C, Sampson M, Ecker JL, Ness RB, et al. Preeclampsia and future cardiovascular disease: potential role of altered angiogenesis and insulin resistance. *J Clin Endocrinol Metab* 2004; 89: 6239–6243.
34. Wikström AK, Larsson A, Eriksson UJ, Nash P, Olovsson M. Early postpartum changes in circulating pro- and anti-angiogenic factors in early-onset and late-onset pre-eclampsia. *Acta Obstet Gynecol Scand.* 2008;87(2):146-153. doi:10.1080/00016340701819262
35. Di Marco GS, Reuter S, Hillebrand U, Amler S, König M, Larger E, et al. The soluble VEGF receptor sFlt1 contributes to endothelial dysfunction in CKD. *J Am Soc Nephrol* 2009; 20: 2235–2245.
36. Robillard PY, Dekker G, Scioscia M, Saito S. Progress in the understanding of the pathophysiology of immunologic maladaptation related to early-onset preeclampsia and metabolic syndrome related to late-onset preeclampsia. *Am J Obstet Gynecol.* 2021. doi: 10.1016/j.ajog.2021.11.019

Figures




Figure 1

Study Flowchart. PE, pre-eclampsia; EOP, early-onset pre-eclampsia; LOP, late-onset pre-eclampsia

the risk persistent hypertensio

ORIGINALITY REPORT

19%

SIMILARITY INDEX

12%

INTERNET SOURCES

16%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 "Urogynecology", Journal of Obstetrics and Gynaecology Research, 2020 1%

Publication
- 2 QiLong Jiang, Yang Liu, Yong Cao, Xinwen Bai, Yu Deng, Chengxiang Yu, Qizhi Song, Yan Li. "Comparison of nonanatomical reduction of femoral neck fractures in young patients treated with femoral neck system: a retrospective cohort study", Research Square Platform LLC, 2023 1%

Publication
- 3 Bianca Covella, Angela Elena Vinturache, Gianfranca Cabiddu, Rossella Attini et al. "A systematic review and meta-analysis indicates long-term risk of chronic and end-stage kidney disease after preeclampsia", Kidney International, 2019 1%

Publication
- 4 Maren Goetz, Mitho Müller, Raphael Gutsfeld, Tjeerd Dijkstra et al. "An observational claims data analysis on the risk of maternal chronic 1%

kidney disease after preterm delivery and preeclampsia", Scientific Reports, 2021

Publication

5

Shaz Ahmad, Reena Singh, Tanvir Arfin, Krishna Neeti. "Fluoride contamination, consequences and removal techniques in water: a review", Environmental Science: Advances, 2022

Publication

1 %

6

Saeedeh Salimi, Farzaneh Farajian-Mashhadi, Ehsan Tabatabaei, Mahnaz Shahrakipour, Minoos Yaghmaei, Mojgan Mokhtari. "Estrogen receptor alpha XbaI GG genotype was associated with severe preeclampsia", Clinical and Experimental Hypertension, 2017

Publication

<1 %

7

A. Perales, J. L. Delgado, M. de la Calle, J. A. García-Hernández et al. "sFlt-1/PIGF for prediction of early-onset pre-eclampsia: STEPS (Study of Early Pre-eclampsia in Spain)", Ultrasound in Obstetrics & Gynecology, 2017

Publication

<1 %

8

Choi, E.Y.. "Effects of Hemoglobin Concentration and Creatinine Clearance in Pro-B-Type Natriuretic Peptide-Based Left Ventricular Filling Pressure Prediction in Patients With Preserved Left Ventricular

<1 %

Systolic Function", The American Journal of Cardiology, 20080201

Publication

9	docplayer.net Internet Source	<1 %
10	emedicine.medscape.com Internet Source	<1 %
11	researchspace.ukzn.ac.za Internet Source	<1 %
12	www.era-online.org Internet Source	<1 %
13	Eva Maria Roes, Maarten TM Raijmakers, Marieke Schoonenberg, Natasha Wanner, Wilbert HM Peters, Eric AP Steegers. "Physical well-being in women with a history of severe preeclampsia", The Journal of Maternal-Fetal & Neonatal Medicine, 2009 Publication	<1 %
14	Alvarado-Esquivel, Cosme, Fernando Vázquez-Alaníz, Ada A Sandoval-Carrillo, José M Salas-Pacheco, Jesús Hernández-Tinoco, Luis Sánchez-Anguiano, and Oliver Liesenfeld. "Lack of association between Toxoplasma gondii infection and hypertensive disorders in pregnancy: a case-control study in a Northern Mexican population", Parasites & Vectors, 2014.	<1 %

15	cyberleninka.org Internet Source	<1 %
16	medcraveonline.com Internet Source	<1 %
17	Vanesa Alonso-Ventura, Yangzhou Li, Vinay Pasupuleti, Yuani M. Roman, Adrian V. Hernandez, Faustino R. Pérez-López. "Effects of preeclampsia and eclampsia on maternal metabolic and biochemical outcomes in later life: a systematic review and meta-analysis", <i>Metabolism</i> , 2020 Publication	<1 %
18	www.clinicaltrials.gov Internet Source	<1 %
19	S Berrettini, A Carabelli, S Sellari - Franceschini, L Bruschini, F Quartieri, F Sconosciuto, A Abruzzese. "Perennial allergic rhinitis and chronic sinusitis: correlation with rhinologic risk factors", <i>Allergy</i> , 2001 Publication	<1 %
20	escholarship.org Internet Source	<1 %
21	www.degruyter.com Internet Source	<1 %

22

Angela Vinturache, Joyce Popoola, Ingrid Watt-Coote. "The Changing Landscape of Acute Kidney Injury in Pregnancy from an Obstetrics Perspective", Journal of Clinical Medicine, 2019

Publication

<1 %

23

synapse.koreamed.org

Internet Source

<1 %

24

Cinthia E. M. Carbonara, Noemi A. V. Roza, Kelcia R. S. Quadros, Renata A. França et al. "Effect of aluminum accumulation on bone and cardiovascular risk in the current era", PLOS ONE, 2023

Publication

<1 %

25

EVA V. TYLDUM. "Maternal left ventricular and endothelial functions in preeclampsia : Cardiac and endothelial functions in PE", Acta Obstetricia Et Gynecologica Scandinavica, 05/2012

Publication

<1 %

26

Heri Suroto, Gestana Retaha Wardana, Julius Albert Sugianto, Dina Aprilya, Steven Samijo. "Time to surgery and myo-d expression in biceps muscle of adult brachial plexus injury: a preliminary study", BMC Research Notes, 2023

Publication

<1 %

27 Misa Sugeno, Hyo Kyojuka, Toki Jin, Fumihiro Ito et al. "Late-onset hypertensive disorders of pregnancy and red blood cell distribution width during late pregnancy among Japanese primipara: A nested case-control study", Research Square Platform LLC, 2023
Publication

28 cjasn.asnjournals.org
Internet Source

29 files.eric.ed.gov
Internet Source

30 jmedsoc.org
Internet Source

31 repub.eur.nl
Internet Source

32 www.ncbi.nlm.nih.gov
Internet Source

33 George J. Mangos. "Markers of cardiovascular disease risk after hypertension in pregnancy :", Journal of Hypertension, 02/2012
Publication

34 Karoline Mayer-Pickel, Christina Stern, Katharina Eberhard, Uwe Lang, Barbara Obermayer-Pietsch, Mila Cervar-Zivkovic. "Comparison of mean platelet volume (MPV) and sFlt-1/PIGF ratio as predictive markers for

preeclampsia", The Journal of Maternal-Fetal & Neonatal Medicine, 2019

Publication

35 Theodora Pittara, Andreas Vyrides, Demetris Lamnisis, Konstantinos Giannakou. <1 %
"Preeclampsia and long - term health outcomes for mother and infant: An umbrella review", BJOG: An International Journal of Obstetrics & Gynaecology, 2021
Publication

36 docksci.com <1 %
Internet Source

37 journals.lww.com <1 %
Internet Source

38 journals.plos.org <1 %
Internet Source

39 pure.rug.nl <1 %
Internet Source

40 www.alliedacademies.org <1 %
Internet Source

41 "Placental Dysfunction and Future Maternal Cardiovascular Disease", Human Placental Trophoblasts, 2015. <1 %
Publication

42 "Poster Day 1 - Oxidative Stress", Hypertension in Pregnancy, 7/2006 <1 %
Publication

43 Erica Shields Hammer, Marilyn J. Cipolla. "Cerebrovascular Dysfunction in Preeclamptic Pregnancies", Current Hypertension Reports, 2015

Publication

<1 %

44 Eva Stekkinger. "Early-Onset Preeclampsia and the Prevalence of Postpartum Metabolic Syndrome :", Obstetrics and Gynecology, 11/2009

Publication

<1 %

45 H. Nisell, H. Lintu, N. O. Lunell, G. Möllerström, E. Pettersson. "Blood pressure and renal function seven years after pregnancy complicated by hypertension", BJOG: An International Journal of Obstetrics & Gynaecology, 2005

Publication

<1 %

46 Marwan Ma'ayeh, Kara M. Rood, Douglas Kniss, Maged M. Costantine. "Novel Interventions for the Prevention of Preeclampsia", Current Hypertension Reports, 2020

Publication

<1 %

47 Morven Caroline Brown, Kate Elizabeth Best, Mark Stephen Pearce, Jason Waugh, Stephen Courtenay Robson, Ruth Bell. "Cardiovascular disease risk in women with pre-eclampsia: systematic review and meta-analysis",

<1 %

48 W. Gyselaers. "Role of dysfunctional maternal venous hemodynamics in the pathophysiology of pre-eclampsia: a review", *Ultrasound in Obstetrics and Gynecology*, 08/2011
Publication

49 oamjms.eu
Internet Source

50 openaccess.sgul.ac.uk
Internet Source

51 pubmed.ncbi.nlm.nih.gov
Internet Source

52 uu.diva-portal.org
Internet Source

53 www.chp.gov.hk
Internet Source

54 www.hindawi.com
Internet Source

55 www.karger.com
Internet Source

56 www.medrxiv.org
Internet Source

57

"Oral/Free Communication Session Abstracts",
International Journal of Gynecology &
Obstetrics, 2018

Publication

<1 %

58

Virginia Dines, Sonja Suvakov, Andrea Kattah,
Jane Vermunt et al. "Preeclampsia and the
Kidney: Pathophysiology and Clinical
Implications", Wiley, 2023

Publication

<1 %

59

Östlund, Eva, Maha Al-Nashi, Rangeen Rafik
Hamad, Anders Larsson, Maria Eriksson,
Katarina Bremme, and Thomas Kahan.
"Normalized endothelial function but
sustained cardiovascular risk profile 11 years
following a pregnancy complicated by
preeclampsia", Hypertension Research, 2013.

Publication

<1 %

60

"Preeclampsia", Springer Nature, 2018

Publication

<1 %

61

Carmen Garrido-Gimenez, Manel Mendoza,
Monica Cruz-Lemini, Laura Galian-Gay et al.
"Angiogenic Factors and Long-Term
Cardiovascular Risk in Women That
Developed Preeclampsia During Pregnancy",
Hypertension, 2020

Publication

<1 %

62

Egle Bytautiene, Fangxian Lu, Esther H. Tamayo, Gary D. V. Hankins, Monica Longo, Karolina Kublickiene, George R. Saade. "Long-term maternal cardiovascular function in a mouse model of sFlt-1-induced preeclampsia", *American Journal of Physiology-Heart and Circulatory Physiology*, 2010

Publication

<1 %

63

Sandra M. Blois, Ralf Dechend, Gabriela Barrientos, Anne Cathrine Staff. "A potential pathophysiological role for galectins and the renin–angiotensin system in preeclampsia", *Cellular and Molecular Life Sciences*, 2014

Publication

<1 %

64

Terry D. Bilhartz. "Making Use of a Natural Stress Test: Pregnancy and Cardiovascular Risk", *Journal of Women s Health*, 03/31/2011

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

the risk persistent hypertensio

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14

PAGE 15